

The impact of defense hormones on the interaction between plants and the soil microbial community

Zhang, J.

Citation

Zhang, J. (2021, May 4). The impact of defense hormones on the interaction between plants and the soil microbial community. Retrieved from https://hdl.handle.net/1887/3166490

Version:	Publisher's Version
License:	<u>Licence agreement concerning inclusion of doctoral thesis in the</u> <u>Institutional Repository of the University of Leiden</u>
Downloaded from:	https://hdl.handle.net/1887/3166490

Note: To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <u>https://hdl.handle.net/1887/3166490</u> holds various files of this Leiden University dissertation.

Author: Zhang, J. Title: The impact of defense hormones on the interaction between plants and the soil microbial community Issue Date: 2021-05-04

The impact of defense hormones on the interaction between plants and the soil microbial community

by Jing Zhang

张 静

Copyright © 2021, Zhang, Jing The impact of defense hormones on the interaction between plants and the soil microbial community PhD thesis Leiden University, The Netherlands Front cover by Hakim Valiton Thesis lay-out by Jing Zhang Printing and binding: Boekendeal.nl ISBN 9789492597694

The impact of defense hormones on the interaction between plants and the soil microbial community

Proefschrift

ter verkrijging van de graad van doctor aan de Universiteit Leiden, op gezag van de rector magnificus prof.dr.ir. H. Bijl, volgens besluit van het college voor promoties te verdedigen op dinsdag 4 mei 2021 klokke 13:45 uur

door

Jing Zhang

Geboren te Jinan, Shandong, China in 1991 Promotores: Prof. Dr. P.G.L. Klinkhamer

Prof. Dr. ir. T.M. Bezemer

Co-promotor: Dr. K. Vrieling

Promotiecommissie: Prof. Dr. G.P. van Wezel

Prof. Dr. R. Offringa

Prof. Dr. J. Memelink

Dr. E. Hannula (Netherlands Institute of Ecology (NIOO-KNAW))

Dr. i.r. Feng Zhu (Chinese Academy of Sciences, China)

Contents

Chapter 17
General introduction
Chapter 2
Activation of hormone-associated plant defense pathways alters the effects of soil microbial communities on plant performance
Chapter 3 61
Activation of SA-associated plant defense pathway alters the composition of soil bacterial communities
Chapter 4 109
Activation of SA-associated plant defense pathway alters the functions of soil microbial communities in four sequential generations
Chapter 5 151
The negative effects of soil microorganisms on plant growth only extends to the first weeks
Chapter 6 175
General discussion
Summary
Nederlandse samenvatting193
Curriculum Vitae197
Acknowledgements 199
Publications

Chapter 1

General introduction

The soil ecosystem consists of the largest reservoir of biodiversity on Earth (Zak et al., 2003; Decaëns, 2010; Bardgett and Van der Putten, 2014). Microbial communities are unseen drivers in soil ecosystems, and they play an important role in determining a wide variety of soil processes in terrestrial ecosystems (Van der Heijden, et al., 2008; Fester et al., 2014; Delgado-Baquerizo et al., 2016; Singh and Gupta, 2018). Soil microbial communities can influence plant performance and can drive plant species composition on a particular soil. Soil microbes are associated with an extensive range of ecosystem processes, such as nitrogen (N) and carbon (C) cycling, organic matter decomposition, soil structural formation and stability and these processes, in turn, affect plant growth (Beare et al., 1992; Ehrenfeld et al., 2005; Schimel and Schaeffer, 2012; Bardgett et al., 2014). While these processes can benefit plant growth, the soil microbial community also harbors microbes that compete with plants for nutrients or are pathogenic and impair plant growth. This leads to the question if plants can manipulate the composition of the soil microbial community to their advantage.

In agriculture, the physical structure of the soil is often altered to improve crop production and this modifies biological components and microbial properties of the soil (Mousavi and Eskandari, 2011; Van der Heijden et al., 2013; Van der Putten, et al., 2013). Moreover, green crop management technologies, such as regulation of soil microbial biodiversity, application of beneficial microbial agents and induction of plant hormonal resistance, are regarded as promising approaches against pests and microbial pathogens (Chung et al., 1988; Kennedy and Smith, 1995; Neher, 1999; Sturz and Christie, 2003). Although many experiments have shown that activation of hormonal signaling pathways can boost a plant's immunity against pathogenic microbial attacks, whether and how these hormonal signaling pathways affect the soil microbial community and consequently plant growth is still poorly understood (Berendsen et al., 2012; Graham, et al., 2016). Therefore, to better understand the roles of soil microbial communities at both taxonomic and functional level, studies from a plant defensive perspective are timely and needed, and this is the main focus of this thesis.

1. The relationship between plants and soil microbial communities

In the early middle ages, under Charles the Great, cropping and fallow rotations were already applied in Europe (van der Putten et al., 2013). Chinese historical books record that cropping and fallow rotations in China began even in the ancient Warring States period, which ran from 475 BC to 221 BC (Zhang and Yu, 2006). In agriculture, all these ancient practical actions aimed to overcome the same problem: a soil becomes less suitable for a crop if this crop is grown in that soil repeatedly.

We have since become aware that the negative impact of soil on plant growth is dependent on the role of soil microbes. Particularly, in the early 2000s, a large number of studies began to emphasize that the relationship between plants and soil microbial communities is bidirectional, rather than unidirectional. Plants can affect the microbial communities in the soil, and in turn, soil microbial communities also influence plant growth. Nowadays it is clear that the interactions between plants and soil microbial communities are extremely complicated (Van der Heijden, et al., 2008; Fester et al., 2014; Singh and Gupta, 2018).

1.1 Effects of soil microbial communities on plants

Generally speaking, plant-microbial interactions can be broadly subdivided into three basic groups of effects. Firstly, in many cases plants and specific microbes do not affect each other strongly. Secondly, there are negative effects on plants through root-associated organisms that form pathogenic relationships with plants. These pathogenic organisms in the rhizosphere include parasitic nematodes, fungi, Archaea, bacteria and invertebrate herbivores. Soil pathogens can reduce plant productivity, thus impacting ecosystem processes. Ecologists are long aware that soil microbes can cause serious reductions in plant growth (Nijjer et al., 2007). Among a myriad of soilborne microbial pathogens, *Phytopathora, Pythium, Fusarium* and *Verticillium* are well-known genera and they have been widely reported to have negative effects on the production of many crops and economically important tree species, such as potato, wheat, radish, pea and oaks (Harman et al., 1980; Nirenberg, 1981; Jung et al., 1999).

Thirdly, root-associated organisms can have positive effects on plant growth. For example, several plant growth-promoting *rhizobacteria* (PGPR), like *Pseudomonas* and *Burkholderia*, residing in the rhizosphere may repress the growth and activity of

soil-borne pathogens and other attackers (Bhattacharyya and Jha, 2012). PGPR commonly reside in the rhizosphere, where they are important regulators involved in numerous biological processes affecting host plants, e.g. solubilizing phosphate, fixing available soil nitrogen, producing siderophores, phytohormones, producing volatile organic compounds (VOCs), inducing host systemic resistance and systemic acquired resistance and stimulating antifungal compounds (Wei et al., 1991; Nelson, 2004; Esitken et al., 2010; Bhattacharyya and Jha, (2012). Arbuscular mycorrhizal fungi (AMF) can act as a natural extension of the host root system, to increase the possibility of plants to obtain resources from the soil, and plants provide carbon (C) to the AMF in exchange (Azcón-Aguilar et al., 1992; Barea, 2000). Moreover, some proteobacteria, such as legume-nodulating Burkholderia strains, and species of the genus Azoarcus and Sinorhizobium meliloti, are well known for their functions in fixing soil atmospheric nitrogen, which also benefits host plants (Reinhold-Hurek et al., 1993; Chen et al., 2003; Hayat et al., 2010; Schlüter et al., 2010). Additionally, root endophytes can play an important role in enhancing both biotic and abiotic stress tolerance in plants (Dimkpa et al., 2009), while some rhizosphere bacteria produce antibiotic compounds or protective biofilms that prevent the plant from attack by pathogenic soil bacteria.

Many studies report that the overall net effect of soil microbial communities on plant performance is negative (Nijjer et al., 2007). Inoculation of soil-borne microbial communities into sterilized soil often causes a reduction in plant growth. Negative effects of the soil microbial community on plant growth can be due to nutrient competition between plants and microbes and due to soil-borne plant-pathogenic microbes. Soil microbes, such as AMF, phosphorus-solubilizing bacteria, proteases and nitrogen-fixing bacteria can assist plants in taking up more nutrients from the surrounding soil. However, plants and microbes also depend largely on the same inorganic nutrients and therefore compete for these nutrients.

Plant-soil feedback studies, mostly show that inoculation of sterilized soil with microbial communities collected from underneath conspecific plants has a stronger negative effect on plant growth than inoculation with microbial communities from other plant species, suggesting that plant species-specific pathogenic or plant growth-

inhibiting microorganisms build up in the rhizosphere (Pendergast et al., 2013; Dawson et al., 2017; Howard et al., 2020). Studies with pure "home soil" and "away soil" also generally report negative effects of "home soil" on plant growth (Manning et al., 2008; Ayres et al., 2009). All these studies suggest that pathogenic effects of micro-organisms play an important role in the interaction between plants and the soil microbial community. These studies furthermore suggest that many of the interactions are species-specific and that plants affect the microbial community in a speciesspecific way and vice versa. Moreover, it is worth noticing that soil microorganisms can also affect the above and belowground defense system of a plant (Huberty et al., 2020). For instance, soil-borne microorganisms affect the composition of pyrrolizidine alkaloids (PAs) and the total PA concentration in the plant Jacobaea vulgaris (Joosten et al., 2009; Kostenko et al., 2012; Kos et al., 2015). Studies that explore the mechanisms behind the interactions between plants and the soil microbial community nowadays are boosted by the molecular tools that make it possible to study the composition and functions of microbial communities. In this thesis, I will concentrate on the effects of the microbial communities on plant growth.

1.2 Effects of plants on rhizosphere microbial communities

The term rhizosphere was introduced by Hiltner in 1904 to describe the layer of soil that was influenced by the root of a plant (Hiltner, 1904). In comparison to root-free soil, the rhizosphere is an area where plant roots and soil microorganisms are mutually interacting. Soil properties (pH, humidity, chemical composition, texture and structure) play an important role in the modulation of rhizosphere microbial communities (Börner, 1960; Bach et al., 2010). However, plants also greatly influence the structure and function and diversity of microbial communities, especially in the rhizosphere (Grayston et al., 1998; Girvan et al., 2003; Nunan et al., 2005; Berg and Smalla, 2009; Dennis et al., 2010). Plant species differ in their effect, and in the strength of this effect, on microbial communities (Zak et al., 2003; Mangan et al., 2010). Other studies showed that bacterial communities in the soil of grass and forb species differ (Hannula et al., 2019) and that the structure and function of soil microbial communities of exotic plant species are different from those of native species (Kourtev et al., 2003). A prediction of how specific crops will influence the

soil microbial community may help to reduce risks and yield losses in agriculture, but, so far, this is poorly understood and remains a long-term challenge.

Plants synthesize a vast array of secondary metabolites (SMs) and more than 100,000 are reported (Dixon, 2001; Quiroga et al., 2001; Bartwal et al., 2013). Many studies have demonstrated that these compounds are involved in the chemical defense of plants against pathogenic microbes (Van Loon, 2007; Boller and He, 2009). In particular, root exudates are key drivers of microbial diversity and composition in the rhizosphere. For example, sugars, organic acids and amino acids are well-known nutrients for microbes (Canarini, et al., 2019). The composition and concentration of these metabolites in the rhizosphere depend upon the plant species and overall environmental conditions (Broeckling et al., 2008; Zahar et al., 2008).

Plant chemical defenses play an important role in plant-pathogenic microbe interactions. It has been suggested that the diversity of defensive compounds has evolved as a result of an evolutionary arms race between the plants and their potential attackers (Ehrlich and Raven, 1964). In particular, those defenses based on molecules with low molecular weight, and long-distance communicating molecules, such as volatile organic compounds (VOCs) (Insam and Seewald, 2010; Frankenberger and Arshad, 2020). Moreover, SMs, such as citronellal, berberine and pyrazines are also functioning in plant defense against pathogenic microbes (Wink, 1988; Tyc et al., 2017).

In addition, plants can regulate the production of protease inhibitors to defend themselves (Lawrence and Koundal, 2002; Habib and Fazili, 2007). Microbial pathogens can secrete extracellular protease enzymes, and those enzymes can digest some proteins in the tissues of plants (Ryan, 1990). Plants can defend themselves from protease-related pathogens through expressing protease inhibitors and also regulate them to accurate and strict concentrations. Researchers have generated some transgenic plants with high expression of protease inhibitors, such as transgenic rice, potato, soybean, and these plants exhibit increased resistance against various pathogens (Cowgill et al., 2002; Rahbé et al., 2003).

Moreover, induced defense responses in the plant influence the chemical composition of root exudates and through that the bacterial community structure in the soil. Salicylic acid (SA), ethylene (ET), abscisic acid (ABA) and methyl jasmonate (MeJA) hormonal signaling pathways can alter the bacterial community composition in the soil (Carvalhais et al., 2015; Lebeis et al., 2015). Van der Meij et al. (2018) showed that application of SA to endophytic actinobacteria stimulates antibiotic production. Altogether these findings suggest that activating hormonal signaling may not only boosts the plant's defense system directly but also can affect the microbial composition on the soil thereby potentially mitigating the negative effects of the soil microbial community on plant growth.

2. Plant hormonal induced defense against soil-borne pathogens

To counteract the effects of microbial pathogens, plants have evolved a broad range of defensive mechanisms, which are partly regulated via hormonal signaling pathways (Fujita et al., 2006). Defense, as an essential and effective strategy for terrestrial plant species against pests and pathogens has been broadly developed in plants (Wesson and Wesson, 1993; Bronstein, 1998; Agrawal, 2011; Turley et al., 2013; Zhang et al., 2015). Induced defenses are defenses that are activated after infection occurs and enhance plant fitness (Boots and Best, 2018). Phytohormones are a group of natural plant compounds with low molecular weight that play an important role in the regulation of plant growth and development and induced plant resistance against pests and pathogens. SA, ET, abscisic ABA, MeJA, auxin, cytokinins (CKs), gibberellic acid (GA) and brassinosteroids (GAs) are commonly studied phytohormones. Besides these, there are also several other compounds (e.g. karrikins, triacontanol and nitric oxide) that can be involved in induced plant defense, but their functions are still under debate.

Each phytohormone has clear functions, however, they can also exhibit strong interactive effects. For instance, JA and SA are well-known for their negative cross-talk (Munné-Bosch and Müller, 2013). Upregulating the SA signaling can lead to downregulation of the JA signaling and vice versa. Such cross talk is one of the mechanisms that can explain why plant pathogens in the soil can e.g. affect herbivory above ground (Aljbory and Chen, 2018). Although most hormones have been 13

implicated to be involved in defense pathways, the key regulator against pathogens and pests, in particular, to defend plants against biotrophic, necrotrophic pathogenes and herbivores, are the phytohormones JA and SA (Bari and Jones, 2009). In the following sections, JA and SA induced resistance in plants against soil-borne pathogens is described separately.

2.1 JA-induced resistance

JA is associated with several biological processes in plants. Specifically, JA can stimulate the germination of seeds, negatively impacts root growth, and invokes tuber formation and fruit ripening. Apart from these functions, JA is well-known for being involved in induced resistance against herbivores and for being a regulator of the activation of induced systemic resistance (ISR) of plants against necrotrophic microbial pathogens.

The biosynthesis of JA has been mostly studied in the model plant species Arabidopsis thaliana and tomato (Lycopersicon esculentum) (Ruan et al., 2019). Both biotic and abiotic stresses can induce the synthesis of JA. In plant tissues JA can be converted into JA-isoleucine conjugate (JA-Ile) by JAR1 (an auxin-induced gene), JA-Ile is a bioactive state of JA. MYC is a family of regulator genes that code for transcription factors and JA-Ile activates the MYC transcription factors by directly binding to the jasmonate zim-domain (JAZ) and a coronatine insensitivel (COI1) protein, which results in the degradation of JAZ transcriptional repressor proteins through the proteasome pathway. These processes result in the activation of transcription factors and the regulation of JA-responsive genes (e.g. MYC2, ERF1 and ORA59), which are associated with plant responses against environmental stresses from pathogens, wounding, and insect herbivory, biosynthesis of secondary metabolites, and with plant growth and development. For example, Carvalhais et al. (2015) demonstrated that the JA signaling pathway affects the composition of root exudates by enhancing the production of ornithine and that ornithine can be used by plant growth-promoting bacteria such as Pseudomonas fluorescens, which in turn has a positive effect on plant growth.

Due to the effectivity of JA-induced resistance against herbivores, in agriculture, foliar application of JA or MeJA has been considered as an alternative approach to control pests rather than using chemical pesticides. This theory has been tested in several crop plant species, including corn, tomato and wheat (Mandal et al., 2006; War et al., 2011; Chen et al., 2018).

In addition, to increased resistance to pests, JA-mediated defense also regulates the plant's response to necrotrophic microbial pathogens, e.g. *Pseudomonas syringae, Fusarium oxysporum and Hyaloperonospora parasitica* (Antico et al., 2012; Wasternack and Strnad, 2018; Li et al., 2019). The activation of JA-signaling pathways in the plant can result in changes in the composition of bacteria in the rhizosphere as was shown for the plant *A. thaliana* (Carvakhais et al., 2013). The mechanisms behind this are not yet fully resolved. JA signaling may directly affect microbial species or through the interaction with SA signaling.

2.2 SA-induced resistance

SA is another well-studied hormonal compound, which plays an important role in the activation of SA-induced resistance against biotrophic microbial pathogens. Hypersensitive response (HR) is a primary manifestation of a plant to pathogenic attack, e.g. due to cell death of the tissues surrounding the infection, to control the spread of pathogens. Cultivars that are highly sensitive to SA are often more tolerant of microbial pathogens (Seskar et al., 1998).

Infection of plant tissues with biotrophic pathogens leads to the accumulation of SA, as well as monomerization of NPR1 via SA-mediated redox changes in the cell. Later, monomeric NPR1 is relocated into the nucleus, at which the monomeric NPR1 interacts with TGA transcription factors, as a result, SA-responsive genes are activated. A large number of WRKY genes are induced by SA, among which some can regulate SA-responsive gene expression (Van der Does et al., 2013).

SAR is associated with the expression of pathogenesis-related proteins (PRPs). PRPs include proteins like β -1, 3-glucanase and chitinases (Van Loon and Van Strien, 1999). Commonly, both chitinases and glucanases show antimicrobial activities. Once PRPs

are induced by SA, they take actions in several ways to assist plants against pathogens. These actions can be direct and indirect (Edreva, 2005). PRPs can direct breakdown or damage pathogens based on their antifungal and antibacterial activities. In addition, PRPs can indirectly boost host defensive abilities through hydrolytic released compounds of fungal cell walls, e.g. ochitin and glucan fragments. These released oligosaccharides could further stimulate a series of defensive responses in host plants (Lawrence et al., 2000; Edreva, 2005).

SA induced resistance can interact with beneficial bacteria and fungi, such as *Pseudomonas*, plant growth-promoting bacteria (PGPR), arbuscular mycorrhiza fungi (AMF). Those beneficial microbes interact with SA-induced resistance through species-specific microbe-associated molecular patterns (MAMPs) (Bittel and Robatzek, 2007; Choi and Klessig, 2016). MAMPs are special components on the surface of general microbes, such as, *Trichoderma*, *Bacillus* and *Pseudomonas* (Pieterse et al., 2014). Once a host plant recognizes these MAMPs, the plant will activate its innate systemic defensive system to be ready to cope with future pathogens.

Exogenous application of SA or Methyl SA to activate SAR is a way to control microbial pathogenic diseases. In many crops, like tomato, pepper and pea exogenous application of SA results in a suppressing effect on microbial pathogens (Esmailzadeh et al., 2008; Barilli et al., 2010; Choi and Hwang, 2011). Overall, in agriculture, improving SA-mediated resistance has become a promising strategy to control microbial pathogens and viruses.

3. Adaptation of microbes to plant defenses

Plants are not the only organisms that can produce hormonal compounds, which stimulate the activation of signaling pathways in the plant. For example, plant growth-promoting rhizobacteria can produce and/or degrade phytohormones, and in this way interfere with the regulation of plant growth (Dodd et al., 2010). Not only beneficial bacteria but also pathogenic microbes can produce hormones or compounds that impact plant growth.

Some microbial pathogens can mimic the production of plant hormones and are able to highjack the plant immune system (Cui et al., 2005; Laurie-Berry et al., 2006; Navarro et al., 2008). Cui et al. (2005), for example, reported that the bacterial pathogen (*Pseudomonas syringae*) activated induced systemic resistance in *A. thaliana* by producing coronatine (COR). This compound can mimic the function of the JA hormone, thereafter induce the JA-related signaling pathway in host plants. Besides, Laurie-Berry et al. (2006) also demonstrated that *P. syringae* could utilize COR to upregulate the JA pathway in host plants and suppress the SA-mediated signaling pathway in tomato plants, making *P. syringae* even more virulent.

Some bacteria can synthesize SA by converting their chorismate through isochorismate synthase (ICS) and isochorismate pyruvate lyase (IPL) (Chen et al., 2009; Dempsey et al., 2011). This has been reported for several bacteria species, like *Pseudomonas aeruginosa* and *P. fluorescens* (Mercado-Blancoet al., 2001; Kerbarh et al., 2005; AI-Mustafaet al., 2009). The production of SA by bacteria can increase the plant's resistance against pathogens. For example, the production of SA by *P. aeruginosa* can enhance the resistance of plants against the bacterial pathogen *Botrytis cinerea*, which causes leaf diseases on bean (De Meyer and Höfte, 1997); *P. fluorescens* enhances plant defense in chickpea against *Fusarium* wilt (Saikia et al., 2003).

4. Contradictory observations of SA signaling on the soil microbial community

Many studies investigated the effects of hormonally induced defenses on single pathogens (El-Khallal 2007; Abo-Elyousr et al., 2009; Mandal et al., 2009). However, our knowledge about the impact of activating plant hormonal signaling pathways on the composition of the soil microbial community is still limited and contradictory (reviewed in Hacquard et al., 2017).

Lebeis et al. (2015) examined the effect of SA on isogenic *A. thaliana* mutants with altered immune systems and found that plants with an altered SA signaling pathway contained rhizospheres that differed in the relative abundance of specific bacterial families as compared to wild type plants. Kniskern et al. (2007) using *A. thaliana* mutants found that activation of SA signaling pathways reduced endophytic bacterial

community diversity, whereas plants that were deficient in JA-mediated defenses experienced greater epiphytic bacterial diversity. When a plant is exposed to microbial pathogens or herbivory, the plant changes the composition of the primary and secondary metabolites that are produced, and this can impact the soil microbial community, and may result in a feedback to the plant (Rolfe et al., 2019).

Several other studies showed that there was no effect of activation of SA signaling on the soil microbial community. For instance, Wang et al. (2015) found that higher concentrations of exogenously added SA inhibited the growth of grape plants, however, there was no direct correlation between the inhibitory effects of SA on plant growth and the diversity of the soil bacterial or fungal community. Similarly, Doornbos et al. (2011) found that chemical activation of JA- or SA-induced resistance did not significantly affect the composition and diversity of the rhizosphere bacterial community in A. thaliana. Hein et al. (2008) compared the effect of SA-induced resistance on the diversity of rhizosphere bacterial communities in several Arabidopsis mutants and found that changes in microbial composition were not caused by the induction of the SA signaling pathway. Even though the previous mentioned study showed that activation of SA-dependent defenses did not change the composition of soil microbial community in A. thaliana and the SA-independent defense was not induced by foliar application of SA, this still provides a great value to understand the interplay of activating SA-signaling pathways and microbial composition (Sonnemann et al., 2002; Doornbos, et al., 2012; Moccia and Lebeis, 2019).

All taken together, the role of induced resistance on the soil microbial community is still being debated, and more research on the impact of upregulated hormonal signaling in plants on the composition and functionality of the soil microbial community is necessary.

5. Characterizing microbial communities

Microbial communities can be characterized using metagenomics tools. Metagenomics aims at determining the microorganisms as a whole and allows us to extract the biological information of all the microbes from the environment directly (Hugenholtz and Tyson, 2008). Up to date, metagenomics has been widely applied in various environments to investigate microbial communities ranging from soils, water, ocean and human gut (Handelsman, 2004; Daniel, 2005). However, the lack of reference sequences and genomes is a major drawback of metagenomics (Krehenwinkel et al., 2019).

The microbial community can be characterized on basis of the species present or on the basis on the genes and their functions that are expressed in the microbial community. While the first is highly relevant to understand the diversity and the dynamics of microbial populations and communities, the second is of great importance if we want to understand the mechanisms behind plant-microbial interactions and how a microbial community adapts to the environment. In other words: taxonomic information helps to answer the most primary question for soil microbial-ecologists: Who is there? Analysis of gene expression helps to understand what they are doing. Amplification of 16S rRNA barcode markers is commonly used to determine bacterial microbial communities and the costs of characterizing part of the genome are much lower than sequencing the whole genome. Pipelines for 16S rRNA gene sequencing and identifying operational taxonomic units (OTUs) by aligning the reads against known public databases (e.g. NCBI, EzBioCloud 16S database and MBGD) are available. Nowadays, high-throughput sequencing has become a vital and cost-effective tool for profiling functions of soil microbial communities. It can generate a high volume of data and long read lengths. Illumina short reads sequencing (up to 250 bp) has a high output and low read errors. In this thesis, the Illumina sequencing platform is used to examine the microbial composition and functional genes of rhizosphere soil microbial communities.

Also to process metatranscriptomics data existing pipelines can be used. These pipelines can be modified and applied to different experimental designs. For example, the IMP pipeline incorporates robust read preprocessing and is suitable for analyzing metagenomic and metatranscriptomics as it provides information on both microbial structure and functional genes (Narayanasamy et al., 2016). MetaTrans is an open-source pipeline developed for a paired-end RNA-Seq analysis (Martinez et al., 2016) while the functional mapping and analysis pipeline (FMAP) provides alignment, gene

family abundance calculations and open-level statistical analysis (Kim et al., 2016). SAMSA2 is a standalone metatranscriptome analysis pipeline and is used on a supercomputing cluster, which is more flexible and reproductive in processing a large volume of sequence data (Ni et al., 2016). In this thesis I used a modified pipeline that can run on a regular computer and is easily customized. The pipeline that was used assists with the transcriptomic tools Trinity and Trinotate. Transcripts generated by Trinity can be annotated with Trinotate and Trinotate allows users to perform functional annotation with several selective methods, such as homology search, protein domain search, or protein peptide domain search. The combination of these two bioinformatic tools enabled us to explore the structure and the functionality of microbial communities (Haas et al., 2013).

6. Research questions

A number of studies have shown that the overall effect of the soil microbial community on the growth of *J. vulgaris* seedlings is negative (Bezemer and van Dam, 2005; Van de Voorde et al., 2012; Wang et al., 2019). In this project, we asked whether this negative effect of the soil microbial community could be mitigated by the activation of Me-JA and SA signaling pathways through an effect on the composition of the soil microbial community.

First, I studied in four plant species (*J. vulgaris, Cirsium vulgare, Trifolium repens* and *Daucus carota*) how the growth of these plants was affected by the presence of a live microbial community in the soil. Then, I examined, for *J. vulgaris* that grew less well in live soil than in sterilized soil and for which the negative effect of the live soil on plant growth was mitigated by exogenous application of SA on leaves of the plant, how the application of SA alters the soil microbial community on both taxonomic and functional levels through a multi-generational experiment where I analyzed mRNA of the soil. In addition, I studied for how long during the plant's life the negative effect of a live soil on plant growth is maintained.

Specifically, the following research questions are addressed in this thesis:

(1) Do the effects of the soil microbial community on plant growth differ among four plant species that occur in the same habitat? Does the foliar application of JA and SA alter the effects of the soil microbial community on plant growth of these four plant species? Does the negative effect of the soil microbial community increase or decrease over successive generations of plant growth in *J. vulgaris* inoculated with the soil of the previous generation, and how is this influenced by SA application?

(2) How does the application of SA on *J. vulgaris* affect the composition of the microbial community in the rhizosphere? How does the soil microbial composition change over plant generations, when in each generation sterilized soil is inoculated with soil from the previous generation for plants that are treated with SA and untreated control plants?

(3) Does the application of SA on *J. vulgaris* alter the gene expression in the rhizosphere? Does the application of SA impact microbial gene expression over generations? Which groups of genes are influenced by SA-treated soil samples compared to control over generations?

(4) How long does the effect of inoculum of 10% soil containing a natural microbial community on plant growth last? Does the timing of inoculation change the effect of soil microbial communities on plant growth in *J. vulgaris*?

7. Thesis outline

Many plant species grow better in sterilized soil than in soil that contains a live microbial community, this could be due to an overall net pathogenic effect of soil microbial communities. To find out if an overall negative effect on plant growth is a common phenomenon in nature, in **Chapter 2**, four plant species were grown in either sterilized soil or sterilized soil containing 10% of live soil. In addition, I exposed plant leaves to two hormonal treatments (jasmonic acid and salicylic acid) to examine if hormonal defense pathways can influence the microbial effects on plants.

In Chapter 3, I sequenced and analyzed the microbial communities from the experiment of Chapter 2, to investigate if SA-induced defense had an impact on the

taxonomic composition of the microbial community in rhizosphere samples using Illumina sequencing. Since the application of SA mitigated the negative effect of soil microbial communities on the growth of *J. vulgaris*, I used this species to study the changes in the composition of the microbial community in response to SA application for four generations of plant growth.

As described in **Chapter 2** and **3**, certain groups of microbial species responded differently to the exogenous application of SA on plant leaves. However, the functions of those microbial species in the rhizosphere are largely unknown. Therefore, in **Chapter 4** I used metatranscriptomics to study the functional genes and clusters in the rhizosphere microbiome of both SA-treated and control samples. The changes in microbial functional genes over four generations were analyzed and compared.

Studies on plant-soil-interactions often address the soil microbial effect with measurements on plant absolute biomass. However, even if there is no difference in the relative growth rate (RGR), the absolute difference in plant growth can still increase. In **Chapter 5**, I studied for *J. vulgaris* how long the negative effect of live soil on plant growth is maintained. Also, I studied if the timing of inoculation affected the RGR of this species. Finally, the results described in this thesis and their implications are discussed in **Chapter 6**.

References

- Abo-Elyousr, K.A., Hussein, M.A.M., Allam, A.D.A. and Hassan, M.H., 2009. Salicylic acid induced systemic resistance on onion plants against *Stemphylium vesicarium*. Archives of *Phytopathology and Plant Protection*, 42(11), pp.1042-1050.
- Agrawal, A.A., 2011. Current trends in the evolutionary ecology of plant defence. *Functional Ecology*, 25(2), pp.420-432.
- Al-Mustafa, M.M., Badran, I.Z., Abu-Ali, H.M., Al-Barazangi, B.A., Massad, I.M. and Al-Ghanem, S.M., 2009. Intravenous dexmedetomidine prolongs bupivacaine spinal analgesia. *Middle East J Anesthesiol*, 20(2), pp.225-231.
- Aljbory, Z. and Chen, M.S., 2018. Indirect plant defense against insect herbivores: a review. *Insect Science*, 25(1), pp.2-23.
- Antico, C.J., Colon, C., Banks, T. and Ramonell, K.M., 2012. Insights into the role of jasmonic acidmediated defenses against necrotrophic and biotrophic fungal pathogens. *Frontiers in Biology*, 7(1), pp.48-56.
- Ayres, E., Steltzer, H., Simmons, B.L., Simpson, R.T., Steinweg, J.M., Wallenstein, M.D., Mellor, N., Parton, W.J., Moore, J.C. and Wall, D.H., 2009. Home-field advantage accelerates leaf litter decomposition in forests. *Soil Biology and Biochemistry*, 41(3), pp.606-610.
- Azcón-Aguilar, C., & Barea, J. M., 1992. Interactions between mycorrhizal fungi and other rhizosphere microorganisms. *Mycorrhizal functioning: an integrative plant-fungal process. Chapman and Hall, New York*, pp.163-198.
- Bach, E.M., Baer, S.G., Meyer, C.K. and Six, J., 2010. Soil texture affects soil microbial and structural recovery during grassland restoration. *Soil Biology and Biochemistry*, 42(12), pp.2182-2191.
- Bardgett, R.D. and Van der Putten, W.H., 2014. Belowground biodiversity and ecosystem functioning. *Nature*, *515*(7528), pp.505-511.
- Bardgett, R.D., Mommer, L. and De Vries, F.T., 2014. Going underground: root traits as drivers of ecosystem processes. *Trends in Ecology & Evolution*, 29(12), pp.692-699.
- Barea, J. M., 2000. Rhizosphere and mycorrhiza of field crops. In *Biological resource management connecting Science and Policy*. Springer, Berlin, Heidelberg, 2000, pp.81-92.
- Bari, R. and Jones, J.D., 2009. Role of plant hormones in plant defence responses. *Plant molecular biology*, 69(4), pp.473-488.
- Barilli, E., Sillero, J.C. and Rubiales, D., 2010. Induction of systemic acquired resistance in pea against rust (Uromyces pisi) by exogenous application of biotic and abiotic inducers. Journal of Phytopathology, 158(1), pp.30-34.
- Bartwal, A., Mall, R., Lohani, P., Guru, S.K. and Arora, S., 2013. Role of secondary metabolites and brassinosteroids in plant defense against environmental stresses. *Journal of plant growth regulation*, 32(1), pp.216-232.
- Beare, M.H., Parmelee, R.W., Hendrix, P.F., Cheng, W., Coleman, D.C. and Crossley Jr, D.A., 1992. Microbial and faunal interactions and effects on litter nitrogen and decomposition in agroecosystems. *Ecological Monographs*, 62(4), pp.569-591.
- Berendsen, R.L., Pieterse, C.M. and Bakker, P.A., 2012. The rhizosphere microbiome and plant health. *Trends in plant science*, *17*(8), pp.478-486.
- Berg, G. and Smalla, K., 2009. Plant species and soil type cooperatively shape the structure and function of microbial communities in the rhizosphere. *FEMS microbiology ecology*, *68*(1), pp.1-13.
- Bezemer, T.M. and van Dam, N.M., 2005. Linking aboveground and belowground interactions via induced plant defenses. *Trends in Ecology & Evolution*, 20(11), pp.617-624.

- Bhattacharyya, P.N. and Jha, D.K., 2012. Plant growth-promoting rhizobacteria (PGPR): emergence in agriculture. *World Journal of Microbiology and Biotechnology*, 28(4), pp.1327-1350.
- Bittel, P. and Robatzek, S., 2007. Microbe-associated molecular patterns (MAMPs) probe plant immunity. *Current opinion in plant biology*, *10*(4), pp.335-341.
- Boller, T. and He, S.Y., 2009. Innate immunity in plants: an arms race between pattern recognition receptors in plants and effectors in microbial pathogens. *Science*, *324*(5928), pp.742-744.
- Boots, M. and Best, A., 2018. The evolution of constitutive and induced defences to infectious disease. *Proceedings of the Royal Society B: Biological Sciences*, 285(1883), p.20180658.
- Börner, H., 1960. Liberation of organic substances from higher plants and their role in the soil sickness problem. *The Botanical Review*, *26*(3), pp.393-424.
- Broeckling, C.D., Broz, A.K., Bergelson, J., Manter, D.K. and Vivanco, J.M., 2008. Root exudates regulate soil fungal community composition and diversity. *Appl. Environ. Microbiol.*, 74(3), pp.738-744.
- Bronstein, J.L., 1998. The contribution of ant-plant protection studies to our understanding of mutualism 1. *Biotropica*, 30(2), pp.150-161.
- Canarini, A., Wanek, W., Merchant, A., Richter, A. and Kaiser, C., 2019. Root exudation of primary metabolites: mechanisms and their roles in plant responses to environmental stimuli. *Frontiers in Plant Science*, *10*, p.157.
- Carvalhais, L.C., Dennis, P.G., Badri, D.V., Tyson, G.W., Vivanco, J.M. and Schenk, P.M., 2013. Activation of the jasmonic acid plant defence pathway alters the composition of rhizosphere bacterial communities. *PLoS One*, 8(2), p.e56457.
- Carvalhais, L.C., Dennis, P.G., Badri, D.V., Kidd, B.N., Vivanco, J.M. and Schenk, P.M., 2015. Linking jasmonic acid signaling, root exudates, and rhizosphere microbiomes. *Molecular Plant-Microbe Interactions*, 28(9), pp.1049-1058.
- Chen, Z., Zheng, Z., Huang, J., Lai, Z. and Fan, B., 2009. Biosynthesis of salicylic acid in plants. *Plant signaling & behavior*, 4(6), pp.493-496.
- Chen, W.M., Moulin, L., Bontemps, C., Vandamme, P., Béna, G. and Boivin-Masson, C., 2003. Legume symbiotic nitrogen fixation by β-proteobacteria is widespread in nature. *Journal of bacteriology*, 185(24), pp.7266-7272.
- Chen, G., Klinkhamer, P.G., Escobar-Bravo, R. and Leiss, K.A., 2018. Type VI glandular trichome density and their derived volatiles are differently induced by jasmonic acid in developing and fully developed tomato leaves: Implications for thrips resistance. *Plant Science*, 276, pp.87-98.
- Choi, H.W. and Hwang, B.K., 2011. Systemic acquired resistance of pepper to microbial pathogens. *Journal of Phytopathology*, 159(6), pp.393-400.
- Choi, H.W. and Klessig, D.F., 2016. DAMPs, MAMPs, and NAMPs in plant innate immunity. *BMC plant biology*, *16*(1), p.232.
- Chung, Y.R., Hoitink, H.A.H. and Lipps, P.E., 1988. Interactions between organic-matter decomposition level and soilborne disease severity. *Agriculture, ecosystems & environment, 24*(1-3), pp.183-193.
- Cowgill, S.E., Wright, C. and Atkinson, H.J., 2002. Transgenic potatoes with enhanced levels of nematode resistance do not have altered susceptibility to nontarget aphids. *Molecular Ecology*, 11(4), pp.821-827.
- Cui, J., Bahrami, A.K., Pringle, E.G. Hernandez-Guzman, G., Bender, C.L., Pierce, N.E. and Ausubel, F.M., 2005. *Pseudomonas syringae* manipulates systemic plant defenses against pathogens and herbivores. *Proceedings of the National Academy of Sciences*, U.S.A, 102(5), pp.1791-1796.
- Daniel, R., 2005. The metagenomics of soil. Nature Reviews Microbiology, 3(6), pp.470-478.

- Dawson, W., Hör, J., Egert, M., van Kleunen, M. and Pester, M., 2017. A small number of lowabundance bacteria dominate plant species-specific responses during rhizosphere colonization. *Frontiers in microbiology*, 8, p.975.
- De Meyer, G. and Höfte, M., 1997. Salicylic acid produced by the rhizobacterium *Pseudomonas aeruginosa* 7NSK2 induces resistance to leaf infection by *Botrytis cinerea* on bean. *Phytopathology*, 87(6), pp.588-593.
- Decaëns, T., 2010. Macroecological patterns in soil communities. *Global Ecology and Biogeography*, 19(3), pp.287-302.
- Delgado-Baquerizo, M., Maestre, F.T., Reich, P.B., Jeffries, T.C., Gaitan, J.J., Encinar, D., Berdugo, M., Campbell, C.D. and Singh, B.K., 2016. Microbial diversity drives multifunctionality in terrestrial ecosystems. *Nature communications*, 7(1), pp.1-8.
- Dempsey, D.M.A., Vlot, A.C., Wildermuth, M.C. and Klessig, D.F., 2011. Salicylic acid biosynthesis and metabolism. *The Arabidopsis book/ American Society of Plant Biologists*, 9.
- Dennis, P.G., Miller, A.J. and Hirsch, P.R., 2010. Are root exudates more important than other sources of rhizodeposits in structuring rhizosphere bacterial communities? *FEMS microbiology ecology*, 72(3), pp.313-327.
- Dimkpa, C., Weinand, T. and Asch, F., 2009. Plant-rhizobacteria interactions alleviate abiotic stress conditions. *Plant, cell & environment, 32*(12), pp.1682-1694.
- Dixon, R.A., 2001. Natural products and plant disease resistance. Nature, 411 (6839), pp.843-847.
- Dodd, I.C., Zinovkina, N.Y., Safronova, V.I. and Belimov, A.A., 2010. Rhizobacterial mediation of plant hormone status. *Annals of Applied Biology*, 157(3), pp.361-379.
- Doornbos, R.F., Geraats, B.P., Kuramae, E.E., Van Loon, L.C. and Bakker, P.A., 2011. Effects of jasmonic acid, ethylene, and salicylic acid signaling on the rhizosphere bacterial community of *Arabidopsis thaliana*. *Molecular plant-microbe interactions*, *24*(4), pp.395-407.
- Doornbos, R.F., van Loon, L.C. and Bakker, P.A., 2012. Impact of root exudates and plant defense signaling on bacterial communities in the rhizosphere. A review. *Agronomy for Sustainable Development*, 32(1), pp.227-243
- Edreva, A., 2005. Pathogenesis-related proteins: research progress in the last 15 years. *Gen Appl Plant Physiol*, *31*(1-2), pp.105-24.
- Ehrenfeld, J.G., Ravit, B. and Elgersma, K., 2005. Feedback in the plant-soil system. *Annu. Rev. Environ. Resour*, 30, pp.75-115.
- Ehrlich, P.R. and Raven, P.H., 1964. Butterflies and plants: a study in coevolution. *Evolution*, pp.586-608.
- El-Khallal, S.M., 2007. Induction and modulation of resistance in tomato plants against *Fusarium* wilt disease by bioagent fungi (arbuscular mycorrhiza) and/or hormonal elicitors (jasmonic acid & salicylic acid): 2-changes in the antioxidant enzymes, phenolic compounds and pathogen related-proteins. *Aust J Basic Appl Sci*, 1(4), pp.717-732.
- Esitken, A., Yildiz, H. E., Ercisli, S., Donmez, M. F., Turan, M., & Gunes, A., 2010. Effects of plant growth promoting bacteria (PGPB) on yield, growth and nutrient contents of organically grown strawberry. *Scientia Horticulturae*, *124*(1), pp.62-66.
- Esmailzadeh, M., Soleimani, M.J. and Rouhani, H., 2008. Exogenous applications of salicylic acid for inducing systemic acquired resistance against tomato stem canker disease. *Journal of Biological Sciences*, 8(6), pp.1039-1044.
- Fester, T., Giebler, J., Wick, L.Y., Schlosser, D. and Kästner, M., 2014. Plant-microbe interactions as drivers of ecosystem functions relevant for the biodegradation of organic contaminants. *Current opinion in biotechnology*, 27, pp.168-175.

- Frankenberger Jr, W.T. and Arshad, M., 2020. *Phytohormones in Soils Microbial Production & Function*. CRC Press.
- Fujita, M., Fujita, Y., Noutoshi, Y., Takahashi, F., Narusaka, Y., Yamaguchi-Shinozaki, K. and Shinozaki, K., 2006. Crosstalk between abiotic and biotic stress responses: a current view from the points of convergence in the stress signaling networks. *Current opinion in plant biology*, 9(4), pp.436-442.
- Girvan, M.S., Bullimore, J., Pretty, J.N., Osborn, A.M. and Ball, A.S., 2003. Soil type is the primary determinant of the composition of the total and active bacterial communities in arable soils. *Appl. Environ. Microbiol.*, 69(3), pp.1800-1809.
- Graham, E.B., Knelman, J.E., Schindlbacher, A., Siciliano, S., Breulmann, M., Yannarell, A., Beman, J.M., Abell, G., Philippot, L., Prosser, J. and Foulquier, A., 2016. Microbes as engines of ecosystem function: when does community structure enhance predictions of ecosystem processes? *Frontiers in microbiology*, 7, p.214.
- Grayston, S.J., Wang, S., Campbell, C.D. and Edwards, A.C., 1998. Selective influence of plant species on microbial diversity in the rhizosphere. *Soil Biology and Biochemistry*, 30(3), pp.369-378.
- Haas, B.J., Papanicolaou, A., Yassour, M., Grabherr, M., Blood, P.D., Bowden, J., Couger, M.B., Eccles, D., Li, B., Lieber, M. and MacManes, M.D., 2013. De novo transcript sequence reconstruction from RNA-seq using the Trinity platform for reference generation and analysis. *Nature protocols*, 8(8), pp.1494-1512.
- Habib, H. and Fazili, K.M., 2007. Plant protease inhibitors: a defense strategy in plants. *Biotechnol Mol Biol Rev*, 2(3), pp.68-85.
- Hacquard, S., Spaepen, S., Garrido-Oter, R. and Schulze-Lefert, P., 2017. Interplay between innate immunity and the plant microbiota. *Annual review of Phytopathology*, 55, pp.565-589.
- Handelsman, J., 2004. Metagenomics: application of genomics to uncultured microorganisms. *Microbiol.* Mol. Biol. Rev., 68(4), pp.669-685.
- Hannula, S.E., Kielak, A.M., Steinauer, K., Huberty, M., Jongen, R., Jonathan, R., Heinen, R. and Bezemer, T.M., 2019. Time after time: temporal variation in the effects of grass and forb species on soil bacterial and fungal communities. *mBio*, 10(6), e02635-19.
- Harman, G.E., Chet, I. and Baker, R., 1980. *Trichoderma hamatum* effects on seed and seedling disease induced in radish and pea by *Pythium spp.* or *Rhizoctonia solani*. *Phytopathology*, 70(12), pp.1167-1172.
- Hayat, R., Ali, S., Amara, U., Khalid, R., & Ahmed, I., 2010. Soil beneficial bacteria and their role in plant growth promotion: a review. *Annals of Microbiology*, 60(4), pp.579-598.
- Hein, J.W., Wolfe, G.V. and Blee, K.A., 2008. Comparison of rhizosphere bacterial communities in *Arabidopsis thaliana* mutants for systemic acquired resistance. *Microbial ecology*, 55(2), pp.333-343.
- Hiltner, L., 1904. Über neuere erfahrungen und probleme auf dem debiete der bo denbakteriologie und unter besonderer berucksichtigung der grundund und brache. *Zbl. Bakteriol*, *2*, pp.14-25.
- Howard, M.M., Muñoz, C.A., Kao-Kniffin, J. and Kessler, A., 2020. Soil microbiomes from fallow fields have species-specific effects on crop growth and pest resistance. *Frontiers in Plant Science*, 11, p.1171.
- Huberty, M., Choi, Y.H., Heinen, R. and Bezemer, T.M., 2020. Above-ground plant metabolomic responses to plant-soil feedbacks and herbivory. *Journal of Ecology*, 2020(108), pp.1703-1712.
- Hugenholtz, P. and Tyson, G.W., 2008. Metagenomics. Nature, 455(7212), pp.481-483.
- Insam, H. and Seewald, M.S., 2010. Volatile organic compounds (VOCs) in soils. *Biology and fertility* of soils, 46(3), pp.199-213.

- Joosten, L., Mulder, P.P., Klinkhamer, P.G. and van Veen, J.A., 2009. Soil-borne microorganisms and soil-type affect pyrrolizidine alkaloids in *Jacobaea vulgaris*. *Plant and Soil*, 325(1-2), p.133.
- Jung, T., Cooke, D.E.L., Blaschke, H., Duncan, J.M. and Oßwald, W., 1999. *Phytophthora quercina* sp. nov., causing root rot of *European oaks*. *Mycological Research*, 103(7), pp.785-798.
- Kennedy, A.C. and Smith, K.L., 1995. Soil microbial diversity and the sustainability of agricultural soils. *Plant and soil*, 170(1), pp.75-86.
- Kerbarh, O., Ciulli, A., Howard, N.I. and Abell, C., 2005. Salicylate biosynthesis: overexpression, purification, and characterization of Irp9, a bifunctional salicylate synthase from *Yersinia enterocolitica. Journal of bacteriology*, 187(15), pp.5061-5066.
- Kim, J., Kim, M.S., Koh, A.Y., Xie, Y. and Zhan, X., 2016. FMAP: functional mapping and analysis pipeline for metagenomics and metatranscriptomics studies. *BMC bioinformatics*, 17(1), p.420.
- Kniskern, J.M., Traw, M.B. and Bergelson, J., 2007. Salicylic acid and jasmonic acid signaling defense pathways reduce natural bacterial diversity on *Arabidopsis thaliana*. *Molecular plant-microbe interactions*, 20(12), pp.1512-1522.
- Kos, M., Tuijl, M.A., de Roo, J., Mulder, P.P. and Bezemer, T.M., 2015. Plant-soil feedback effects on plant quality and performance of an aboveground herbivore interact with fertilisation. *Oikos*, 124(5), pp.658-667.
- Kostenko, O., van de Voorde, T.F., Mulder, P.P., van der Putten, W.H. and Martijn Bezemer, T., 2012. Legacy effects of aboveground-belowground interactions. *Ecology Letters*, *15*(8), pp.813-821.
- Kourtev, P.S., Ehrenfeld, J.G. and Häggblom, M., 2003. Experimental analysis of the effect of exotic and native plant species on the structure and function of soil microbial communities. *Soil Biology and Biochemistry*, 35(7), pp.895-905.
- Krehenwinkel, H., Pomerantz, A. and Prost, S., 2019. Genetic biomonitoring and biodiversity assessment using portable sequencing technologies: current uses and future directions. *Genes*, 10(11), p.858.
- Laurie-Berry, N., Joardar, V., Street, I.H. and Kunkel, B.N., 2006. The Arabidopsis thaliana JASMONATE INSENSITIVE 1 gene is required for suppression of salicylic acid-dependent defenses during infection by Pseudomonas syringae. Molecular Plant-Microbe Interactions, 19(7), pp.789-800.
- Lawrence, C.B., Singh, N.P., Qiu, J., Gardner, R.G. and Tuzun, S., 2000. Constitutive hydrolytic enzymes are associated with polygenic resistance of tomato to *Alternaria solani* and may function as an elicitor release mechanism. *Physiological and Molecular Plant Pathology*, 57(5), pp.211-220.
- Lawrence, P.K. and Koundal, K.R., 2002. Plant protease inhibitors in control of phytophagous insects. *Electronic Journal of Biotechnology*, 5(1), pp.5-6.
- Lebeis, S.L., Paredes, S.H., Lundberg, D.S., Breakfield, N., Gehring, J., McDonald, M., Malfatti, S., Del Rio, T.G., Jones, C.D., Tringe, S.G. and Dangl, J.L., 2015. Salicylic acid modulates colonization of the root microbiome by specific bacterial taxa. *Science*, 349(6250), pp.860-864.
- Li, N., Han, X., Feng, D., Yuan, D. and Huang, L.J., 2019. Signaling crosstalk between salicylic acid and ethylene/jasmonate in plant defense: do we understand what they are whispering? *International Journal of Molecular Sciences*, 20(3), p.671.
- Mandal, M.K., Pandey, D., Purwar, S., Singh, U.S. and Kumar, A., 2006. Influence of jasmonic acid as potential activator of induced resistance against Karnal bunt in developing spikes of wheat. *Journal of biosciences*, 31(5), pp.607-616.
- Mandal, S., Mallick, N. and Mitra, A., 2009. Salicylic acid-induced resistance to *Fusarium oxysporum* f. sp. lycopersici in tomato *lant Physiology and Biochemistry*, 47(7), pp.642-649.

- Mangan, S.A., Schnitzer, S.A., Herre, E.A., Mack, K.M., Valencia, M.C., Sanchez, E.I. and Bever, J.D., 2010. Negative plant-soil feedback predicts tree-species relative abundance in a tropical forest. *Nature*, 466(7307), pp.752-755.
- Manning, P., Morrison, S.A., Bonkowski, M. and Bardgett, R.D., 2008. Nitrogen enrichment modifies plant community structure via changes to plant-soil feedback. *Oecologia*, 157(4), pp.661-673.
- Martinez, X., Pozuelo, M., Pascal, V., Campos, D., Gut, I., Gut, M., Azpiroz, F., Guarner, F. and Manichanh, C., 2016. MetaTrans: an open-source pipeline for metatranscriptomics. *Scientific reports*, 6, p.26447.
- Mercado-Blanco, J., van der Drift, K.M., Olsson, P.E., Thomas-Oates, J.E., van Loon, L.C. and Bakker, P.A., 2001. Analysis of the pmsCEAB gene cluster involved in biosynthesis of salicylic acid and the siderophore pseudomonine in the biocontrol strain *Pseudomonas fluorescens* WCS374. *Journal of Bacteriology*, 183(6), pp.1909-1920.
- Moccia, K.M. and Lebeis, S.L., 2019. Microbial ecology: how to fight the establishment. *Current Biology*, 29(24), pp.R1320-R1323.
- Mousavi, S.R. and Eskandari, H., 2011. A general overview on intercropping and its advantages in sustainable agriculture. *Journal of Applied Environmental and Biological Sciences*, *1*(11), pp.482-486.
- Munné-Bosch, S. and Müller, M., 2013. Hormonal cross-talk in plant development and stress responses. *Frontiers in plant science*, *4*, p.529.
- Narayanasamy, S., Jarosz, Y., Muller, E.E., Heintz-Buschart, A., Herold, M., Kaysen, A., Laczny, C.C., Pinel, N., May, P. and Wilmes, P., 2016. IMP: a pipeline for reproducible referenceindependent integrated metagenomic and metatranscriptomic analyses. *Genome biology*, 17(1), pp.1-21.
- Navarro, L., Bari, R., Seilaniantz, A., Nemri, A. and Jones, J.D., 2008. Roles of plant hormones in plant resistance and susceptibility to pathogens. In *Genomics of Disease*. Springer, New York, NY, pp.1-10.
- Neher, D.A., 1999. Soil community composition and ecosystem processes: comparing agricultural ecosystems with natural ecosystems. Agroforestry Systems, 45(1-3), pp.159-185.
- Nelson, L.M., 2004. Plant growth promoting rhizobacteria (PGPR): Prospects for new inoculants. Crop management, 3(1), pp.1-7.
- Ni, Y., Li, J. and Panagiotou, G., 2016. COMAN: a web server for comprehensive metatranscriptomics analysis. BMC genomics, 17(1), p.622.
- Nijjer, S., Rogers, W.E. and Siemann, E., 2007. Negative plant-soil feedbacks may limit persistence of an invasive tree due to rapid accumulation of soil pathogens. *Proceedings of the Royal Society B: Biological Sciences*, 274(1625), pp.2621-2627.
- Nirenberg, H.I., 1981. A simplified method for identifying *Fusarium* spp. occurring on wheat. *Canadian Journal of Botany*, 59(9), pp.1599-1609.
- Nunan, N., Daniell, T.J., Singh, B.K., Papert, A., McNicol, J.W. and Prosser, J.I., 2005. Links between plant and rhizoplane bacterial communities in grassland soils, characterized using molecular techniques. *Appl. Environ. Microbiol*, 71(11), pp.6784-6792.
- Pendergast IV, T.H., Burke, D.J. and Carson, W.P., 2013. Belowground biotic complexity drives aboveground dynamics: a test of the soil community feedback model. *New Phytologist*, 197(4), pp.1300-1310.
- Pieterse, C.M., Zamioudis, C., Berendsen, R.L., Weller, D.M., Van Wees, S.C. and Bakker, P.A., 2014. Induced systemic resistance by beneficial microbes. *Annual review of phytopathology*, 52, pp.347-375.

- Quiroga, E.N., Sampietro, A.R. and Vattuone, M.A., 2001. Screening antifungal activities of selected medicinal plants. *Journal of ethnopharmacology*, 74(1), pp.89-96.
- Rahbé, Y., Deraison, C., Bonadé-Bottino, M., Girard, C., Nardon, C. and Jouanin, L., 2003. Effects of the cysteine protease inhibitor oryzacystatin (OC-I) on different aphids and reduced performance of *Myzus persicae* on OC-I expressing transgenic oilseed rape. *Plant science*, 164(4), pp.441-450.
- Reinhold-Hurek, B., Hurek, T., Gillis, M., Hoste, B., Vancanneyt, M., Kersters, K. and De Ley, J., 1993. Azoarcus gen. nov., nitrogen-fixing proteobacteria associated with roots of Kallar grass (Leptochloa fusca (L.) Kunth), and description of two species, Azoarcus indigens sp. nov. and Azoarcus communis sp. nov. International Journal of Systematic and Evolutionary Microbiology, 43(3), pp.574-584.
- Rolfe, S.A., Griffiths, J. and Ton, J., 2019. Crying out for help with root exudates: adaptive mechanisms by which stressed plants assemble health-promoting soil microbiomes. *Current opinion in microbiology*, 49, pp.73-82.
- Ruan, J., Zhou, Y., Zhou, M., Yan, J., Khurshid, M., Weng, W., Cheng, J. and Zhang, K., 2019. Jasmonic acid signaling pathway in plants. *International journal of molecular sciences*, 20(10), p.2479.
- Ryan, C.A., 1990. Protease inhibitors in plants: genes for improving defenses against insects and pathogens. Annual review of phytopathology, 28(1), pp.425-449.
- Saikia, R., Singh, T., Kumar, R., Srivastava, J., Srivastava, A.K., Singh, K. and Arora, D.K., 2003. Role of salicylic acid in systemic resistance induced by *Pseudomonas fluorescens* against *Fusarium* oxysporum f. sp. ciceri in chickpea. *Microbiological Research*, 158(3), pp.203-213.
- Schimel, J. and Schaeffer, S.M., 2012. Microbial control over carbon cycling in soil. Frontiers in microbiology, 3, p.348.
- Schlüter, J.P., Reinkensmeier, J., Daschkey, S., Evguenieva-Hackenberg, E., Janssen, S., Jänicke, S., Becker, J.D., Giegerich, R. and Becker, A., 2010. A genome-wide survey of sRNAs in the symbiotic nitrogen-fixing alpha-proteobacterium *Sinorhizobium meliloti. BMC* genomics, 11(1), p.245.
- Seskar, M., Shulaev, V. and Raskin, I., 1998. Endogenous methyl salicylate in pathogen-inoculated tobacco plants. *Plant physiology*, 116(1), pp.387-392.
- Singh, J.S. and Gupta, V.K., 2018. Soil microbial biomass: a key soil driver in management of ecosystem functioning. *Science of the Total Environment*, 634, pp.497-500.
- Sonnemann, I., Finkhaeuser, K. and Wolters, V., 2002. Does induced resistance in plants affect the belowground community? *Applied Soil Ecology*, 21(2), pp.179-185.
- Sturz, A.V. and Christie, B.R., 2003. Beneficial microbial allelopathies in the root zone: the management of soil quality and plant disease with rhizobacteria. *Soil and Tillage Research*, 72(2), pp.107-123.
- Turley, N.E., Godfrey, R.M. and Johnson, M.T., 2013. Evolution of mixed strategies of plant defense against herbivores. New Phytologist, 197(2), pp.359-361.
- Tyc, O., Song, C., Dickschat, J.S., Vos, M. and Garbeva, P., 2017. The ecological role of volatile and soluble secondary metabolites produced by soil bacteria. *Trends in microbiology*, 25(4), pp.280-292.
- Van der Does, D., Leon-Reyes, A., Koornneef, A., Van Verk, M.C., Rodenburg, N., Pauwels, L., Goossens, A., Körbes, A.P., Memelink, J., Ritsema, T. and Van Wees, S.C., 2013. Salicylic acid suppresses jasmonic acid signaling downstream of SCFCOII-JAZ by targeting GCC promoter motifs via transcription factor ORA59. *The Plant Cell*, 25(2), pp.744-761.

- Van der Heijden, M.G., Bardgett, R.D. and Van Straalen, N.M., 2008. The unseen majority: soil microbes as drivers of plant diversity and productivity in terrestrial ecosystems. *Ecology letters*, *11*(3), pp.296-310.
- Van der Heijden, M.G. and Wagg, C., 2013. Soil microbial diversity and agro-ecosystem functioning. *Plant and soil*, 363(1-2), pp.1-5.
- Van der Meij, A., Willemse, J., Schneijderberg, M.A., Geurts, R., Raaijmakers, J.M. and van Wezel, G.P., 2018. Inter-and intracellular colonization of *Arabidopsis* roots by endophytic actinobacteria and the impact of plant hormones on their antimicrobial activity. *Antonie Van Leeuwenhoek*, 111(5), pp.679-690.
- Van der Putten, W.H., Bardgett, R.D., Bever, J.D., Bezemer, T.M., Casper, B.B., Fukami, T., Kardol, P., Klironomos, J.N., Kulmatiski, A., Schweitzer, J.A. and Suding, K.N., 2013. Plant-soil feedbacks: the past, the present and future challenges. *Journal of Ecology*, 101(2), pp.265-276.
- Van de Voorde, K., Paauwe, J. and Van Veldhoven, M., 2012. Employee well-being and the HRMorganizational performance relationship: a review of quantitative studies. *International Journal of Management Reviews*, 14(4), pp.391-407.
- Van Loon, L.C. and Van Strien, E.A., 1999. The families of pathogenesis-related proteins, their activities, and comparative analysis of PR-1 type proteins. *Physiological and molecular plant pathology*, 55, pp.85-97.
- Van Loon, L.C., 2007. Plant responses to plant growth-promoting rhizobacteria. In New perspectives and approaches in plant growth-promoting Rhizobacteria research. Springer, Dordrecht, pp.243-254.
- Wang, B., Guo, X.W., Li, K., Han, X., Xu, S.J., Liu, Z.D., Guo, Y.S. and Xie, H.G., 2015. Effects of Salicylic acid on grape plants and the soil microbial community. *Allelopathy Journal*, 36(1), pp.49-61.
- Wang, M., Ruan, W., Kostenko, O., Carvalho, S., Hannula, S.E., Mulder, P.P., Bu, F., van der Putten, W.H. and Bezemer, T.M., 2019. Removal of soil biota alters soil feedback effects on plant growth and defense chemistry. *New Phytologist*, 221(3), pp.1478-1491.
- War, A.R., Paulraj, M.G., War, M.Y. and Ignacimuthu, S., 2011. Jasmonic acid-mediated-induced resistance in groundnut (*Arachis hypogaea* L.) against *Helicoverpa armigera* (Hubner) (Lepidoptera: Noctuidae). Journal of Plant Growth Regulation, 30(4), pp.512-523.
- Wasternack, C. and Strnad, M., 2018. Jasmonates: News on occurrence, biosynthesis, metabolism and action of an ancient group of signaling compounds. *International journal of molecular sciences*, 19(9), p.2539.
- Wei, G., Kloepper, J.W. and Tuzun, S., 1991. Induction of systemic resistance of cucumber to *Colletotrichum orbiculare* by select strains of plant growth-promoting rhizobacteria. *Phytopathology*, 81(11), pp.1508-1512.
- Wesson, R.G. and Wesson, R., 1993. Beyond natural selection. MIT press.
- Wink, M., 1988. Plant breeding: importance of plant secondary metabolites for protection against pathogens and herbivores. *Theoretical and applied genetics*, 75(2), pp.225-233.
- Zahar Haichar, F., Marol, C., Berge, O., Rangel-Castro, J.I., Prosser, J.I., Balesdent, J., Heulin, T. and Achouak, W., 2008. Plant host habitat and root exudates shape soil bacterial community structure. *The ISME journal*, 2(12), pp.1221-1230.
- Zak, D.R., Holmes, W.E., White, D.C., Peacock, A.D. and Tilman, D., 2003. Plant diversity, soil microbial communities, and ecosystem function: are there any links? *Ecology*, 84(8), pp.2042-2050.
- Zhang, M. and Yu, JR, 2006. History of Chinese science and technology. Jilin Literature and History Press.

Zhang, S., Zhang, Y. and Ma, K., 2015. The equal effectiveness of different defensive strategies. *Scientific reports*, 5, p.13049.

Chapter 2

Activation of hormone-associated plant defense pathways alters the effects of live soils on plant performance

> Jing Zhang^{1*} Klaas Vrieling¹ Peter G.L. Klinkhamer¹ T. Martijn Bezemer^{1,2}

 Plant Science and Natural Products, Institute of Biology Leiden (IBL), Leiden University, Sylviusweg 72, 2333BE Leiden, The Netherlands

2) The Netherlands Institute of Ecology (NIOO-KNAW), Department of Terrestrial Ecology, Droevendaalsesteeg 10, 6708 PB, Wageningen, The Netherlands

Abstract

Many plant species grow better in sterilized soil than in soil that contains a live microbial community. One hypothesis to explain this phenomenon is that the overall net pathogenic effect of soil microbial communities reduces plant performance. Induced plant defenses triggered by the application of the plant hormones jasmonic acid (JA) and salicylic acid (SA) may help to mitigate this pathogenic effect of live soil. However, little is known about how such hormonal application to the plant affects the soil and how this, in turn, impacts plant growth. We grew four plant species in sterilized and inoculated live soil and exposed their leaves to two hormonal treatments (JA and SA). Two species (Jacobaea vulgaris and Cirsium vulgare) were negatively affected by soil inoculation. In these two species foliar application of SA led to higher plant growth in live soil but not in sterilized soil. Two other species (Trifolium repens and Daucus carota) were not affected by soil inoculation and for these two species foliar application of SA reduced plant growth in both the sterilized and live soil. Application of JA reduced plant growth in both soils for all species. We subsequently carried out a multiple generation experiment for one of the plant species, J. vulgaris. In each generation, the live soil was a mixture of 10% soil from the previous generation and 90% sterilized soil and the same hormonal treatments were applied. The negative effects of live soil on plant growth were similar in all four generations, and this negative effect was mitigated by the application of SA. Our research suggests that the application of SA can mitigate the negative effects of live soil on plant growth. However, although the inoculum of soil containing a natural live soil microbial community had a strong negative effect on the growth of J. vulgaris, we found no evidence for an increase in the negative plant-soil feedback in either the control or the SA treated plants as plant performance did not decrease consistently with succeeding generations.

Keywords

Plant-soil interactions, Plant-soil feedback, Induced resistance, Rhizosphere soil, Salicylic acid, Jasmonic acid
Introduction

The interactions between plants and soil microorganisms have long been recognized for their importance in terrestrial ecological systems (Bever 1994; van der Heijden et al., 2008). Although the effects may vary depending on the plant species and the soils tested, in the majority of cases the soil microbial community has a negative effect on plant growth (Kulmatiski et al., 2008). Plants also affect the composition of the soil microbial community, which, in turn, will impact plant growth. The process is called plant-soil feedback (Bever et al., 1997; Van Breemen and Finzi, 1998). Most plant species exhibit negative conspecific soil feedbacks. This means that they grow worse in soil, in which the same species has been grown than in soil where other species have grown (Kulmatiski et al., 2008). From natural situations and agriculture, it is well-known that soil can become less suitable for a species if this species is grown in the same soil for multiple generations. This negative effect is thought to be caused by soil pathogens or root herbivores, allelopathy, nutrient immobilization or nutrient depletion (Miki, 2012). In some cases, plants also cause positive plant-soil feedbacks and these can be mediated by plant promoting rhizobacteria, mycorrhizal fungi or other unknown mechanisms (Revillini et al., 2016; van der Putten 2017).

Plant-induced resistance has been regarded as a promising defense strategy against pathogens or herbivores (Haney and Ausubel, 2015; Lebeis et al., 2015; Yang et al., 2015). In nature, plants are exposed to complex selection pressures, involving both abiotic and biotic stresses. Plants are under constant attack by a myriad of pathogens and pests and have to compete with neighboring plants. As a result, plants have evolved a wide range of responses to cope with biotic stresses. The abilities of plants to respond to different biotic stresses are regulated through sophisticated interacting hormonal signaling networks (Bezemer and van Dam, 2005; Fujita et al., 2006; Arnaud and Hwang, 2015). Phytohormones are a group of natural plant compounds with low molecular weights. Salicylic acid (SA), Jasmonic acid (JA), ethylene (ET), abscisic acid (ABA), methyl jasmonate (MeJA), auxin, cytokinins (CKs), gibberellins and brassinosteroids are commonly studied phytohormones. Plant hormones have been implicated to be involved in defense pathways, the key regulator against pathogens and pests, are the phytohormones JA and SA (Bari and Jones, 2009). Experimental

evidence indicates that application of SA to plant leaves, activates systemic acquired resistance (SAR) against pathogens (Reymond and Farmer, 1998; Mandal et al., 2009). JA, in turn, activates induced defenses against herbivores and necrotrophic pathogens (Nahar et al., 2011). Although to some extent, the SA or JA-induced hormonal signaling pathway could interact with other phytohormones, such as CKs, ET, ABA and auxins, they do show clear effects on the plant's defense system when applied as single hormones (Fujita et al., 2006; Yang et al., 2015; Berens et al., 2019). The crosstalk between SA or JA and other hormones is still not fully understood.

A still uncharted territory is how plant hormone-activated signaling pathways impact soil microbial communities and how these, in turn, affect plant growth. Here we restricted ourselves to two prime hormones involved in activating defense pathways, SA and JA. We aimed to quantify the effect of induced SA or JA resistance on the soil microbial communities that affect plant growth. If the negative effect of the soil containing a live soil microbial community on plant growth is caused by an overall pathogenic effect we expect that activating SA signaling by exogenous application mitigates these negative effects. As a result, we expect that the effect of SA application on plant growth differs between plants in sterile soil and in live soil. Exogenous application of JA typically induces resistance against herbivores and necrotrophic pathogens (Pieterse et al., 2009; van Dam and Oomen, 2014; Carvalhais et al., 2017; Palmer et al., 2017). JA signaling can exhibit negative crosstalk with SA signaling (Leon-Reves et al., 2010). One can therefore hypothesize that activating JA signaling will reduce the ability of plants to cope with pathogens (which causes induction of the SA pathway in the plant) and thus will increase the overall negative effect of the soil microbial community on plant growth. The responses of plants, after activating hormonal defense pathways, to an inoculum containing a live soil microbial community are, as yet, not well studied and understood. Moreover, the evidence for the existence of such effects is contradictory. Activation of JA and SA signaling pathways did not affect the resident soil microflora in several studies (Doornbos et al., 2012; Berendsen et al., 2012; Rashid and Chung, 2017), but a more recent study showed that SA modulates colonization of the root microbiome by specific bacterial taxa (Lebeis et al., 2015).

2

If the induction of signaling pathways in the plant leads to changes in the composition of the soil microbial community, its effect is likely to extend over time or plant generations. Potentially this could lead to the selection of more beneficial soil microbial communities either by suppressing pathogens or by promoting beneficial microbes. As far as we are aware, the effects of plant hormones through plants on soils containing a live microbial community over multiple generations have not been studied so far, despite its potential to select for more beneficial soils containing plant growth-promoting microbial communities in agriculture.

In a preliminary experiment, we found strong evidence for negative effects of soil that consisted of a mixture of 90% sterilized soil and 10% live soil on the growth of common ragwort (*Jacobaea vulgaris*), compared to sterilized soil. After treating plants with SA, this negative effect diminished. Based on these findings, we grew four different plant species individually in both sterilized soil and live soil. For *J. vulgaris*, the species which showed the strongest negative effect towards the live soil, and for which this negative effect was mitigated by foliar application of SA, we grew plants for three more generations. For each generation, sterilized soil was inoculated with live soil from the previous generation from the same treatment. We addressed four questions: (1) Do the effects of live soil on plant growth differ among plant species? (2) Does the foliar application of JA and SA alter the effects of the live soil? (3) Does the negative effect of live soil change in four successive generations of *J. vulgaris* for control plants and plants treated with SA and SA or JA.

Materials and methods

Plant material and seeds germination

Jacobaea vulgaris (common ragwort), Cirsium vulgare (bull thistle), Trifolium repens (white clover) and Daucus carota (wild carrot), were chosen because they are common native species at the dune area where we collected soil. We collected seeds at the dunes for J. vulgaris, C. vulgare, and D. carota. T. repens seeds were bought from Cruydt-Hoeck a seed company that sells seeds of wild plant species (Nijeberkoop, The Netherlands). Prior to seed germination, all seeds were shaken for

2 min in 70% ethanol, then washed with sterilized water, put for 12 min in 2% bleach, and finally rinsed four times with sterilized water to minimize influences of seedborne microbes.

Soil material

The soil was collected at Meijendel, a calcareous sandy area from a coastal dune area north of The Hague, The Netherlands (52°11'N, 4°31'E). The topsoil was collected to a depth of 15 cm after removing the grassland vegetation and the organic layer of the surface. The soil was sieved using a 5 mm sized mesh, homogenized with a concrete mixer, and then stored into 20-liter plastic bags (Nasco Whirl-Pak Sample Bag). Bags were either sterilized by 35-K Gray gamma-irradiation (Synergy Health Company, Ede, The Netherlands) or kept at 4°C for inoculation.

Plant growth and foliar application of hormones

Surface sterilized seeds of the four species (J. vulgaris, C. vulgare, T. repens and D. carota) were germinated in sterile Petri dishes on filter paper. After one week, 60 seedlings per species were planted individually in 500 ml pots containing either sterilized soil or inoculated live soil. The live soil consisted of a mixture of 90% sterilized soil and 10% live soil. Nutrient availability often increases after sterilization of the soil, and we therefore inoculated the sterilized soil rather than using pure live soil, to enable comparison of the two types of soil. Sterilized soil and live soil were kept in bags and left in the climate room for 14 days to enable the establishment of microbial communities in the inoculated soil before potting. Before potting, the soil in each bag was mixed. After planting the seedlings, pots were randomly distributed over a climate room (relative humidity 70%, light 16h at 20°C, dark 8h at 20°C). Plants were watered regularly with Milli-Q water. Five ml Steiner nutrient solution was added per plant on day seven. Ten ml Steiner nutrient solution was added on day 13, and 20 ml Steiner nutrient solution was added on days 19, 28, 37, 42. The Steiner nutrient solution (Steiner, 1980) was prepared from seven different stock solutions (106.2 g Ca(NO₃)₂·4H₂O, 29.3 g KNO₃, 13.6 g KH₂PO₄, 49.2 g MgSO₄·7H₂O, 25.2 g K₂SO₄, 2.24 g KOH and 3.29 g Fe-EDTA added to 1 liter demineralized water, and a stock solution with micro elements (a mixed solution of 0.181 g MnCl₂·4H₂O, 0.286

g H₃BO₃, 0.022 g ZnSO₄·7H₂O, 0.0078 g CuSO₄·5H₂O and 0.0126 g NaMoO₄·2H₂O added to 1 liter demineralized water). Ten ml of each stock solution was diluted in 1 liter of demineralized water before use.

The pots for each species were divided over six treatments: two soil treatments (sterilized soil and live soil) and three hormonal treatments (JA, SA and control (only solvent)). Each treatment was replicated 10 times. The experiment, therefore, consisted of 240 pots (4 species \times 2 soil treatments \times 3 hormonal treatments \times 10 replicates). The plant hormones JA and SA were applied through foliar application three times a week for four consecutive weeks. The first application was given when plants were 14 days old. Either 0.75 ± 0.05 ml of 100 μ M JA or SA was sprayed on the leaves while carefully avoiding spillover to the soil. One week later the treatment was repeated with 1.50 ± 0.05 ml of 100 μ M JA or SA. In the next week, the treatment was repeated with 2.25 ± 0.05 ml of 100 μ M JA or SA. The JA-solution was prepared by adding 105.135 µl JA stock solution into Milli-Q water until a final volume of 500 ml. The JA stock solution was prepared by adding 500 mg JA to 5 ml ethanol. JA was purchased from Cayman Chemical Company (product number: 88300). SA (purchased from Sigma-Aldrich, \geq 99.0%) was made by dissolving 6.9055 mg in 69.055 µl of ethanol to which Milli-Q water was added until a final volume of 500 ml. Control plants were sprayed with sterile water with the same solvent (85 µl ethanol in 500 ml Milli-Q water).

Harvesting plants and soil samples

Fifty-four days after planting, all plants were harvested, except for *C. vulgare*. *C. vulgare* plants were considerably larger than the other species and were therefore harvested after 45 days to prevent pot size becoming limit growth. Plants were gently removed from the pots. Shoots were separated from roots with a scissor just above the root crown, and roots were rinsed with water and then put into paper bags. Harvested plant parts were oven-dried at 60°C for approximately one week. The dry weight of roots and shoots was determined until the nearest 0.1 mg. The rhizosphere soil was harvested individually from each pot by gently shaking the roots and soil three times to remove the loosely adhering soil, after which rhizosphere soil samples were collected onto a sterile filter paper by removing the remnant soil from the roots

with a fine sterilized brush. Finally, all the labeled soil samples were transferred to a 4°C room and stored for the multiple generation experiment.

Multi-generation experiment

J. vulgaris was chosen for the multiple generation selection experiment to examine if the observed effect on plant biomass of the first generation would increase further over later generations. For *J. vulgaris* we grew the plants from each of the six treatments (sterilized and live soils, two hormone treatments and control) for another three generations under the same conditions as described for the first generation. The only difference being that each time, the soil inoculate was derived from the previous generation from the same treatment, 100% sterilized soil was used as control. A schematic drawing of the experiment is presented in Fig. 1. Fourteen days after mixing the sterilized and live soil, a single *J. vulgaris* seedling was planted into each pot. All replicate rhizosphere soils from a single treatment were mixed before inoculation to avoid a selection of particular microbial species in individual pots. All treatments were carried out as described above.



Fig. 1 Experimental design of the multigeneration experiment with *J. vulgaris*. Soil used for the 1^{st} generation was a mixture of 90% sterilized soil and 10% live soil both collected from the dunes. Soil used for the 2^{nd} , 3^{rd} and 4^{th} generations was a mixture of 10% rhizosphere soil collected from the previous generation from the same treatment and 90% sterilized soil collected from the dunes. In each generation we tested two hormonal treatments in inoculated and 100% sterilized soil. JA denotes foliar application of jasmonic acid, SA denotes foliar application of salicylic acid and

C denotes control. In each treatment 10 replicates were used even though only three are depicted.

Statistical analysis

Data were first checked for homogeneity of variance and normal distribution of errors and data were transformed when necessary. To test whether the effect of the live soil was different among the four species we performed a three-way ANOVA on the total data set of the first experiment with soil (sterilized and live, 2 levels)", hormones (3 levels) and species (4 levels) as fixed factors, Relative plant dry mass was used as a dependent variable and was arcsine square-root transformed prior to analysis. Relative plant dry mass was calculated as 100 times the dry mass of a plant divided by the average dry mass of the control plants from the same species in the sterilized soil. In this way, the average dry mass of the control plants in the sterilized soil was set at 100 for each of the four species. By doing so we removed species-specific size differences enabling to make the data more comparable among species. This analysis showed a significant soil \times species interaction (see results section). On basis of this we divided the data set in two groups. One group for the two species that were negatively affected by the lives soil and one group for the two species that were not. We did this because we expected the effect of the hormonal treatments to be only present for the species that were affected by the live soil. To answer the question if the effect of the live soil was affected by foliar application of hormones, we performed four three-way ANOVAs (for the two groups of species and the two hormonal treatments) with plant mass as dependent variable and species (2 levels), soil (2 levels) and hormonal treatment (2 levels) as fixed factors. Usually, the negative effects of live soils on plant biomass are stronger in the roots than the shoots, thus we also carried out three-way ANOVA analysis for shoot-root ratios of the four plant species.

To answer the question whether the negative effect of live soil changes in four successive generations of *J. vulgaris* for control plants and plants treated with SA or JA we used a three-way ANOVA with soil (2 levels), generation (4 levels) and hormones (3 levels) as a fixed factor, and log-transformed plant dry mass or shoot-root ratio as dependent variables. We furthermore compared the effects of the two hormones separately using a three-way ANOVA with log-transformed plant dry mass

as a dependent variable and soil (2 levels), hormone (2 levels) and generation (4 levels) as fixed factors. Differences between treatments were tested with a Tukey post-hoc test.

We used a linear regression model to estimate the effects of SA and JA on the growth of J. vulgaris over four consecutive generations in both sterilized and live soil. In the regression model, the dry mass of plants of the SA or JA treatment divided by dry mass of control plants was the dependent variable and generation was the independent variable. Since we could not pair the pots (SA or JA/control) and we only had 10 replicates for each treatment, we used a Monte-Carlo simulation to test if the linear regression model differed from y = 1. Each time we randomly paired one plant of the hormone treatment and one plant of the control to calculate the ratio of the dry mass of treated and dry mass of control. Then we repeated this procedure 1000 times, to obtain 1000 ratios of each generation per soil. Then we took the mean of 1000 ratios per generation to fit linear regression models for the two soils, respectively. To test whether the linear regressions in sterilized and live soils differed from y = 1, we calculated the 95% confidence intervals (CFI) of the slopes for both soils. We also tested whether the two linear regression models differed between sterilized and live soils with ANCOVA (analysis of covariance) analysis. All analyses were performed in IBM SPSS Statistics 25.

Results

Do the effects of live soil on plant growth differ among plant species?

For *J. vulgaris* and *C. vulgare*, biomass in live soils was about half that in sterilized soils. This negative effect of the live soil was present irrespective of the hormonal treatment. For the other two species (*T. repens* and *D. carota*) biomass was not significantly different in live and sterilized soils (Fig. 2, Table 1). The difference in response to live soils among the four species is reflected by the highly significant interaction term (species \times soil) in the ANOVA (Table 1)



Fig. 2 Mean (+ SE) relative plant dry mass (%) of *J. vulgaris, C. vulgare, T. repens* and *D. carota* plants treated with JA and SA in sterilized soil and live soil. C represents the control treatment. Note: within species different letters above bars indicate significant differences between treatments based on a Tukey post-hoc test for each single species. N=10.

Table 1 Three-way ANOVA of arcsine square-root transformed relative plant dry mass of *J. vulgaris*, *C. vulgare*, *T. repens* and *D. carota* in live and sterilized soil for plants treated with JA or SA and for control plants. df = degrees of freedom.

Source of variations	df	F-value	Р
species	3, 239	53.67	***
soil	1, 239	147.78	***
hormone	2, 239	27.17	***
species × soil	3, 239	59.81	***
species × hormone	6, 239	0.45	ns
soil × hormone	2, 239	4.75	**
species × soil × hormone	6, 239	1.48	ns

** P < 0.01, *** P < 0.001, ns not significant.

Does the foliar application of JA and SA alter the effects of live soil on plant growth for those species that were negatively affected by live soil?

Salicylic acid.

For the two species (*J. vulgaris and C. vulgare*) that were negatively affected by the live soil, foliar application of SA reduced the biomass for plants grown in the sterilized soil while it increased the biomass for plants grown in the live soil (Fig. 2). As a result, the main effect of SA in the ANOVA was not significant (Table 2). Although by itself the differences between the SA treatment and the control were not significant (Fig. 2), the effect of the SA treatment, as we hypothesized, depended strongly on soil type as is reflected by the significant soil × hormone interaction term in the ANOVA (Table 2). For the two species (*T. repens, D. carota*) that were not negatively affected by the live soil foliar application of SA reduced plant biomass in both soils, although this effect was not significant (Fig. 2, Table 2).

Jasmonic acid

Foliar application of JA decreased plant mass in all plant species in both sterilized and live soils. For the two species that were negatively affected by live soil the negative effect of JA was stronger in sterilized soils than in live soil (Fig. 2). This difference in response between plants grown in the two soils was significant as reflected by the soil × hormone interaction term in the ANOVA (Table 2). For the two species that did not grew less well in the live soils, such a difference in the response to JA application in the two soils was not found (Table 2). **Table 2** Three-way ANOVAs of arcsine square-root transformed relative plant dry mass for species that grew less well in live soil compared to sterilized soil (upper part) and for species that were not negatively affected by the live soil (lower part). Left: hormonal treatment is foliar application of SA. Right: hormonal treatment is foliar application of JA. Species, soil (live or sterilized), and hormone treatment were used as fixed factors. df = degrees of freedom.

		SA	SA treatment			JA treatment		
Species respond to soil effect	Source of variations	df	F- value	Р	df	F- value	Р	
	species	1, 79	10.00	*	1, 79	5.25	*	
	soil	1, 79	190.26	**	1, 79	191.88	***	
Yes	hormone	1, 79	2.87	ns	1, 79	21.04	***	
(I miloavis	species × soil	1, 79	11.17	**	1, 79	12.57	**	
(J. vulgaris C. vulgare)	species × hormone	1, 79	0.35	ns	1, 79	0.08	ns	
0 /	soil × hormone	1, 79	8.20	**	1, 79	8.49	**	
	species \times soil \times hormone	1, 79	0.05	ns	1, 79	0.00	ns	
	species	1, 79	7.56	**	1, 79	5.74	*	
	soil	1, 79	0.92	ns	1, 79	1.97	ns	
No	hormone	1, 79	0.35	ns	1, 79	32.94	***	
(T ronous	species × soil	1, 79	0.48	ns	1, 79	0.36	ns	
D. carota)	species × hormone	1, 79	1.10	ns	1, 79	1.86	ns	
	soil \times hormone	1, 79	1.21	ns	1, 79	1.58	ns	
	species \times soil \times hormone	1, 79	0.08	ns	1, 79	1.44	ns	

* P < 0.05, ** P < 0.01, *** P < 0.001, ns not significant.

The shoot-root ratio of plants differed among species soils and hormone treatments (Table S1). Except for *T. repens*, JA application increased the shoot-root ratio. We found no significant effects of SA application on the shoot-root ratio. The effects of hormone application on the shoot-root ratio varied among species and soils. In all species, the shoot-root ratio was on average higher in live soils (Fig. S1).

Does the negative effect of live soil change in four successive generations of *J*. *vulgaris* for control plants and plants treated with SA or JA?

The effect of the live soil across generations.

As in generation 1, in all three subsequent generations plants grew less well in the live soil than in the sterilized soil. Although the strength of this effect varied among generations there was no clear trend across subsequent generations (Fig. 3, Table 3).



Fig. 3 Mean (+ SE) plant dry mass of *J. vulgaris* during four successive generations treated with JA and SA in sterilized soil and live soil. C represents the control treatment. For each generation soil from the previous generation and originating from the same treatment was used as an inoculum. Within each generation, different letters above bars indicate significant differences between treatment groups based on a Tukey post-hoc test. N=10.

Source of variation	df	F-value	Р
soil	1,250	569.88	***
hormone	2,250	39.83	***
generation	3, 250	68.36	***
soil × hormone	2,250	8.17	***
soil \times generation	3, 250	57.96	***
hormone × generation	6,250	1.88	ns
soil \times hormone \times generation	6,250	0.68	ns

Table 3 Three-way ANOVA of log-transformed plant dry mass of *J. vulgaris* during four generations in live and sterilized soils after JA, SA or control treatment. df = degrees of freedom.

*** P < 0.001, ns not significant

The effect of foliar application of SA across generations

Again, as in generation 1, in all three subsequent generations foliar application of SA reduced plant biomass in sterilized soil and increased plant biomass in live soils (Fig. 3, Table 4). Although within generations and soils these differences were not significant, plants responded clearly different to the SA treatment in the two soils as is reflected by the significant soil x hormone interaction term in the ANOVA (Table 4, left part). The effect of foliar SA application did not differ among generations as was reflected by the non-significant interaction term in the ANOVA (Table 4). To examine if the effect of hormone application in live and sterilized soils showed a trend over generations in more detail, we regressed the ratio between the dry mass of SAtreated and control plants in both sterilized and live soils against generations (Fig. 4). This ratio was higher than 1 for all generations in live soils while it was close to 1 in sterilized soils. This difference between the two soils was significant (ANCOVA df =(1, 7), F = 20.18, P < 0.01, Fig. 4A). The slopes of the regressions for both sterilized and live soils did not significantly differ from 0 (for sterilized soil the lower and upper 95% CFIs are -0.15 and 0.19; for live soil the lower and upper 95% CFIs are -0.13 and (0.33) The latter results indicate that there is no significant trend in the effect of foliar application of SA over generations.

The effect of foliar application of JA across generations

As in generation 1, in all three subsequent generations foliar application of JA reduced plant biomass in both sterilized and live soil (Fig. 3, Table 4). This reduction was less strong in live soils, as is reflected by the significant soil \times hormone interaction term in the ANOVA (Table 4, right part). The effect of foliar JA application did not differ among generations as was reflected by the non-significant interaction term in the ANOVA (Table 4, right part). To examine if the effect of JA application in live and sterilized soils showed a trend over generations in more detail, we regressed the ratio between the dry mass of JA-treated and control plants in both sterilized and live soils against generations (Fig. 4). This ratio was lower than 1 for all generations in both live soils and sterilized soils. The ratios did not differ between the two soils (ANCOVA df = (1, 7), F = 0.01, P > 0.05, Fig. 4B). The latter result is somewhat surprising given the significant interaction we found between the effects of JA application and soil type in Table 4. The slopes of the regressions for both sterilized and live soils did not significantly differ from 0 (for sterilized soil the 95% CFI is -0.31 to 0.21; for live soil the 95% CFI is -1.8 to 0.24). The latter results indicate that there is no significant trend in the effect of foliar application of JA over generations.

Table 4 Three-way ANOVAs of plant dry mass of *J. vulgaris* during four generations in live and sterilized soils with soil (live and sterilized soils) generation, hormone (control and SA or JA) as fixed factors. df = degrees of freedom.

	SA treatment			JA treatment			
Source of variations	df	F-value	Р	df	F-value	Р	
soil	1, 164	241.79	***	1, 170	307.05	***	
hormone	1, 164	0.28	ns	1, 170	39.11	***	
generation	3, 164	8.98	***	3, 170	11.07	***	
soil × hormone	1, 164	8.75	**	1, 170	11.36	**	
soil \times generation	3, 164	7.98	***	3, 170	7.12	***	
hormone × generation	3, 164	0.17	ns	3, 170	0.14	ns	
soil \times hormone \times generation	3, 164	0.50	ns	3, 170	0.70	ns	

** *P* < 0.01, *** *P* < 0.001, ns not significant.



Fig. 4 The ratio of dry mass of hormone treated *J. vulgaris* plants divided by control plants in both sterilized and live soil for four generations. (A) SA treated plants (B) JA treated plants. C represents control treatment. The dashed line indicates y = 1. The data points are the average of 1000 ratios of dry mass of SA or JA and dry mass of control for each generation, the error bars represent the 95% confidence intervals of 1000 ratios for each generation (see material and methods for details).

The shoot-root ratio of *J. vulgaris* differed among generations and was affected by soil and hormone treatments (Table S2). While the effects of hormone application on the shoot-root ratio did not vary among generations, the effects of the hormone treatments differed among soils. In general, in both sterilized and live soils, application of JA increased shoot-root ratios relative to the control and the SA treatments except for the third generation in live soil (Fig. S2). Application of SA did not affect the shoot-root ratio across generations in the sterilized and live soils.

Discussion

In this study, we examined how exogenous application of the plant signaling hormones SA and JA interacts with the effects of inoculation of soil on plant growth of different plant species and how those effects altered plant performance during multiple consecutive generations. In two of the four species, plant biomass was lower in live soil than in sterilized soil. We found that foliar application of SA mitigated the negative effects of the live soil on plant performance for these species. We then grew *J. vulgaris* for three additional generations and found that SA application mitigated the negative effect of live soil on plant growth in all four generations, and that this overall effect was significant.

Our results show that the effect of the live soil on plant growth strongly varied among plant species although all plants received the same soil inoculum containing a natural soil microbial community and the growth conditions were identical for all four species. Species-specific effects have also been found in other plant-soil feedback experiments that showed that *J. vulgaris* and *C. vulgare* responded negatively to soil conditioning by conspecifics, while this is not the case for *T. repens* and *D. carota* (Klironomos, 2002; Joosten et al., 2009; Harrison and Bardgett, 2010; Wang et al., 2019). Together these results show that the responses of plant species to live soil are highly species-specific. Other studies have suggested that net positive or negative plant-soil feedback effects are related to the capacity of plants to cope with biotic or abiotic stresses, to influence soil nutrients, or to the way they impact soil microbial communities (Bezemer et al., 2006; van der Heijden et al., 2008; Eisenhauer et al., 2011). The soil microbial community present in live soil might have pathogenic effects on plant growth. This is in line with previous studies that indicate that soil sterilization enhanced plant growth by killing soil-borne pathogens in crops (Li et al., 2019).

The effect of SA application also varied among plant species. Interestingly, a positive effect of SA on plant growth in live soils occurred in *J. vulgaris* and *C. vulgare*, the two species that responded negatively to exposure to the live soil, and not in the other two species, *T. repens* and *D. carota* that were unresponsive to the soil with a live soil inoculum. These results strongly suggest that the negative impact of the live soil on plant growth is driven by pathogens. The difference in response between the four species can have different non-exclusive causes. The pathogenic effect of the live soils itself may differ among plant species due to specificity of the soil microbial species in the live soil inoculum, or due to inherent plant characteristics. We started with our hormone application when seedlings were 14 days old. In retrospect, we should have started earlier. The negative effects of the live soil on plant growth are most apparent during the first few weeks of plant growth (Jing et al., Chapter 5; Bezemer et al., 2018). If we would have applied the exogenous SA earlier, effects may therefore have been sol

stronger because plants in our study may have outgrown the negative effects of the live soil e.g. by upregulating their defense system (Vernooij et al., 1994; Métrauxs, 2001). In addition, in this paper, we used only one concentration of the phytohormones. Plant species may have a different sensitivity to the foliar application of these hormones, and the species that did not show a response may have responded to higher concentrations. It is important to note that, in this paper, we performed experiments with two phytohormones. Other plant hormones like auxins and cytokinins have been reported to play a role in fighting off the potentially pathogenic bacteria in the live soil via changing physiological and morphological features of plants (Hamill, 1993; Clarke et al., 2000). They may interact with JA or SA signaling pathways; however, this is still not fully understood. Applying combinations of different phytohormones would present a next logical step. To find a clear effect of SA and JA on plant growth against the pathogenic effect caused by live soils is the base for carrying out more extensive experiments. For example, in further tests, different plant hormones and their crosstalk effects could be tested.

Importantly, application of SA mitigated the negative effects of the live soil on the growth of J. vulgaris in all four generations. Sterilization of the soil resulted in higher plant growth, indicating an overall pathogenic effect due to soil-borne pathogens, and SA-induced resistance may help to mitigate this pathogenic effect caused by soil pathogens. Activation of SA-dependent signaling pathway leads to the expression of pathogenesis-related proteins (PRP) contributing to resistance, by limiting pathogen growth, the access of pathogens to water and nutrients in the plant, or by changing the composition of the cell wall of the plant (O'Donnell, et al., 2001; Heil, 2002; Glazebrook, 2005; Spoel et al., 2007). All this can result in higher plant mass in SAtreated plants than in control plants in live soil. In addition, activation of SA pathway regulates a myriad of compounds and enzymes, for example, peroxidase (POD), polyphenol oxidase (PPO), superoxide dismutase (SOD) etc., and those compounds play an important role in plant SA-induced defense against biotic stresses caused by pathogens (Achuo et al., 2004; War et al., 2011; Bakker et al., 2018). The effects of SA in the first generation were similar to those observed in the second or later generations indicating that SA application did not result in selection for more beneficial soil microbial communities over time. In part, this may be an artefact of the

experimental set-up. For each generation, we used an inoculum, which means that we placed a subset of the microbial community in a sterile background. This may have led to selection for microbes with similar particular characteristics in each of the four generations. However, we urge not to overemphasize the conclusion that application of SA results in a change in the effects of live soils on plant growth over generations (see chapters 3 and 4). Future studies should also include a comparison between the growth of SA-treated plants and control plants grown in soils that are conditioned by either SA-treated plants and control plants in a full factorial design.

JA-induced defenses are activated in response to herbivore attack, the infection of necrotrophic organisms or nematodes (Pieterse et al., 2009; Nahar, et al., 2011; van Dam and Oomen, 2014; Carvalhais et al., 2017; Palmer et al., 2017). In our study, the foliar JA application did not clearly mitigate the negative effects of the live soil on plant performance. Instead, it led to a significant negative effect on plant growth. This exemplifies that hormonal signaling is costly for plants (Baldwin, 1998; Vos et al., 2013).

In conclusion, our study suggests that negative effects in live soil on plant growth can be mitigated with foliar applications of SA. Sterilization benefited plant growth for two of the four species we investigated, suggesting the microbial community in live soils contains pathogens. For *J. vulgaris*, the plant species that responded most strongly to SA application, we did not observe an increasingly stronger effect on plant growth over further plant generations, but instead, the effect was stable over time. To better understand what caused the positive effect of SA application on plant growth in live soil, we examined changes in the diversity and functional role of the soil microbial community in live soil in Chapters 3 and 4.

Acknowledgements

We would like to thank Karin van der Veen-van Wijk for assistance at the final biomass harvest, Jan Vink for driving us to the field site to collect soil used for the inoculation experiment, Gang Chen for collecting soil, and the China Scholarship Council for financial support.

References

- Achuo, E.A., Audenaert, K., Meziane, H. and Höfte, M., 2004. The salicylic acid-dependent defence pathway is effective against different pathogens in tomato and tobacco. *Plant Pathology*, 53(1), pp.65-72.
- Arnaud, D. and Hwang, I., 2015. A sophisticated network of signaling pathways regulates stomatal defenses to bacterial pathogens. *Molecular plant*, 8(4), pp.566-581.
- Bakker, M.G., Chaparro, J.M., Manter, D.K. and Vivanco, J.M., 2015. Impacts of bulk soil microbial community structure on rhizosphere microbiomes of *Zea mays. Plant and Soil*, 392(1-2), pp.115-126.
- Baldwin, I.T., 1998. Jasmonate-induced responses are costly but benefit plants under attack in native populations. *Proceedings of the National Academy of Sciences*, U.S.A, 95(14), pp.8113-8118.
- Bari, R. and Jones, J.D., 2009. Role of plant hormones in plant defence responses. *Plant molecular biology*, 69(4), pp.473-488.
- Berendsen, R.L., Pieterse, C.M. and Bakker, P.A., 2012. The rhizosphere microbiome and plant health. *Trends in plant science*, 17(8), pp.478-486.
- Berens, M.L., Wolinska, K.W., Spaepen, S., Ziegler, J., Nobori, T., Nair, A., Krüler, V., Winkelmüller, T.M., Wang, Y., Mine, A. and Becker, D., 2019. Balancing trade-offs between biotic and abiotic stress responses through leaf age-dependent variation in stress hormone crosstalk. *Proceedings of the National Academy of Sciences*, U.S.A, 116(6), pp.2364-2373.
- Bever, J.D., 1994. Feeback between plants and their soil communities in an old field community. *Ecology*, 75(7), pp.1965-1977.
- Bever, J.D., Westover, K.M. and Antonovics, J., 1997. Incorporating the soil community into plant population dynamics: the utility of the feedback approach. *Journal of Ecology*, pp.561-573.
- Bezemer, T.M. and van Dam, N.M., 2005. Linking aboveground and belowground interactions via induced plant defenses. *Trends in Ecology & Evolution*, 20(11), pp.617-624.
- Bezemer, T.M., Lawson, C.S., Hedlund, K., Edwards, A.R., Brook, A.J., Igual, J.M., Mortimer, S.R. and Van der Putten, W.H., 2006. Plant species and functional group effects on abiotic and microbial soil properties and plant-soil feedback responses in two grasslands. *Journal of Ecology*, 94(5), pp.893-904.
- Bezemer, T.M., Jing, J., Bakx-Schotman, J.T. and Bijleveld, E.J., 2018. Plant competition alters the temporal dynamics of plant-soil feedbacks. *Journal of Ecology*, 106(6), pp.2287-2300.
- Carvalhais, L.C., Schenk, P.M. and Dennis, P.G., 2017. Jasmonic acid signalling and the plant holobiont. *Current opinion in microbiology*, *37*, pp.42-47.
- Clarke, S.F., Burritt, D.J., Jameson, P.E. and Guy, P.L., 2000. Effects of plant hormones on white clover mosaic potexvirus double-stranded RNA. *Plant Pathology*, 49(4), pp.428-434.
- Doornbos, R.F., van Loon, L.C. and Bakker, P.A., 2012. Impact of root exudates and plant defense signaling on bacterial communities in the rhizosphere. A review. *Agronomy for Sustainable Development*, 32(1), pp.227-243.
- Eisenhauer, N., Milcu, A., Sabais, A.C., Bessler, H., Brenner, J., Engels, C., Klarner, B., Maraun, M., Partsch, S., Roscher, C. and Schonert, F., 2011. Plant diversity surpasses plant functional groups and plant productivity as driver of soil biota in the long term. *PloS one*, 6(1), p.e16055.
- Fujita, M., Fujita, Y., Noutoshi, Y., Takahashi, F., Narusaka, Y., Yamaguchi-Shinozaki, K. and Shinozaki, K., 2006. Crosstalk between abiotic and biotic stress responses: a current view from the points of convergence in the stress signaling networks. *Current opinion in plant biology*, 9(4), pp.436-442.

- Glazebrook, J., 2005. Contrasting mechanisms of defense against biotrophic and necrotrophic pathogens. Annu. Rev. Phytopathol., 43, pp.205-227.
- Hamill, J.D., 1993. Alterations in auxin and cytokinin metabolism of higher plants due to expression of specific genes from pathogenic bacteria: a review. *Functional Plant Biology*, 20(5), pp.405-423.
- Haney, C.H. and Ausubel, F.M., 2015. Plant microbiome blueprints. Science, 349(6250), pp.788-789.
- Harrison, K.A. and Bardgett, R.D., 2010. Influence of plant species and soil conditions on plant-soil feedback in mixed grassland communities. *Journal of Ecology*, *98*(2), pp.384-395.
- Heil, M. and BOSTOCK, R.M., 2002. Induced systemic resistance (ISR) against pathogens in the context of induced plant defences. *Annals of botany*, 89(5), pp.503-512.
- Heinen, R., van der Sluijs, M., Biere, A., Harvey, J.A. and Bezemer, T.M., 2018. Plant community composition but not plant traits determine the outcome of soil legacy effects on plants and insects. *Journal of Ecology*, 106(3), pp.1217-1229.
- Joosten, L., Mulder, P.P., Klinkhamer, P.G. and van Veen, J.A., 2009. Soil-borne microorganisms and soil-type affect pyrrolizidine alkaloids in Jacobaea vulgaris. *Plant and Soil*, 325(1-2), p.133.
- Klironomos, J.N., 2002. Feedback with soil biota contributes to plant rarity and invasiveness in communities. *Nature*, 417(6884), pp.67-70.
- Kos, M., Tuijl, M.A., de Roo, J., Mulder, P.P. and Bezemer, T.M., 2015. Species-specific plant-soil feedback effects on above-ground plant-insect interactions. *Journal of Ecology*, 103(4), pp.904-914.
- Kulmatiski, A., Beard, K.H., Stevens, J.R. and Cobbold, S.M., 2008. Plant-soil feedbacks: a metaanalytical review. *Ecology letters*, 11(9), pp.980-992.
- Lebeis, S.L., Paredes, S.H., Lundberg, D.S., Breakfield, N., Gehring, J., McDonald, M., Malfatti, S., Del Rio, T.G., Jones, C.D., Tringe, S.G. and Dangl, J.L., 2015. Salicylic acid modulates colonization of the root microbiome by specific bacterial taxa. *Science*, 349(6250), pp.860-864.
- Leon-Reyes, A., Van der Does, D., De Lange, E.S., Delker, C., Wasternack, C., Van Wees, S.C., Ritsema, T. and Pieterse, C.M., 2010. Salicylate-mediated suppression of jasmonate-responsive gene expression in *Arabidopsis* is targeted downstream of the jasmonate biosynthesis pathway. *Planta*, 232(6), pp.1423-1432.
- Li, K., DiLegge, M.J., Minas, I.S., Hamm, A., Manter, D. and Vivanco, J.M., 2019. Soil sterilization leads to re-colonization of a healthier rhizosphere microbiome. *Rhizosphere*, *12*, p.100176.
- Mandal, S., Mallick, N. and Mitra, A., 2009. Salicylic acid-induced resistance to *Fusarium oxysporum* f. sp. lycopersici in tomato. *Plant physiology and Biochemistry*, 47(7), pp.642-649.
- Métrauxs, J.P., 2001. Systemic acquired resistance and salicylic acid: current state of knowledge. *European Journal of plant pathology*, 107(1), pp.13-18.
- Miki, T., 2012. Microbe-mediated plant-soil feedback and its roles in a changing world. *Ecological Research*, 27(3), pp.509-520.
- Nahar, K., Kyndt, T., De Vleesschauwer, D., Höfte, M. and Gheysen, G., 2011. The jasmonate pathway is a key player in systemically induced defense against root knot nematodes in rice. *Plant physiology*, 157(1), pp.305-316.
- O'Donnell, P.J., Jones, J.B., Antoine, F.R., Ciardi, J. and Klee, H.J., 2001. Ethylene-dependent salicylic acid regulates an expanded cell death response to a plant pathogen. *The Plant Journal*, 25(3), pp.315-323.
- Palmer, I.A., Shang, Z. and Fu, Z.Q., 2017. Salicylic acid-mediated plant defense: Recent developments, missing links, and future outlook. *Frontiers in biology*, 12(4), pp.258-270.
- Pieterse, C.M., Leon-Reyes, A., Van der Ent, S. and Van Wees, S.C., 2009. Networking by smallmolecule hormones in plant immunity. *Nature chemical biology*, 5(5), pp.308-316.

- Rashid, M. and Chung, Y.R., 2017. Induction of systemic resistance against insect herbivores in plants by beneficial soil microbes. *Frontiers in plant science*, *8*, p.1816.
- Revillini, D., Gehring, C.A. and Johnson, N.C., 2016. The role of locally adapted mycorrhizas and rhizobacteria in plant-soil feedback systems. *Functional Ecology*, *30*(7), pp.1086-1098.
- Reymond, P. and Farmer, E.E., 1998. Jasmonate and salicylate as global signals for defense gene expression. *Current opinion in plant biology*, *1*(5), pp.404-411.
- Spoel, S.H., Johnson, J.S. and Dong, X., 2007. Regulation of tradeoffs between plant defenses against pathogens with different lifestyles. *Proceedings of the National Academy of Sciences*, U.S.A, 104(47), pp.18842-18847.
- Steiner, A.A., 1979, September. The selective capacity of plants for ions and its importance for the composition and treatment of the nutrient solution. In Symposium on Research on Recirculating Water Culture, 98, pp. 87-98.
- Van Breemen, N. and Finzi, A.C., 1998. Plant-soil interactions: ecological aspects and evolutionary implications. *Biogeochemistry*, 42(1-2), pp.1-19.
- van Dam, N.M. and Oomen, M.W., 2008. Root and shoot jasmonic acid applications differentially affect leaf chemistry and herbivore growth. *Plant Signaling & Behavior*, *3*(2), pp.91-98.
- Van der Heijden, M.G., Bardgett, R.D. and Van Straalen, N.M., 2008. The unseen majority: soil microbes as drivers of plant diversity and productivity in terrestrial ecosystems. *Ecology letters*, 11(3), pp.296-310.
- Van der Putten, W.H., Bardgett, R.D., Bever, J.D., Bezemer, T.M., Casper, B.B., Fukami, T., Kardol, P., Klironomos, J.N., Kulmatiski, A., Schweitzer, J.A. and Suding, K.N., 2013. Plant-soil feedbacks: the past, the present and future challenges. *Journal of Ecology*, 101(2), pp.265-276.
- Van der Putten, W.H., 2017. Belowground drivers of plant diversity. Science, 355(6321), pp.134-135.
- Vernooij, B., Friedrich, L., Morse, A., Reist, R., Kolditz-Jawhar, R., Ward, E., Uknes, S., Kessmann, H. and Ryals, J., 1994. Salicylic acid is not the translocated signal responsible for inducing systemic acquired resistance but is required in signal transduction. *The Plant Cell*, 6(7), pp.959-965.
- Vos, I.A., Pieterse, C.M. and Van Wees, S.C., 2013. Costs and benefits of hormone-regulated plant defences. *Plant Pathology*, 62, pp.43-55.
- War, A.R., Paulraj, M.G., War, M.Y. and Ignacimuthu, S., 2011. Role of salicylic acid in induction of plant defense system in chickpea (*Cicer arietinum* L.). *Plant signaling & behavior*, 6(11), pp.1787-1792.
- Wang, M., Ruan, W., Kostenko, O., Carvalho, S., Hannula, S.E., Mulder, P.P., Bu, F., van der Putten, W.H. and Bezemer, T.M., 2019. Removal of soil biota alters soil feedback effects on plant growth and defense chemistry. *New Phytologist*, 221(3), pp.1478-1491.
- Yang, Y.X., J Ahammed, G., Wu, C., Fan, S.Y. and Zhou, Y.H., 2015. Crosstalk among jasmonate, salicylate and ethylene signaling pathways in plant disease and immune responses. *Current Protein and Peptide Science*, 16(5), pp.450-461.

Supplementary data

Table S1 Three-way ANOVA of shoot to root ratio of *J. vulgaris*, *C. vulgare*, *T. repens* and *D. carota* plants in live and sterilized soils after JA, SA application and control plants. Degrees of freedom, F- and P values are shown. df = degrees of freedom.

Source of variations	df	F-value	Р
species	3, 239	35.82	***
soil	1, 239	5.16	*
hormone	2, 239	20.02	***
species × soil	3, 239	1.88	ns
species × hormone	6, 239	9.28	***
soil × hormone	2, 239	4.28	*
species × soil × hormone	6, 239	2.67	*

* *P* < 0.05, *** *P* < 0.001, ns not significant.

Table S2 Three-way ANOVA of shoot-root ratios of *J. vulgaris* grown in live and sterilized soil after JA, SA application and control plants over four generations. Degrees of freedom, F- and P values are shown. df = degrees of freedom.

Source of variations	df	F-value	Р
soil	1,250	18.88	***
hormone	2,250	20.95	***
generation	3, 250	37.10	***
soil × hormone	2,250	1.41	*
soil × generation	3, 250	5.34	ns
hormone × generation	6,250	1.27	ns
soil \times hormone \times generation	6,250	1.70	ns

* *P* < 0.05, *** *P* < 0.001, ns not significant.



Fig. S1 Shoot-root ratio (+ SE) of *J. vulgaris, C. vulgare, T. repens* and *D. carota* plants treated with JA and SA in sterilized soil and live soil. C represents control treatment. Within species different letters indicate significant differences between treatments based on a Tukey post-hoc test for each species separately.



Fig. S2. Shoot-root ratio (+ SE) over four generations of *J. vulgaris* plants foliar treated with SA or JA and control plants grown in sterilized soil and live soil. C represents the control treatment. Within generations different letters indicate significant differences between treatment based on a Tukey post-hoc test for each generation separately.

Chapter 3

Activation of the SA-associated plant defense pathway alters the composition of soil bacterial communities

Jing Zhang^{1*} Peter G.L. Klinkhamer¹ Klaas Vrieling¹ T. Martijn Bezemer^{1,2}

 Plant Science and Natural Products, Institute of Biology Leiden (IBL), Leiden University, Sylviusweg 72, 2333BE Leiden, The Netherlands

2) The Netherlands Institute of Ecology (NIOO-KNAW), Department of Terrestrial Ecology, Droevendaalsesteeg 10, 6708 PB, Wageningen, The Netherlands

Abstract

Many plant species grow better in sterilized than in live soil. Foliar application of SA mitigated this negative effect of live soil on the growth of the plant Jacobaea vulgaris, as described in Chapter 2. This "SA-effect" on plant growth in live soils did not change over further cycles (generations), neither did the negative effect of live soils. To examine what causes the positive effect of SA application on plant growth in live soils, in this chapter we analyzed the effects of SA application on the composition of active rhizosphere bacteria in the live soil and how this change over time using RNA sequencing of the microbial communities in the rhizosphere of Jacobaea vulgaris. Our study shows that the composition of the rhizosphere bacterial communities of J. vulgaris greatly differed among generations. Application of SA resulted in both increases and decreases in a number of bacterial genera in the rhizosphere soil, but the genera that were affected by the treatment differed among generations. In the first generation, there were no genera that were significantly affected by the SA treatment, indicating that induction of the SA defense pathway in plants does not lead to immediate changes in the soil microbial community. 89 species out of the total 270 (32.4%) were present in all generations in all soils of SA-treated and control plants suggesting that these make up the "core" microbiome. On average in each generation, 72.9% of all genera were present in both soils. Application of SA to plants significantly up-regulated genera of Caballeronia, unclassified Cytophagaceae, Crinalium and Candidatus Thermofonsia Clade 2, and down-regulated genera of Thermomicrobiales, unclassified Rhodobacterales, Paracoccus and Flavihumibacter. While the functions of many of these bacteria are poorly understood, bacteria of the genus Caballeronia play an important role in fixing nitrogen and promoting plant growth, and hence this suggests that activation of the SA signaling pathway in J. vulgaris plants may select for bacterial genera that are beneficial to the plant. Further studies should examine how activation of the SA signaling pathway in the plant changes the functional genes of the rhizosphere soil bacterial community. Overall, our study shows that aboveground activation of defenses in the plant affects soil

microbial communities and as soil microbes can greatly influence plant performance, this implies that induction of plant defenses, can lead to complex above-belowground feedbacks.

Keywords

Metatranscriptomics, Soil microbial community, Taxonomy, Plant-soil interactions, Microbial diversity, Rhizosphere soil, Salicylic acid

Introduction

Plants encounter a myriad of threats from the surrounding environment, including both abiotic and biotic stresses (Suzuki et al., 2014). Biotic stresses are mostly due to herbivory and pathogen infestation both below- and above-ground (Pieterse and Dicke, 2007; Adair and Douglas, 2017). Microbes in the soil can have a beneficial, pathogenic or neutral effect on the host plant. For example, soil bacteria such as *Rhizoctonia species*, often strongly negative affect plant growth and survival (Issac et al., 1971). On the other hand, plant growth-promoting rhizobacteria (PGPR), such as *Pseudomonas* and *Burkholderia* species are beneficial for the plant, e.g. via suppressing the growth of soil-borne pathogens or increasing nutrient availability (Bhattacharyya and Jha, 2012). However, the overall net effect of soil microbial communities on plant growth is often negative (Nijjer et al., 2007). Most plant species grow less well in soils that contain a natural microbial community than in sterilized soils. This might be due e.g. competition between plants and microbes for available nutrients or due to soil-borne plant pathogens (Callaway et al., 2004; Berendsen et al., 2012; Mazzoleni et al., 2015; Cesarano et al., 2017).

Systemic acquired resistance (SAR) is one of the most common defensive strategies of plants against biotrophic pathogenic microbes. Foliar application of salicylic acid to plant tissues can activate SAR and boost the innate immune system of a plant (Reymond and Farmer, 1998). Cultivars with a higher sensitivity to SA are often better defended against the pathogens. For example, in tomato, exogenous application of SA can be effective against the pathogens *Oidium neolycopersici* and *Botrytis cinerea*, which cause powdery mildew and grey mould diseases (Seskar et al., 1998; Achuo et al., 2004). In agriculture, application of SA is now used to suppress pathogenic microbial effects in e.g. tomato, pepper and pea crops (Esmailzadeh et al., 2008; Barilli et al., 2010; Choi and Hwang, 2011). How SA application to the plant affects the microbial community in the soil is not fully uncovered.

Because plants alter the composition of the microbial community in the soil in which they grow, and SAR protects plants against pathogens, an important question is how activation of SAR alters the plant's effect on the soil microbial community. Several studies have demonstrated that the activation of SAR indeed altered the composition of soil microbial community and that SA can play a key role as a regulator in shaping root bacterial communities (Kniskern et al., 2007; Hein et al., 2008; Lebeis et al., 2015). However, several other studies reported that foliar application of SA did not affect the bacterial composition in the soil (Doornbos et al., 2011; Wang et al., 2015). These studies on the effects of SAR on soil bacterial composition were mostly limited to the model plant species *Arabidopsis*. As plant species differ greatly in the way and magnitude in which they influence the soil bacterial community (Wubs and Bezemer, 2018; Hannula et al., 2019; Pineda et al., 2020), we may expect that the effects of SA application on the soil microbial community also differ among plant species.

Several studies have shown that the composition of the soil bacterial microbial community varies greatly over time (e.g. Hannula et al., 2019). In a study on temporal variation in three land-use types, the number of taxa present in the soil showed strong temporal variability, and these changes over time were considerably larger than the variation associated with land-use types (Lauder et al., 2013). In contrast, Shade et al. (2012) demonstrated that soil microbial communities have clear successional trajectories. If generally true this would imply that application of SA to plants could also cause directed changes in the soil microbial community over time. An important question is therefore how activation of SAR will alter the soil microbial community over time.

In Chapter 2, we showed that inoculation of a sterilized soil with natural, live soil, reduced plant growth in comparison with that in sterilized soil for the plant species *Jacobaea vulgaris*. Interestingly, applying SA to the leaves mitigated these negative effects. This implies that activation of SA-induced resistance may potentially suppress microbial pathogens present in live soil. If this is the case, an important question is whether the repeated foliar application of SA during consecutive generations of plant growth will increase this effect and hence, whether there is a selection for a more beneficial bacterial community. Conceptually, the temporal dynamics of foliar application of SA and the effect of different generations do not alter the soil bacterial composition (Fig. 1-i). Second, foliar application of SA may lead to different bacterial communities independent of time (Fig. 1-ii). Third, bacterial communities may differ among generations but are not influenced by the SA

application (Fig. 1-iii). Fourth, foliar application of SA may influence bacterial communities but these effects may differ among generations (Fig. 1-iv).

In this study, we sequenced the mRNA from rhizosphere soil samples of both SAtreated and control plants during four consecutive generations of growth of *J. vulgaris*. In each consecutive generation soil from the previous plant growth period was used. Using mRNA instead of DNA or rRNA enabled us to focus on the active soil microbial community (Gilbert et al., 2008). In this study, we focus on the bacterial community. Twenty-four rhizosphere soils were sequenced with an Illumina sequencing platform. The goal of this study is to answer the following questions: (1) How does the foliar application of SA in *J. vulgaris* affect the bacterial composition in the rhizosphere and is there a time effect or an interactive time x SA effect on the bacterial community? (2) What is the "core" bacterial community in the soils of plants exposed to the SA treatment and of control plants? (3) How does the application of SA influence the bacterial community in each generation? Are the SA effects consistent over time?



Fig. 1 Conceptual figure showing the potential effects of SA application on *J. vulgaris* over four generations.

i) No effect of SA and time. The bacterial community does not differ between SAtreated plants and control, and does not differ over time. ii) SA effect only. The bacterial community is affected by SA application but the effect does not differ over time. iii) Time effect only. The bacterial community changes over time, but is not affected by the SA treatment. iv) SA \times time effect. The bacterial community is affected by SA-application but these effects differ among generations.

Materials and methods

Multi-generational plant growth experiment

In Chapter 2, we report the effects of foliar application of SA on plant growth in inoculated and sterilized soils. The current chapter focuses on the effect of foliar SA application on the composition of the bacterial community in the rhizosphere in the inoculated soil. Details of the experiment are described below.

J. vulgaris (common ragwort) seeds were collected at the dunes of Meijendel (a calcareous sandy area from a coastal dune area north of The Hague, The Netherlands, 52°11′N, 4°31′E). Prior to germination, all seeds were surface sterilized (shaken for 2 min in 70% ethanol, then rinsed with sterilized water, put for 12 min in 2% bleach, and then rinsed again four times with sterilized water to minimize influences of seedborne microbes (Bakker et al., 2015). The soil was also collected at Meijendel. The topsoil was collected to a depth of 15 cm after removing the grassland vegetation and the organic layer of the surface. The soil was sieved using a 5 mm sized mesh, homogenized with a concrete mixer, and then stored into 20-liter plastic bags (Nasco Whirl-Pak Sample Bag). Bags were either sterilized by 35-K Gray gamma-irradiation (Synergy Health Company, Ede, The Netherlands) or kept at 4°C for inoculation.

Surface sterilized seeds were germinated in sterile Petri dishes on filter paper. After one week, seedlings were randomly transferred individually to 500 ml pots consisting of a mixture of 90% sterilized soil and 10% live soil. Prior to potting but after mixing, the soil was kept in bags and left in the climate room for 14 days so that the mixed

soil could settle and microbial communities could colonize the sterilized soil. After potting the seedlings, pots were randomly distributed over a climate room (relative humidity 70%, light 16h at 20°C, dark 8h at 20°C). Plants were watered regularly with Milli-Q and 5 ml Steiner nutrient solution was added per plant on day 7 after planting, 10 ml Steiner nutrient solution (Steiner, 1979) was added on day 13, and 20 ml Steiner nutrient solution was added on days 19, 28, 37, 42. The Steiner nutrient solution was prepared from 7 different stock solutions (106.2 g Ca(NO₃)₂·4H₂O, 29.3 g KNO₃, 13.6 g KH₂PO₄, 49.2 g MgSO₄·7H₂O, 25.2 g K₂SO₄ and 2.24 g KOH, 3.29 g Fe-EDTA added to 1 liter demineralized water, and a stock solution with micro elements (a mixed solution of 0.181 g MnCl₂·4H₂O, 0.286 g H₃BO₃, 0.022 g ZnSO₄·7H₂O, 0.0078 g CuSO₄·5H₂O and 0.0126 g NaMoO₄·2H₂O added to 1 liter demineralized water). Ten ml of each stock solution was diluted in 1 liter of demineralized water before use.

Plants were allocated to either a hormonal treatment (SA) or served as control (only solvent). Both treatments were replicated 10 times. During plant growth, the phytohormone SA was applied through foliar application three times a week for four consecutive weeks. The first application was given when plants were 14 days old. About 0.75 ml of 100 μ M SA was sprayed on the leaves while carefully avoiding spillover to the soil. One week later the treatment was repeated with 1.50 ml of SA. In the next week, the treatment was repeated with 2.25 ml of SA. SA solvent (purchased from Sigma-Aldrich, \geq 99.0%) was made by dissolving 6.91 mg in 69.10 μ l of ethanol. Milli-Q water was then added until a final volume of 500 ml. Control plants were sprayed with sterile water with the same solvent (ethanol in Milli-Q water).

After six weeks, plants were gently removed from the pots. The rhizosphere soil for each treatment was harvested for each pot individually by gently shaking three times to remove the loosely adhering soil, after which rhizosphere soil samples were collected onto a sterile filter paper by removing the remnant soil with a fine sterilized brush. Rhizosphere soil samples were put in a 2 ml Eppendorf tube and stored at - 80°C for further RNA extraction. The remaining rhizosphere soil and adhering soil of the ten pots were mixed and used as inoculum (live soil) for the next generational of plant growth. The inoculum soil (10%) was mixed with 90% sterilized soil.

The set-up was repeated for another three generations under the same conditions as described above so that there were four generations of plant growth. For the second, third, and fourth generation, the soil inoculum was derived from the previous generation from the same treatment and was a mixture of rhizosphere soil and the loosen adhering soil surrounding the roots. Again, after mixing, the soil was kept in bags and left in the climate room for 14 days. Hereafter, pots were filled with soil and a *J. vulgaris* seedling was planted into each pot. All replicate soils from the SA or control treatment were mixed before the inoculation. The SA treatment was carried out as described above in each generation. Fifty-four days after planting, all plants were harvested each time.

RNA extraction and metatranscriptomic sequencing

For each treatment, the three successively labeled samples (No. 1, 2, 3, No. 4, 5, 6 and No. 7, 8, 9) were mixed and used as one composed replicate, Hence, three replicates were generated and used for RNA extraction for each treatment in each generation and a total of 24 soil samples were used for RNA extraction (3 replicates x 2 treatments x 4 generations). Total RNA was extracted with the RNeasy PowerSoil Total RNA kit (Qiagen). RNA concentration and quality were assessed by running 1µl of the extracted raw RNA on the 4200 TapeStation (Agilent). Subsequently, unwanted DNA, salts and buffers were removed with the RNeasy minElute Cleanup Kit (Qiagen). Later, the Ribo-Zero Magnetic kit for bacteria (Illumina) was used for mRNA enrichment. In the end, a RNA Clean & Concentrator kit (Zymoresearch) was used to clean additional buffers and proteins of the rRNA-depleted RNA. All the steps in extracting and cleaning RNA were according to the supplier's instructions. Doublestranded cDNA was generated from the cleaned RNA obtained in the final step. Library preparation (Illumina Nextera XT DNA library), processing and sequencing were performed by FG Technologies (Leiden, The Netherlands) with paired-end (PE) 150 bp templates. Twenty-four metatranscriptomic libraries were generated, the size of each library was indicated in Table S1 and Fig. S1.

Bioinformatics processing

Trimmomatic 0.39 was used for the removal of adapters of paired-end raw reads (Bolger et al., 2014). FastQC was applied to check the qualities, the bases with a threshold lower than 30 were cut off with Trimmomatic (Andrews, 2010). Ribosomal RNAs of all 24 metatranscriptomic libraries were filtered with SortMeRNA (Sorting ribosomal RNA) (Kopylova et al., 2012). Eight rRNA representative databases (silvabac-16s-id90, silva-arc-16s-id95, silva-euk-18s-id95, silva-bac-23s-id98, silva-arc-23s-id98, silva-euk-28s-id98 rfam-5s-id98, rfam-5.8s-id98) were derived from the SILVA SSU and LSU databases (release 119) and the RFAM databases with HMMER 3.1b1 and SumaClust v1.0.00 were used for fast filtering of rRNA from eukaryote, prokaryote and archaea. Then, all reads of the 24 metatranscriptomic libraries were combined into one set, which was the input of a de novo assembly. Trinity with default parameters was used for the metatranscriptomic assembly (Haas et al., 2013). Later, the quality of assembled contigs was assessed with Trinity scripts. The CD-HIT-EST algorithm was used to remove the duplicates of each transcript and reads with shorter than 300 bps were removed with a homemade script (Li and Godzik, 2006), after which reads of each library were mapped back to transcriptome with Bowtie2 (Langmead and Salzberg, 2012). The isoform IDs per sample were extracted with Seqkit (Shen et al., 2016). Contigs of each sample were generated and then aligned against the NCBI NR (non-redundant) database by DIAMOND with a cut off e-value at 1e-5 (Buchfink et al., 2015). The closest match with an identity higher than 80% was kept for mapping. The output file of Blastx was further analyzed with the lowest common ancestor (LCA) algorithm in MEGAN (version 6.0) with all default parameters (Camon et al., 2005; Huson et al., 2016). MEGAN helped to compute and explore our data at different taxonomic levels and in this process NCBI taxonomy was employed for summarizing and outputting results, the detailed workflow is referred to Huson et al. (2007). A count table of microbial species was obtained with read counts assigned directly to taxon for the 24 samples. The number of assigned reads per taxa was extracted at species, genus, family and phylum levels respectively. The number of identified phyla, families, genera and species were counted, and the composition and the percentage of reads used for each classification level were calculated.
Statistical analyses

Differences in the numbers of the total reads and the numbers of the non-rRNA reads over four generations were presented as mean \pm SD. A Heinrich's triangle figure was generated to visualize microbial composition at different phylogenetic levels of all the identified microbes from the 24 rhizosphere soil samples. Log10 transformed hit numbers of each genus were plotted as a function of ranked genus abundance numbers including all species and a cut-off was performed with an abundance larger than 0.01% of the total reads. A Shapiro-Wilk test was used to test for differences between the distribution of abundance between the SA and control treatment.

The Shannon-diversity index was calculated for the 24 samples and differences between the Shannon-diversity of soils of SA treated plants and soils of control plants were tested with a student t-test. Subsequently, abundance at genus level was used for to construct NMDS (nonmetric multidimensional scaling), PCA (Principal component analysis), OPLS-DA (orthogonal projection to latent structures discriminant analysis), and Venn diagrams, and Pearson distance and the Ward clustering algorithm statistical analysis was calculated since most of the reads were identified at the genus level.

Two-factor Venn diagrams were constructed to illustrate the numbers of unique and common genera in soil samples within each generation for the SA and control treatments, and a four-factor Venn diagram including all generations was performed for the SA and the control treatment separately (Heberle et al., 2015).

PCA and OPLS-DA were performed with SIMACA 15.0 using relative abundance at genus level. The relative abundance was calculated using the absolute abundance number of one genus divided by the total abundance of all bacterial genera in the sample. Before performing OPLS-DA analysis, we checked that our data fitted the model with a cross-validated residual (CV)-ANOVA significance testing (n = 270, P < 0.02).

To visualize the compositional changes among different treatment and time categories, a NMDS using the Bray-Curtis index as a measure of dissimilarity was generated using relative abundances. To verify changes in composition due to the SA treatment and time effect, a PERMANOVA test was performed using the *Adonis* function (number of permutations = 999) in R within the "vegan" package.

Local "immigration" and "extinction" in the rhizosphere soil of SA-treated or control plants over generations at genus level was calculated and the numbers were presented in Venn diagram. A Student's t-test was used to identify bacterial genera that were significantly enriched in soil samples of SA-treated or control plants. *P* values were adjusted for false discovery rates (FDR).

Spearman's rank correlation without multiple comparison tests were performed to identify the genera that were significantly positively or negatively correlated with generation within the SA or control treatment. Genera with P values smaller than 0.05 were selected to create a heatmap for all the 24 samples. Hierarchical clustering analysis was done for the 24 samples together, based on the relative abundance to show the similarity. The row-centered relative abundance of each genus was used to construct the color key (Chong et al., 2018). Heatmaps for only SA and only control treatments were generated in addition.

Results

Metatranscriptomic sequence data

A total of 898.4 million raw sequence reads were obtained from the 24 metatranscriptomic libraries, the smallest and largest library contained 25.0 and 52.0 million raw sequence reads, respectively (supplementary data Table S1). 846.9 million reads were kept after removing adapters and quality filtering control with FastQC. In total, 775.3 million reads were removed with the SortMeRNA program as ribosomal RNA (rRNA) reads when aligning them against eight rRNA representative databases (silva-bac-16s-id90, silva-arc-16s-id95, silva-euk-18s-id95, silva-bac-23s-id98, silva-arc-23s-id98, silva-euk-28s-id98 rfam-5s-id98, rfam-5.8s-id98), and 71.6 million reads were used as non-rRNA reads for further de novo assembly with Trinity (Fig. S1), of which the smallest library contained 1.5 million reads and the largest library 5.9 million reads. Reads for de novo assembly were normalized with Trinity

in silico normalization algorithm. The average guanine-cytosine (GC) content for the 24 libraries was 60.10%. After assembly, 0.99 million contigs were removed because their length was shorter than 300 bps. A total of 1.3 million unique contigs were identified after removing duplicates with CD-HIT-EST. In total, 392.4 million bases were assembled. After we checked the quality of the contigs in all samples by realigning all contigs back to the assemblies using Bowtie2, the average mapping rate for proper pairs was 45.41%.

Overview of the assigned reads at differential microbial classification levels

When we aligned the 1.3 million unique contigs against the NR (non-redundant) database with DIAMOND and MEGAN 6.0, 0.39 million contigs were taxonomically classified, while the others did provide a match with the available taxonomic information. Based on the analysis in MEGAN, the identified contigs were assigned at different classification levels. 22 different bacterial phyla were identified, 283 families and 382 bacterial genera and 1081 bacterial species (Fig. S2). At the phylum, family, genus and species level 23.4%, 23.4%, 20.4% and 14.9% of the total number of contigs were assigned, respectively. Bacteria were the most prevalent in the microbial community taking up 98.3 % of the total number of reads (Fig. S3a). Eukaryotes, with algae taking the largest proportion, were the second dominant, but Eukaryotes only covered 1.5% of the total number of reads (Fig. S3b).

SA application and time effects on bacterial community diversity and composition

From the total of 408 bacterial genera, 270 genera were included in the analysis (contigs with more than 0.01% of the total number of reads Fig. S5). The genera in both soils showed significantly different abundance curves (Shapiro-Wilk test, df = 407, P < 0.0001; Fig. S5), the abundance curve in the SA soil is lower than that in the control soil. Application of SA did not significantly increase or decrease the Shannon diversity at genus level within each generation (t-test for the 1st generation: t = -0.63, df = 5, P = 0.27; 2nd generation: t = 0.07, df = 5, P = 0.47; 3rd generation: t = 0.67, df = 5, P = 0.26; 4th generation: t = 0.50, df = 5, P = 0.31).

The NMDS plot showed that the bacterial communities of the same generation clustered together (Fig. 2a), PERMANOVA $R^2 = 0.30$, P = 0.001). The SA and control separated in the NMDS plot (Fig. S6) but this was not significant (PERMANOVA $R^2 = 0.05$, P = 0.18). Similar patterns were observed in a principal component analysis (PCA; Fig. S7). The OPLS-DA analysis showed clusters for replicates within each generation, and clear separation for the SA effect but only in the 2nd, 3rd and 4th generation (Fig. 2b). However, the generation effect was more evident than the SA effect.



Fig. 2 Multivariate analysis of the bacterial community in soil samples from SAtreated and control plants grown in four generations. Shown are sample scores from a Nonmetric Multi-Dimensional Scaling (NMDS) plot (a) and an Orthogonal

Projections to Latent Structures Discriminant analysis (OPLS-DA) plot (b) from the 24 rhizosphere soil samples.

Core bacterial community

89 species out of the total of 270 (32.4%) were present in all generations in at least two out of the three replicates of the soils of SA-treated and control plants suggesting that these make up the "core" microbiome (Fig. 3a). On average in each generation, 72.9% of all the genera were present in both soils (Fig. 3b). In the first generation, both soils shared about 74.2% of the genera while 7.7% only occurred in the SAtreatment and 18.0% only in the control (Fig. 3b-1). The percentage of shared genera by the two soils in the 2nd, 3rd and 4th generation was 67.6%, 72.9% and 76.8% (Fig. 3b-2, 3, 4). For soils of the control treatment, 49.5% of the genera were shared over all four generations; while 45.1% of genera were shared in soils of the SA treated plants over four generations (Fig. S2c; Table S2). Immigration was somewhat higher in the SA treatment (on average 42 new genera) than in the control (on average 34 new genera) while the opposite was true for extinction rates (on average 31 genera in the SA treatment and 33 in the control treatment; Fig. 4). The information of Archaea, virus and eukaryote is listed in supplementary Fig. S4.



Fig. 3 Venn diagrams showing the unique and shared genera of bacteria in the rhizosphere soil samples of SA-treated and control *J. vulgaris* plants. The diagram in (a) is based on an analysis of genera that occur in all growth generations of the SA treatment, in (b) each generation is analyzed separately, (c) shows the diagram for all generations combined for the SA and control treatment.



Fig. 4 Local "immigration" and "extinction" of bacterial genera in the rhizosphere soil of SA-treated and control plants over time. For each two consecutive generations, shown are the number of genera present only in the first of those generations (i.e., representing genera that go extinct), present in both generations, and present only in the second of those generations (i.e., representing generate that immigrate). Genera were considered present in a treatment when present in at least two of the tree replicates. 1, 2, 3 and 4 represent the 1st, 2nd, 3rd and 4th generation.

SA selection of soil bacteria

When analyzed per generation, in total eight genera differed among the SA treatment and control (Fig. 5). No genus was significantly affected in more than one generation and no genera were significantly affected in the first generation. Most of the significant genera were only present in either the control or SA treatment. A Spearman's rank correlation showed that 41 (out of 240) genera in the rhizosphere soil of SA-treated plants were significantly increasing or 31 genera were decreasing over generations. For the control soils these numbers were 47 and 27, respectively out of a total of 239 genera (Table S3). The heatmap including all 24 samples showed a clear generation effect, but no clear SA effect (Fig. 6). A heatmap representing the patterns of all identified genera in the 12 rhizosphere soils of SA-treated plants showed that replicates within a generation clustered and that the 2nd, 3rd and 4th generation showed a higher similarity than the 1st generation (Fig S8a). For the control plants, the samples from the 1st generation differed from the three other generations (Fig S8b).



Fig. 5 Bar chart showing relative abundance (%) (mean \pm SE) of the significant up or down regulated genera in the rhizosphere soils by SA-treated *J. vulgaris* plants. The significance is based on a student t-test with a false discovery rate (FDR) adjusted *P* values (< 0.05).



Fig. 6 Heatmap with a hierarchical clustering analysis of all the bacterial genera of rhizosphere soil of SA-treated *J. vulgaris* plants and control plants in the 24 samples. The hierarchical clustering was calculated with Pearson distance and the Ward clustering algorithm based on the relative abundance of the reads of each genus. The color code represents the row-centered relative abundance. SA1, SA2, SA3 and SA4 represent SA treatments from the 1st generation, 2nd generation of plant growth, 3rd generation and 4th generation. Control 1, control2, control3 and control4 represent

control treatments from the 1^{st} generation, 2^{nd} generation, 3^{rd} generation and 4^{th} generation.

Discussion

In this study, we examined how the activation of SA-induced resistance in the plant impacts the microbial composition in the rhizosphere, and how this change over generations of plant growth. Our study shows that the composition of rhizosphere bacteria communities of *J. vulgaris* changed significantly over generations, *but* that neither the effects of activation of SA-associated plant defense pathways nor the interaction between generation number and SA on the bacterial composition was significant. Within generations the application of SA selected for different bacterial genera in the rhizosphere soil, but these selected genera differed from generation to generation. There were no SA-mediated changes in active bacterial genera in the first generation, suggesting that there are no immediate effects of activation of the SA defense pathway on the soil microbial composition. The majority (76.1%) of the bacterial genera that we detected was present in all soils and represents the "core" bacterial microbiome.

Our study showed that aboveground activation of SA-associated plant defense pathways influenced different bacterial genera in the second, third and fourth generations. Effects of SA-induced resistance on the soil microbial community have been reported in several other studies. For example, Hein et al. (2008) compared the effect of SA application on the composition of rhizosphere bacterial communities in several *Arabidopsis* mutants with terminal restriction fragment length polymorphism (T-RFLP) analysis. They found that SA-induced resistance changed the structure of bacterial communities in the rhizosphere. In addition, Lebeis et al. (2015) demonstrated that SA application modulates colonization of the root microbiome by specific bacterial taxa. SA in plants is associated with the expression of pathogenesis-related proteins (PRPs). These PRPs possess antimicrobial activities resulting in suppression of microbial pathogens, consequently changing the microbial composition (Yalpani et al., 1991; Van Loon and Van Strien, 1999). Alternatively, hormonal-induced resistance in the plant may promote beneficial bacteria and fungi.

However, the impact of SA-induced resistance on soil microbial communities is still debated. For instance, Wang et al. (2015) and Doornbos et al. (2011) both demonstrated that activation of SA-induced resistance did not significantly affect the composition and diversity of the rhizosphere bacterial community.

Even though the experimental conditions and plant genotypes remained the same throughout the experiment, the effects of SA application on the bacterial community differed among generations. In this context, it is important to note that for each generation we used an inoculum, which means that we placed a subset of the microbial community in a sterile background. This may explain why we saw so much variation temporally as in each generation a different subset of the microbial community may have been activated. It is also possible that the composition of the bacterial community is variable over time within each generation and as a consequence also among generations (Gilbert et al., 2009; Hickey et al., 2013; Lauber et al., 2013; Hannula et al., 2019).

Of the four potential models, our data confirmed the third hypothesis (Fig. 1-iii), showing that the bacterial communities did differed among generations but were not strongly influenced by SA application. This is line with studies showing that the composition of the soil bacterial microbial community exhibits large fluctuations over time (Hannula et al., 2019; Lauder et al., 2013). Moreover, our data also shows that the application of SA selects for different bacterial genera in the rhizosphere soil but that these selected genera differ from generation to generation. This suggests that the effects of SA application to plants on the soil microbial community are not consistent over time and that it will be difficult to predict the effects of activation of plant defenses on soil microbes, and ultimately how this will influence the interactions between plants and microbes in the rhizosphere.

Interestingly, in soils of SA-treated plants, we found an increase of *Caballeronia*, unclassified *Cytophagaceae*, *Crinalium* and *Candidatus Thermofonsia Clade* 2. The *Caballeronia* genus is often reported as playing an important role in fixing nitrogen and promoting plant growth. Species in this genus are predominantly endophytic diazotrophic bacteria and N-fixing bacteria (Padda et al., 2018; Puri et al., 2018; Puri et al., 2018; Puri et al., 2020). This suggests that activation of SA signaling pathways in *J. vulgaris*

plants benefited bacteria that were more beneficial to plant growth, but further studies are needed to confirm this. The functions of the other species of which their abundance differentially increases are poorly understood. It is noteworthy though that *Crinalium* is a genus that is often isolated from sandy dune soils so it not surprising that we detected this genus as we used dune soils in our experiment. Further studies should extract the information of these detected genera at the species level.

In conclusion, we provide evidence that the composition of bacterial communities in the rhizosphere significantly differed between plant cycles (generation), but we found no evidence that application of SA altered this pattern. However, application of SA influenced different bacterial genera in the rhizosphere, but the responsive genera varied between generations. No bacterial genera were detected that responded to SA application in the first generation suggesting that there are no immediate responses of bacteria in the rhizosphere to SA application to plants. This would question the socalled 'cry for help" hypothesis (Biere and Bennett, 2013; Rasmann et al., 2017; Pineda et al., 2013), but further studies are required before we can make firm conclusions about this. Our results provide a new perspective on the effects of plant hormones on temporal changes in the soil microbial community.

Acknowledgements

We would like to thank Connor Philippo for his practical assistance in extracting RNA and quality control of the raw reads, Bing Xie for his generous supporting in sharing his private server and help in troubleshooting, Fons Verbeek for sharing public resource of the server in the Leiden institute of advanced computer science (LIACS). We also thank Martine Huberty for discussing PCA statistical analysis. We thank Robin van Schendel for assisting in the Trinity assembly on Shark cluster, Meciael Liem and Alexander Kloosterman for troubleshooting coding errors in Linux, Niels van Tol for helping introducing people who can help us with performing Trinity assembly. Jing Zhang would also like to thank the China Scholarship Council for financial support.

References

- Achuo, E.A., Audenaert, K., Meziane, H. and Höfte, M., 2004. The salicylic acid-dependent defence pathway is effective against different pathogens in tomato and tobacco. *Plant Pathology*, 53(1), pp.65-72.
- Adair, K.L. and Douglas, A.E., 2017. Making a microbiome: the many determinants of host-associated microbial community composition. *Current Opinion in Microbiology*, 35, pp.23-29.
- Andrews, S., 2010. FastQC: a quality control tool for high throughput sequence data. http://www.bioinformatics.babraham.ac.uk/projects/fastqc/
- Bakker, M.G., Chaparro, J.M., Manter, D.K. and Vivanco, J.M., 2015. Impacts of bulk soil microbial community structure on rhizosphere microbiomes of *Zea mays. Plant and Soil*, 392(1-2), pp.115-126.
- Barilli, E., Sillero, J.C. and Rubiales, D., 2010. Induction of systemic acquired resistance in pea against rust (*Uromyces pisi*) by exogenous application of biotic and abiotic inducers. *Journal of Phytopathology*, 158(1), pp.30-34.
- Berendsen, R.L., Pieterse, C.M. and Bakker, P.A., 2012. The rhizosphere microbiome and plant health. *Trends in plant science*, 17(8), pp.478-486.
- Bhattacharyya, P.N. and Jha, D.K., 2012. Plant growth-promoting rhizobacteria (PGPR): emergence in agriculture. *World Journal of Microbiology and Biotechnology*, 28(4), pp.1327-1350.
- Biere, A. and Bennett, A.E., 2013. Three-way interactions between plants, microbes and insects. *Functional Ecology*, 27(3), pp.567-573.
- Bolger, A.M., Lohse, M. and Usadel, B., 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics*, 30(15), pp.2114-2120.
- Buchfink, B., Xie, C. and Huson, D.H., 2015. Fast and sensitive protein alignment using DIAMOND. *Nature methods*, 12(1), p.59.
- Callaway, R.M., Thelen, G.C., Rodriguez, A. and Holben, W.E., 2004. Soil biota and exotic plant invasion. *Nature*, 427(6976), pp.731-733.
- Camon, E.B., Barrell, D.G., Dimmer, E.C., Lee, V., Magrane, M., Maslen, J., Binns, D. and Apweiler, R., 2005. An evaluation of GO annotation retrieval for BioCreAtIvE and GOA. *BMC bioinformatics*, 6(S1), p.S17.
- Cesarano, G., Zotti, M., Antignani, V., Marra, R., Scala, F. and Bonanomi, G., 2017. Soil sickness and negative plant-soil feedback: A reappraisal of hypotheses. *Journal of Plant Pathology*, 99(3), pp.545-570.
- Choi, H.W. and Hwang, B.K., 2011. Systemic acquired resistance of pepper to microbial pathogens. *Journal of Phytopathology*, 159(6), pp.393-400.
- Chong, J., Soufan, O., Li, C., Caraus, I., Li, S., Bourque, G., Wishart, D.S. and Xia, J., 2018. MetaboAnalyst 4.0: towards more transparent and integrative metabolomics analysis. *Nucleic acids research*, 46(W1), pp. W486-W494.
- Doornbos, R.F., Geraats, B.P., Kuramae, E.E., Van Loon, L.C. and Bakker, P.A., 2011. Effects of jasmonic acid, ethylene, and salicylic acid signaling on the rhizosphere bacterial community of *Arabidopsis thaliana*. *Molecular plant-microbe interactions*, 24(4), pp.395-407.
- Esmailzadeh, M., Soleimani, M.J. and Rouhani, H., 2008. Exogenous applications of salicylic acid for inducing systemic acquired resistance against tomato stem canker disease. *Journal of Biological Sciences*, 8(6), pp.1039-1044.
- Gilbert, J.A., Field, D., Huang, Y., Edwards, R., Li, W., Gilna, P. and Joint, I., 2008. Detection of large numbers of novel sequences in the metatranscriptomes of complex marine microbial communities. *PloS one*, 3(8), p.e3042.

- Haas, B.J., Papanicolaou, A., Yassour, M., Grabherr, M., Blood, P.D., Bowden, J., Couger, M.B., Eccles, D., Li, B., Lieber, M. and MacManes, M.D., 2013. De novo transcript sequence reconstruction from RNA-seq using the Trinity platform for reference generation and analysis. *Nature protocols*, 8(8), p.1494.
- Hannula, S.E., Kielak, A.M., Steinauer, K., Huberty, M., Jongen, R., Jonathan, R., Heinen, R. and Bezemer, T.M., 2019. Time after time: temporal variation in the effects of grass and forb species on soil bacterial and fungal communities. *MBio*, 10(6), e02635-19.
- Heberle, H., Meirelles, G.V., da Silva, F.R., Telles, G.P. and Minghim, R., 2015. InteractiVenn: a webbased tool for the analysis of sets through Venn diagrams. *BMC bioinformatics*, *16*(1), p.169.
- Hein, J.W., Wolfe, G.V. and Blee, K.A., 2008. Comparison of rhizosphere bacterial communities in *Arabidopsis thaliana* mutants for systemic acquired resistance. *Microbial ecology*, 55(2), pp.333-343.
- Hickey, R.J., Abdo, Z., Zhou, X., Nemeth, K., Hansmann, M., Osborn III, T.W., Wang, F. and Forney, L.J., 2013. Effects of tampons and menses on the composition and diversity of vaginal microbial communities over time. *BJOG: An International Journal of Obstetrics & Gynaecology*, 120(6), pp.695-706.
- Huson, D.H., Auch, A.F., Qi, J. and Schuster, S.C., 2007. MEGAN analysis of metagenomic data. *Genome research*, 17(3), pp.377-386.
- Huson, D.H., Beier, S., Flade, I., Górska, A., El-Hadidi, M., Mitra, S., Ruscheweyh, H.J. and Tappu, R., 2016. MEGAN community edition-interactive exploration and analysis of large-scale microbiome sequencing data. *PLoS computational biology*, *12*(6), p.e1004957.
- Isaac, I., Fletcher, P. and Harrison, J.A.C., 1971. Quantitative isolation of *Verticillium spp*. from soil and moribund potato haulm. *Annals of Applied Biology*, 67(2), pp.177-183.
- Kniskern, J.M., Traw, M.B. and Bergelson, J., 2007. Salicylic acid and jasmonic acid signaling defense pathways reduce natural bacterial diversity on *Arabidopsis thaliana*. *Molecular plant-microbe interactions*, 20(12), pp.1512-1522.
- Kopylova, E., Noé, L. and Touzet, H., 2012. SortMeRNA: fast and accurate filtering of ribosomal RNAs in metatranscriptomic data. *Bioinformatics*, 28(24), pp.3211-3217.
- Langmead, B. and Salzberg, S.L., 2012. Fast gapped-read alignment with Bowtie 2. Nature methods, 9(4), p.357.
- Lebeis S. L., Paredes S. H., Lundberg D. S., Breakfield N., Gehring J., McDonald M., Malfatti SRTGD., Jones C. D., Tringe S. G., Dang I. J. L., 2015. Salicylic acid modulates colonization of the root microbiome by specific bacterial taxa. *Science* 349, PP.860-864.
- Li, W. and Godzik, A., 2006. Cd-hit: a fast program for clustering and comparing large sets of protein or nucleotide sequences. *Bioinformatics*, 22(13), pp.1658-1659.
- Lauber, C.L., Ramirez, K.S., Aanderud, Z., Lennon, J. and Fierer, N., 2013. Temporal variability in soil microbial communities across land-use types. *The ISME journal*, 7(8), pp.1641-1650.
- Li, W. and Godzik, A., 2006. Cd-hit: a fast program for clustering and comparing large sets of protein or nucleotide sequences. *Bioinformatics*, 22(13), pp.1658-1659.
- Mazzoleni, S., Bonanomi, G., Incerti, G., Chiusano, M.L., Termolino, P., Mingo, A., Senatore, M., Giannino, F., Cartenì, F., Rietkerk, M. and Lanzotti, V., 2015. Inhibitory and toxic effects of extracellular self-DNA in litter: a mechanism for negative plant-soil feedbacks? *New Phytologist*, 205(3), pp.1195-1210.
- Nijjer, S., Rogers, W.E. and Siemann, E., 2007. Negative plant-soil feedbacks may limit persistence of an invasive tree due to rapid accumulation of soil pathogens. *Proceedings of the Royal Society B: Biological Sciences*, 274(1625), pp.2621-2627.

- Padda, K.P., Puri, A. and Chanway, C.P., 2018. Isolation and identification of endophytic diazotrophs from lodgepole pine trees growing at unreclaimed gravel mining pits in central interior British Columbia, Canada. *Canadian Journal of Forest Research*, 48(12), pp.1601-1606.
- Pieterse, C.M. and Dicke, M., 2007. Plant interactions with microbes and insects: from molecular mechanisms to ecology. *Trends in plant science*, *12*(12), pp.564-569.
- Pineda, A., Dicke, M., Pieterse, C.M. and Pozo, M.J., 2013. Beneficial microbes in a changing environment: are they always helping plants to deal with insects? *Functional Ecology*, 27(3), pp.574-586.
- Pineda, A., Kaplan, I., Hannula, S.E., Ghanem, W. and Bezemer, T.M., 2020. Conditioning the soil microbiome through plant-soil feedbacks suppresses an aboveground insect pest. *New Phytologist*, 226(2), pp.595-608.
- Puri, A., Padda, K.P. and Chanway, C.P., 2018. Evidence of endophytic diazotrophic bacteria in lodgepole pine and hybrid white spruce trees growing in soils with different nutrient statuses in the West Chilcotin region of British Columbia, Canada. *Forest Ecology and Management*, 430, pp.558-565.
- Puri, A., Padda, K.P. and Chanway, C.P., 2020. Can naturally-occurring endophytic nitrogen-fixing bacteria of hybrid white spruce sustain boreal forest tree growth on extremely nutrient-poor soils? Soil Biology and Biochemistry, 140, p.107642.
- Rasmann, S., Bennett, A., Biere, A., Karley, A. and Guerrieri, E., 2017. Root symbionts: Powerful drivers of plant above-and belowground indirect defenses. *Insect science*, 24(6), pp.947-960.
- Reymond, P. and Farmer, E.E., 1998. Jasmonate and salicylate as global signals for defense gene expression. *Current opinion in plant biology*, *1*(5), pp.404-411.
- Seskar, M., Shulaev, V. and Raskin, I., 1998. Endogenous methyl salicylate in pathogen-inoculated tobacco plants. *Plant physiology*, *116*(1), pp.387-392
- Shade, A., Peter, H., Allison, S.D., Baho, D., Berga, M., Bürgmann, H., Huber, D.H., Langenheder, S., Lennon, J.T., Martiny, J.B. and Matulich, K.L., 2012. Fundamentals of microbial community resistance and resilience. *Frontiers in microbiology*, 3, p.417.
- Shen, W., Le, S., Li, Y. and Hu, F., 2016. SeqKit: a cross-platform and ultrafast toolkit for FASTA/Q file manipulation. *PloS one*, *11*(10), p.e0163962.
- Steiner, A.A., 1979. The selective capacity of plants for ions and its importance for the composition and treatment of the nutrient solution. In *Symposium on Research on Recirculating Water Culture*, 98, pp.87-98.
- Suzuki, N., Rivero, R.M., Shulaev, V., Blumwald, E. and Mittler, R., 2014. Abiotic and biotic stress combinations. *New Phytologist*, 203(1), pp.32-43.
- Van Loon, L.C. and Van Strien, E.A., 1999. The families of pathogenesis-related proteins, their activities, and comparative analysis of PR-1 type proteins. *Physiological and molecular plant pathology*, 55, pp.85-97.
- Wang, B., Guo, X.W., Li, K., Han, X., Xu, S.J., Liu, Z.D., Guo, Y.S. and Xie, H.G., 2015. Effects of Salicylic acid on grape plants and the soil microbial community. *Allelopathy Journal*, 36(1).
- Wubs, E.J. and Bezemer, T.M., 2018. Plant community evenness responds to spatial plant–soil feedback heterogeneity primarily through the diversity of soil conditioning. *Functional Ecology*, 32(2), pp.509-521.
- Yalpani, N., Silverman, P., Wilson, T.M., Kleier, D.A. and Raskin, I., 1991. Salicylic acid is a systemic signal and an inducer of pathogenesis-related proteins in virus-infected tobacco. *The Plant Cell*, 3(8), pp.809-818.

3

Supplementary data

Table S1. Summary of Illumina sequencing of the 24 libraries. Total RNA was extracted from 24 soil samples collected from *J. vulgaris* rhizospheres (2 treatments (control/SA application) \times 4 generations \times 3 replicates). The mapped reads and the percentage of properly paired reads that mapped back to the assembled metatranscriptome by Bowie2 are presented.

	Treatment	No Raw	No. filtered	No. rRNA	No. Non	Mapped
Gene.	(replicate)	reads	reads	reads	rRNA reads	reads
	Con 1	22 210 026	(percentage)	(percentage)	(percentage)	(percentage)
	Coll-1	55,519,920	51,005,490	29,951,407	1,932,091	901,492
	Com 2	17 161 076	(93.0)	(95.9)	(0.1)	(49.5)
	Coll-2	47,404,970	45,550,642	43,005,095	2,34/,14/	(40.5)
	Com 2	20 074 024	(90.0)	(94.4)	(3.0)	(49.3)
	Coll-3	36,074,624	(04.5)	(02.4)	2,300,270	(45.2)
1^{st}	So 1	26 801 026	(94.3)	(95.4)	(0.0)	(43.5)
	5a-1	30,891,920	33,337,342 (06 4)	33,148,711	2,408,831	1,046,220
	S- 2	20 695 279	(90.4)	(93.2)	(0.8)	(43.4)
	5a-2	29,085,578	28,040,038	27,055,158	1,580,900	044,792 (40.6)
	S- 2	41 (00 759	(90.3)	(94.3)	(3.3)	(40.6)
	5a-5	41,099,738	39,323,110	30,398,300	2,920,744	1,445,924
	Care 1	51 522 069	(94.3)	(92.0)	(7.4)	(49.4)
	Con-1	51,525,968	40,908,512	43,407,004	3,440,908	1,082,552
	Com 2	26 022 206	(91.0)	(92.7)	(7.3)	(48.9)
	Coll-2	30,922,500	55,600,604	30,217,009	3,383,833	1,196,910
	C 2	26 009 466	(91.0)	(89.4)	(10.0)	(33.3)
	Con-3	30,098,400	34,487,094	51,087,990	2,799,098	1,300,320
2 nd	So 1	24 409 252	(95.5)	(91.9)	(8.1)	(48.0)
	5a-1	54,498,252	55,458,050	50,528,949 (01.2)	2,929,087	1,122,032
	5. 2	20 828 060	(97.0)	(91.2)	(0.0)	(30.3)
	5a-2	50,828,900	29,287,510	23,012,404	5,474,620	(25.0)
	5.2	20 241 270	(95.0)	(00.1)	(11.9)	(33.0)
	5a-5	50,241,570	29,244,304	(02.0)	2,037,107	(25.2)
	Con 1	32 336 630	(90.7)	(92.9)	2 462 747	1 652 708
	Coll-1	52,550,050	(06.3)	(02.1)	(7.9)	(67.1)
	Con 2	37 877 082	(10.5)	(52.1)	5 905 460	2 000 760
	C0II-2	52,877,082	(96.4)	(81.4)	(18.6)	(35.4)
	Con 3	31 458 002	(90.4)	(01.4)	5 310 500	(33.4)
2 rd	C0II-3	51,458,902	(05.8)	(82.4)	(17.7)	(41.0)
3	Sa 1	25 035 684	24 001 844	(02.4)	1 500 830	866 602
	5a-1	25,055,084	(05.0)	(03.3)	(67)	(54.2)
	Sa-2	29 313 124	(<i>JJ.J.)</i> 27 003 208	24 721 628	2 371 670	(3+.2) 1 293 100
	5a-2	27,313,124	(97.4)	(91.2)	(8.8)	(54.5)
	So 3	52 080 670	(32.4) 18 177 381	(31.2)	(0.0)	2 907 272
	-3a-3	52,000,070	+0,+/2,304	+3,300,073	+,712,309	2,907,372

SA defense pathway an	d the composition	of soil bacterial	communities
-----------------------	-------------------	-------------------	-------------

			(93.1)	(89.9)	(10.1)	(59.2)
	Con-1	40,952,998	37,197,048	33,471,054	3,725,994	1,727,084
			(90.8)	(90.0)	(10.0)	(46.4)
	Con-2	40,803,576	37,914,816	34,101,981	3,812,835	1,641,020
			(92.9)	(89.9)	(10.1)	(43.0)
	Con-3	46,249,750	42,340,292	39,136,920	3,203,372	1,100,132
∕th			(91.6)	(92.4)	(7.6)	(34.3)
4	Sa-1	39,495,864	36,597,750	34,868,480	1,729,270	814,560
			(92.7)	(95.3)	(4.8)	(47.1)
	Sa-2	41,678,588	39,339,896	36,363,915	2,975,981	1,552,592
			(94.4)	(92.4)	(7.6)	(52.2)
	Sa-3	38,834,084	36,852,408	35,344,564	1,507,844	580,832
			(94.9)	(95.9)	(4.1)	(38.5)

Table S2. A list of all bacterial genera presents in different generations in the SA and
control treatment. "+" represents present and "-" absent. When a genus is present in
at least two out of three replicates within a treatment it is recorded as "+".

	1SA	2SA	3SA	4SA	1C	2C	3C	4C
Acidovorax	+	+	+	+	+	+	+	+
Acinetobacter	+	+	+	+	+	+	+	+
Aeromicrobium	+	+	+	+	+	+	+	+
Afipia	+	+	+	+	+	+	+	+
Aminobacter	+	+	+	+	+	+	+	+
Aquabacterium	+	+	+	+	+	+	+	+
Arenimonas	+	+	+	+	+	+	+	+
Arthrobacter	+	+	+	+	+	+	+	+
Azohydromonas	+	+	+	+	+	+	+	+
Bosea	+	+	+	+	+	+	+	+
Bradyrhizobium	+	+	+	+	+	+	+	+
Caenimonas	+	+	+	+	+	+	+	+
Cellvibrio	+	+	+	+	+	+	+	+
Chryseolinea	+	+	+	+	+	+	+	+
Cupriavidus	+	+	+	+	+	+	+	+
Curvibacter	+	+	+	+	+	+	+	+
Devosia	+	+	+	+	+	+	+	+
Dongia	+	+	+	+	+	+	+	+
Ensifer	+	+	+	+	+	+	+	+
environmental Bacteria	+	+	+	+	+	+	+	+
Flavobacterium	+	+	+	+	+	+	+	+
Fluviicola	+	+	+	+	+	+	+	+
Gemmatimonas	+	+	+	+	+	+	+	+
Herbaspirillum	+	+	+	+	+	+	+	+
Herminiimonas	+	+	+	+	+	+	+	+
Hydrogenophaga	+	+	+	+	+	+	+	+
Ideonella	+	+	+	+	+	+	+	+
Janthinobacterium	+	+	+	+	+	+	+	+
Lacibacter	+	+	+	+	+	+	+	+
Lacunisphaera	+	+	+	+	+	+	+	+
Luteimonas	+	+	+	+	+	+	+	+
Lysobacter	+	+	+	+	+	+	+	+
Marmoricola	+	+	+	+	+	+	+	+
Massilia	+	+	+	+	+	+	$^+$	+
Mesorhizobium	+	+	+	+	+	+	$^+$	+
Methylibium	+	+	+	+	+	+	+	+
Methylotenera	+	+	+	+	+	+	+	+
Microbacterium	+	+	+	+	+	+	+	+
Mycobacterium	+	+	+	+	+	+	+	+
Niastella	+	+	+	+	+	+	$^+$	+
Nitrospira	+	+	+	+	+	+	+	+
Nocardioides	+	+	+	+	+	+	+	+
Noviherbaspirillum	+	+	+	+	+	+	+	+
Opitutus	+	+	+	+	+	+	+	+
Pelomonas	+	+	+	+	+	+	+	+
Phenylobacterium	+	+	+	+	+	+	+	+

Phycicoccus	+	+	+	+	+	+	+	+
Polaromonas	+	+	+	+	+	+	+	+
Pseudarthrobacter	+	+	+	+	+	+	+	+
Pseudomonas	+	+	+	+	+	+	+	+
Pseudoxanthomonas	+	+	+	+	+	+	+	+
Ramlibacter	+	+	+	+	+	+	+	+
Revranella	+	+	+	+	+	+	+	+
Rhizobacter	+	+	+	+	+	+	+	+
Rhizobium	+	+	+	+	+	+	+	+
Rhodoferar	+	+	+	+	+	+	+	+
Rivibacter	+	+	+	+	+	+	+	+
Rubrivivar	+	+	+	+	+	+	+	+
Solimonas	+	+	+	+	+	+	+	+
Solimonus						1		
Springomonas	+	+	+	+	+	+	+	+
Spningopyxis	+	+	+	+	+	+	+	+
Sporichthya	+	+	+	+	+	+	+	+
Staphylococcus	+	+	+	+	+	+	+	+
Streptomyces	+	+	+	+	+	+	+	+
unclassified Acidobacteria	+	+	+	+	+	+	+	+
unclassified Actinobacteria (class) (miscellaneous)	+	+	+	+	+	+	+	+
unclassified Alphaproteobacteria (miscellaneous)	+	+	+	+	+	+	+	+
unclassified Bacteroidetes (miscellaneous)	+	+	+	+	+	+	+	+
unclassified Betaproteobacteria (miscellaneous)	+	+	+	+	+	+	+	+
unclassified Burkholderiaceae	+	+	+	+	+	$^+$	+	+
unclassified Burkholderiales (miscellaneous)	+	+	+	+	+	+	+	+
unclassified Chitinophagaceae	+	+	+	+	+	+	+	+
unclassified Chloroflexi (miscellaneous)	+	+	+	+	+	+	+	+
unclassified Comamonadaceae	+	+	+	+	+	+	+	+
unclassified Cytophagaceae	+	+	+	+	+	+	+	+
unclassified Deltaproteobacteria (miscellaneous)	+	+	+	+	+	+	+	+
unclassified Gammanrateabacteria (miscellaneous)	+	+	+	+	+	+	+	+
unclassified Gammatimonadalas	+	+	+	+	+	+	+	+
unclassified Gammatimonadatas	+	+	+	+	+	+	+	+
unclassified Managagagalag (miggallangaga)						1		
unclassified <i>Myxococcales</i> (<i>miscellaneous</i>)	- T	- -	- T	- T	- T	т ,	т	- -
	+	+	+	+	+	+	+	+
unclassified Proteobacteria	+	+	+	+	+	+	+	+
unclassified Rhizobiales (miscellaneous)	+	+	+	+	+	+	+	+
unclassified Sphingobacteriaceae	+	+	+	+	+	+	+	+
unclassified Sphingobacteriales	+	+	+	+	+	+	+	+
unclassified Verrucomicrobia (miscellaneous)	+	+	+	+	+	+	+	+
unclassified Xanthomonadaceae	+	+	+	+	+	+	+	+
Variovorax	+	+	+	+	+	+	+	+
Vulcaniibacterium	+	+	+	+	+	+	+	+
Azospira	+	+	+	+	-	+	+	+
Candidatus Thermofonsia Clade 2	+	+	+	+	-	+	+	+
Clostridioides	+	+	+	+	-	+	+	+
Collimonas	+	+	+	+	-	+	+	+
Escherichia	+	+	+	+	-	-	+	+
Hassallia	+	+	+	+	+	+	-	+
Ohtaekwangia	+	+	+	+	+	+	-	+
Pedobacter	+	+	+	+	+	+	-	+
Pedosphaera	+	+	+	+	+	+	-	+
Ralstonia	+	+	+	+	+	+	-	+
Rhodobactar	, +	+	+	+	I	- -		, +
MIDHODUCIEI	T		E E	T	-	- F	-	F

Shigella	+	+	+	+	+	-	-	+
Sinorhizobium	+	+	+	+	+	-	-	+
Tabrizicola	+	+	+	+	+	+	+	-
unclassified Acidimicrobiaceae	+	+	+	+	+	+	+	-
unclassified Flavobacteriales (miscellaneous)	+	+	+	+	+	-	+	-
unclassified Oxalobacteraceae	+	+	+	+	+	-	+	-
unclassified Sphingomonadales	+	+	+	+	+	-	-	-
Acidobacterium	-	+	+	+	+	+	+	+
Algoriphagus	-	+	+	+	+	$^+$	+	+
Altererythrobacter	-	+	+	+	+	+	+	+
Paeniglutamicibacter	+	-	+	+	+	+	+	+
Piscinibacter	+	-	+	+	+	+	+	+
Rhodococcus	+	-	+	+	+	+	+	+
Rhodoplanes	+	+	-	+	+	+	+	+
Sphingobium	+	+	+	-	+	+	+	+
unclassified Angerolineacege	+	+	+	-	+	+	+	+
unclassified Opitutae	+	+	-	-	+	+	+	+
unclassified Sinobacteraceae	+	+	-	-	+	+	+	+
unclassified Xanthomonadales	+	-	-	-	+	+	+	+
Actinomycetales	+	-	+	+	-	+	+	+
Actinoplanes	+	-	+	+	_	+	+	+
Agromyces	+	-	+	+	+	_	+	+
Alistines	+	-	+	+	+	_	+	_
Angerobutvricum	+	-	+	+	+	-	_	-
Anaerostines	+	-	+	+	+	_	_	_
Aquimonas	+	-	+	+	+	_	_	_
Aquincola	+	-	+	+	+	_	_	_
Azotobacter	+	+	_	+	+	+	_	+
Racteriovorax	+	+	_	+	+	+	_	+
hacterium	+	+	_	+	+	+	_	+
Racteroides	+	+	_	+	+	+	_	-
Batrachochytrium	+	+	_	+	-	_	_	_
Rdellovihrio	+	+	_	+	-	-	_	-
Rifidobacterium	+	_	_	+	+	+	_	+
Rlautia	+	-	_	+	+	_	_	+
Brewindimonas	+	-	_	+	+	-	_	_
Bryohacter	+	+	+	_	_	_	+	+
Burkholderia	+	+	+	_	-	+	_	+
Caballeronia	+	+	+	_	+	+	+	_
Candidatus Kaiserbacteria	+	+	+	_	+	+	_	_
Candidatus Kanabacteria	+	_	+	-	+	+	_	_
Candidatus Nitrosocosmicus	+	-	+	_	_	_	_	-
Candidatus Rokubacteria	+	+	_	-	+	+	_	+
Catellatospora	+	+	_	-	+	+	_	+
Chitinonhaga	+	+	_	-	+	+	_	+
Clostridium	+	+	_	_	+	+	+	_
Collinsella	+	+	_	_	+	+	_	-
Comamonas	+	+	_	_	+	_	_	_
Coprococcus	+	+	_	-	+	_	_	_
Crinalium	+	+	-	-	+	-	-	-
Crocinitomix	+	+	_	-	+	-	-	-
Cutibacterium	+	+	-	-	_	-	-	-
Dorea	+	+		-	-	-	-	_
Duganella	+	+	_	-	+	_	_	_
					•			

Dyadobacter	+	-	-	-	-	-	+	-
Dyella	+	-	-	-	+	+	-	-
environmental samples <bacteria, phylum<="" td=""><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td></bacteria,>								
Gemmatimonadetes>	+	-	-	-	+	+	-	-
environmental samples <crenarchaeotes,phylum< td=""><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td></crenarchaeotes,phylum<>								
Crenarchaeota>	+	-	-	-	+	+	-	-
environmental samples < firmicutes, phylum								
<i>Firmicutes></i>	+	-	-	-	+	-	-	-
environmental samples <gns bacteria,phylum<="" td=""><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td></gns>								
Chloroflexi>	+	-	-	-	+	-	-	-
Erythrobacter	+	-	-	-	+	-	-	-
Eubacterium	+	-	-	-	+	-	-	-
Faecalibacterium	+	-	-	-	+	-	-	-
Flavihumibacter	+	-	-	-	-	-	-	-
Fontimonas	+	-	-	-	-	-	-	-
Gemmatirosa	+	-	-	-	-	-	-	-
Gemmohacter	-	+	+	+	-	+	+	+
Glaciecola	-	+	+	+	-	+	+	+
Hernetosinhon	-	+	+	+	_	+	+	+
Holdemanella	-	+	+	+	_	+	+	+
Hylemonella	_	+	+	+	-	+	+	+
Hyphomicrobium	_	+	+	+	_	_	+	+
Ilumatobacter	_	+	+	+	_	+	-	+
Inhalla	-	+	+	+	-	+	-	+
Klabsiella	-	+	+	+	-	'	-	+
Knoollig	-	I	, 	- -	-	-	-	- -
Knoella	-	-	- -	- -	-	т 	т 	т
Kouleollin ix	-	-	, 	- -	-	- -	- -	- -
	-	-	- -	- T	-	т	т ,	т ,
	-	-	+	+	Ŧ	-	+	+
Leptolyngbya	-	-	+	+	-	-	+	+
Leptotnrix	-	-	+	+	-	-	+	+
Leptothrix	-	-	+	+	-	-	-	+
Limnobacter	-	-	+	+	-	+	+	-
Limnohabitans	-	-	+	+	-	-	+	-
Listeria	-	+	-	+	-	-	+	+
Longispora	-	+	-	+	-	-	+	+
Methylobacteriaceae	-	+	-	+	+	+	-	+
Microcoleus	-	+	-	+	-	+	-	+
Micromonospora	-	+	-	+	+	-	-	+
Nitrobacter	-	+	-	+	+	-	-	+
Nitrosomonas	-	+	-	+	+	-	-	+
Nitrososphaera	-	+	-	+	+	-	-	+
Novosphingobium	-	+	-	+	+	-	-	+
Oscillochloris	-	+	-	+	+	-	-	+
Paenarthrobacter	-	+	-	+	+	-	-	+
Panacagrimonas	-	+	-	+	+	-	-	+
Parabacteroides	-	+	-	+	+	-	-	+
Paracoccus	-	+	-	+	+	-	-	+
Paucibacter	-	+	-	+	+	-	-	+
Paucimonas	-	+	-	+	+	-	-	+
Phormidium	-	+	-	+	+	-	-	+
Phyllobacterium	-	+	-	+	+	-	-	+
Pirellula	-	+	-	+	+	-	-	+
Planomicrobium	-	+	-	+	-	-	-	+

Prevotella	-	+	-	+	-	-	-	+
Pseudolabrvs	-	+	-	+	-	+	+	-
Pseudonocardia	-	+	_	+	+	-	-	-
Pseudorhodobacter	-	+	_	+	+	-	-	-
Rheinheimera	-	+	_	+	-	-	-	-
Rickettsia	_	_	-	+	_	+	+	+
Rosenteles	_	-	_	+	-	+	+	+
Roseburia	_	-	_	+	+	_	+	+
Roseiflerus	_	_	_	+	-	_	+	+
Ruminococcus	_	-	_	+	-	_	+	+
Sandaracinus	_	_	_	+	_	_	+	+
Sadiminibactorium	_	-	_	+	-	-	_	+
Simplicispira	-	-	-	+	-	-	-	+
Simplicispiru	-	-	-	- -	-	-	-	, ,
Sinimarinibacierium Sabia zashah dua	-	-	-	т	-	-	-	- -
Springornabaus	-	-	-	+	-	-	-	Ŧ
Sporocytopnaga	-	-	-	+	-	-	+	-
Stella	-	-	-	+	-	-	+	-
Stenotrophomonas	-	-	-	+	+	-	-	-
Streptococcus	-	-	-	+	-	-	-	-
Streptosporangiaceae	-	-	-	+	-	-	-	-
Subdoligranulum	-	-	-	+	-	-	-	-
Thermomonas	-	-	-	+	-	-	-	-
Thermomonosporaceae	-	-	-	+	-	-	-	-
unclassified Acidobacteriia	-	-	-	+	-	-	-	-
unclassified Actinobacteria	-	+	+	-	-	-	-	-
unclassified Anaerolineae	-	+	+	-	-	-	-	-
unclassified Bradyrhizobiaceae	-	-	+	-	-	+	+	+
unclassified Caulobacteraceae	-	-	+	-	+	-	+	+
unclassified Clostridiales (miscellaneous)	-	-	+	-	-	-	+	+
unclassified Crocinitomicaceae	-	-	+	-	+	$^+$	+	-
unclassified Cyanobacteria (miscellaneous)	-	-	+	-	-	-	-	-
unclassified Cyclobacteriaceae	-	-	+	-	-	-	-	-
unclassified Firmicutes sensu stricto								
(miscellaneous)	-	+	-	-	-	-	-	+
unclassified Frankiales (miscellaneous)	-	+	-	-	+	+	-	-
unclassified Hyphomicrobiaceae	-	+	-	-	-	+	-	-
unclassified Ignavibacteriae	-	+	-	-	-	-	-	-
unclassified Lachnospiraceae	-	-	_	-	-	-	-	+
unclassified Nitrosomonadales	-	-	_	-	-	-	-	+
unclassified Nitrosonumilales	_	-	-	_	_	_	_	+
unclassified Parcubacteria group	_	-	_	_	-	-	_	+
unclassified Phyllobacteriaceae	_	_	_	_	_		+	-
unclassified Pseudomonadales	_	_	_	_	_	_	+	_
unclassified Rhodobactaracaaa	_						+	
unclassified Rhodobacterales	_	-	_	-	-	-	+	-
unclassified Rhodospirillagage	-	-	-	-	-	-	- -	-
unclassified Rhodospirillalos (miscollanoous)	-	-	-	-	-	-	т 1	-
unclassified Rhouospirillales (miscellaneous)	-	-	-	-	-	-	т	-
unclassified Subjection and a subject of the subjec	-	-	-	-	-	+	-	-
unclassified They manch and a function of the	-	-	-	-	-	+	-	-
unclassified <i>I naumarchaeota</i> (miscellaneous)	-	-	-	-	+	-	-	-
unclassified <i>I hermomicrobiales</i>	-	-	-	-	+	-	-	-
unclassified <i>Verrucomicrobia subdivision 3</i>	-	-	-	-	+	-	-	-
unclassified Verrucomicrobiaceae	-	-	-	-	+	-	-	-
Undibacterium	-	-	-	-	+	-	-	-

SA defense pathway and the composition of soil bacterial communities

Xanthomonas	-	-	-	-	+	-	-	-
Xenophilus	-	-	-	-	+	-	-	-

Table S3. Genera, of which the relative abundance was significantly positively or negatively correlated with generation number within the SA or control treatment. The correlation is based on a Spearman's rank correlation test. R_s represents Spearman's rank correlation coefficient. *P* values less than 0.05 were selected.

	SA		Control		
Genera Name	R _s value	Р	R _s value	Р	
Agromyces	0.85	***	-0.68	*	
Arthrobacter	0.93	***	-0.84	**	
Candidatus Kaiserbacteria	-0.74	**	0.59	*	
Crocinitomix	0.81	**	-0.83	**	
Ensifer	0.89	***	-0.62	*	
environmental samples < <i>crenarchaeotes,</i> <i>Crenarchaeota</i> >	0.77	**	-0.82	*	
Erythrobacter	-0.83	**	0.70	*	
Flavihumibacter	-0.81	**	0.86	***	
Flavobacterium	0.73	**	-0.64	*	
Gemmobacter	-0.83	**	0.69	*	
Lacibacter	-0.60	*	0.68	*	
Leptothrix	0.77	**	-0.72	*	
Listeria	0.60	*	-0.82	**	
Lysobacter	-0.76	**	0.64	*	
Mycobacterium	-0.74	**	0.74	**	
Opitutus	0.84	**	-0.82	**	
Phyllobacterium	0.75	**	-0.62	*	
Piscinibacter	0.95	***	-0.81	**	
Polaromonas	-0.76	**	0.62	*	
Pseudorhodobacter	-0.63	*	0.90	***	
Pseudoxanthomonas	-0.73	**	0.67	*	
Ramlibacter	0.81	**	-0.77	**	
Rheinheimera	-0.87	***	0.62	*	
Rhodobacter	0.82	**	-0.85	***	
Rhodococcus	0.78	**	-0.69	*	
Rhodoferax	0.60	*	-0.82	**	
Rickettsia	-0.70	*	0.76	**	
Sporichthya	-0.89	***	0.66	*	

Tabrizicola	-0.60	*	0.59	*
unclassified Acidobacteria	-0.73	**	0.80	***
unclassified Alphaproteobacteria (miscellaneous)	-0.78	**	0.72	*
unclassified Betaproteobacteria (miscellaneous)	-0.71	*	0.73	*
unclassified Chloroflexi (miscellaneous)	-0.95	***	0.76	***
unclassified Deltaproteobacteria (miscellaneous)	-0.63	*	0.79	**
unclassified Gemmatimonadales	-0.70	*	0.64	*
unclassified Myxococcales (miscellaneous)	-0.81	**	0.80	**
unclassified Parcubacteria group	-0.77	**	0.72	*
unclassified Rhodobacteraceae	0.63	*	-0.72	*
unclassified Rhodobacterales	-0.69	*	0.69	*
Undibacterium	-0.69	*	0.66	*
Vulcaniibacterium	0.67	*	-0.61	*
Afipia	-0.82	**	0.57	ns
Aquabacterium	0.82	**	-0.42	ns
Arenimonas	-0.81	**	0.55	ns
Azotobacter	-0.80	**	0.59	ns
Bosea	-0.78	**	0.58	ns
Bradyrhizobium	0.78	**	-0.31	ns
Bryobacter	0.73	**	-0.40	ns
Caenimonas	0.72	*	-0.40	ns
Dyella	-0.72	*	0.54	ns
environmental Bacteria	-0.60	*	-0.02	ns
Gemmataceae	-0.71	*	-0.08	ns
Hassallia	0.78	**	-0.31	ns
Herpetosiphon	0.67	*	-0.36	ns
Inhella	-0.69	*	-0.36	ns
Limnobacter	0.67	*	-0.36	ns
Luteimonas	-0.65	*	0.55	ns
Methylotenera	-0.65	*	0.48	ns
Novosphingobium	0.65	*	-0.32	ns
Oscillochloris	0.64	*	-0.53	ns
Paracoccus	0.63	*	-0.54	ns
Pirellula	-0.60	*	0.35	ns
Pseudarthrobacter	-0.60	*	0.32	ns
Rhizobium	-0.60	*	0.29	ns

Rhodoplanes	-0.60	*	0.13	ns
Roseiflexus	-0.60	*	0.40	ns
Sandaracinus	0.95	***	-0.59	ns
Sphingobium	0.87	***	-0.50	ns
unclassified Actinobacteria	0.86	***	-0.43	ns
unclassified Cytophagaceae	-0.60	*	0.25	ns
unclassified Flavobacteriales (miscellaneous)	0.60	*	-0.14	ns
unclassified Rhodospirillaceae	-0.60	*	-0.02	ns
unclassified Xanthomonadaceae	0.61	*	-0.50	ns
Variovorax	-0.60	*	0.08	ns
Acidovorax	-0.59	ns	0.88	***
Algoriphagus	0.36	ns	-0.85	***
Azohydromonas	0.19	ns	-0.83	**
Bdellovibrio	0.10	ns	-0.82	**
Brevundimonas	-0.36	ns	-0.82	**
Burkholderia	-0.41	ns	0.82	**
Candidatus Nitrosocosmicus	0.02	ns	0.81	**
Candidatus Thermofonsia Clade 2	0.56	ns	0.80	**
Collimonas	-0.48	ns	0.78	**
Cupriavidus	-0.41	ns	0.78	**
Dongia	-0.37	ns	0.78	**
Herbaspirillum	-0.07	ns	-0.77	**
Hyphomicrobium	-0.37	ns	0.76	**
Klebsiella	-0.50	ns	0.76	**
Mesorhizobium	0.30	ns	-0.75	**
Microbacterium	-0.52	ns	0.73	**
Nitrososphaera	-0.23	ns	-0.72	*
Noviherbaspirillum	-0.04	ns	-0.72	*
Paeniglutamicibacter	-0.36	ns	0.71	*
Parabacteroides	-0.17	ns	0.69	*
Pedosphaera	-0.24	ns	0.69	*
Pelomonas	-0.29	ns	0.68	*
Planomicrobium	0.35	ns	-0.67	*
Sediminibacterium	-0.21	ns	0.67	*
Sinorhizobium	0.40	ns	-0.67	*
Stenotrophomonas	-0.58	ns	0.66	*
Streptomyces	-0.45	ns	0.65	*

unclassified Anaerolineae	0.32	ns	-0.65	*
unclassified Caulobacteraceae	-0.47	ns	0.65	*
unclassified Comamonadaceae	-0.52	ns	0.64	*
unclassified Gemmatimonadetes	-0.16	ns	0.63	*
unclassified Nitrosopumilales	-0.17	ns	-0.62	*
unclassified Opitutae	0.17	ns	-0.61	*
unclassified Oxalobacteraceae	-0.41	ns	0.61	*
unclassified Phyllobacteriaceae	0.45	ns	-0.61	*
unclassified Planctomycetes	-0.38	ns	0.61	*
unclassified Sphingomonadales	0.26	ns	-0.61	*
unclassified Thaumarchaeota (miscellaneous)	-0.26	ns	0.60	*
unclassified Thermomicrobiales	0.39	ns	0.60	*
unclassified Verrucomicrobia (miscellaneous)	-0.57	ns	0.59	*
unclassified Verrucomicrobiaceae	0.00	ns	-0.59	*

* *P* < 0.05, ** *P* < 0.01, *** *P* < 0.001, ns not significant. n=3.



Fig. S1 Number of total raw reads (mean \pm SE) (left) and non-rRNA reads (mean \pm SE) (right) in the metatranscriptomic datasets over four generations in SA treatments and control. N=3.



Fig. S2 Heinrich's triangle visualizing information on microbial composition at different classification levels of all 24 rhizosphere soil samples identified with MEGAN against NCBI. The numbers of identified phyla, families, genera and species are shown and the percentage of reads used for each classification level is noted. At phylum level, 2 Archaea, 22 bacteria, 22 eukaryotes and 2 viruses were found; at family level, 3 Archaea, 134 bacteria, 69 eukaryote (only fungi were counted) and 2 viruses were found; at genus level, 4 Archaea, 382 bacteria, 83 eukaryotes (9 fungi, 27 nematodes, 27 algae and 20 others) and 3 viruses were found: at species level, 12 Archaea, 1086 bacteria, 12 eukaryotes and 3 viruses were found. In the Heinrich's triangle, only the information of bacteria is listed, this includes numbers of bacteria and their read percentage against all the reads in that taxonomic level.



Fig. S3 The percentage of reads at different taxonomic levels. (a) The percentage of reads mapping to bacteria, archaea, eukaryote and virus reads of the total mapped reads. (b) The percentage of reads mapping to fungi, nematodes, algae and others in the total eukaryote reads.



Fig. S4 Venn diagram showing the immigration and extinction numbers at genus level for Archaea, eukaryotes and viruses at each generation in the rhizosphere soil samples of SA-treated or control *J. vulgaris* plants.



Fig. S5 Scatter plot showing log $_{10}$ transformed read counts of all 408 genera in rhizosphere soil samples of SA-treated and control plants against genus abundance. Rank number was based on the values of total read counts of all genera from the sum of the reads in SA and control treatments. The genus with highest abundance corresponds to the rank number 1. The cut-off line is based on < 0.01% of the total reads.



Fig. S6 Nonmetric multidimensional scaling (NMDS) plot based on Bray-Curtis distance representing the taxonomic information from the bacterial genera of 24 rhizosphere soil samples. For each treatment combination, the centroid is connected to the three replicates.



Fig. S7 Principal component analysis (PCA) representing the taxonomic information on the genera of 24 rhizosphere soil samples of SA-treated and control *J. vulgaris* plants. PCA scores are based on relative abundance at genus level.




Fig. S8 Heatmap with a hierarchical clustering analysis of 264 genera of rhizosphere soil of SA-treated plants (a) and 270 genera of soil samples of control plants (b). The hierarchical clustering analysis was calculated with Pearson distance and the Ward clustering algorithm based on the relative hit numbers of each genus. The color code represents the values of log2 transformed row-centered relative hit numbers. SA1, SA2, SA3 and SA4 represent SA treatments from the 1st generation, 2nd generation, 3rd generation and 4th generation. Control 1, control2, control3 and control4 represent control treatments from the 1st generation, 3rd generation and 4th generation is represented by three replicates indicated as 01, 02 and 03 respectively. E.g. the code 1C_01 represent the first generation control's first replicate.

Chapter 4

Activation of the SA-associated plant defense pathway alters the functions of soil microbial communities in four sequential generations

> Jing Zhang^{1*} T. Martijn Bezemer^{1, 2} Peter G.L. Klinkhamer¹ Klaas Vrieling¹

 Plant Science and Natural Products, Institute of Biology Leiden (IBL), Leiden University, Sylviusweg 72, 2333BE Leiden, The Netherlands

2) The Netherlands Institute of Ecology (NIOO-KNAW), Department of Terrestrial Ecology, Droevendaalsesteeg 10, 6708 PB, Wageningen, The Netherlands

Abstract

Systemic acquired resistance (SAR) is an immune response of plants that regulates plant hormonal signaling pathways and strengthens the ability of the plant to withstand pathogenic microbes. Aboveground application of salicylic acid (SA) to the plant can induce SAR and we showed that it mitigates negative effects of the soil microbial community on the performance of the plant Jacobaea vulgaris. How SAinduced resistance affects the expression of functional genes and gene ontology in the rhizosphere and how this phenomenon extends over multiple generations is not well studied. In this study, a meta-transcriptomics approach was used to characterize gene expression profiles of microbial communities in 24 soil samples of SA-treated and control plants over 4 generations. 71.6 million reads were used for de-novo assembly of the microbial transcriptome, after which a total of 1.3 million unique contigs (genes) were identified. Multivariate analysis revealed that the SA treatment, generation and the interaction between these two affected the functional genes of the rhizosphere microbial communities of J. vulgaris. In general, the effect of the SA treatment on microbial gene expression was lowest in the first generation and strongest in the fourth generation. Microbes in soil samples of SA-treated and control plants showed 1663 differentially expressed genes. In the first generation only two genes differed significantly in gene expression between microbes from soils of SA treated and control plants while in the fourth generation 361 genes were differentially expressed between microbes from soils of SA treated and control plants. None of the significantly expressed SA-downregulated genes were present in all four generations, while only one SA-upregulated gene was observed in all four generations. Gene ontology (GO) analysis showed that soil microbial communities in rhizosphere soil of SA-treated plants increased the expression of thirteen GO terms in the second, third and fourth generation. These increased GO terms were mostly related to viral RNA genome replication, to interactions with host cells, to organelles of the host cells and to RNA polymerase activities. There were six GO terms of which the expression decreased in the second, third and fourth generation, and these were associated with processing nitrogen and macromolecules. Overall, our results show that aboveground activation of defenses in the plant affects the expression of functional genes in the soil microbial communities belowground. This suggests that plants may recruit functional

rhizosphere microbiomes that improve plant health and crop production in agriculture.

Keywords

Meta-transcriptomics, Soil microbial community, Functional genes, Plant-soil interactions, Induced resistance, Rhizosphere soil, Salicylic acid

Introduction

Plants can alter the microbiome of the soil in which they grow, and in turn, microorganisms can influence plant performance. The rhizosphere microbiome, defined as the microbial community established near or on plant roots, can have negative, positive and neutral effects on the growth of a host plant (Van Wees et al., 2008; Raaijmakers, et al., 2009; Berendsen et al., 2012). Microbes such as plant growth-promoting bacteria (PGPB) and arbuscular mycorrhizal fungi (AMF) are typically characterized as plant beneficial, because of their contribution to plant health and nutrient uptake (Jeffries et al., 2003; Compant et al., 2010). In contrast, pathogenic microbes typically reduce plant growth and trigger defense mechanisms in the plant (Pieterse et al., 2001). However, the overall net effect of soil microbial communities on plant growth is often negative (Nijjer et al., 2007; Wardle et al., 2011). This might be due to e.g. competition between plants and microbes for available nutrients or soil pathogens (Berendse, 1994; Callaway et al., 2004; Mazzoleni et al., 2015; Cesarano et al., 2017). In response, plants have evolved hormone-driven defensive strategies to suppress these pathogenic impacts, such as systemic acquired resistance (SAR) and induced systemic resistance (ISR) (Bruce and Pickett, 2007; Berendsen et al., 2012; Huang et al., 2014; Ökmen and Doehlemann, 2014).

Systemic acquired resistance (SAR) is a distinct transduction pathway, which is involved in the biological processes that enhance the plant's immune system and defense against microbial pathogens (Reymond and Farmer, 1998; Walters and Heil, 2007; Pieterse et al., 2014; Haney and Ausubel, 2015). An infection caused by a pathogenic microbe can induce SAR, in which plants enhance their immune system by expressing genes coding for pathogenic-proteins (PR) in infested and uninfected tissues (Kachroo and Robin, 2013; Shah and Zeier, 2013; Gao et al., 2015). Apart from local induction by pathogenic microbes, SAR can also be induced by foliar sprays of the phytohormone salicylic acid (SA) (Reymond and Farmer, 1998). Applying a low concentration of SA directly to leaf tissues results in the activation of SA signaling pathways and this has been considered an effective way to activate defense signals in many plant species (Reymond and Farmer, 1998; Pozo and Azcón-Aguilar, 2007; Vlot et al., 2009).

In Chapter 2, we showed that the application of SA mitigates the negative effects of soil microbes on the growth of J. vulgaris although this effect did not increase further in subsequent generations of plant growth. A number of studies have examined the expression of functional genes in soil microbial communities. For example, Xue et al. (2016) showed that changing the temperature of soil significantly altered the gene expression in soil microbial communities and these genes were related to maintaining carbon and nitrogen stability in the soil, resulting in higher plant growth. Moreover, Castro et al. (2019) recently demonstrated that plants can change the expression of functional genes (i.e., carbon metabolic genes) in the soil microbial community in response to environmental changes such as drought. Here we hypothesize that application of SA to plants can also cause changes in the expression of functional genes in the soil microbial community and we hypothesize that the altered gene expression is related to the suppression of soil microbial pathogens of plants (Maurhofer et al., 1998; Verberne et al., 2000; Tanaka et al., 2015). Moreover, we expect, that the gene expression difference in the rhizosphere microbial community of control and SA treated plants will increased over generations of plant growth.

Previously, we analyzed the changes in the composition of the microbial community in the rhizosphere soil upon foliar application with SA and showed that the composition of rhizosphere bacterial communities differed among four plant generations of *J. vulgaris* and between soils from SA treated and control plants. However, the composition differed strongly among generations (Chapter 3). Functions of the soil microbial community can be performed by different microbial taxa (Burke et al., 2011; Liu et al., 2018; Liu et al., 2020) and hence we expect that there is functional redundancy in the soil microbial community and a consistent effect of SA application on gene expression in the microbial community.

In this study we ask the following questions: (1) Does the application of SA on leaves of *J. vulgaris* significantly alter the gene expression of the microbial community in the rhizosphere? (2) Does the effect differ between generations or is there an interaction between the SA treatment and generation on the gene expression in the microbial communities? (3) Which groups of genes or gene ontology pathways in the rhizosphere microbiome are influenced by SA-application over generations?

Materials, methods and bioinformatics processing

The multi-generation growth experiment with *J. vulgaris* has been described in Chapter 3. In short, *J. vulgaris* plants were grown for four generations on soils inoculated with soil from the previous generation from the same treatment with a foliar SA application treatment and a control treatment. Each treatment had 10 replicates. For each treatment, the three successively labeled replicates (No. 1, 2, 3, No. 4, 5, 6 and No. 7, 8, 9) were mixed and used as one pooled replicate, Hence, the three pooled replicates were used for RNA extraction for each treatment in each generation and a total of 24 soil samples were used for RNA extraction (3 replicates x 2 treatments x 4 generations). RNAseq was carried out using the Illumina platform.

Processing of the data included quality control of raw reads (FastQC), data trimming (Trimmomatic 0.39), filtering out ribosomal RNAs (SortMeRNA), de novo assembly of reads (Trinity), remove duplicates (CD-HIT-EST algorithm), mapping back to the transcriptome (Bowtie2). For a detailed description see Chapter 3. Gene ontology enrichment was performed using Trinotate and Goseq against the SwissProt, NR (non-redundant) and Pfam databases (Bryant et al., 2017; Bateman, 2019; El-Gebali et al., 2019).

Statistical analyses

Prior to analysis, the raw data were normalized. TMM (trimmed mean of M-values) normalization was used for read counts among all 24 samples (Robinson and Oshlack, 2010). A principal component analysis (PCA) was employed using the normalized number of genes to examine the composition of rhizosphere soil samples of SA-treated and control plants for the four generations. A PERMANOVA test was performed using the *adonis* function (number of permutations = 999) in R within the "vegan" package to verify the effects of the SA treatment and time on the composition of all expressed genes. To compare similarities among samples of treatment SA and control over four generations, a Pearson correlation for pairwise sample comparison based on the normalized raw read counts of all replicates in the control and SA treatments was performed in R and a heatmap was produced.

Differential gene expression (DE) analysis was performed for all possible combinations of replicates of sets of 8 samples (2 treatments x 4 generations) with EdgeR with raw read counts as input. EdgeR normalizes the data to TMM before further processing. After DE analysis in EdgeR, for all differentially expressed genes of the 8 samples Volcano plots were made for the contrast between SA-treated and control samples per generation. Log2 (FC) values were used as x-variable and -log10 (FDR) for the y-variable to produce a volcano plot of differentially expressed genes between control and SA-treated soil samples per generation. Genes that were significantly differentially expressed between SA-treated and control soil samples that could be annotated were listed. A clustered heatmap based on Euclidean distances (Danielsson, 1980) of gene expression derived from EdgeR per treatment after Z-scored transformation was generated in R using the package "pheatmap" (Kolde and Kolde, 2015).

To visualize the gene expression changes among different hormonal treatments and time categories, an NMDS (nonmetric multidimensional scaling) plot using the Bray-Curtis index as a measure of dissimilarity was generated using TMM normalized read counts. To verify changes in the composition of the 1663 expressed genes due to the SA treatment and time effect, a PERMANOVA test was performed using the adonis function (number of permutations = 999) in R within the "vegan" package.

Gene ontology (GO) enrichment was performed with "GoSeq" for each generation separately. Gene functional classification was determined for three categories: biological processes, cellular components and molecular functions. GO terms affiliated to Eukaryotes (e.g. mitochondria) were removed. The rich factor was calculated as the number of differentially expressed genes in the ontology divided by the number of all genes that were used as a background gene list.

Results

Comparing read counts between generations and treatments

A total of 898,4 million raw sequencing reads were obtained from the 24 metatranscriptomic libraries. The details of the library size and basic information

about read quality were described in Chapter 3. A principal component analysis (PCA) using log2 transformed normalized CPM showed that the read counts of contigs in the microbial community of rhizosphere soil of the *J. vulgaris* samples among generations were well separated (Fig. 1), this was in line with the permutation test (PERMANOVA $R^2 = 0.22$, F = 19.6, $df_1 = 3$, $df_2 > 999$, p < 0.01). In addition, the effect of SA application was significant (PERMANOVA $R^2 = 0.07$, F = 6.3, $df_1 = 1$, $df_2 > 999$, p < 0.05). Gene expression patterns of SA-treated *J. vulgaris* and control samples were better separated in the third and fourth generation than in the first and second generation (Fig. 1). In the correlation matrix for all sample replicates generated with PtR (a tool for comparing sample replicates in Trinity) (Fig. 1), samples within treatments were positively correlated with each other and also there was a positive correlation between samples within generations especially for the first generation. The heatmap showed clear clustering of treatments within generations except for generation 1. The separation between the SA and the control treatment became more distinct over generations (Fig. 2).



Fig. 1 Scatter plot from a principal component analysis (PCA) of TMM normalized CPM representing the overall gene expression patterns of different rhizosphere soil



samples of SA-treated and control *J. vulgaris* plants over generations. Shapes represent the treatments and colors represent generations.

Fig. 2 Clustered heatmap visualizing the Pearson correlation matrix for pairwise sample comparisons based on TMM normalized read counts per million. The heat map shows the correlation in microbial gene expression in all paired replicates between rhizosphere soil samples of SA-treated and control *J. vulgaris* plants over four generations. The dendrogram illustrates the relationship-distance between samples and is calculated based on a Pearson correlation coefficient. The color key represents the z-score of log2 CPM. The legends on the sides represent: Generation (1-4), treatment (SA/Control) and replicate number (01-03).

Differential gene expression

In total, 0.36 million genes were detected. Of those genes, 1663 were differentially expressed between all possible combinations of replicates of sets of 8 samples (2 treatments x 4 generations). Hierarchical clustering on CPM for 1663 differentially expressed genes was performed to explore the patterns of gene expression of the microbial communities between all pairwise combinations of all the samples among SA and control treatments over four generations (Fig. 3). Except for the first generation, SA and control samples were separated from each other in different clusters (Fig. 3). However, among generations, different clusters of genes were differentially grouped. Differences were most pronounced between on the one hand, the first and second generation, and on the other hand, the third and fourth generation.



Fig. 3 Heatmap showing 1663 differentially expressed genes (FDR < 0.05 and fold change ≥ 2) between all possible combinations of replicates of rhizosphere soil samples of SA-treated and control *J. vulgaris* plants over four generations based on TMM normalized CPMs. The color key represents the z-score of log2 CPM. The dendrogram on the x-axis illustrates the hierarchal clustering of relationship-distance between replicates using TMM normalized log2-transformed CPM. The legend on the bottom represents: generation (1-4), treatment (SA/control) and replicate number (01-03).

The NMDS plot showed that the 1663 differentially expressed microbial genes detected with EdgeR were differentially expressed in the different generations 119

(Fig. 4, PERMANOVA $R^2 = 0.63$, F = 21.8, $df_1 = 3$, $df_2 = 1662$, p < 0.01) and also that genes were differently expressed between the SA treatment and the control (PERMANOVA $R^2 = 0.07$, F = 7.0, $df_1 = 1$, $df_2 = 1662$, p < 0.01). The effect of the SA treatment was not the same in each generation as indicated by the significant interaction (PERMANOVA $R^2 = 0.15$, F = 5.2, $df_1 = 3$, $df_2 = 1662$, p < 0.001).



Fig. 4 Multivariate analysis of 1663 differentially expressed microbial genes between all replicates in rhizosphere samples from SA-treated and control *J. vulgaris* plants grown in four generations. Shown are sample scores from a nonmetric multidimensional scaling (NMDS) plot.

To identify the numbers of significantly down- or up-regulated genes in the SA treatment in each generation in the rhizosphere microbial community, volcano plots were made (Fig. 5). In the first generation, no downregulated genes were observed and only two upregulated genes were detected (Fig. 5a). This increased to 59 and 76 in the second, 89 and 26 in the third, and 187 and 174 in the fourth generation, respectively (Fig. 5b, c, d). Among all the significant differentially expressed genes, no genes were found that were downregulated after SA application in all four generations, while only one gene was observed that was upregulated in SA in all four generations (Fig. 6). Circa 90% of the genes that were significantly altered by the SA

treatment could not be annotated. Among all the annotated genes, only two of the significant differentially expressed microbial genes were detected in three generations and eight genes were detected in two generations (Fig. 6). Not all the genes could be matched with a function in the database. Detailed information of successfully annotated genes was listed in Table S1.



Fig. 5 Volcano plots of 1663 differentially expressed genes of the microbial community in rhizosphere samples of SA-treated and control *J. vulgaris* plants per generation. The x-axes show log2 fold changes of read counts of the genes of the SA treatment compared to the control, and the y-axes show the -log10 adjusted for FDR 121

values. SA upregulated genes are presented in purple, and SA downregulated genes are displayed in red, while non-significant genes are shown as light grey dots. 1st, 2nd, 3rd and 4th represent the different generations. The numbers inside each box represent the number of significantly up/down-regulated genes. The two vertical dashed lines represent the positive or negative log2 fold changes in the number of readcounts in the SA treatment compared to the control in the generation when -log10(FDR) is 2 as presented by the horizontal dashed lines.



Fig. 6 Venn diagrams showing the number of shared and unique up and down-regulated microbial genes over generations in the rhizosphere of *J. vulgaris*. The numbers represent the significantly differently expressed genes from the volcano plot (Fig. 5). 1^{st} , 2^{nd} , 3^{rd} and 4^{th} represent the different generations.

Gene ontology (GO) analysis

To profile differentially expressed pathways, we performed a gene ontology (GO) analysis for the soil samples of SA-treated and control plants for each generation (Fig. 7, Table S2). No significantly upregulated or downregulated GO terms were observed in the first generation (Fig. 7a, 7b). In the second, third and fourth generations, genes from classes of the GO categories "biological processes", "cellular components" and "molecular functions" were differentially expressed (Fig. 7a). 13 GO terms were upregulated in the SA treatment in three generations, while 18 GO terms were upregulated in one or two generations (Fig. 7a). Of the 13 GO terms upregulated in

three generations three belonged to the GO category "biological processes", and these GO terms were all related to viral RNA genome replication, seven belonged to the GO category "cellular components" and these GO terms were related to interactions with host cells and to organelles of the host cells and finally three belonged to the GO category "molecular function" and these GO terms were all related to RNA polymerase activity.

Only six GO terms were downregulated in the second, third and fourth generation in the rhizosphere of SA treated plants, while 58 GO terms were downregulated in one or two generations only (Fig. 7b). The six GO terms downregulated in three generations fell all in the GO category "biological processes" and the GO terms were related the localization of processes, to nitrogen processing and to processes involving macromolecules. None of GO terms involved in cellular components and molecular functions were present in these three generations.



SA-up regulated GO terms per generation



SA-down regulated GO terms per generation

Fig. 7 Gene ontology (GO) enrichment analysis of significantly differentially expressed genes in the microbial community in the rhizosphere. A bubble chart shows

Generation

enrichment of differentially expressed GO terms. The Y-axis label lists the GO terms, the size of the bubbles represents the rich factor (= amount of differentially expressed genes enriched in the ontology/total amount of all genes in the background gene set) in different generations. Gene classification of the annotated GO terms was grouped in three categories. Colors of the bubbles represent the significance level of enrichment as calculated with Goseq. a and b represent up and down-regulated GO terms in the SA treatment, respectively. Note: in the first generation, there were no up-or down-regulated GO terms.

Discussion

In this study, a high-throughput metatranscriptomic sequencing approach was used to examine how the aboveground application of SA to the plant impacts the functional gene expression of the microbial communities in the rhizosphere over four subsequent generations of plant growth. Our study shows that the activation of the SA-associated plant defense pathways significantly affected the gene expression of the microbial communities in the rhizosphere, but this effect differed over four generations. Notably, the numbers of differentially expressed genes increased over generations, and there was almost no overlap of in the genes that were significantly expressed in the four generations. Moreover, foliar application of SA caused upregulation of genes of the microbial community related to GO terms associated to viral RNA genome replication, to interactions with host cells, to organelles of the host cells and to RNA polymerase activities, while downregulated GO terms of the microbial community were associated biosynthetic processes involving nitrogen and metabolic processes.

Our study shows that application of SA to plants changed the functional gene expression in the rhizosphere microbial community. This complements previous studies, which report that effects of different abiotic factors alter the expression of functional genes in the soil community (Xue et al., 2016; Castro et al., 2019). Interestingly, in our study, the highest number of significantly expressed genes was recorded in the fourth generation, which suggests that the effect of SA on gene expression becomes more pronounced over time. We did not find a selection-effect of SA on the rhizosphere bacterial community over multiple generations (results in Chapter 3). Hence, we cannot conclude that the increase in the number of significant

expressed genes in our study was due to a specific rhizosphere bacterial community that became increasingly active.

Our finding that the expression of functional genes differed strongly among generations is in line with the previous findings that different taxonomic groups are present in the rhizosphere of SA treated J. vulgaris plants in each generation (Chapter 3). However, this clearly contrasts our prediction that there will be functional redundancy in the microbial community. In the same experiment also plant biomass was measured (Chapter 2) and SA treated plants in all generations did better than the control plants showing that from the plant's perspective different microbial taxa with different gene expressions in the rhizosphere provided similar functions. Our findings are in contrast to studies (e.g. Burke et al., 2011; Liu et al., 2018; Liu et al., 2020) that mention that particular functions of the soil microbial community are often distributed across multiple microbial taxa and more closely resemble other studies that show that environmental changes can cause selection of both different taxa and functions in the soil microbial communities (Haggerty and Dinsdale, 2017). It is important to note that in our study, in each generation we placed a subset of the microbial community in a sterile background. This may have led to selection for microbes and consequently different functions in each generation.

At the gene ontology level, we mapped 13 SA-upregulated and six SA-downregulated GO terms that were expressed in the second, third and fourth generation. The proportion of significantly expressed GO terms was high, compared to the proportion of significantly expressed genes. This is because most of the functional genes in this study could not be annotated, while at the ontology level more reads were matched with a function. As the taxa significantly selected by SA differed strongly from generation to generation, it is notable that there we detected many significant GO terms that were found in multiple generations.

Our results show that activating SA resistance in the plant drives gene expression in the rhizosphere microbiome. However, whether SA application to the plant suppressed soil pathogenic microbes remains unproven in our study. SA induced resistance is often reported to play an important role in resistance to a broad range of microbial pathogens, such as bacteria, fungi and viruses (Murphy et al., 1999; 126 Gilliland et al., 2003; Mayers et al., 2005; Kundu et al., 2011; Li et al., 2019; Yuan et al., 2019). Interestingly, at the ontology level, we found up-regulated GO terms that were involved in viral (RNA) genome replication and viral processes, and these GO terms increased in importance over generations. These results indicate that viruses in the soil may play a role in SA-induced resistance of host plants against soil microbes. It is well known that the soil contains bacteriophages as well as virus controlling microbial pathogens (Duckworth and Gulig, 2002; Svircev et al., 2018; Jamal et al., 2019; Kortright et al., 2019; Rehman et al., 2019). However, their exact role in the rhizosphere microbiome is still poorly understood and further studies should examine these virus-microbe-plant interactions in more detail.

In conclusion, our study shows that application of SA to the plant *J. vulgaris* causes differential gene expression in the rhizosphere microbial community. However, our data also shows that these effects vary among plant generations. Plant-defense-soil microbe interactions may be regulated by viruses or viral phages.

Acknowledgements

We would like to thank Karin van der Veen-van Wijk for assistance at the final biomass harvest, Connor Philippo for his practical assistance in extracting RNA and performing the quality control of raw sequencing reads, Bing Xie for his generous supporting in sharing his private server and help in troubleshooting, Fons Verbeek for sharing public resource of the server in the Leiden Institute of advanced computer science (LIACS). We also thank Yangan Chen for discussing general transcriptomic knowledge, and Martine Huberty for PCA statistical analysis. Jing Zhang would also like to thank the China Scholarship Council for financial support.

References

- Anantharaman, V. and Aravind, L., 2003. Evolutionary history, structural features and biochemical diversity of the NlpC/P60 superfamily of enzymes. *Genome biology*, 4(2), p.R11.
- Andersson, C.E., Lagerbäck, P. and Carlson, K., 2010. Structure of bacteriophage T4 endonuclease II mutant E118A, a tetrameric GIY-YIG enzyme. *Journal of molecular biology*, 397(4), pp.1003-1016.
- Antoine, R., Jacob-Dubuisson, F., Drobecq, H., Willery, E., Lesjean, S. and Locht, C., 2003. Overrepresentation of a gene family encoding extracytoplasmic solute receptors in *Bordetella. Journal of bacteriology*, 185(4), pp.1470-1474.
- Aravind, L. and Koonin, E.V., 1998. The HD domain defines a new superfamily of metal-dependent phosphohydrolases. *Trends in biochemical sciences*, 23(12), pp.469-472.
- Bateman, A., 2019. Uniprot: A Universal Hub of Protein Knowledge. In Protein Science, 28, pp. 32-32.
- Beatson, S.A., Minamino, T. and Pallen, M.J., 2006. Variation in bacterial flagellins: from sequence to structure. *Trends in microbiology*, 14(4), pp.151-155.
- Berendse, F., 1994. Competition between plant populations at low and high nutrient supplies. *Oikos*, pp.253-260.
- Berendsen, R.L., Pieterse, C.M. and Bakker, P.A., 2012. The rhizosphere microbiome and plant health. *Trends in plant Science*, *17*(8), pp.478-486.
- Bonocora, R.P. and Shub, D.A., 2001. A novel group I intron-encoded endonuclease specific for the anticodon region of tRNAfMet genes. *Molecular microbiology*, *39*(5), pp.1299-1306.
- Boonrod, K., Galetzka, D., Nagy, P.D., Conrad, U. and Krczal, G., 2004. Single-chain antibodies against a plant viral RNA-dependent RNA polymerase confer virus resistance. *Nature biotechnology*, 22(7), pp.856-862.
- Braun, T.F., Khubbar, M.K., Saffarini, D.A. and McBride, M.J., 2005. Flavobacterium johnsoniae gliding motility genes identified by mariner mutagenesis. Journal of bacteriology, 187(20), pp.6943-6952.
- Briers, Y., Schmelcher, M., Loessner, M.J., Hendrix, J., Engelborghs, Y., Volckaert, G. and Lavigne, R., 2009. The high-affinity peptidoglycan binding domain of *Pseudomonas* phage endolysin KZ144. *Biochemical and biophysical research communications*, 383(2), pp.187-191.
- Bruce, T.J. and Pickett, J.A., 2007. Plant defence signalling induced by biotic attacks. *Current opinion* in plant biology, 10(4), pp.387-392.
- Bryant, D.M., Johnson, K., DiTommaso, T., Tickle, T., Couger, M.B., Payzin-Dogru, D., Lee, T.J., Leigh, N.D., Kuo, T.H., Davis, F.G. and Bateman, J., 2017. A tissue-mapped axolotl de novo transcriptome enables identification of limb regeneration factors. *Cell reports*, 18(3), pp.762-776.
- Burd, C.G. and Dreyfuss, G., 1994. Conserved structures and diversity of functions of RNA-binding proteins. Science, 265(5172), pp.615-621.
- Burke, C., Steinberg, P., Rusch, D., Kjelleberg, S. and Thomas, T., 2011. Bacterial community assembly based on functional genes rather than species. *Proceedings of the National Academy of Sciences*, U.S.A, 108(34), pp.14288-14293.
- Callaway, R.M. and Ridenour, W.M., 2004. Novel weapons: invasive success and the evolution of increased competitive ability. *Frontiers in Ecology and the Environment*, 2(8), pp.436-443.
- Castro, S.P., Cleland, E.E., Wagner, R., Al Sawad, R. and Lipson, D.A., 2019. Soil microbial responses to drought and exotic plants shift carbon metabolism. *The ISME journal*, *13*(7), pp.1776-1787.

- Cesarano, G., De Filippis, F., La Storia, A., Scala, F. and Bonanomi, G., 2017. Organic amendment type and application frequency affect crop yields, soil fertility and microbiome composition. Applied Soil Ecology, 120, pp.254-264.
- Cho, K.H. and Salyers, A.A., 2001. Biochemical analysis of interactions between outer membrane proteins that contribute to starch utilization by Bacteroides thetaiotaomicron. Journal of Bacteriology, 183(24), pp.7224-7230.
- Compant, S., Clément, C. and Sessitsch, A., 2010. Plant growth-promoting bacteria in the rhizo-and endosphere of plants: their role, colonization, mechanisms involved and prospects for utilization. Soil Biology and Biochemistry, 42(5), pp.669-678.
- Convery, M.A., Rowsell, S., Storehouse, N.J., Ellington, A.D., Hirao, I., Murray, J.B., Peabody, D.S., Phillips, S.E. and Stockley, P.G., 1998. Crystal structure of an RNA aptamer-protein complex at 2.8 Å resolution. *Nature structural biology*, 5(2), pp.133-139.
- D'Amico, S., Collins, T., Marx, J.C., Feller, G., Gerday, C. and Gerday, C., 2006. Psychrophilic microorganisms: challenges for life. EMBO reports, 7(4), pp.385-389.
- De Mot, R., Schoofs, G., Roelandt, A., Declerck, P., Proost, P., Damme, J.V. and Vanderleyden, J., 1994. Molecular characterization of the major outermembrane protein OprF from plant rootcolonizing Pseudomonas fluorescens. Microbiology, 140(6), pp.1377-1387.

Duckworth, D.H. and Gulig, P.A., 2002. Bacteriophages. BioDrugs, 16(1), pp.57-62.

- Dunin-Horkawicz, S., Feder, M. and Bujnicki, J.M., 2006. Phylogenomic analysis of the GIY-YIG nuclease superfamily. BMC genomics, 7(1), p.98.
- El-Gebali, S., Mistry, J., Bateman, A., Eddy, S.R., Luciani, A., Potter, S.C., Qureshi, M., Richardson, L.J., Salazar, G.A., Smart, A. and Sonnhammer, E.L.L., 2019. The Pfam protein families database in 2019. Nucleic acids research, 47(D1) pp.D427-D432.
- Felix, G., Duran, J.D., Volko, S. and Boller, T., 1999. Plants have a sensitive perception system for the most conserved domain of bacterial flagellin. The Plant Journal, 18(3), pp.265-276.
- Fernando Gil, J., Wibberg, D., Eini, O., Savenkov, E.I., Varrelmann, M. and Liebe, S., 2020. Comparative transcriptome analysis provides molecular insights into the interaction of beet necrotic yellow vein virus and beet soil-borne mosaic virus with their host sugar beet. Viruses, 12(1), p.76.
- Moscat, J., Diaz-Meco, M.T., Albert, A. and Campuzano, S., 2006. Cell signaling and function organized by PB1 domain interactions. Molecular cell, 23(5), pp.631-640.
- Gao, Q.M., Zhu, S., Kachroo, P. and Kachroo, A., 2015. Signal regulators of systemic acquired resistance. Frontiers in plant Science, 6, p.228.
- Gilliland, A., Singh, D.P., Hayward, J.M., Moore, C.A., Murphy, A.M., York, C.J., Slator, J. and Carr, J.P., 2003. Genetic modification of alternative respiration has differential effects on antimycin A-induced versus salicylic acid-induced resistance to Tobacco mosaic virus. Plant Physiology, 132(3), pp.1518-1528.
- Haggerty, J.M. and Dinsdale, E.A., 2017. Distinct biogeographical patterns of marine bacterial taxonomy and functional genes. Global Ecology and Biogeography, 26(2), pp.177-190.
- Haney, C.H. and Ausubel, F.M., 2015. Plant microbiome blueprints. Science, 349(6250), pp.788-789.
- Hardy, S.F., German, T.L., Loesch-Fries, L.S. and Hall, T.C., 1979. Highly active template-specific RNA-dependent RNA polymerase from barley leaves infected with brome mosaic virus. Proceedings of the National Academy of Sciences, U.S.A, 76(10), pp.4956-4960.
- Henry, J.T. and Crosson, S., 2011. Ligand-binding PAS domains in a genomic, cellular, and structural context. Annual review of microbiology, 65, pp.261-286.
- Hong, Y., Cole, T.E., Brasier, C.M. and Buck, K.W., 1998. Evolutionary relationships among putative RNA-dependent RNA polymerases encoded by a mitochondrial virus-like RNA in the Dutch

elm disease fungus, *Ophiostoma novo-ulmi*, by other viruses and virus-like RNAs and by the *Arabidopsis* mitochondrial genome. *Virology*, 246(1), pp.158-169.

- Hoppe, J., Ünal, C.M., Thiem, S., Grimpe, L., Goldmann, T., Gaßler, N., Richter, M., Shevchuk, O. and Steinert, M., 2017. PilY1 promotes *Legionella pneumophila* infection of human lung tissue explants and contributes to bacterial adhesion, host cell invasion, and twitching motility. *Frontiers in cellular and infection microbiology*, 7, p.63.
- Huang, X.F., Chaparro, J.M., Reardon, K.F., Zhang, R., Shen, Q. and Vivanco, J.M., 2014. Rhizosphere interactions: root exudates, microbes, and microbial communities. *Botany*, 92(4), pp.267-275.
- Ito, T., Matsui, Y., Ago, T., Ota, K. and Sumimoto, H., 2001. Novel modular domain PB1 recognizes PC motif to mediate functional protein-protein interactions. *The EMBO journal*, 20(15), pp.3938-3946.
- Jamal, M., Bukhari, S.M., Andleeb, S., Ali, M., Raza, S., Nawaz, M.A., Hussain, T., Rahman, S.U. and Shah, S.S., 2019. Bacteriophages: an overview of the control strategies against multiple bacterial infections in different fields. *Journal of basic microbiology*, 59(2), pp.123-133.
- Jeffries, P., Gianinazzi, S., Perotto, S., Turnau, K. and Barea, J.M., 2003. The contribution of arbuscular mycorrhizal fungi in sustainable maintenance of plant health and soil fertility. *Biology and fertility of soils*, 37(1), pp.1-16.
- Johnson, E.R. and McKay, D.B., 1999. Crystallographic structure of the amino terminal domain of yeast initiation factor 4A, a representative DEAD-box RNA helicase. *Rna*, *5*(12), pp.1526-1534.
- Jung, A.L., Stoiber, C., Herkt, C.E., Schulz, C., Bertrams, W. and Schmeck, B., 2016. Legionella pneumophila-derived outer membrane vesicles promote bacterial replication in macrophages. PLoS pathogens, 12(4), p.e1005592.
- Kachroo, A. and Robin, G.P., 2013. Systemic signaling during plant defense. Current Opinion in Plant Biology, 16(4), pp.527-533.
- Kolde, R. and Kolde, M.R., 2015. Package 'pheatmap'. R Package, 1(7), p.790.
- Kortright, K.E., Chan, B.K., Koff, J.L. and Turner, P.E., 2019. Phage therapy: a renewed approach to combat antibiotic-resistant bacteria. *Cell host & microbe*, *25*(2), pp.219-232.
- Krishna, S.S., Majumdar, I. and Grishin, N.V., 2003. Structural classification of zinc fingers: survey and summary. *Nucleic acids research*, 31(2), pp.532-550.
- Krogh, S., Jørgensen, S.T. and Devine, K.M., 1998. Lysis genes of the Bacillus subtilis defective prophage PBSX. Journal of bacteriology, 180(8), pp.2110-2117.
- Kundu, S., Chakraborty, D. and Pal, A., 2011. Proteomic analysis of salicylic acid induced resistance to Mungbean yellow mosaic India virus in Vigna mungo. *Journal of proteomics*, 74(3), pp.337-349.
- Kuo, W.T., Chin, K.H., Lo, W.T., Wang, A.H.J. and Chou, S.H., 2008. Crystal structure of the C-terminal domain of a flagellar hook-capping protein from *Xanthomonas campestris*. *Journal of molecular biology*, 381(1), pp.189-199.
- Li, T., Huang, Y., Xu, Z.S., Wang, F. and Xiong, A.S., 2019. Salicylic acid-induced differential resistance to the Tomato yellow leaf curl virus among resistant and susceptible tomato cultivars. *BMC plant biology*, 19(1), pp.1-14.
- Lindquist, J.A. and Mertens, P.R., 2018. Cold shock proteins: from cellular mechanisms to pathophysiology and disease. *Cell Communication and Signaling*, 16(1), p.63.
- Liu, Y.C., Machuca, M.A., Beckham, S.A., Gunzburg, M.J. and Roujeinikova, A., 2015. Structural basis for amino-acid recognition and transmembrane signalling by tandem Per-Arnt-Sim (tandem PAS) chemoreceptor sensory domains. *Acta Crystallographica Section D: Biological Crystallography*, 71(10), pp.2127-2136.

- Liu, Y.R., Delgado-Baquerizo, M., Bi, L., Zhu, J. and He, J.Z., 2018. Consistent responses of soil microbial taxonomic and functional attributes to mercury pollution across China. *Microbiome*, 6(1), pp.1-12.
- Liu, Y.R., Delgado-Baquerizo, M., Yang, Z., Feng, J., Zhu, J. and Huang, Q., 2020. Microbial taxonomic and functional attributes consistently predict soil CO2 emissions across contrasting croplands. *Science of The Total Environment*, 702, p.134885.
- Maurhofer, M., Reimmann, C., Schmidli-Sacherer, P., Heeb, S., Haas, D. and Défago, G., 1998. Salicylic acid biosynthetic genes expressed in *Pseudomonas fluorescens* strain P3 improve the induction of systemic resistance in tobacco against tobacco necrosis virus. *Phytopathology*, 88(7), pp.678-684.
- Mayers, C.N., Lee, K.C., Moore, C.A., Wong, S.M. and Carr, J.P., 2005. Salicylic acid-induced resistance to *Cucumber mosaic virus* in squash and *Arabidopsis thaliana*: contrasting mechanisms of induction and antiviral action. *Molecular plant-microbe interactions*, 18(5), pp.428-434.
- Mazzoleni, S., Bonanomi, G., Incerti, G., Chiusano, M.L., Termolino, P., Mingo, A., Senatore, M., Giannino, F., Cartenì, F., Rietkerk, M. and Lanzotti, V., 2015. Inhibitory and toxic effects of extracellular self-DNA in litter: a mechanism for negative plant-soil feedbacks? *New Phytologist*, 205(3), pp.1195-1210.
- Mitchell, P., Petfalski, E., Shevchenko, A., Mann, M. and Tollervey, D., 1997. The exosome: a conserved eukaryotic RNA processing complex containing multiple 3'→ 5' exoribonucleases. *Cell*, 91(4), pp.457-466.
- Murphy, A.M., Chivasa, S., Singh, D.P. and Carr, J.P., 1999. Salicylic acid-induced resistance to viruses and other pathogens: a parting of the ways? *Trends in plant science*, *4*(4), pp.155-160.
- Nar, H., Huber, R., Meining, W., Schmid, C., Weinkauf, S. and Bacher, A., 1995. Atomic structure of GTP cyclohydrolase I. *Structure*, 3(5), pp.459-466.
- Nandhagopal, N., Simpson, A.A., Gurnon, J.R., Yan, X., Baker, T.S., Graves, M.V., Van Etten, J.L. and Rossmann, M.G., 2002. The structure and evolution of the major capsid protein of a large, lipid-containing DNA virus. *Proceedings of the National Academy of Sciences*, U.S.A, 99(23), pp.14758-14763.
- Nijjer, S., Rogers, W.E. and Siemann, E., 2007. Negative plant-soil feedbacks may limit persistence of an invasive tree due to rapid accumulation of soil pathogens. *Proceedings of the Royal Society B: Biological Sciences*, 274(1625), pp.2621-2627.
- Ökmen, B. and Doehlemann, G., 2014. Inside plant: biotrophic strategies to modulate host immunity and metabolism. *Current opinion in plant biology*, 20, pp.19-25.
- Pao, G.M. and Saier, M.H., 1995. Response regulators of bacterial signal transduction systems: selective domain shuffling during evolution. *Journal of Molecular Evolution*, 40(2), pp.136-154.
- Pieterse, C.M., Van Pelt, J.A., Van Wees, S.C., Ton, J., Léon-Kloosterziel, K.M., Keurentjes, J.J., Verhagen, B.W., Knoester, M., Van der Sluis, I., Bakker, P.A. and Van Loon, L.C., 2001. Rhizobacteria-mediated induced systemic resistance: triggering, signalling and expression. *European Journal of Plant Pathology*, 107(1), pp.51-61.
- Pieterse, C.M., Zamioudis, C., Berendsen, R.L., Weller, D.M., Van Wees, S.C. and Bakker, P.A., 2014. Induced systemic resistance by beneficial microbes. *Annual review of phytopathology*, 52, pp.347-375.
- Pozo, M.J. and Azcón-Aguilar, C., 2007. Unraveling mycorrhiza-induced resistance. Current opinion in plant biology, 10(4), pp.393-398.

- Price, M.N., Wetmore, K.M., Waters, R.J., Callaghan, M., Ray, J., Liu, H., Kuehl, J.V., Melnyk, R.A., Lamson, J.S., Suh, Y. and Carlson, H.K., 2018. Mutant phenotypes for thousands of bacterial genes of unknown function. *Nature*, 557(7706), pp.503-509.
- Raaijmakers, J.M., Paulitz, T.C., Steinberg, C., Alabouvette, C. and Moënne-Loccoz, Y., 2009. The rhizosphere: a playground and battlefield for soilborne pathogens and beneficial microorganisms. *Plant and soil*, 321(1-2), pp.341-361.
- Ramos, H.C., Rumbo, M. and Sirard, J.C., 2004. Bacterial flagellins: mediators of pathogenicity and host immune responses in mucosa. *Trends in microbiology*, 12(11), pp.509-517.
- Rehman, S., Ali, Z., Khan, M., Bostan, N. and Naseem, S., 2019. The dawn of phage therapy. *Reviews in medical virology*, 29(4), p.e2041.
- Reymond, P. and Farmer, E.E., 1998. Jasmonate and salicylate as global signals for defense gene expression. *Current opinion in plant biology*, *1*(5), pp.404-411.
- Robinson, M.D. and Oshlack, A., 2010. A scaling normalization method for differential expression analysis of RNA-seq data. *Genome biology*, 11(3), pp.1-9.
- Saier, M.H., 2000. A functional-phylogenetic classification system for transmembrane solute transporters. *Microbiology and molecular biology reviews*, 64(2), pp.354-411.
- Schwach, F., Vaistij, F.E., Jones, L. and Baulcombe, D.C., 2005. An RNA-dependent RNA polymerase prevents meristem invasion by potato virus X and is required for the activity but not the production of a systemic silencing signal. *Plant physiology*, 138(4), pp.1842-1852.
- Sganga, M.W., Aksamit, R.R., Cantoni, G.L. and Bauer, C.E., 1992. Mutational and nucleotide sequence analysis of S-adenosyl-L-homocysteine hydrolase from *Rhodobacter capsulatus*. *Proceedings* of the National Academy of Sciences, U.S.A 89(14), pp.6328-6332.
- Shah, J. and Zeier, J., 2013. Long-distance communication and signal amplification in systemic acquired resistance. *Frontiers in Plant Science*, *4*, p.30.
- Shen, C., Shi, Y., Ni, Y., Deng, Y., Van Nostrand, J.D., He, Z., Zhou, J. and Chu, H., 2016. Dramatic increases of soil microbial functional gene diversity at the treeline ecotone of Changbai Mountain. *Frontiers in microbiology*, 7, p.1184.
- Sharma, M., Ellis, R.L. and Hinton, D.M., 1992. Identification of a family of bacteriophage T4 genes encoding proteins similar to those present in group I introns of fungi and phage. *Proceedings* of the National Academy of Sciences, U.S.A, 89(14), pp.6658-6662.
- Steen, A., Buist, G., Leenhouts, K.J., El Khattabi, M., Grijpstra, F., Zomer, A.L., Venema, G., Kuipers, O.P. and Kok, J., 2003. Cell wall attachment of a widely distributed peptidoglycan binding domain is hindered by cell wall constituents. *Journal of Biological Chemistry*, 278(26), pp.23874-23881.
- Stuwe, T., Hothorn, M., Lejeune, E., Rybin, V., Bortfeld, M., Scheffzek, K. and Ladurner, A.G., 2008. The FACT Spt16 "peptidase" domain is a histone H3-H4 binding module. *Proceedings of the National Academy of Sciences*, U.S.A, 105(26), pp.8884-8889.
- Sumimoto, H., Kamakura, S. and Ito, T., 2007. Structure and function of the PB1 domain, a protein interaction module conserved in animals, fungi, amoebas, and plants. *Science's STKE*, 2007(401), pp.re6-re6.
- Svircev, A., Roach, D. and Castle, A., 2018. Framing the future with bacteriophages in agriculture. Viruses, 10(5), p.218.
- Tanaka, S., Han, X. and Kahmann, R., 2015. Microbial effectors target multiple steps in the salicylic acid production and signaling pathway. *Frontiers in Plant Science*, 6, p.349.
- Van Wees, S.C., Van der Ent, S. and Pieterse, C.M., 2008. Plant immune responses triggered by beneficial microbes. *Current opinion in plant biology*, 11(4), pp.443-448.

- Verberne, M.C., Verpoorte, R., Bol, J.F., Mercado-Blanco, J. and Linthorst, H.J., 2000. Overproduction of salicylic acid in plants by bacterial transgenes enhances pathogen resistance. *Nature biotechnology*, 18(7), pp.779-783.
- Vlot, A.C., Dempsey, D.M.A. and Klessig, D.F., 2009. Salicylic acid, a multifaceted hormone to combat disease. *Annual review of phytopathology*, 47, pp.177-206.
- Wardle, D.A., Bardgett, R.D., Callaway, R.M. and Van der Putten, W.H., 2011. Terrestrial ecosystem responses to species gains and losses. *Science*, 332(6035), pp.1273-1277.
- Walters, D. and Heil, M., 2007. Costs and trade-offs associated with induced resistance. *Physiological and Molecular Plant Pathology*, 71(1-3), pp.3-17.
- Westman, J., Hube, B. and Fairn, G.D., 2019. Integrity under stress: Host membrane remodelling and damage by fungal pathogens. *Cellular microbiology*, 21(4), p.e13016.
- Wilkens, S., Zhang, Z. and Zheng, Y., 2005. A structural model of the vacuolar ATPase from transmission electron microscopy. *Micron*, *36*(2), pp.109-126.
- Williams, M.A., Taylor, E.B. and Mula, H.P., 2010. Metaproteomic characterization of a soil microbial community following carbon amendment. *Soil Biology and Biochemistry*, 42(7), pp.1148-1156.
- Witherell, G.W., Gott, J.M. and Uhlenbeck, O.C., 1991. Specific interaction between RNA phage coat proteins and RNA. Progress in nucleic acid research and molecular biology, 40, pp.185-220.
- Xue, K., Xie, J., Zhou, A., Liu, F., Li, D., Wu, L., Deng, Y., He, Z., Van Nostrand, J.D., Luo, Y. and Zhou, J., 2016. Warming alters expressions of microbial functional genes important to ecosystem functioning. *Frontiers in microbiology*, 7, p.668.
- Yao, Z., Zou, C., Peng, N., Zhu, Y., Bao, Y., Zhou, Q., Wu, Q., Chen, B. and Zhang, M., 2020. Virome identification and characterization of *Fusarium sacchari* and *F. andiyazi*: causative agents of *Pokkah boeng* disease in sugarcane. *Frontiers in microbiology*, 11, p.240.
- Yeh, L.C.C. and Lee, J.C., 1998. Yeast ribosomal proteins L4, L17, L20, and L25 exhibit different binding characteristics for the yeast 35S precursor rRNA. *Biochimica et Biophysica Acta* (BBA)-Gene Structure and Expression, 1443(1-2), pp.139-148.
- Younas, F., Soltanmohammadi, N., Knapp, O. and Benz, R., 2018. The major outer membrane protein of *Legionella pneumophila* Lpg1974 shows pore-forming characteristics similar to the human mitochondrial outer membrane pore, hVDAC1. *Biochimica et Biophysica Acta (BBA)-Biomembranes*, 1860(8), pp.1544-1553.
- Yuan, W., Jiang, T., Du, K., Chen, H., Cao, Y., Xie, J., Li, M., Carr, J.P., Wu, B., Fan, Z. and Zhou, T., 2019. Maize phenylalanine ammonia-lyases contribute to resistance to Sugarcane mosaic virus infection, most likely through positive regulation of salicylic acid accumulation. *Molecular plant pathology*, 20(10), pp.1365-1378.
- Zhang, G. and Darst, S.A., 1998. Structure of the *Escherichia coli* RNA polymerase α subunit aminoterminal domain. *Science*, 281(5374), pp.262-266.

Supplementary data

Table S1 Log2 (FC) of 70 differentially expressed annotated genes, the expression of which are significantly altered by SA treatments in at least one generation in the rhizosphere of *J. vulgaris* plants in four generations. When Log2 (FC) is > 0, the gene is up-regulated in the SA treatment and when Log2 (FC) is < 0, the gene is down-regulated in the SA treatment. '-' indicates that the gene was not detected in the treatment; 'ns' represents the gene is not significantly altered by the SA treatment, but it is present. 1,2,3 and 4 represent the four generations.

Isoform ID	Functional gene name	—	2	ω	4	SA down- /up- regulatio n	Go category	GO term	Homologous organisms	Function	Literature
TRINITY_DN								adhesion of		Major outer membrane protein; The attachment of a symbiont to a host cell via adhesion molecules, general	Jung et al., 2016; Hoppe et al., 2017;
_i1	nella_OMP	•	-7.5		•	down	process	host cell	bacteria	indirectly.	2018.
TRINITY DN								adhesion of		Major outer membrane protein; The attachment of a symbiont to a host cell via adhesion molecules opperal	Jung et al., 2016; Hoppe et al 2017 [.]
17KINITY_DN 237954_c0_g1	PF05150.12//Legio						Biological	adhesion of symbiont to		via adhesion molecules, general stickiness etc., either directly or	al., 2017; Younas et al.,
.i1	nella OMP	x	-6.3	,	3	down	process	host cell	bacteria	indirectly.	2018.
TRINITY DN	PF00687.21^Ribos							cellular metabolic		Ribosomal protein L1 is the largest protein from the large ribosomal subunit, also can involve moleclar function of RNA binding and cellular	
2161 c0 g4 i1	omal L1	ns	ns	-5.1	4.7	down	Biological proces	s process	bacteria	component of large ribosomal subunit.	
										Involing with multi-organism process in which a virus is a participant. The other participant is the host. Includes infection	
										of a host cell, replication of the viral genome, and assembly of progeny virus particles. In some cases the viral genetic material may integrate into the host	
TRINITY_DN 116491_c0_g1	PF00729.18 ^{\V} iral						Biological	virual		genome and only subsequently, under particular circumstances, 'complete' its life cycle. Viral genome integration into	Nandhagopal
il	coat		-6.7	-	ł,	down	process Biological	process	virus	host DNA. The process by which a virion attaches	et al., 2002.
15	PF03863.13^Phag e_mat-A	1	-7.5	'	ns	down	cellular component	virion	bacteriophag e	host cell surface. The complete fully infectious extracellular virus particle.	Witherell et al., 1991.
TRINITY_DN 342313_c0_g1 i2	PF03863.13^Phag e mat-A	ns	1	ns	-7.9	down	Biological process or cellular component	virion	bacteriophag e	The process by which a virion attaches to a host cell by binding to a pilus on the host cell surface. The complete fully infectious extracellular virus particle.	Witherell et al., 1991.
TRINITY_DN 497_c0_g1_i6	PF03863.13^Phag e_mat-A	ns	ns	ns	-2.8	down	Biological process or cellular component	virion	bacteriophag e	The process by which a virion attaches to a host cell by binding to a pilus on the host cell surface. The complete fully infectious extracellular virus particle.	Witherell et al., 1991.

TRINITY_DN 4472_c0_g1_i5	TRINITY_DN 4077 cl g2 il	TRINITY_DN 1559755_c0_g 1_il	TRINITY_DN 21112_c0_g1_i 1	TRINITY_DN 5762_c0_g1_i2	TRINITY_DN 4152_c3_g1_i1	TRINITY_DN 418 c0 g1 i14	TRINITY_DN 2233_c0_g1_i1
PF00281.19^Ribo omal_L5	PF00573.22^Ribo omal L4	PF00564.24/PB1	PF03863.13^Phag	PF11645.8^PDDE XK_5	PF05736.11^OprF	PF00460.20'Flg_1 b_rod	PF03863.13^Phage e_mat-A
	- ⁵⁰	r;	•	IIS		ns	ns
ns	4.7		•	ns	ns	ns	ns
ns	IIS	-61	ns	,	43	ns	ns
-6.8	IIS	ns	-5.6	-3.7	ns	4.6	-1.5
down	down	down	down	down	down	down	down
Cellular component	Cellular component	Celhilar	Cellular component	Cellular component	Cellular component	Cellular component	Biological process or cellular component
ribosome	ribosome	membrane	membrane	integral component of membrane	cell outer membrane	bacterial-type flagellum basal body, rod	virion
	eubacteria		bacteria	bacteria	bacteria	bacteria	bacteriophag e
	This family includes Ribosomal L4/L1 from eukaryotes and archaebacteria and L4 from eubacteria. L4 from yeast has been shown to bind rRNA.	Phox and Bem1 (PB1) domains contain approximately 80 amino acids and are found in a number of cytoplasmic signaling proteins. A PB1 domain may form heterodimers with a paired PB1 domain, although not all PB1 domains will associate with one another. A highly conserved internal sequence known as OPR, PC or AID motifs is necessary for PB1 domain function. Regions outside the OPR, PC and AID help confer specificity, for binding.	Flagellans polymeruse to form bacterial flagella. This family includes flagellins and hook associated protein 3. Structurally this family forms an extended helix that interacts with PF00700.	This family of endonucleases includes a group I intron-encoded endonuclease. This family belongs to the PD (D/E)XK superfamily	This domain represents the presumed membrane-spanning region of the OprF proteins. This region is involved in channel formation and is thought to form an 8-stranded beta-barrel.	The central portion of the bacterial-type flagellar basal body, which spans the periplasm and threads through the rings.	The process by which a virion attaches to a host cell by binding to a pilus on the host cell surface. The complete fully infectious extracellular virus particle.
	Yeh and Lee, 1998.	Ito et al., 2001; Moscat et al., 2006; Sumimoto et al., 2007.	Felix et al., 1999 ; Ramos et al., 2004; Beatson et al., 2006.	Bonocora and Shub, 2001.	De Mot et al., 1994.		Witherell et al., 1991.

TRINITY_DN 332299_c0_g1 PF1 i1 xyp	TRINITY_DN PF0 1498 c0 g1 i2 ind	TRINITY_DN 24155_c0_g2_i PFc 1	TRINITY_DN 30386_c0_g2_i PF1 1	TRINITY_DN 41896_c0_g1_i PFc 1. Ho	TRINITY_DN 164781_c0_g2 PFG i2 e P
13620.6^Carbo hepD_reg	01471.18^PG_b ing_1 -	01471.18^PG_b	12081.8^GłdM	05221.17^Ado 7ase	13725.15°RNas
ns		ns		ns	118
-s	د		5	د	ы
.9 down	.2 down	.0 down	.7. down	5 down	8 down
Molecular function	Molecular function	Molecular function	Molecular	Molecular function	Molecular
carboxypepti dase activity	binding	binding	ATP binding	adenosylhom ocy steinase activity	5-3 exoribonucle ase activity
	bacetria	bacetria	bacteria		bacteria
	Putative peptidoglycan binding domain. It is found at the N or C terminus of a variety of enzymes involved in bacterial cell wall degradation.	Putative peptidoglycan binding domain. It is found at the N or C terminus of a variety of enzymes involved in bacterial cell wall degradation	This domain is found in bacteria at the N-terminus of the GldM protein. This domain is typically between 169 to 182 amino acids in length. This domain has two completely conserved residues (Y and N) that may be functionally important. GldM is named for the member from <i>Cytoplagg jolnsonae</i> (<i>Flavobacterium jolnsoniae</i>), which is required for a type of rapid gliding motility found in certain members of the <i>Bacteriodeles</i> .	AdoHcyase is an enzyme of the activated methyl cycle, responsible for the reversible hydration of S-adenosyl-L- homocysteine into adenosine and homocysteine.	ecoribonucleases. Ribonuclease PH contains a single copy of this domain, and removes nucleotide residues following the -CCA terminus of tRNA. Polyribonucleotide nucleotidyltransferase (PNPase) contains two tandem copies of the domain. PNPase is involved in nRNA degradation in a 3'-5' direction. The ecosome is a 3'-5' direction. The ecosome is a 3'-5' ecoribonuclease complex that is required for 3' processing of the 5.8S rRNA
	Krogh et al., 1998; Steen et al., 2003; Briers et al., 2009.	Krogh et al., 1998; Steen et al., 2003; Briers et al., 2009.	Braun et al., 2005.	Sganga et al., 1992	Mitchell et al.,

TRINITY_DN 6228 c0_g1_i3	TRINITY DN 1070 el al il	TRINITY_DN 2301_c0_g4_i1	TRINITY_DN 4160 c0 <u>c</u> 2 i3	TRINITY_DN 16479_c0_g1_i 8	TRINITY_DN 664942_c0_ <u>9</u> 2
PF00989.25^PAS	PF13609.6°Porin 4	PF00006.25^ATP	PF01541.24^GIY YIG	PF01541.24^GIY YIG	PF00270.29^DE4 D
1 1	5.7	· ·5.1	64	- ns -7.0	
-6.5	46	r	ns n	-7.1	r
5.9 dov	1.0 dov	doy	s dov	7.1 dov	5.4 dov
vn	SI .	M	Ϋ́Π	Ϋ́Π	1 1 1
Molecular function	Molecular function	Molecular function	Molecular function	Molecular function	Molecular
protein binding	por in activity	hydrolase activity or RNA binding	endonuclease activity	endonuclease activity	DEAD/H- box RNA helicase binding
bacteria	bacteria	bacteria, archaea eukaryotes	bacteriophag e	bacteriophag e	
L1 3are positioned at the amino terminus of signaling proteins such as sensor histidine kinases, cyclic-di-CBAP synthases and hydrolases, and methyl- accepting chemotaxis proteins.	Enables the transfer of substances, sized less than 1000 Da, from one side of a membrane to the other. The transmembrane portions of porins consist exclusively of beta-strands, which form a beta-barrel. They are found in the outer membranes of Gram- negative bacteria, mitochondria, plastids and possibly acid-fast Gram-positive bacteria	Catalysis of the hydrolysis of various bonds, e.g. C-Q, C-N, C-C, phosphoric anhydride bonds, etc. Hydrolase is the systematic name for any enzyme of EC class 3 or Interacting selectively and non-covalently with an RNA molecule or a portion thereof.	It is involved in degradation of host DNA, permitting scavenging of host- derived nucleotides for phage DNA synthesis; in enzymes involved in DNA repair and recombination.	It is involved in degradation of host DNA, permitting scavenging of host- derived nucleotides for phage DNA synthesis; in enzymes involved in DNA repair and recombination.	The DEAD/DEAH lox helicases are a family of proteins whose purpose is to unwind nucleic acids. The DEAD box helicases are involved in various aspects of RNA metabolism, including nuclear transcription, pre nRNA splicing, ribosome biogenesis, nucleocytoplasmic transport, translation, RNA decay and organellar gene expression.
Henry and Crosson, 2011; Liu et al., 2015.	Saier, 2000	Wilkens et al., 2005.	Sharma et al., 1992; Dunin- Horkawicz et al., 2006; Andersson et al., 2010.	Sharma et al., 1992; Dunin- Horkawicz et al., 2006; Andersson et al., 2010	Johnson and McKay, 1999

TRINITY_DN 243032_c0_g1 i2	TRINITY_DN 19010_c0_g4_i 3	TRINITY_DN 34032_c0_g1_i 1	TRINITY_DN 5435_c0_g1_i4	TRINITY_DN 109372_c0_g1 i2	TRINITY_DN 264317_c0_g1 i1	TRINITY_DN 20104_c0_g1_i 6	TRINITY_DN 135_c0_g1_i7
PF05919.11^Mito vir RNA. pol	PF00978.21^RdRP	PF05919.11^Mito vir RNA_pol	PF05919.11^Mito vir RNA. pol	PF05919.11^Mito vir RNA pol	PF03431.13^RNA .replicase B	PF05919.11^Mito vir RNA pol	PF13360.6^PQQ_2
1	ns		'	ns	•	ns	ns
'	-6.3	ns	-5.0	-5.5	ns	-7.0	ns
		-7.2		IIS	-8.0	ns	ns
-6.3		-6.6	-7.3	-7.2	ns	-6.8	-2.5
down	down	down	down	down	down	down	down
Biological process	Molecular	Molecular function	Molecular function	Molecular function	Molecular function	Molecular function	Molecular function
RNA- directed 5'-3' RNA polymerase activity	viral replication or RNA- directed RNA polymerase activity	RNA- directed 5'-3' RNA polymerase activity	RNA- directed 5'-3' RNA polymerase activity	RNA- directed 5'-3' RNA polymerase activity	RNA- directed 5'-3' RNA polymerase activity	RNA- directed 5'-3' RNA polymerase activity	pyrroloquinol ine quinone binding
virus	vinus	virus	virus	vins	virus	vinus	
Mitovinus RNA-dependent RNA polymerase proteins. The family also contains fragment matches in the mitochondria of Arabidopsis theliana.	KNA-dependent KNA polymerase or RNA replicase is an enzyme that catalyzes the replication of RNA from an RNA template. Specifically, it catalyses synthesis of the RNA strand complementary to a given RNA template.	Mitovirus RNA-dependent RNA polymerase proteins. The family also contains fragment matches in the mitochondria of <i>Arabidopsis theliana</i> .	Mitovinus RNA-dependent RNA polymerase proteins. The family also contains fragment matches in the mitochondria of <i>Arabidopsis theliana</i> .	Mitovinus RNA-dependent RNA polymerase proteins. The family also contains fragment matches in the mitochondria of <i>Arabidopsis thaliana</i> .	This family is of Leviviridae RNA replicases. The replicase is also known as RNA dependent RNA polymerase.	Mitovinus RNA-dependent RNA polymerase proteins. The family also contains fragment matches in the mitochendria of Arabidopsis theliana.	Interacting selectively and non- covalently with pyrroloquinoline quinone, PQQ, the coenzyme or the prosthetic group of certain alcohol deliydrogenases and ghucose deliydrogenases.
Hong et al., 1998; Yao et al., 2020	Hardy et al., 1979; Boonrod et al., 2004; Schwan et al., 2005.	Hong et al., 1998; Yao et al., 2020.	Hong et al., 1998; Yao et al., 2020	Hong et al., 1998; Yao et al., 2020.	Fernando et al., 2020.	Hong et al., 1998; Yao et al., 2020	

1	endonucleases.		hydrolysis	process	чр	5.9	,	,	•	73n	3358 c0 g1 i2
on of a number of	zinc-binding loops regi much longer chain hon		pnospnotiest er bond	Biological					-1	PF05551.11^F25:F	TRINITY_DN
on of a number of ing main is the short	zinc-binding loops regi much longer chain hon endonucleases. This do		nucleic acid								
n of homing ain is the short	Zinc-binding loop region endonuclease. This dom										
	endonucleases.		hydrolysis	process	цр	5.8	•	ns		His Me endon	_i3
ŭ	much longer chain homi		er bond	Biological						PF05551.11^zf-	473649_c0_g1
1 of a number of	zinc-binding loops region		phosphodiest								TRINITY DN
of homing in is the short	Zinc-binding loop region endonuclease. This doma		mucleic acid								
domains.	of one or more cold shock		translation,	process	цр	2	ns	4.4	ns	shock protein	1057 c0 g1 i4
the presence	proteins, characterized by		transcription,	Biological						COG1278/Cold	TRINITY_DN
DNA binding	are multifunctional RNA/		regulation of								
nock proteins	for that organism Cold sh		like								
al temperature	stimulus below the optim		DLOCGESSES								
on, cu. j as a	regult of a cold stimulus :		hiological								
yme	movement, secretion, enz		Manu								
erms of	a cell or an organism (in t										
e or activity of	results in a change in state										
ocess that	Response to stress, any pi										
domains. I	of one or more cold shock		translation,	process	ų	ns	ns	3.8	ns	shock protein	306 c0 g2 i5
the presence	proteins, characterized by		transcription,	Biological						COG1278/Cold	TRINITY_DN
DNA binding	are multifunctional RNA/		regulation of								
nock proteins	for that organism. Cold sh		like								
a temperature	stimulus below the ontim		DIOLOGICAL								
on, etc.) as a	production, gene expressi		Many								
yme	movement, secretion, enz										
erms of	a cell or an organism (in t										
te or activity of	results in a change in stat										
rocess that	Response to stress, any p										
	PF00700.	bacteria	membrane	component	down	-1.7	ns	ns	ns	llin N	2369 c0 g1 i4
cts with	extended helix that intera			Cellular						PF00669.20/Flage	TRINITY DN
orms an	Structurally this family for										
n 3. Ű	and hook associated protei										
ides flagellins	flagella. This family inch										
	1 1. 1										

1994.	ribosome.		ribosome	function	dn	•	5 INS	.4	S3 C -	omal	924 c0 $\overline{g_3}$ il
Drevfuss	to the structural integrity of the	-f	constituent o	molecular					89 20'R ihos	PF001	TRINITY DN
Burd and	The action of a molecule that contributes		structural	process or							
	polypeptide chain from the ribosome or		process or	Biological							
	Translation ends with the release of a		metabolic								
	ribosome and an mRNA or circRNA.		cellular								
	associates with the small subunit of the										
	initiation factor 2, which subsequently										
	initiator methionine tRNA, GTP, and										
	ternary complex between aminoacylated										
	and begins with the formation of a										
	Translation is mediated by the ribosome,										
	amino acids in a polypeptide chain.										
	molecule to specify the sequence of										
	of a mature mixing or circking										
	a protein is formed, using the sequence										
	a protein is formed using the sequence										
1001.	The cellular metabolic process in which	č	VII OI	component	up		110			C 1100	0022 00 52 0
1001	infactions extracellular virus narticle	0		commonent	I	د د د	ne	10	0	e mat	1 00 00 00
Witherell et al.	host cell surface. Or The complete fully	bacteriophag		cellular					863.13 [^] Phag	PF038	TRINITY DN
	to a host cell by binding to a pilus on the			process or							
	The process by which a virion attaches			Biological							
1775.	COLUMN		LI GIISGICLIOII	TIOCOS	цр	0.7	115		-	OTISC 1	10
1005	domain		transduction	Drocess		л 2	ne				
Pao and Saier.	N-terminal to a DNA binding effector		signal	Biological)72.24^Resp	PF000	220624 c0 gl
	component systems. It is usually found										TRINITY DN
	sensor partner in bacterial two-										
	This domain receives the signal from the										
	Kesponse regulator receiver domain.										
2000.		Dacteria	proteorysis	process	цр	4.0	115	5 115	11	00	1
2000 ct ut.,	TTITOD like bets bened dennein			The Store		2			100.0 1 8U_1	11 100	
Kiin et al	shows that it is inserted within a			Biological					RU UVEIOD !	PF138	1442 c0 ol il
	protein. THe structure for this domain										TRINITY DN
	FlgD protein the flagellar hook capping										
	beta sandwich fold. It is found in the										
	This domain has an immunoglobulin like										
1999.	membrane	bacteria	transport	process	цр	3.3	ns	ns	-	ExbB	il
Braun et al.,	large molecules across the outer bacterial		protein	Biological					518.16 MotA	PF016	422335_c0_g1
	the inner bacterial membrane to transport										TRINITY_DN
	complex uses the proton gradient across										
	transduction complex. The Tomb										
	is part of the round-upperformed										
	is nort of the TonD demondent										
	notational motion in the flaveller ExhB										
	uses a proton gradient to generate										
	component of the flageller motor that										
	proton channels. MotA is an essential										
	a membrane. These proteins are probably										
	involved translocation of proteins across										
	membrane proteins that appear to be										
	This family groups together integral										

	polypeptide.		ribosome	component	dn c	ns 4	ns	'	omal L10	468/ c0 g1 12
	terminal or penultimate peptide bond at the C-terminal end of a peptide or			Cellular	,				PF00466.20'Ribos	TRINITY_DN
	Catalysis of the hydrolysis of the				T.					
	nolvnentide		ribosome	component	9 Im	ns 2	ns		mal L12 N	653 c7 g1 i1
	terminal or penultimate peptide bond at the C-terminal end of a peptide or			Cellular					PF16320.5^Riboso	TRINITY DN
	Catalysis of the hydrolysis of the									
	other proteins involved in secretion.	prokaryotes	of membrane	component	dn s	3.9 n		,	ethyl	1598_c0_g1_i1
	found at the N-terminus of pilins and		component	Cellular					PF07963.12/N m	TRINITY DN
	phenylalanine residue. It is most often		integral							
	methylation of the conserved		membrane or							
	Prokaryotic N-terminal methylation									
2018.	sequences.	bacteria	membrane	component	.6 up	ns 5	'	a.	126	3530 c0 g1 i1
Price et al.,	domain is always found as pair in such			Cellular					PF03458.13//UPF0	TRINITY_DN
	medicted to be obvine transportant. This									
	in obvine utilization and thus are									
	have since heen shown to be immortant									
	channel. Proteins containing this domain									
	glycine, which are suggestive of an ion									
	transmembrane helices, with conserved									
	This domain contains three									
Antoine et al., 2003.	group of families, distinct from the ABC and TRAP-T families	bacteria	membrane	Cellular component	.2 up	- 2		•	PF03401.14 [^] TctC	TRINITY_DN 429 c0 g2 i2
	binding receptor-dependent transporter									
	receptors of the extracytoplasmic solute									
	of bacterial tripartite tricarboxy late									
	IOTHERIY KNOWN as Bug - BOTUERENA									
	formerly because and the formerly because the formerly because and the former of the formerly because the formerly									
	several heta-protechacteria This family									
	These probable extra-cytoplasmic solute									
2006.	PF00700.	bacteria	membrane	component	dn s	ns n	3.9	ns	llin N	-i3
Beatson et al.,	extended helix that interacts with			Cellular					PF00669.20/Flage	297806 c0 g2
et al., 2004;	Structurally this family forms an									TRINITY DN
1999; Ramos	and hook associated protein 3.									
Felix et al.,	flagella. This family includes flagellins									
	Flagellins polymerise to form bacterial									
Salyers, 2001.	PF07715 and PF00593.	eukaryotes	of membrane	component	.7 up	ns 2	ns	•	pepD_reg_2	520_c0_g2_i1
Cho and	The family is found in association with	archaea or	component	Cellular					PF13715.6//Carbo	TRINITY DN
	approximately 90 amino acids in length.	bacteria,	integral							
	archaea and eukaryotes, and is									
	This domain family is found in bacteria,									
	it to associate with the outer membrane.		membrane	component	.4 up	ns 2	'	•	like_2	5169 c0 g1 i4
	with an N-terminal lipid tail that allows		cell outer	Cellular					PF12771.7^SusD-	TRINITY_DN
	SusD is a secreted starch-binding protein									
TRINITY_DN 1632 c0 gl il	TRINITY_DN 193314_c0_g1 i1	TRINITY_DN 6187 c0 g1 i1	TRINITY_DN 12161_c0_g4_i 3	TRINITY_DN 35 c0 g1 i1	TRINITY_DN 64210_c0_g2_i 1	TRINITY_DN 3461_c0_g1_i1	TRINITY_DN 3461_c0_g3_i1	TRINITY_DN 3585 c0 g1 i4	TRINITY_DN 2568 c0 gl il	
--	---	--	---	--	--	--	--	--	---	
PF13328.6^HD 4	PF00557.24^Pepti dase M24	COG0302^GTP cyclohydrolase i	PF01 193.24^RNA 	PF01471.18 [^] PG_b inding_1	PF01471.18^PG_b inding_1	PF01471.18 [^] PG_b inding_1	PF01471.18 ^{PG_b}	COG0402^deamin ase	PF00466.20'Ribos omal_L10	
•		E.	10		L.					
ns		•		IIS	•	7.2	3.5	1	•	
ns	ř.	ns		ns	6.2	6.6	•	IIS	•	
2.2	5.7	3.4	5.4	5.7	•	ns	5.7	3.8	5.4	
ų	Ð	Ð	8	Ð	ŧ	Ð	Ð	Ð	Ð	
Molecular function	Molecular function	Molecular function	Molecular function	Molecular function	Molecular function	Molecular function	Molecular function	Molecular function	Cellular component	
hydrolase activity	lydrolase activity	GTP cyclohydrola se I activity	DNA- directed 5'-3' RNA polymerase activity or protein dimerization activity	binding	binding	binding	binding	deaminase ac tivity	ribosome	
	bacteria	bacteria		bacetria	bacetria	bacetria	bacetria			
HD domains are metal dependent phospholydrolases.	This family contains metallopeptidases. It also contains non-peptidase homologues such as the N terminal domain of Spt16, which is a histone H3- H4 binding module.	This family includes GTP cyclohydrolase enzymes and a family of related bacterial proteins	The prokaryotic equivalent of the Rpb3/Rpb11 platform is the alpha-alpha dinner. The dimerisation domain of the alpha subunit/Rpb3 is interrupted by an insert domain (PF01000). Some of the alpha subunits also contain iron-sulphur binding domains	Putative peptidoglycan binding domain. It is found at the N or C terminus of a variety of enzymes involved in bacterial cell wall degradation	Putative peptidoglycan binding domain. It is found at the N or C terminus of a variety of enzymes involved in bacterial cell wall degradation	Putative peptidoglycan binding domain. It is found at the N or C terminus of a variety of enzymes involved in bacterial cell wall degradation	Putative peptidoglycan binding domain. It is found at the N or C terminus of a variety of enzymes involved in bacterial cell wall degradation	Catalysis of the removal of an amino group from a substrate, producing ammonia	Catalysis of the hydrolysis of the terminal or penultimate peptide bond at the C-terminal end of a peptide or polypeptide.	
Aravind and Koonin, 1998.	Stuwe et al., 2008.	Nar et al., 1995.	Zhang and Darst, 1998	Krogh et al., 1998; Steen et al., 2003; Briers et al., 2009.	Krogh et al., 1998; Steen et al., 2003; Briers et al., 2009.	Krogh et al., 1998; Steen et al., 2003; Briers et al., 2009.	Krogh et al., 1998; Steen et al., 2003; Briers et al., 2009.			

	Domain of unknown function				dn	3.4	ns	ns	OUF	PF01595.20^I 21	TRINITY_DN 8253_c0_g1_i9
	Domain of unknown function				ų	3.3	ns	•	UF2 -	PF09984.9^DI 222	TRINITY_DN 17868_c1_g1_i 9
Convey et al., 1998.	The Leivivins coat protein forms the bacteriophage coat that encapsidates the viral RNA. 180 copies of this protein form the virion shell. The MS2 bacteriophage coat protein controls two distinct processes: sequence-specific RNA encapsidation and repression of replicase translation-by binding to an RNA stem-loop structure of 19 nucleotides containing the initiation codon of the replicase gene. The binding of a coat protein dimer to this hairpin shuts off synthesis of the viral replicase, switching the viral replication cycle to virion assembly rather than continued replication	bacteriophag	viral capsid	Molecular function	Ð	3 1	20	su	evi_	PF01819.17/T coat	TRINITY DN 1761 c0 g2 il
Fernando et al., 2020.	This family is of Leviviridae RNA replicases. The replicase is also known as RNA dependent RNA polymerase.	VINS	RNA- directed 5'-3' RNA polymerase activity	Molecular function	Ð	5	8.2	ns	RNA -	PF03431.13^F replicase B	TRINITY_DN 1905 c1 g2 i1
Wilkens et al., 2005.	Catalysis of the hydrolysis of various bonds, e.g. C-O, C-N, C-C, phosphoric anhydride bonds, etc. Hydrolase is the systematic name for any enzyme of EC class 3 or Interacting selectively and non-covalently with an RNA molecule.	bacteria, archaea eukaryotes	hydrolase activity or RNA binding	Molecular function	Æ	ns	3.9	ns	ATP-	PF02874.23^A synt_ab	TRINITY_DN 9091_c0_g1_i1
Anantharaman and Aravind, 2003.	The function of this domain is unknown. It is found in several lipoproteins.		hydrolase activity	Molecular function	ųp	3.6	ns		. Ч	COG0791^NL P60 protein	TRINITY_DN 2812 c0 g1 i4

Table S2 The definition of GO terms in Fig. 7.

		SA		
GO ID	GO term	regulation	Catergory	Definition
				Interaction between organisms physiological interaction
	multi-organism		Biological	between organisms physiological interaction with another
GO:0051704	cellular process	up	process	organism
				A process carried out by gene products in an organism that
				enable the organism to engage in a symbiotic relationship, a
			Biological	more or less intimate association, with another organism.
GO:0044403	symbiotic process	up	process	Microscopic symbionts are often referred to as endosymbionts.
	multi-organism	ŧ	Biological	Any process that is carried out at the cellular level, which
GO:0044764	cellular process	up	process	involves another organism of the same or different species.
	viral genome		Biological	Any process involved directly in viral genome replication.
GO:0019079	replication	110	process	including viral nucleotide metabolism.
	viral RNA genome		Biological	
GO:0039694	replication	110	process	The replication of a viral RNA genome.
			Biological	The cellular metabolic process in which a cell duplicates one or
GO:0039703	RNA replication	110	process	more molecules of RNA
00.00007700			Process	A multi-organism process in which a virus is a participant. The
				other participant is the host Includes infection of a host cell
				replication of the viral genome and assembly of progeny virus
				narticles. In some cases, the viral genetic material may
			Biological	integrate into the bost genome and only subsequently under
GO:0016032	wiral process	100	process	naticular circumstances 'complete' its life cycle
00.0010032	viral process	цр	process	Lateral transfer of an intron to a homologous allele that lacks
				the intron mediated by a gite medific endenuelesse encoded
			Dialogical	within the mehile intern. It investigates with collular
CO.0006214	intern banning		Biological	walling the moone multiplice and multiplice and metabolice process
60:0006514	muon nonning	up	process	A match alia ana ana ana fuciele acid metabolic process
				A metabolic process - chemical reactions and pailways,
	1.1		Distantiant	line anatomisma transformation of a state transformation
CO.0044022	muid-organism		Biological	inving organisms transform chemical substances, which
GO:0044033	metabolic process	up	process	involves more than one organism.
	RNA-protein covalent		Biological	The formation of a covalent cross-link between RNA and a
GO:0018144	cross-linking	up	process	protein. It involves in cellular protein modification.
				The process by which a virion attaches to a host cell by binding
				to a pilus on the host cell surface. Pili are retractile filaments
	N 6 705 6 875			that protrude from gram-negative bacteria. Filamentous viruses
~~~~~	virion attachment to		Biological	can attach to the pilus tip, whereas icosahedral viruses can
GO:0039666	host cell pilus	up	process	attach to the pilus side.
	a ni 200 ni		Cellular	Any constituent part of a host cell. The host is defined as the
GO:0033643	host cell part	up	Component	larger of the organisms involved in a symbiotic interaction.
			Cellular	Any constituent part of a secondary organism with which the
GO:0044217	other organism part	up	Component	first organism is interacting.
				Any constituent part of the living contents of a host cell; the
				matter contained within (but not including) the plasma
				membrane, usually taken to exclude large vacuoles and masses
				of secretory or ingested material. In eukaryotes it includes the
			Cellular	nucleus and cytoplasm. The host is defined as the larger of the
GO:0033646	host intracellular part	up	Component	organisms involved in a symbiotic interaction.
				Any constituent part of the host cell cytoplasm, all of the
				contents of a cell excluding the plasma membrane and nucleus,
				but including other subcellular structures. The host is defined
	host cell cytoplasm		Cellular	as the larger of the organisms involved in a symbiotic
GO:0033655	part	up	Component	interaction.
			Cellular	Any constituent part of a virion, a complete fully infectious
GO:0044423	virion part	up	Component	extracellular virus particle.
				The protein coat that surrounds the infective nucleic acid in
			Cellular	some virus particles. It comprises numerous regularly arranged
GO:0019028	viral capsid	up	Component	subunits, or capsomeres.
				Organized structure of distinctive morphology and function.
				occurring within the host cell. Includes the nucleus.
	host intracellular		Cellular	mitochondria, plastids, vacuoles, vesicles, ribosomes and the
GO:0033647	organelle	110	Component	cytoskeleton. Excludes the plasma membrane. The host is

				1.0 1.4.1 04
				defined as the larger of the organisms involved in a symbiotic
				Organized structure of distinctive mombology and function as
				found in host cells, bounded by a single or double lipid bilayer
				membrane and occurring within the cell. Includes the nucleus,
	host intracellular			mitochondria, plastids, vacuoles, and vesicles. Excludes the
00 0033640	membrane-bounded		Cellular	plasma membrane. The host is defined as the larger of the
GO:0033648	organelle	up	Component	organisms involved in a symbiotic interaction.
				cells that occur in varying numbers shapes and sizes in the
	host cell		Cellular	cell cytoplasm. The host is defined as the larger of the
GO:0033650	mitochondrion	up	Component	organisms involved in a symbiotic interaction.
				The protein coat that surrounds the infective nucleic acid in
				some virus particles; the subunits are arranged to form a
CO:0010020	balical rinal consid		Cellular	protein helix with the genetic material contained within.
GO:0019029	nencai virai capsid	up	Component	The protein coat that surrounds the infective nucleic acid in
				some virus particles: the subunits are arranged to form an
				icosahedron, a solid with 20 faces and 12 vertices. Icosahedral
				capsids have 12 pentamers plus 10(T-1) hexamers, where T is
	ic osahedral viral		Molecular	the triangulation number. Tobacco satellite necrosis virus has
GO:0019030	capsid	up	function	such a capsid structure.
				The protein coat that surrounds the infective nucleic acid in some virus particles where the submits (consomerse) and
				arranged to form an icosabedron with T=3 symmetry. The T=3
	T=3 icosahedral viral		Molecular	capsid is composed of 12 pentameric and 20 hexameric
GO:0039617	capsid	up	function	capsomeres.
				Catalysis of the reaction: nucleoside triphosphate + RNA (n) =
			21212 2	diphosphate + RNA (n+1); the synthesis of RNA from
00 002 40 (2	5'-3' RNA polymerase		Molecular	ribonucleotide triphosphates in the presence of a nucleic acid
GO:0034062	activity	up	nunction	Catalysis of the reaction: nucleoside trinhosphate + $RNA(n) =$
				diphosphate + RNA $(n+1)$ : the synthesis of RNA from
	RNA polymerase		Molecular	ribonucleotide triphosphates in the presence of a nucleic acid
GO:0097747	activity	up	function	temp late.
	nucleotidyltransferase		Molecular	Catalysis of the transfer of a nucleotidyl group to a reactant.
GO:0016779	activity	up	function	The upper group belongs to trasnfeerase activety.
	PNA directed 5' 2'			Catalysis of the reaction: nucleoside triphosphate + $RNA(n) =$
	RNA nolymerase		Molecular	catalysis of RNA-template-directed extension of the 3'-end of
GO:0003968	activity	up	function	an RNA strand by one nucleotide at a time.
			Molecular	Interacting selectively and non-covalently with any component
GO:0001848	complement binding	up	function	or product of the complement cascade.
				Interacting selectively and non-covalently with an opsonin,
			Malandar	such as a complement component or antibody, deposited on the
GO:0001846	onsonin hinding	110	function	surface of a bacteria, virus, infinitine complex, or other
50.001640	complement		menon	
	component C3b		Molecular	Interacting selectively and non-covalently with the C3b product
GO:0001851	binding	up	function	of the complement cascade.
				Any process, in which a cell, a substance, or a cellular entity,
				such as a protein complex or organelle, is transported, tethered
			Biological	to or outerwise maintained in a specific location. In the case of substances, localization may also be achieved via selective
GO:0051179	localization	down	Drocess	degradation
30.0001119				Any process that localizes a substance or cellular component.
	establishment of		Biological	This may occur via movement, tethering or selective
GO:0051234	localization	down	process	degradation.
				The directed movement of substances (such as
				macromolecules, small molecules, ions) or cellular components
			Biological	or between cells, or within a multicellular organism by means
GO:0006810	transport	down	process	of some agent such as a transporter, pore or motor protein.
				The chemical reactions and pathways involving small
	small molecule		Biological	molecules, any low molecular weight, monomeric, non-
GO:0044281	metabolic process	down	process	encoded molecule.

	organic cyclic			
00 100120	compound		Biological	The chemical reactions and pathways resulting in the formation
GO:1901362	biosynthetic process	down	process	of organic cyclic compound.
	1.02.00000010		Distantion	The chemical reactions and pathways resulting in the formation
CO.0010120	heterocycle	- dimension	Biological	of heterocyclic compounds, those with a cyclic molecular
GO:0018130	biosynthetic process	down	process	structure and at least two different atoms in the ring (or rings).
				The chemical reactions and pathways resulting in the formation
			Dislogical	of substances, typically, the energy-requiring part of
CO.000058	hi agenthatia neasaga	damm	Biological	metabolism in which simpler substances are transformed into
00.0009038	biosynatetic process	uown	process	The chemical reactions and nathurwa resulting in the formation
	organic substance		Biological	of an organic substance, any molecular entity containing
GO-1901576	biosynthetic process	down	process	carbon
00.1901970	cellular biosynthetic	down	Biological	The chemical reactions and nathways resulting in the formation
GO:0044249	process	down	process	of substances, carried out by individual cells
	organonitrogen			
	compound metabolic		Biological	The chemical reactions and pathways involving organonitrogen
GO:1901564	process	down	process	compound.
••••••	organonitrogen			
	compound		Biological	The chemical reactions and pathways resulting in the formation
GO:1901566	biosynthetic process	down	process	of organonitrogen compound.
	organic substance		Biological	The chemical reactions and pathways involving an organic
GO:0071704	metabolic process	down	process	substance, any molecular entity containing carbon.
	cellular metabolic		Biological	The chemical reactions and pathways by which individual cells
GO:0044237	process	down	process	transform chemical substances.
	cellular nitrogen			
	compound		Biological	The chemical reactions and pathways resulting in the formation
GO:0044271	biosynthetic process	down	process	of organic and inorganic nitrogenous compounds.
	carbohydrate			
~~	derivative metabolic		Biological	The chemical reactions and pathways involving carbohydrate
GO:1901135	process	down	process	derivative.
00 0010530	protein metabolic	10. • 10.0 IV/20	Biological	The chemical reactions and pathways involving a protein.
GO:0019538	process	down	process	Includes protein modification.
				The chemical reactions and pathways involving those
	primary metabolia		Dialogical	compounds, which are formed as a part of the normal anabolic and establic processes. These processes take place in post if
GO-0044238	printary metabolic	down	biological	and catabolic processes. These processes take place in most, if
00.0044238	nitrogen compound	down	Biological	The chemical reactions and nathways involving organic or
GO:0006807	metabolic process	down	process	in organic compounds that contain nitrogen
00.0000007	metabolic process	down	process	The chemical reactions and nathways resulting in the formation
				of a macromolecule, any molecule of high relative molecular
				mass, the structure of which essentially comprises the multiple
	macromolecule		Biological	repetitions of units derived, actually or conceptually, from
GO:0009059	biosynthetic process	down	process	molecules of low relative molecular mass.
				The chemical reactions and pathways resulting in the formation
				of a macromolecule, any molecule of high relative molecular
				mass, the structure of which essentially comprises the multiple
	cellular			repetition of units derived, actually or conceptually, from
	macromolecule		Biological	molecules of low relative molecular mass, carried out by
GO:0034645	biosynthetic process	down	process	individual cells.
			100 N N N	The chemical reactions and pathways resulting in the formation
	aromatic compound		Biological	of aromatic compounds, any substance containing an aromatic
GO:0019438	biosynthetic process	down	process	carbon ring.
				Metabolic process resulting in cell growth   metabolism
00.00001 55			Biological	metabolism resulting in cell growth   multicellular organism
GO:0008152	metabolic process	down	process	metabolic process   single-organism metabolic process
	organic cyclic		Distant	The sharping postions and path-
CO-10012 CO	compound metabolic	darm	Biological	The chemical reactions and pathways involving organic cyclic
00:1901300	process	uown	process	The chemical reactions and nathering involving assessti-
				compounds any organic compound characterized by one or
	cellular aromatic			more planar rings, each of which contains conjugated double
	compound metabolic		Biological	bonds and delocalized pi electrons as carried out by individual
GO:0006725	process	down	process	cells.

	aallular nitraa oo			The chemical constiant and nathrana involving various accessio
	compound metabolic		Biological	and inorganic nitrogenous compounds, as carried out by
GO:0034641	process	down	process	individual cells
00.0051011	process	down	process	The chemical reactions and pathways involving heterocyclic
	heterocy cle metabolic		Biological	compounds, those with a cyclic molecular structure and at least
GO:0046483	process	down	process	two different atoms in the ring (or rings).
				The chemical reactions and pathways involving
				macromolecules, any molecule of high relative molecular mass,
				the structure of which essentially comprises the multiple
	macromolecule		Biological	repetitions of units derived, actually or conceptually, from
GO:0043170	metabolic process	down	process	molecules of low relative molecular mass.
				The chemical reactions and pathways resulting in the
			Distant	break down of substances, including the break down of carbon
00.000056		4	Biological	compounds with the liberation of energy for use by the cell or
60.0009030	catabolic process	down	process	The chemical reactions and nathuraus involving
				macropolecules, any polecule of high relative polecular mass
				the structure of which essentially comprises the multiple
	cellular			repetition of units derived actually or conceptually from
	macromolecule		Biological	molecules of low relative molecular mass, as carried out by
GO:0044260	metabolic process	down	process	individual cells.
	·····		Biological	Any process that modulates a measurable attribute of any
GO:0065007	biological regulation	down	process	biological process, quality or function.
	nucleobase-containing			
	compound metabolic		Biological	Any cellular metabolic process involving nucleobases,
GO:0006139	process	down	process	nucleosides, nucleotides and nucleic acids.
				Any constituent part of the living contents of a host cell; the
				matter contained within (but not including) the plasma
				membrane, usually taken to exclude large vacuoles and masses
			~ !! !	of secretory or ingested material. In eukaryotes it includes the
00.0022646	· · · · · · · · · · · · · · · · · · ·	4	Cellular	nucleus and cytoplasm. The host is defined as the larger of the
GO:0033646	intracentilar part	down	Component	organisms involved in a symplotic interaction.
CO:0016020	momhron o	down	Cenular	A lipid bilayer along with all the proteins and protein
60.0010020	memorane	down	Component	All of the contents of a cell excluding the plasma membrane
GO:0005737	cytonlasm	down	Component	and nucleus but including other subcellular structures
00.0003737	cytopiasii	down	Component	The membrane surrounding a cell that separates the cell from
			Cellular	its external environment. It consists of a phospholipid bilayer
GO:0005886	plasma membrane	down	Component	and associated proteins.
	ribonucleoprotein		Cellular	A macromolecular complex containing both protein and RNA
GO:1990904	complex	down	Component	molecules.
				A stable assembly of two or more macromolecules, i.e.,
				proteins, nucleic acids, carbohydrates or lipids, in which at
	protein-containing	24	Cellular	least one component is a protein and the constituent parts
GO:0032991	complex	down	Component	function together.
				The component of a membrane consisting of the gene products
				having some covalently attached portion, for example part of a
			Cellular	peptide sequence or some other covalently attached group such
CO.0021224	intrinsic component	damm	Centuar	as a GP1 anchor, which spans or is embedded in one or boun
GO:0051224	of memorane	down	Component	The component of a membrane consisting of the gene products
				and protein complexes having at least some part of their
	integral component of		Celhilar	peptide sequence embedded in the hydrophobic region of the
GO:0016021	membrane	down	Component	membrane.
				Organized structure of distinctive morphology and function.
	non-membrane-		Cellular	not bounded by a lipid bilayer membrane. Includes ribosomes,
GO:0043228	bounded organelle	down	Component	the cytoskeleton and chromosomes.
			Cellular	All of the contents of a cell excluding the plasma membrane
GO:0005737	cytoplasmic part	down	Component	and nucleus, but including other subcellular structures.
				Organized structure of distinctive morphology and function,
	non-membrane-		Cellular	not bounded by a lipid bilayer membrane. Includes ribosomes,
GO:0043228	bounded organelle	down	Component	the cytoskeleton and chromosomes.
			<u></u>	An intracellular organelle, about 200 A in diameter, consisting
00 00000			Cellular	of RNA and protein. It is the site of protein biosynthesis
GO:0005840	ribosome	aown	Component	resulting from translation of messenger RNA (mRNA). It

				consists of two subunits, one large and one small, each containing only protein and RNA.
				Organized structure of distinctive morphology and function.
				Includes the nucleus, mitochondria, plastids, vacuoles, vesicles,
				ribosomes and the cytoskeleton, and prokaryotic structures
			Cellular	such as anammoxosomes and pirellulosomes. Excludes the
GO:0043226	organelle	down	Component	plasma membrane.
			••••••	Organized structure of distinctive morphology and function,
				occurring within the cell. Includes the nucleus, mitochondria,
			Cellular	plastids, vacuoles, vesicles, ribosomes and the cytoskeleton.
GO:0043229	intracellular organelle	down	Component	Excludes the plasma membrane.
			Molecular	Interacting selectively and non-covalently with anions, charged
GO:0043168	anion binding	down	function	atoms or groups of atoms with a net negative charge.
			Molecular	Interacting selectively and non-covalently with ions, charged
GO:0043167	ion binding	down	function	atoms or groups of atoms.
	carbohydrate		Molecular	Interacting selectively and non-covalently with a carbohydrate
GO:0097367	derivative binding	down	function	derivative.
				Interacting selectively and non-covalently with a
				ribonucleotide, any compound consisting of a ribonucleoside
	ribonucleotide		Molecular	that is esterified with (ortho) phosphate or an oligophosphate at
GO:0032553	binding	down	function	any hydroxyl group on the ribose moiety.
				Interacting selectively and non-covalently with a purine
				ribonucleotide, any compound consisting of a purine
	purine ribonucleotide		Molecular	ribonucleoside that is esterified with (ortho) phosphate or an
GO:0032555	binding	down	function	oligophosphate at any hydroxyl group on the ribose moiety.
				Interacting selectively and non-covalently with a pyrimidine
	pyrimidine		200200	ribonucleotide, any compound consisting of a pyrimidine
	ribonucleotide	10. · · · · · · · · · · · · · · · · · · ·	Molecular	ribonucleoside that is esterified with (ortho) phosphate or an
GO:0032557	binding	down	function	oligophosphate at any hydroxyl group on the ribose moiety.
				Interacting selectively and non-covalently with a purine
				ribonucleoside triphosphate, a compound consisting of
~~ ~~ ~~ ~	purine ribonucleoside	20 <b>•</b> 100000000	Molecular	a purine base linked to a ribose sugar esterified
GO:0035639	triphosphate binding	down	function	with triphosphate on the sugar.
				Any molecular function by which a gene product interacts
00 00000	DATA 1 ' I'		Molecular	selectively and non-covalently with DNA (deoxyribonucleic
GO:00036//	DNA binding	down	nunction	acid).
			Malandar	Catalysis of the hydrolysis of various bonds, e.g., C-O, C-N, C-
CO.0016797	has donal and a set in site.		forecular	C, phosphoric annydride bonds, etc. Hydrolase is the
GO:0010/8/	ny drotase activity	down	nunction	systematic name for any enzyme of EC class 5.
CO:0016919	pnospnorus-	dorra	function	Catalysis of the hydrolysis of any acid annyaride, which
60.0010818	containing annyurides	uown	minemon	Catalway of the hydrohysis of a symphoce has deal between
	average based atops		Malaanlar	two phoenbate groups, logving and phoenbate on each of the
CO:0016462	pyrophosphatase	down	function	two phosphate groups, leaving one phosphate on each of the
00.0010402	mucleogide	uown	Italiction	two inaginetits.
	triphognhatage		Molecular	Catalysis of the reaction: a nucleoside triphosphate $\pm$ H2O =
GO:0017111	activity	down	function	mucleoside dinhombate + phombate
00.001/111	activity	uown	Molecular	The selective non covalent often stoichiometric interaction of
GO:0005488	hinding	down	function	a molecule with one or more specific sites on another molecule
00.0000400	omang	down	miletion	Interacting selectively and non-covalently with an organic
	organic cyclic		Molecular	cyclic compound any molecular entity that contains carbon
GO:0097159	compound hinding	down	function	arranged in a cyclic molecular structure
00.0077107	heterocyclic	donn	Molecular	Interacting selectively and non-covalently with heterocyclic
GO:1901363	compound binding	down	function	compound
	*			Catalysis of a biochemical reaction at physiological
				temperatures. In biologically catalyzed reactions, the reactants
				are known as substrates, and the catalysts are naturally
				occurring macromolecular substances known as enzymes.
				Enzymes possess specific binding sites for substrates, and are
				usually composed wholly or largely of protein, but RNA that
			Molecular	has catalytic activity (ribozyme) is often also regarded as
GO:0003824	catalytic activity	down	function	enzymatic.
				Interacting selectively and non-covalently with a nucleotide,
				any compound consisting of a nucleoside that is esterified with
			Molecular	(ortho) phosphate or an oligophosphate at any hydroxyl group
GO:0000166	nucleotide binding	down	function	on the ribose or deoxyribose.

GO:1901265	nucleoside phosphate binding	down	Molecular function	Interacting selectively and non-covalently with nucleoside phosphate.
			Molecular	Interacting selectively and non-covalently with cations,
GO:0043169	cation binding	down	function	charged atoms or groups of atoms with a net positive charge.
			Molecular	
GO:0046872	metal ion binding	down	function	Interacting selectively and non-covalently with any metal ion.

# **Chapter 5**

The negative effect of soil microorganisms on plant growth only extends to the first weeks

> Jing Zhang^{1*} Peter G.L. Klinkhamer¹ Klaas Vrieling¹ T. Martijn Bezemer^{1, 2}

1) Plant Science and Natural Products, Institute of Biology Leiden (IBL), Leiden University, Sylviusweg 72, 2333BE Leiden, The Netherlands

2) The Netherlands Institute of Ecology (NIOO-KNAW), Department of Terrestrial Ecology, Droevendaalsesteeg 10, 6708 PB, Wageningen, The Netherlands

#### Abstract

Soil biotic communities can strongly impact plant performance. Many plant species grow worse in live soil than in sterilized soil. So far, most studies on plant-soilinteractions have estimated the effect of the soil microbial community on plant mass after a fixed duration of plant growth. However, these interactions may change over time and several studies have argued that plant-soil interactions are more important for young seedlings than for older plants. In this paper we ask the question: how longlasting is the effect of the soil microbial community on plant growth and we focus on relative plant growth rates at three stages: early growth (0-21 days), mid growth (22 to 42 days) and late growth (43 to 63 days). This is important as a plant with a reduced relative growth rate early in life, due to negative effects of the soil microbial community, may increase less in biomass for a much longer period, even though the relative growth rates do not differ anymore. We performed two growth experiments with Jacobaea vulgaris lasting 49 and 63 days. Plants were grown in sterilized soil or in sterilized soil inoculated with natural dune soil. In both experiments, differences in biomass of plants grown in sterilized soil and inoculated soil (live soil) increased throughout the experiment. Interestingly, linear regression models testing the relationship between *ln* transformed dry weight and time for younger plants and for older plants in sterilized soil and live soil, respectively, showed that the relative growth rate of plants in the sterilized soil was only significantly higher than that of plants in live soil in the first 2-3 weeks. After that period, there was no negative effect of live soil on plant relative growth rate anymore. In a third experiment, we examined the effect of the timing of soil inoculation prior to planting on the relative growth rate of J. vulgaris plants with four different timing treatments. Plant biomass was reduced in all inoculated soils compared to the sterilized soil. With increasing time between inoculation and planting, plant biomass decreased. Again, in all inoculated soils the negative effect of the soil microbial community on plant growth disappeared during the first weeks after planting. Overall, our results show that plants grow less well in live soil than in sterilized soil. The negative effects of soil inoculation on plant mass appear to extend over the whole growth period but arise from the negative effects on relative growth rates that occur in the first weeks after planting when plants have only less than 5% of the mass they obtained after 42 days. Our study highlights the

importance of examining relative growth rates rather than final biomass to estimate the effects of soil microbial communities on plants.

# Keywords

Dry plant biomass, Growth analysis, *Jacobaea vulgaris*, Live soil, Plant performance, Plant-soil interactions, Pathogenic soil microbial community, Relative growth rate, Sterilized soil

## Introduction

Interactions between plants and soil microbial communities are vital in mediating the balance and functioning of terrestrial ecosystems (Bever, 1994; Churchland and Grayston, 2014; Teste et al., 2017; Erktan et al., 2018). The soil microbiome is an important driver of plant performance. Soil microbial species e.g. pathogenic organisms, plant-growth-promoting rhizobacteria (PGPR, such as *Pseudomonas* and *Burkholderia*) and arbuscular mycorrhizal fungi (AMF), play an active role in modifying the development of plants (Johnson et al., 1997; Arora and Mishra, 2016; Artursson et al., 2016; Gil-Martinez et al., 2018). The effect of the soil microbial community on plant growth in laboratory experiments is often negative (Mangan et al., 2010; van de Voorde et al., 2012; Cortois et al., 2016).

One potential explanation for the negative effect of soil microbes on plant performance is that microbes and plants compete for nutrients. Alternatively, pathogens may accumulate in the soil over time, eventually resulting in a negative overall effect on plant performance (Dobson and Crawley, 1994; Wardle et al., 2004; Raaijmakers et al., 2009; Mordecai, 2011; van der Putten et al., 2013; Jacoby et al., 2017). In the previous chapters, we showed in experiments with ample nutrient supply that the negative effect of the soil microbial community on plant growth is mitigated if the plant's defense system is activated by foliar application of salicylic acid. This led us to hypothesize that the negative effect of the soil microbial community on plant growth in our system is due to an overall pathogenic effect of the soil microbial community. Although this effect was consistent, we did not find this effect to increase over several generations of plant growth (Chapter 2). An important question is therefore how long the negative effects of the soil microbial community on plant growth lasts.

So far, most studies on the effect of the soil microbial community on plant growth are conducted in pots (Hodge and Fitter, 2013). In such experiments, the negative effects of any treatment on plant mass often decline after some period of plant growth (typically six to eight weeks) (Bezemer et al., 2018; Dudenhöffer et al., 2018). This is often attributed to restricted root growth due to limitations in pot size, or to a decline in nutrient availability, and therefore considered an artefact of the experimental design 154

(Smith and Reynolds, 2012; Van de Voorde et al., 2012; Jing et al., 2015). It is also possible, however, that the pathogenic effect of the soil microbial community only last for a short period because (1) only seedlings are susceptible or (2) because over time plants alter the composition of the microbial community in the soil in which they grow so that it becomes less harmful (Bezemer et al., 2018; Dudenhöffer et al., 2018).

Previous studies on plant-soil-interactions typically focus on the effect of the soil microbial community on final plant biomass (van de Voorde et al., 2012; Bezemer et al., 2013; Anacker et al., 2014). It is important to note, however, that the effects of the soil microbial community on plant growth depend on the life stages of the plant (Arrigoni et al., 2018; Bezemer et al., 2018; Dudenhöffer et al., 2018). Seedlings are often highly vulnerable and susceptible to pathogenic microbes in the soil (Packer and Clay, 2000). In contrast, older plants with a more developed root system are typically less vulnerable (Kardol et al., 2013; Bezemer et al., 2018).

Effects on plant growth that occur during early life stages can still affect plant size and plant phenology in late life stages. When plants after some period grow with a similar relative growth rate, differences in absolute plant mass will still continue to increase. In Fig. 1 it is assumed that plants in sterilized soil grow with a constant relative growth rate (red line). Plants in live soil either grow with a constant relative growth rate lower than that of the plants in the live soil (green line) or they first grow with a lower relative growth rate but after an initial period ( $t_1$ ) their relative growth rate becomes similar to that of plants in the sterilized soil (blue line). In the latter case, although the effect of the soil microbial community only is present until  $t_1$ , differences in plant mass still continue to increase (Fig. 1b). Hence, to study the effect of soil microbes on plants, it is important to analyze relative growth rates.

In this study, we used linear regression models and *ln* transformed biomass data from repeated harvests to estimate relative growth rates in sterilized and live soil. We hypothesized that i) relative growth rates of plants are smaller in live soil than in sterilized soils ii) the negative effect on relative growth rates lasts only for a short period during the early plant life stages; and iii) the differences in plant mass between

plants grown in live soils and sterilized soils will continue to increase during the experiment.

We used *Jacobaea vulgaris* to test these hypotheses. *J. vulgaris* is native to The Netherlands. In a former experiment, we found that the plant mass of *J. vulgaris* growing in soil containing a live microbial community was 66% lower than when plants were grown in sterilized soil (Jing et al., Chapter 2). This negative effect of live soil on plant growth is in line with previous findings (e.g. van de Voorde et al., 2012; Kos et al., 2015; Wang et al., 2019). In the present study, to avoid nutrient limitation during the growth of *J. vulgaris*, nutrients were supplied regularly according to estimates of nutrient demand obtained from previous experiments (Steiner, 1980; Joosten et al., 2009). We carried out growth experiments with multiple harvesting points to estimate changes in relative growth rates in live and sterilized soils. Additionally, we grew *J. vulgaris* plants in soil that had been inoculated with live soil at varying time points before planting to manipulate the abundance of the microbial community in the soil. With the latter experiment we aimed to examine how the timing of inoculation into sterilized soil impacts the growth of *J. vulgaris*.



**Fig. 1** Conceptual figures showing relative growth rate and plant mass of *J. vulgaris* in both sterilized soil and live soil over time. (a) The relative growth rate in sterilized soil (red line) is higher than that in live soil (green line) (hypothesis 1) and this difference is maintained during the entire plant growth period. The blue line indicates

an initial lower relative growth rate of plants in the live soil but at  $t_1$  these plants obtain an equal relative growth rate as plants in the live soil (hypothesis 2). In Fig. 1a the yaxis denotes *ln* transformed plant mass. In Fig. 1b the y-axis denotes absolute plant mass. Note that even when relative growth rates become equal after an initial difference in relative growth rate in the early stage of life (the red line and blue line in Fig. 1a) the difference in absolute biomass continues to increase after that period (the red line and blue line in Fig. 1b).

#### Materials and methods

*J. vulgaris* (common ragwort) was used as plant species. We chose this species because it is a common species in The Netherlands that is strongly affected by plantsoil interactions (van de Voorde et al., 2011; van de Voorde et al., 2012; Bezemer et al., 2013). Seeds and soil were collected from Meijendel, a calcareous sandy dune area north of The Hague, The Netherlands (52°11′N, 4°31′E).

#### Seeds

Before seed germination, all seeds were shaken for 2 min in 70% ethanol, then washed with sterilized water, put for 12 min in 2% bleach, and finally rinsed four times with sterilized water to avoid influences of seed-borne microbes. The surface-sterilized seeds were then placed in standard Petri dishes containing filter paper, which was moistened with Milli-Q water. Afterwards, all Petri dishes containing seeds were placed in plastic zip-lock bags and stored in a climate room (relative humidity 70%, light 16 h at 20°C, dark 8 h at 20°C) for the duration of germination.

#### Soil

At Meijendel, the topsoil was collected to a depth of 15 cm after removing the grassland vegetation and the organic layer of the surface. The soil was sieved using a 5 mm sized mesh to remove plant roots and various soil fauna, homogenized with a concrete mixer and then stored into 20-liter plastic bags (Nasco Whirl-Pak Sample Bag). Bags were either sterilized by 35-K Gray gamma-irradiation (Synergy Health Company, Ede, The Netherlands) or kept at 4°C for inoculation. Potting soil

(Slingerland potgrond, Zoeterwoude, The Netherlands) was also sterilized by 35-K Gray gamma-irradiation.

#### **Plant growth**

After germination, seedlings were randomly transferred individually to 500 ml pots containing either "sterilized soil" or "live soil". The live soil treatment consisted of a mixture of 87.5% sterilized dune soil, 2.5% sterilized potting soil and 10% live soil. The sterilized soil treatment contained 97.5% of sterilized dune soil and 2.5% of sterilized potting soil. Sterilized potting soil was added to all pots to increase the organic matter content of the soil. Sterilized soil and live soil were kept in bags and left in the climate room for 14 days (relative humidity 70%, light 16h at 20°C, dark 8h at 20°C) to enable the establishment of microbial communities in the inoculated soil before potting. Before filling the pots, the soil in each bag was mixed. After filling, pots were randomly distributed over the climate room. Plants were watered regularly with Milli-Q water and 5 ml Steiner nutrient solution was added per plant on day 7, 10 ml Steiner nutrient solution was added on day 13, and 20 ml Steiner nutrient solution was added on days 19, 28, 37, and 42. The Steiner nutrient solution (Steiner, 1980) was prepared from 7 different stock solutions (106.2 g Ca(NO₃)₂·4H₂O, 29.3 g KNO₃, 13.6 g KH₂PO₄, 49.2 g MgSO₄·7H₂O, 25.2 g K₂SO₄ and 2.24 g KOH, 3.29 g Fe-EDTA added to 1 liter demineralized water, and a stock solution with micro elements (a mixed solution of 0.181 g MnCl₂·4H₂O, 0.286 g H₃BO₃, 0.022 g  $ZnSO_4 \cdot 7H_2O$ , 0.0078 g CuSO₄  $\cdot 5H_2O$  and 0.0126 g NaMoO₄  $\cdot 2H_2O$  added to 1 liter demineralized water). Ten ml of each stock solution was diluted in 1 liter of demineralized water before use.

#### The effect of live soil on the growth of *J. vulgaris*

**Experiment 1:** An experiment to measure the growth of *J. vulgaris* overtime was performed starting with 1-week-old seedlings, two soil treatments and eight harvesting time points over seven weeks. The harvests were on days 0 (1-week-old seedlings), 7, 14, 21, 28, 35, 42 and 49 after planting. Pots were randomly labeled and allocated to the different harvests. Ten replicates were used for each treatment resulting in 2 treatments × 8 harvesting points × 10 replicates = 160 plants. Harvested

plants (shoots and roots) were oven-dried at 60°C for approximately one week and dry mass was determined.

**Experiment 2:** The growth experiment was repeated using the same soil treatments, but with more harvests during the first 3 weeks. In this experiment, plants were harvested at day 0, 3, 6, 9, 12, 15, 18, 21, 28, 35, 42, 49, 56 and 63 after planting. Ten plants per soil treatment were harvested at each harvesting time point thus resulting in 2 treatments × 14 harvest points × 10 replicates = 280 plants. In this experiment, at each harvest, the plants were gently removed from the pot. Shoots were separated from roots with a pair of scissors just above the root crown, and roots were cleaned with water and then put into aluminum foil. Then, all the harvested plant parts were freeze-dried for approximately one week, and dry mass was determined.

#### The effect of time of inoculation on the growth of J. vulgaris

To examine the effect of the timing of soil inoculation on the relative growth rate of *J. vulgaris* plants, sterilized soils were inoculated at different time points prior to planting the seedlings. In this experiment, 1-week-old seedlings were planted into 500 ml pots containing either "sterilized soil" or four different "live soil" treatments. For these four treatments, a mixture of 10% of live soil was mixed with 90% sterilized soil, and then the mixed soil was kept in the climate room for 0, 1, 2 and 4 weeks (relative humidity 70%, light 16h at 20°C, dark 8h at 20°C) to enable different build-up times for the microbial community in the soil at the time of planting. The live soil treatments were labeled as "live-0", "live-1", "live-2" and "live-4" respectively. Seedlings were randomly distributed over the five soil treatments and nine harvests over six weeks. Plants were harvested on days 0 (as seedlings), 4, 8, 12, 16, 20, 28, 35 and 42. Eight replicates were used per treatment combination, resulting in 5 treatments × 9 harvests × 8 replicates totaling 360 plants. Fresh weight was recorded, because leaves were frozen immediately as we intended to quantify the levels of SA in the plant material. However, due time limitations these data have not been collected.

#### Calculations and statistical analyses

Biomass was plotted against time for plants grown in sterilized and live soil. A student t-test was then performed to test for differences between dry plant mass in sterilized and live soils at each time point. *Ln* transformed biomass was also plotted against time. Plant growth was divided into three stages: early growth (0-21 days), mid growth (22 to 42 days) and late growth (43 to 63 days). For each experiment, a separate line was then fitted through the dry plant mass data for these different periods. Late growth was only measured in experiment 2. Because this division in two time periods is somewhat arbitrary, we backed this analyses up with a sequential backward regression approach for the entire growth period for each experiment. We started this analysis with the two latest harvesting points and then sequentially added the previous data point. In this way we could test for which time periods differences in relative growth rate were significant. For each regression the slope and standard error (SE) of the slope were determined and differences between the slopes for the linear regression models in sterilized and live soil were then tested with a t-test in Excel, t =Slope1-Slope2  $SQRT(SE1^2+SE2^2)$ 

#### Results

#### The effect of live soil on the growth of J. vulgaris

**Experiment 1:** Soil inoculation had a strong negative effect on plant dry mass throughout the experiment (Fig. 2a). The difference in plant dry mass between the sterilized and live soil treatments increased during the entire experiment. From day 21 onward, the dry plant mass of *J. vulgaris* in sterilized soil was significantly larger than the dry mass of plants grown in live soil (Fig. 2). For young plants (0-21 days) the relative growth rate (slope in Fig. 2b, c) in sterilized soil was significantly larger than that for live soil while relative growth rates did not differ for mid-aged plants (22-49 days, Fig. 2b, c). This result was backed up by the sequential backward regression that showed that the relative growth rates were not significantly different for the periods between 22 and 49 days (Table S1).



**Fig. 2** Experiment 1. (a) Mean ( $\pm$  SE) biomass of *J. vulgaris* in sterilized and live soil over 49 days. For each time point, differences between the biomass of the plants in the two soils were tested for significance with a t-test, * indicates a significant difference (P < 0.05). (b) Two linear regression models (early: day 7-21, and mid: day 28-49) of *ln* transformed biomass of *J. vulgaris* in both sterilized and live soil. The extrapolated dashed parts of the lines are based on the linear regression models for day 28-49. (c) Slopes (mean $\pm$  SE) of the regression lines in (b). Differences between

the slopes for live-soil and control soil were tested for significance with a-test. *** indicates P < 0.001.

**Experiment 2:** The first experiment was repeated with more harvesting points during the first 21 days and an extended growth period. Again, the effect of live soil on plant growth was negative (Fig. 3a). The difference in absolute plant biomass increased until day 56. Young plants (0-21 days) had significantly higher relative growth rates in sterilized soil, mid-aged plants (22-42 days) had similar relative growth rates; while for older plants (49-63 days) the relative growth rates were even higher in live soil (Fig. 3b, c). Backward regression showed that the relative growth rate was higher for the plants in live soil for the period 63-28 days. If younger ages are included in the analysis, differences are no longer significant (Table S2).



**Fig. 3** Experiment 2. (a) Mean ( $\pm$  SE) of dry plant mass of *J. vulgaris* in sterilized and live soil over 63 days. For each time point, differences between the biomass of the plants in the two soils were tested for significance with a t-test, * indicates a significant difference at *P* < 0.05. (b) Three linear regression models (early: day 0-19, mid: 22-42, late: 49-63) of *ln* transformed biomass of *J. vulgaris* in both sterilized and live soil. The two wztrapolated dashed parts of the lines are based on the linear regression models for day 0-19 and 49-63 (c) Mean slope ( $\pm$  SE) of the linear

regression lines in (b). Differences between the slopes for live soil and sterilized soil were tested for significance with a t-test. * indicates P < 0.05.

#### The effect of time of inoculation before planting on the growth of J. vulgaris

**Experiment 3:** Plants produced less biomass in inoculated soils than in sterilized soil (Fig. 4a). For young plants (0-21 days) the relative growth rate in sterilized soil was significantly larger than that for live-0, live-1, live-2 or live-4 soil. Relative growth rates did not differ for mid-aged plants between live-0, live-1 and live-2 soil. Interestingly, relative growth rates of plants from live-4 soil for the mid-aged period were significantly higher than the relative growth rate of plants in sterilized soil (Fig. 3c; Table S3). Timing of the inoculation did affect the relative growth rates of plants in the early phase (0-21 days). The longer the time between inoculation and planting the lower the relative growth rate of young plants was ( $R^2 = 0.99$ , P < 0.05, df = 3). This was no longer true for the mid-aged period ( $R^2 = 0.71$ , P = 0.15, df = 3). These results were largely backed up by the backward sequential regression, which showed that relative growth rates were only higher for plants grown in the sterilized soil if very young plant ages were included. Especially for the live-4 soil the relative growth rate was even higher for plants grown in live soil when only older plants were included (Table S3).





models (early: 0-20, mid: 28-42) of *ln* transformed fresh biomass of *J. vulgaris* in sterilized soil and four live soils. The extrapolated dashed parts of the lines are based on the linear regression models of day 28-42. (c) Mean slope ( $\pm$  SE) of linear lines in (b). Differences between the slopes for live-soil and sterilized soil were tested for significance with a t-test. *** indicates significant difference at *P* < 0.001; ** indicates significant difference at *P* < 0.01.

# Discussion

In this study, we report the results of three experiments in which we measured the growth of *J. vulgaris* to test how the effects of soil microbial communities on plant growth change over time. We found a consistent negative effect of the soil microbial community in all three experiments. Biomass was larger in sterilized soil than in live soil. However, analyses of the *ln* transformed data, show that the relative growth rates were significantly higher in sterilized soil than in live soil only for young plants, and not for mid-aged plants. Moreover, in experiment 2, which was continued for a longer period, older plants even had a higher relative growth rate in the live soil. Hence, all data sets showed that the negative effects of soil inoculation on plant mass appear to extend during a long period, but arise from the negative effects that occur in the first weeks after planting when plants have only obtained less than 5% of the mass they obtain after 42 days.

We observed a consistent negative effect of live soil containing a natural soil microbial community in all three experiments. It is plausible that this was due to a net pathogenic effect of the soil microbial community on plant growth (Klironomos, 2002; Joosten et al., 2009; Harrison and Bardgett, 2010; Cortois et al., 2015). This hypothesis has been widely verified in other studies. For example, bacterial microbes such as *Ralstonia solanacearum, Agrobacterium tumefaciens, Erwinia amylovora and Streptomyces scabies* have been frequently isolated from natural soils (Curl et al., 1998; Michel et al., 1998; Gómez et al., 2017; Sharifazizi et al., 2017). These pathogenic microbes can adversely affect plant health and production (Huang et al., 2013; Cesarano et al., 2017). Several studies have indicated that soil microbes

compete with plants for available nutrients in the soil, and this could also result in negative effects on plant growth in inoculated soil (Bardgett et al., 2003; Fontaine et al., 2003; Dunn et al., 2006). However, in our study, we grew plants in a nutrient-rich environment by supplying a nutrient solution, and hence we argue that it is unlikely that the negative effect of live soil on plant growth was due to plant-microbe competition for nutrients. In Chapter 2, application of SA mitigated the negative effects of the live soil on the growth of *J. vulgaris*, in combination with the fact that activation of SA-dependent signaling pathways leads to the expression of pathogenesis-related proteins (PRP) contributing to resistance (Glazebrook, 2005; Spoel et al., 2007), this together suggests that the negative soil effect on plant growth was due to microbial pathogens.

Our study exemplifies that the negative effects of soil inoculation on plant mass can extend over the entire growth period, even though the differences are due to negative effects that occur during the first weeks after planting. There are several explanations for the observation that older plants do not exhibit a negative response to live soils. First, younger plants or seedlings may be more vulnerable and susceptible to pathogenic microbes in the soil than older plants with well-developed root systems (Packer and Clay, 2000). Root development plays an important role for plants in suppressing soil-borne pathogens (Watt et al., 2006; Emmett et al., 2014), and is correlated with soil abiotic or biotic characteristics (Kardol et al., 2013; Arrigoni et al., 2018; Bezemer et al., 2018). Herms and Mattson (1992) demonstrated that plants have to invest in their roots first before they can defend themselves against biotic stress. Hence, it may take a while for plants to build-up their defense systems (Raaijmakers et al., 2009; Hayat et al., 2010). Alternatively, it is well established that plants influence the soil microbial community during growth and hence, it is also possible that the differences in the response of younger and older plants to live soil is due to changes that have occurred in the soil microbial community. Previous work with the same plant species, J. vulgaris, where seedlings were planted in soil in which plants of the same species had been grown first, showed that the differences between responses of young and old plants are likely related to the sensitivity of plant stages and not due to changes in the soil community. Young J. vulgaris exhibited a strong negative conspecific feedback, but this effect diminished over time and became neutral in older plants (Bezemer et al., 2018).

Interestingly we observed that the longer ago the soil was inoculated the stronger the negative effect of the inoculum on plant growth. This also indicates that the negative effects of live soil on plant growth that are commonly observed for this plant species are mediated by the soil microbial community and the variation that is typically observed in plant growth experiments may result from the different densities of soil-borne microbes. We expect that the oldest inoculated live soil contained the highest density of pathogenic microbes, leading to a stronger negative effect on plant growth (Pernilla et al., 2010; Dudenhöffer, et al., 2018). However, in this study, we did neither quantify the microbial density in the soil nor measure plant defense-related compounds such as salicylic acid, or pyrrolizidine alkaloids, and we suggest future work should focus on these two aspects.

In conclusion, our results indicate that live soil negatively affected plant growth. In most cases the difference between plant biomass of plants grown in sterilized soil and live soil increased during the entire experiment. However, the relative growth rates of plants in the sterilized soil and live soil only differed for young plants. Moreover, there was a negative correlation between the time of soil inoculation before planting and the relative growth rate of *J. vulgaris* plants, but for all incubation periods the negative effects were only present for young plants. Hence, our results suggest that young plants ( $\leq 21$  days) or seedlings are most sensitive to soil pathogens while older plants ( $\geq 22$  days) are no longer affected. Our study highlights the importance of examining relative growth rates rather than final biomass to estimate the effects of soil microbial communities on plants.

#### Acknowledgements

We would like to thank Karin van der Veen-van Wijk for assistance during the final biomass harvest, Jan Vink for driving us to the field site to collect soil used for the inoculation experiment, and Jorian van Kampen, Gang Chen for collecting soil. We would also like to thank the China Scholarship Council for financial support.

#### References

- Anacker, B.L., Klironomos, J.N., Maherali, H., Reinhart, K.O. and Strauss, S.Y., 2014. Phylogenetic conservatism in plant-soil feedback and its implications for plant abundance. *Ecology Letters*, 17(12), pp.1613-1621.
- Arora, N.K. and Mishra, J., 2016. Prospecting the roles of metabolites and additives in future bioformulations for sustainable agriculture. *Applied Soil Ecology*, 107, pp.405-407.
- Arrigoni, E., Antonielli, L., Pindo, M., Pertot, I. and Perazzolli, M., 2018. Tissue age and plant genotype affect the microbiota of apple and pear bark. *Microbiological research*, 211, pp.57-68.
- Artursson, V., Finlay, R.D. and Jansson, J.K., 2006. Interactions between arbuscular mycorrhizal fungi and bacteria and their potential for stimulating plant growth. *Environmental microbiology*, 8(1), pp.1-10.
- Bardgett, R.D., Streeter, T.C. and Bol, R., 2003. Soil microbes compete effectively with plants for organic-nitrogen inputs to temperate grasslands. *Ecology*, 84(5), pp.1277-1287.
- Bever, J.D., 1994. Feedback between plants and their soil communities in an old field community. *Ecology*, 75(7), pp. 1965-1977.
- Bezemer, T.M., van der Putten, W.H., Martens, H., van de Voorde, T.F., Mulder, P.P. and Kostenko, O., 2013. Above-and below-ground herbivory effects on below-ground plant-fungus interactions and plant-soil feedback responses. *Journal of Ecology*, 101(2), pp.325-333.
- Bezemer, T.M., Jing, J., Bakx-Schotman, J.T. and Bijleveld, E.J., 2018. Plant competition alters the temporal dynamics of plant-soil feedbacks. *Journal of Ecology*, 106(6), pp.2287-2300.
- Cesarano, G., Zotti, M., Antignani, V., Marra, R., Scala, F. and Bonanomi, G., 2017. Soil sickness and negative plant-soil feedback: A reappraisal of hypotheses. *Journal of plant pathology*, 99(3), pp.545-570.
- Churchland, C. and Grayston, S.J., 2014. Specificity of plant-microbe interactions in the tree mycorrhizosphere biome and consequences for soil C cycling. *Frontiers in microbiology*, *5*, p.261.
- Cortois, R., Schröder-Georgi, T., Weigelt, A., van der Putten, W.H. and De Deyn, G.B., 2016. Plant-soil feedbacks: role of plant functional group and plant traits. *Journal of Ecology*, *104*(6), pp.1608-1617.
- Curl, E.A., Lartey, R. and Peterson, C.M., 1988. Interactions between root pathogens and soil microarthropods. Agriculture, ecosystems & environment, 24(1-3), pp.249-261.
- Dobson, A. and Crawley, M., 1994. Pathogens and the structure of plant communities. *Trends in Ecology* & *Evolution*, *9*(10), pp.393-398.
- Dudenhöffer, J.H., Ebeling, A., Klein, A.M. and Wagg, C., 2018. Beyond biomass: Soil feedbacks are transient over plant life stages and alter fitness. *Journal of Ecology*, 106(1), pp.230-241.
- Dunn, R.M., Mikola, J., Bol, R. and Bardgett, R.D., 2006. Influence of microbial activity on plantmicrobial competition for organic and inorganic nitrogen. *Plant and Soil*, 289(1-2), pp.321-334.
- Emmett, B., Nelson, E.B., Kessler, A. and Bauerle, T.L., 2014. Fine-root system development and susceptibility to pathogen colonization. *Planta*, 239(2), pp.325-340.
- Erktan, A., McCormack, M.L. and Roumet, C., 2018. Frontiers in root ecology: recent advances and future challenges. *Plant and Soil*, 424, pp.1-9.
- Fontaine, S., Mariotti, A. and Abbadie, L., 2003. The priming effect of organic matter: a question of microbial competition? Soil Biology and Biochemistry, 35(6), pp.837-843.
- Gil-Martínez, M., López-García, Á., Domínguez, M.T., Navarro-Fernández, C.M., Kjøller, R., Tibbett,M. and Marañón, T., 2018. Ectomycorrhizal fungal communities and their functional traits

mediate plant-soil interactions in trace element contaminated soils. *Frontiers in Plant Science*, *9*, p.1682.

- Gómez Expósito, R., De Bruijn, I., Postma, J. and Raaijmakers, J.M., 2017. Current insights into the role of rhizosphere bacteria in disease suppressive soils. *Frontiers in Microbiology*, 8, p.2529.
- Glazebrook, J., 2005. Contrasting mechanisms of defense against biotrophic and necrotrophic pathogens. *Annu. Rev. Phytopathol*, 43, pp.205-227.
- Harrison, K.A. and Bardgett, R.D., 2010. Influence of plant species and soil conditions on plant-soil feedback in mixed grassland communities. *Journal of Ecology*, 98(2), pp.384-395.
- Hayat, R., Ali, S., Amara, U., Khalid, R. and Ahmed, I., 2010. Soil beneficial bacteria and their role in plant growth promotion: a review. *Annals of microbiology*, 60(4), pp.579-598.
- Herms, D.A. and Mattson, W.J., 1992. The dilemma of plants: to grow or defend. *The quarterly review* of biology, 67(3), pp.283-335.
- Hodge, A. and Fitter, A.H., 2013. Microbial mediation of plant competition and community structure. *Functional Ecology*, 27(4), pp.865-875.
- Huang, L.F., Song, L.X., Xia, X.J., Mao, W.H., Shi, K., Zhou, Y.H. and Yu, J.Q., 2013. Plant-soil feedbacks and soil sickness: from mechanisms to application in agriculture. *Journal of chemical ecology*, 39(2), pp.232-242.
- Jacoby, R., Peukert, M., Succurro, A., Koprivova, A. and Kopriva, S., 2017. The role of soil microorganisms in plant mineral nutrition-current knowledge and future directions. *Frontiers* in plant science, 8, p.1617.
- Jing, J., Bezemer, T.M. and van der Putten, W.H., 2015. Complementarity and selection effects in early and mid-successional plant communities are differentially affected by plant-soil feedback. *Journal of Ecology*, 103(3), pp.641-647.
- Johnson, N.C., Graham, J.H. and Smith, F.A., 1997. Functioning of mycorrhizal associations along the mutualism-parasitism continuum. *The New Phytologist*, 135(4), pp.575-585.
- Joosten, L., Mulder, P.P., Klinkhamer, P.G. and van Veen, J.A., 2009. Soil-borne microorganisms and soil-type affect pyrrolizidine alkaloids in *Jacobaea vulgaris*. *Plant and Soil*, 325(1-2), p.133.
- Kardol, P., De Deyn, G.B., Laliberté, E., Mariotte, P. and Hawkes, C.V., 2013. Biotic plant-soil feedbacks across temporal scales. *Journal of Ecology*, 101(2), pp.309-315.
- Klironomos, J.N., 2002. Feedback with soil biota contributes to plant rarity and invasiveness in communities. *Nature*, 417(6884), pp.67-70.
- Kos, F., Poland, D., Simmons-Duffin, D. and Vichi, A., 2015. Bootstrapping the O (N) archipelago. Journal of High Energy Physics, 2015(11), p.106.
- Mangan, S.A., Schnitzer, S.A., Herre, E.A., Mack, K.M., Valencia, M.C., Sanchez, E.I. and Bever, J.D., 2010. Negative plant-soil feedback predicts tree-species relative abundance in a tropical forest. *Nature*, 466(7307), pp.752-755.
- Michel, V.V. and Mew, T.W., 1998. Effect of a soil amendment on the survival of *Ralstonia* solanacearum in different soils. *Phytopathology*, 88(4), pp.300-305.
- Mordecai, E.A., 2011. Pathogen impacts on plant communities: unifying theory, concepts, and empirical work. *Ecological Monographs*, 81(3), pp.429-441.
- Packer, A. and Clay, K., 2000. Soil pathogens and spatial patterns of seedling mortality in a temperate tree. *Nature*, 404(6775), pp.278-281.
- Pernilla Brinkman, E., Van der Putten, W.H., Bakker, E.J. and Verhoeven, K.J., 2010. Plant-soil feedback: experimental approaches, statistical analyses and ecological interpretations. *Journal* of Ecology, 98(5), pp.1063-1073.

- Raaijmakers, J.M., Paulitz, T.C., Steinberg, C., Alabouvette, C. and Moënne-Loccoz, Y., 2009. The rhizosphere: a playground and battlefield for soilborne pathogens and beneficial microorganisms. *Plant and soil*, 321(1-2), pp.341-361.
- Sharifazizi, M., Harighi, B. and Sadeghi, A., 2017. Evaluation of biological control of *Erwinia amylovora*, causal agent of fire blight disease of pear by antagonistic bacteria. *Biological Control*, 104, pp.28-34.
- Smith, L.M. and Reynolds, H.L., 2012. Positive plant-soil feedback may drive dominance of a woodland invader, *Euonymus fortunei*. *Plant Ecology*, 213(5), pp.853-860.
- Spoel, S.H., Johnson, J.S. and Dong, X., 2007. Regulation of tradeoffs between plant defenses against pathogens with different lifestyles. *Proceedings of the National Academy of Sciences*, U.S.A, 104(47), pp.18842-18847.
- Steiner, A.A., 1980. The selective capacity of plants for ions and its importance for the composition and treatment of the nutrient solution. *Acta Horticulturae*, *98*, pp.87-98.
- Teste, F.P., Kardol, P., Turner, B.L., Wardle, D.A., Zemunik, G., Renton, M. and Laliberté, E., 2017. Plant-soil feedback and the maintenance of diversity in Mediterranean-climate shrublands. *Science*, 355(6321), pp.173-176.
- van der Putten, W.H., Bardgett, R.D., Bever, J.D., Bezemer, T.M., Casper, B.B., Fukami, T., Kardol, P., Klironomos, J.N., Kulmatiski, A., Schweitzer, J.A. and Suding, K.N., 2013. Plant-soil feedbacks: the past, the present and future challenges. *Journal of Ecology*, 101(2), pp.265-276.
- van de Voorde, T.F., van der Putten, W.H. and Martijn Bezemer, T., 2011. Intra-and interspecific plantsoil interactions, soil legacies and priority effects during old-field succession. *Journal of Ecology*, 99(4), pp.945-953.
- van de Voorde, T.F., van der Putten, W.H. and Bezemer, T.M., 2012. Soil inoculation method determines the strength of plant-soil interactions. *Soil Biology and Biochemistry*, 55, pp.1-6.
- Wang, M., Ruan, W., Kostenko, O., Carvalho, S., Hannula, S.E., Mulder, P.P., Bu, F., van der Putten, W.H. and Bezemer, T.M., 2019. Removal of soil biota alters soil feedback effects on plant growth and defense chemistry. *New Phytologist*, 221(3), pp.1478-1491.
- Wardle, D.A., Bardgett, R.D., Klironomos, J.N., Setälä, H., Van Der Putten, W.H. and Wall, D.H., 2004. Ecological linkages between aboveground and belowground biota. *Science*, 304(5677), pp.1629-1633.
- Watt, M., Silk, W.K. and Passioura, J.B., 2006. Rates of root and organism growth, soil conditions, and temporal and spatial development of the rhizosphere. *Annals of botany*, *97*(5), pp.839-855.

# Supplementary data

**Table S1.** Sequential regression analysis of *ln* transformed dry plant mass of *J. vulgaris* in experiment 1 in sterilized and live soil. The slopes were calculated backward sequentially. Slope1 and slope2 represent slopes in sterilized soil and live soil, respectively. SE indicates standard error, *df* means degrees of freedom. A t-test was used to determine significance between the two slopes. *** represents P < 0.001.

	Day range	slope1	SE1	slope2	SE2	df	<i>t</i> -value	Р
	42-49	0.11	0.02	0.16	0.03	36	1.60	ns
Slopes for	35-49	0.12	0.01	0.13	0.01	53	1.31	ns
sequential	28-49	0.16	0.01	0.13	0.01	73	-2.86	ns
backward	21-49	0.14	0.01	0.12	0.01	93	-1.55	ns
regression	14-49	0.15	0.01	0.12	0.01	113	-4.54	***
	7-49	0.16	0.00	0.12	0.00	131	-6.85	***

**Table S2.** Sequential regression analysis of *ln* transformed dry plant mass of *J. vulgaris* in experiment 2 in sterilized and live soil. The slopes were calculated backward sequentially. Slope1 and slope2 represent slopes in sterilized soil and live soil. SE indicates standard error, *df* means degrees of freedom. A t-test was used to determine significance between the two slopes. * represents P < 0.05.

	Day	slope1	SE1	slope2	SE2	df	<i>t</i> -value	Р
	Tunge							
	56-63	0.02	0.02	0.11	0.02	28	-2.61	*
	49-63	0.04	0.01	0.09	0.01	44	0.01	*
	42-63	0.08	0.01	0.11	0.01	60	-2.32	*
	35-63	0.09	0.01	0.11	0.01	76	-1.99	*
	28-63	0.10	0.00	0.12	0.00	92	-2.05	*
Slopes for	22-63	0.11	0.00	0.12	0.00	108	-1.20	ns
sequential backward	19-63	0.12	0.00	0.13	0.00	124	-1.62	ns
regression	16-63	0.13	0.00	0.13	0.00	140	-1.34	ns
	13-63	0.13	0.00	0.14	0.00	155	1.59	ns
	10-63	0.14	0.00	0.14	0.00	171	1.02	ns
	7-63	0.14	0.00	0.15	0.00	187	0.52	ns
	4-63	0.15	0.00	0.15	0.00	205	-0.19	ns
	0-63	0.17	0.00	0.17	0.00	223	-0.59	ns

**Table S3.** Sequential regression analysis of *ln* transformed fresh plant mass of *J. vulgaris* in experiment 3 in sterilized and inoculated soil 0, 1, 2 or 4 weeks before planting (live-0, live-1, live-2 and live-4). The slopes were calculated backward sequentially. Slope1 and slope2 represent slopes in sterilized soil and live soil respectively. SE indicates standard error, *df* degrees of freedom. A t-test was used to determine significance between the two slopes for each combination of sterilized and live soil. * represents P < 0.05, ** represents P < 0.01, *** represents P < 0.001. Note that the slopes for the sterilized soil are used for comparison with the slopes of the live-0, live-1, live-2 and live-4 and are thus represented in the table 4 times. Backward sequential slope calculation

	Day range	slope1	SE1	slope2	SE2	df	<i>t</i> -value	Р
	35-42	0.04	0.04	0.07	0.02	28	0.48	ns
	28-42	0.08	0.02	0.10	0.01	44	1.09	ns
	20-42	0.09	0.01	0.12	0.01	60	2.77	*
	16-42	0.10	0.01	0.13	0.01	76	2.64	*
	12-42	0.11	0.01	0.12	0.01	92	1.57	ns
	8-42	0.12	0.01	0.13	0.00	108	1.14	ns
	4-42	0.13	0.01	0.13	0.00	124	0.53	ns
Live-0	0-42	0.15	0.01	0.15	0.00	140	-0.31	ns
	35-42	0.04	0.04	0.12	0.04	28	1.35	ns
	28-42	0.08	0.02	0.09	0.02	44	0.49	ns
	20-42	0.09	0.01	0.10	0.01	60	0.86	ns
	16-42	0.10	0.01	0.10	0.01	76	0.05	ns
	12-42	0.11	0.01	0.11	0.01	92	-0.04	ns
	8-42	0.12	0.01	0.11	0.00	108	-1.22	ns
	4-42	0.13	0.01	0.12	0.00	124	-2.04	*
Live-1	0-42	0.15	0.01	0.13	0.00	140	-2.32	*
	35-42	0.04	0.04	0.10	0.03	28	1.02	ns
	28-42	0.08	0.02	0.10	0.01	44	0.86	ns
	20-42	0.09	0.01	0.11	0.01	60	1.51	ns
	16-42	0.10	0.01	0.10	0.01	76	0.69	ns
	12-42	0.11	0.01	0.11	0.00	92	-0.21	ns
	8-42	0.12	0.01	0.11	0.00	108	-1.75	ns
	4-42	0.13	0.01	0.11	0.00	124	-2.99	**
Live-2	0-42	0.15	0.01	0.13	0.00	140	-3.04	**
	35-42	0.04	0.04	0.16	0.05	26	1.67	ns
	28-42	0.08	0.02	0.17	0.02	40	2.94	**
	20-42	0.09	0.01	0.15	0.02	55	3.59	***
	16-42	0.10	0.01	0.13	0.01	71	2.34	*
	12-42	0.11	0.01	0.13	0.01	87	1.47	ns
	8-42	0.12	0.01	0.11	0.01	103	-0.97	ns
	4-42	0.13	0.01	0.12	0.01	119	-1.14	ns
Live-4	0-42	0.15	0.01	0.13	0.01	135	-2.19	*

# Chapter 6

General discussion

# **General discussion**

Plants can alter the microbial community in their rhizosphere, and in turn, the microbial community influences plant growth and development (Bever 1994; van der Heijden et al., 2008; Hahl et al., 2020). In general, the relationship between plants and soil microbes can be neutral, positive and negative (Nijjer et al., 2007; Bhattacharyya and Jha, 2012). However, often plants grow less well in soil that contains a live microbial community (live soil) than when growing in sterilized soil (Kulmatiski et al., 2008). An overall net effect of soil micorbial pathogenic microbes is one of the many mechanisms behind reduced plant growth in live soils (Cesarano et al., 2017).

In nature, plants have developed many defensive strategies, e.g. via hormone signaling, against microbial pathogens or herbivores. In agriculture, inducing hormonal signaling pathways has become a promising strategy to increase plant resistance against these enemies (Haney and Ausubel, 2015; Yang et al., 2015), and this is now applied to control soil microbial pathogens (Fujita et al., 2006; Yang et al., 2015; Berens et al., 2019). Exogenous application of SA to plant leaves activates systemic acquired resistance in the plant associated with the production of pathogenrelated proteins against microbial pathogens (Reymond and Farmer, 1998), while foliar application of JA or MeJA activates induced systemic defenses against herbivores and necrotrophic pathogens (Nahar et al., 2011). If the reduction of plant growth in live soil is caused by an overall pathogenic effect from the soil microbial community, we would expect that plant defense signaling hormones play a role in the interaction between the soil microbial community and plants. In particular, we would expect the negative effect to be mitigated after the application of SA as this would increase the defense of plants against microbes (Maurhofer et al., 1998; Berendsen et al., 2012; Wang et al., 2013).

Although a number of studies assessed the positive effects of hormonal signaling pathways on a plant's immunity against pathogenic microbes, whether and how these hormonal signaling pathways affect the soil microbial community and subsequently plant growth is still poorly understood, and whether activation of SA-induced resistance could potentially select a more beneficial soil microbial community over time is not known.

In this thesis, I examined how harnessing the plants' immune system affects the relationship between plants and the soil microbial community. I tested the hypothesis that the negative effect of live soil on plant growth is due to changes in the microbial community belowground and that activation of SA-induced resistance will potentially mitigate the negative effect of live soil on plant growth through altering the rhizosphere microbial composition and the expression of functional genes. Furthermore, I investigated how long-lasting the effect of the soil microbial community on plant growth is and if the effect changed at different plant growth stages.

In this chapter, I discuss the findings of this thesis and compare them with the results from several recently published studies in this research field, aiming to provide a broader perspective of my research findings within the field of above-belowground plant-soil interactions.

## Activation of hormonal induced defenses in plants growing in live soil

Many plant species produce more biomass in sterilized soil than in soil that contains a live microbial community. This could be due to an overall net pathogenic effect of the soil microbial community (Kulmatiski et al., 2008; Miki 2012). In Chapter 2 I studied the effect of live soil on plant growth of four plant species to investigate whether the overall negative soil effect is a common phenomenon among these plant species. Interestingly, the live soil only negatively affected two (J. vulgaris and C. vulgare) out of the four tested plant species, while for the other two species (T. repens and D. carota) we found no effect. This finding is in line with previous studies showing that interactions between plant species and soil microbial communities are highly species-specific (Klironomos, 2002; Joosten et al., 2009; Harrison and Bardgett, 2010; Wang et al., 2019). Plant genotype, diversity and neighboring-species can all influence these interactions, and the soil microbial community (reviewed in Bever et al., 2010), and this effect is mostly likely related to secondary metabolites exuded by the plants (Smith et al., 2018; Zhu et al., 2019; Dror et al., 2020). For example, pyrrolizidine alkaloids (PAs) are a group of secondary metabolites of the species J. vulgaris that are known to affect soil microbial pathogens. Genotypes of J. *vulgaris* vary in the concentration of Pas that they contain and presumably exude, and

this can influence soil microbial communities (Kowalchuk et al., 2006; Joosten et al., 2009; Kirk et al., 2010; Kostenko et al., 2012). Other studies have shown that rootemitted volatile compounds influence the composition of soil microbial communities (Delory et al., 2016; Massalha et al., 2017; Bailly 2020).

In addition, I examined if the live soil effect can be altered by the foliar application of SA or JA. Overall, activation of hormonal-induced resistance itself for a plant is costly (Vos et al., 2013). This is exemplified in my work where the foliar application of JA and SA to plant leaves resulted in reduced plant growth when plants were grown in sterilized soil for all four species (Chapter 2). Notably, for the two species in which the live soil had a negative effect on plant growth, we found that this effect was mitigated by the application of SA. In the other two species, the application of SA did not affect plant growth. Plants respond to biotic stresses (i.e., microbial pathogens) through regulation of sophisticated hormonal signaling networks (Fujita et al., 2006; Arnaud and Hwang, 2015). In my study, induced plant defenses triggered by foliar application of JA did not have a positive effect on plant growth in all treatments and all species. This is probably related to the functions of these hormones; SA-induced resistance targets microbial pathogens (Reymond and Farmer, 1998), while JA-induced resistance targets herbivores and necrotrophic pathogens (Nahar et al., 2011).

For *J. vulgaris*, we then continued by studying the effect of the live soil and SA application during four subsequent generations. The negative effect of the live soil was observed in all generations but did not increase or decline over time (Chapter 2). The mitigating effect of SA on the negative effects of the live soil on plant growth also did not change over generations. The reduced plant growth in live soils can be caused by nutrient competition between plants and soil microbes or by an overall pathogenic effect of soil microbial community (Hodge et al., 2013; Cesarano et al., 2017; Trivedi et al., 2020). However, in our experiment, we fertilized the plants and hence we expect that competition for nutrients was not important and that an overall microbial pathogenic effect is the most likely explanation for the plant growth reduction in live soil. Further, we hypothesized that we can select for a more beneficial community over time because the foliar application of SA mitigates the
negative live soil effect in *J. vulgaris* but we did not observe that the mitigating effect increased over generations.

Little is known about how such hormonal pathways affect the inoculated live soils and how this, in turn, impacts plant growth. Several studies argue that the 'SAmitigated effect' can be due to (1) a boosted immune system in the plant itself (Chen et al., 2020; Koo et al., 2020), or (2) changes in the plant-microbes interaction (Nishad et al., 2020; Kumar 2020). However, evidence for the second hypothesis in soil environments is still contradictory. For instance, Berendsen et al. (2012) and Doornbos et al. (2011) demonstrated that activation of JA and SA signaling pathways did not affect the resident soil microflora, while a recent study showed that SA modulates colonization of the root microbiome by specific bacterial taxa (Lebeis et al., 2015). In our study, JA application did not affect the relationship between the soil microbial community and plant growth, and this indicates that in our experiments there was no strong cross-talk between SA and JA.

#### SA-induced defenses and soil microbial composition

Based on the results of Chapter 2 we studied if the SA-mitigated effect on the soil microbial community was accompanied by a shift in the composition of the microbial community. In Chapter 3, we studied the composition of the rhizosphere microbial community of *J. vulgaris* over four generations. We found that the composition of the soil microbial community in the rhizosphere soil changed across generations, but not in a consistent manner. This may have resulted from the experimental design that we selected. For each generation, we used an inoculum, which means that we placed a subset of the microbial community in a sterile background. This may explain why we saw so much variation temporally, as in each generation a different subset of the microbial community may have been activated.

Although we did find an overall effect of SA on the total microbial composition, the direction of these changes was different in each generation. Application of SA selected for different bacterial genera in the rhizosphere soil, but these selected genera differed from generation to generation. This suggests that the effects of SA application to plants on the soil microbial community are not consistent over time. It is also

possible that bacterial microbial community composition is variable over time (Gilbert et al., 2009; Hickey et al., 2013; Lauber et al., 2013; Hannula et al., 2019). The impact of SA-induced resistance on soil microbial communities is still debated. For example, Hein et al. (2008) found that SA-induced resistance in *Arabidopsis* mutants changed the structure of bacterial communities in the rhizosphere. Wang et al. (2015) and Doornbos et al. (2011) both demonstrated that activation of SA-induced resistance did not significantly affect the composition and diversity of the rhizosphere bacterial community. As the SA effect on the microbial composition varied from generation to generation, it is difficult to predict the effects of activation of plant defenses on soil microbes. This may also explain why there was no selection for more beneficial communities over generations.

Application of SA to plants significantly up-regulated genera of *Caballeronia*, unclassified *Cytophagaceae*, *Crinalium* and *Candidatus Thermofonsia Clade* 2, and down-regulated the genera of *Thermomicrobiales*, unclassified *Rhodobacterales*, *Paracoccus* and *Flavihumibacter*. While the functions of many of these bacteria are poorly understood, bacteria of the genus *Caballeronia* are often reported to play an important role in fixing nitrogen and promoting plant growth, and species in this genus are predominantly endophytic diazotrophic bacteria and N-fixing bacteria (Puri et al., 2018; Padda et al., 2018; Puri et al., 2020). Hence, this suggests that activation of the SA signaling pathway in *J. vulgaris* plants may select for bacterial genera that are beneficial to the plant.

### SA-induced defenses and soil microbial functional genes

In Chapter 3, we analyzed the changes of microbial taxonomy in the rhizosphere soil and found that the effects of SA on the rhizosphere bacterial communities of J. *vulgaris* were inconsistent over generations. We hypothesized that we would see a common functional gene expression in the same soil samples, because the functions of the soil microbial community are often distributed across microbial taxa (Burke et al., 2011; Liu et al., 2018; Liu et al., 2020). One of the explanations can be that the composition of the soil microbial community shows a great redundancy concerning the functioning of microbial species and that changes in microbial diversity are not always consistent with changes in functional gene expression in soil microbial communities.

In our study, we found that the functional genes of rhizosphere microbial communities of *J. vulgaris* were affected by the SA treatment, by generation and by the interplay between SA treatment and generation. However, none of the significantly SA-downregulated genes was present in all four generations, while only one SA-upregulated gene was observed in all four generations. To date, information about the effects of phytohormone application to plants on the functions of rhizosphere microbiomes are limited (Anderson et al., 2004; Carvalhais et al., 2013). To our knowledge, the work presented in this thesis is among the first to study how activation of SA induced resistance affects natural soil microbiomes at the functional gene level. SA induced resistance is often reported to play an important role in resistance to a broad range of microbial pathogens, such as bacteria, fungi and viruses. Concerning viruses, SA has been reported to act as an elicitor in various plant species, such as tobacco, cucumber, Vigna mungo, tomato, sugarcane (Murphy et al., 1999; Gilliland et al., 2003; Mayers et al., 2005; Kundu et al., 2011; Li et al., 2019; Yuan et al., 2019).

A limitation in the current work is that not all detected genes could be annotated with known functions. Interestingly, at a gene ontology level, we found that soil microbial communities in the rhizosphere soil of SA-treated plants utilized several gene ontology processes. For the increased GO terms, they were mostly related to viral RNA genome replication, to interactions with host cells, to organelles of the host cells and to RNA polymerase activities; while for the decreased GO terms, they were associated with processing nitrogen and macromolecules. However, it still remains unproven that if those processes are associated with infection processes of the host plant and are potentially linked to suppression of pathogenic infections. Interestingly, up-regulated GO terms that were involved in viral (RNA) genome replication and viral processes were frequently found in our study in the soil of SA-treated plants. As it is well-reported that viral-phage therapy uses viruses or bacteriophages to control pathogens. A viral phage first attaches to the surface of a pathogenic bacteria, then injects its genome into the cells, self-replicates in the bacteria, and eventually kills the bacteria by causing them to burst or lyse (Duckworth and Gulig, 2002; Svircev et al., 2018; Jamal et al., 2019; Kortright et al., 2019; Rehman et al., 2019). This has recently 181

been brought up as an alternative for the usage of pesticides to control bacterial pathogens in agriculture (Rehman et al., 2019). Therefore, it is important to note that virus-microbe-plant interactions should be taken into.account in future studies.

### Plant growth stages and negative plant-soil effects

In our experiments, we placed a subset of the microbial community in a sterile background and this may have led temporal variation in the soil microbial community in each generation. Most studies on plant-soil-interactions have examined the effect of the soil microbial community on plant mass after a fixed duration of plant growth (Smith and Reynolds, 2012; Hodge and Fitter 2013; Dudenhöffer et al., 2018). However, these interactions may change over time (Bezemer et al., 2018). In Chapter 5, we examined how long-lasting the effect of the soil microbial community on plant growth is and we established relative plant growth rates at different growth stages (early, mid and late plant growth).

We found in all experiments that we carried out to study these temporal effects, that differences in dry plant mass between the plants grown in sterilized soil and inoculated soil (live soil) increased over the course of the experiment. Interestingly, linear regression models with *ln* transformed dry plant mass against time at the early stage and later stage in sterilized soil and live soil, respectively, showed that the relative growth rate of plants in the sterilized soil and live soil only differed in the first weeks and that there were no significant differences in relative growth rates during the late stage. Our study exemplifies that the negative effects of soil inoculation on plant mass can extend over the whole growth period, but that these differences are due to negative effects that occur in the first weeks after planting. This might be because younger plants or seedlings are more vulnerable and susceptible to pathogenic microbes in the soil than older plants with well-developed root systems (Packer and Clay, 2000). Root development plays an important role for plants in suppressing soil microbial pathogens (Watt et al., 2006; Emmett et al., 2014), and is often correlated with soil abiotic or biotic characteristics (Kardol et al., 2013; Arrigoni et al., 2018; Bezemer et al., 2018). Our findings are in line with previous work (Bezemer et al., 2018) on the same plant species, J. vulgaris, where seedlings were planted again in soil that had been conditioned by other plants of the same species. 182

Their findings show that the differences between responses of young and old plants are likely related to the sensitivity of plant stages and not due to temporal changes in the soil community.

### Concluding remarks and future perspective

The outcomes of this thesis contribute to our understanding of how harnessing of the plant immune system affects the relationship between plants and the soil microbial community. From this work, we can conclude that the effect of live soil on plant growth is species-specific. Moreover, we conclude that application of SA can mitigate the negative effect of live soil on plant growth and we hypothesize that the negative effect of live soil on plant performance is driven by microbial pathogens in the soil. Further, from the multi-generational experiment, we conclude that activation of SAassociated plant defense pathways alters the composition of soil microbial communities of J. vulgaris but that these effects vary over time. We found no evidence that activation of SA signaling pathways in plants results in the selection of bacteria that are more beneficial to plant growth. The functions of the majority of the significantly affected genera by SA-induced resistance in our experiment are not wellknown. SA-induced resistance, against soil microbial pathogens in J. vulgaris may be through the regulation of virus or viral related pathways. Last but not least, we concluded that negative effects of live soil on plant growth may appear consistent over time, but may only be caused by negative effects on plant growth that occur during the first few weeks. Overall, our study exemplifies that aboveground induction of plant defenses, can lead to complex above-belowground feedbacks.

#### References

- Anderson, J.P., Badruzsaufari, E., Schenk, P.M., Manners, J.M., Desmond, O.J., Ehlert, C., Maclean, D.J., Ebert, P.R. and Kazan, K., 2004. Antagonistic interaction between abscisic acid and jasmonate-ethylene signaling pathways modulates defense gene expression and disease resistance in *Arabidopsis. The Plant Cell*, 16(12), pp.3460-3479.
- Arnaud, D. and Hwang, I., 2015. A sophisticated network of signaling pathways regulates stomatal defenses to bacterial pathogens. *Molecular plant*, 8(4), pp.566-581.
- Arrigoni, E., Antonielli, L., Pindo, M., Pertot, I. and Perazzolli, M., 2018. Tissue age and plant genotype affect the microbiota of apple and pear bark. *Microbiological research*, 211, pp.57-68.
- Bailly, A., 2020. How plants might recognize rhizospheric bacterial volatiles. In *Bacterial Volatile* Compounds as Mediators of Airborne Interactions (pp. 139-165). Springer, Singapore.
- Berendsen, R.L., Pieterse, C.M. and Bakker, P.A., 2012. The rhizosphere microbiome and plant health. *Trends in plant science*, *17*(8), pp.478-486.
- Berens, M.L., Wolinska, K.W., Spaepen, S., Ziegler, J., Nobori, T., Nair, A., Krüler, V., Winkelmüller, T.M., Wang, Y., Mine, A. and Becker, D., 2019. Balancing trade-offs between biotic and abiotic stress responses through leaf age-dependent variation in stress hormone crosstalk. *Proceedings of the National Academy of Sciences, U.S.A, 116*(6), pp.2364-2373.
- Bever, J.D., 1994. Feeback between plants and their soil communities in an old field community. *Ecology*, 75(7), pp.1965-1977.
- Bever, J.D., Dickie, I.A., Facelli, E., Facelli, J.M., Klironomos, J., Moora, M., Rillig, M.C., Stock, W.D., Tibbett, M. and Zobel, M., 2010. Rooting theories of plant community ecology in microbial interactions. *Trends in ecology & evolution*, 25(8), pp.468-478.
- Bezemer, T.M., Jing, J., Bakx-Schotman, J.T. and Bijleveld, E.J., 2018. Plant competition alters the temporal dynamics of plant-soil feedbacks. *Journal of Ecology*, 106(6), pp.2287-2300.
- Bhattacharyya, P.N. and Jha, D.K., 2012. Plant growth-promoting rhizobacteria (PGPR): emergence in agriculture. *World Journal of Microbiology and Biotechnology*, 28(4), pp.1327-1350.
- Burke, C., Steinberg, P., Rusch, D., Kjelleberg, S. and Thomas, T., 2011. Bacterial community assembly based on functional genes rather than species. *Proceedings of the National Academy of Sciences*, U.S.A, 108(34), pp.14288-14293.
- Carvalhais, L.C., Dennis, P.G., Tyson, G.W. and Schenk, P.M., 2013. Rhizosphere metatranscriptomics: challenges and opportunities. *Molecular microbial ecology of the rhizosphere*, 2, pp.1137-1144.
- Cesarano, G., De Filippis, F., La Storia, A., Scala, F. and Bonanomi, G., 2017. Organic amendment type and application frequency affect crop yields, soil fertility and microbiome composition. *Applied Soil Ecology*, *120*, pp.254-264.
- Chen, J., Clinton, M., Qi, G., Wang, D., Liu, F. and Fu, Z.Q., 2020. Reprogramming and remodeling: transcriptional and epigenetic regulation of salicylic acid-mediated plant defense. *Journal of Experimental Botany*, 71(17), pp.5256-5268.
- Delory, B.M., Delaplace, P., Fauconnier, M.L. and Du Jardin, P., 2016. Root-emitted volatile organic compounds: can they mediate belowground plant-plant interactions? *Plant and Soil*, 402(1-2), pp.1-26.
- Doornbos, R.F., van Loon, L.C. and Bakker, P.A., 2012. Impact of root exudates and plant defense signaling on bacterial communities in the rhizosphere. A review. *Agronomy for Sustainable Development*, 32(1), pp.227-243.

- Dror, B., Jurkevitch, E. and Cytryn, E., 2020. State-of-the-art methodologies to identify antimicrobial secondary metabolites in soil bacterial communities-A review. *Soil Biology and Biochemistry*, p.107838.
- Duckworth, D.H. and Gulig, P.A., 2002. Bacteriophages. BioDrugs, 16(1), pp.57-62.
- Dudenhöffer, J.H., Ebeling, A., Klein, A.M. and Wagg, C., 2018. Beyond biomass: Soil feedbacks are transient over plant life stages and alter fitness. *Journal of Ecology*, *106*(1), pp.230-241.
- Emmett, B., Nelson, E.B., Kessler, A. and Bauerle, T.L., 2014. Fine-root system development and susceptibility to pathogen colonization. *Planta*, 239(2), pp.325-340.
- Fujita, M., Fujita, Y., Noutoshi, Y., Takahashi, F., Narusaka, Y., Yamaguchi-Shinozaki, K. and Shinozaki, K., 2006. Crosstalk between abiotic and biotic stress responses: a current view from the points of convergence in the stress signaling networks. *Current opinion in plant biology*, 9(4), pp.436-442.
- Gilbert, J.A., Field, D., Huang, Y., Edwards, R., Li, W., Gilna, P. and Joint, I., 2008. Detection of large numbers of novel sequences in the metatranscriptomes of complex marine microbial communities. *PloS one*, 3(8), p.e3042.
- Gilliland, A., Singh, D.P., Hayward, J.M., Moore, C.A., Murphy, A.M., York, C.J., Slator, J. and Carr, J.P., 2003. Genetic modification of alternative respiration has differential effects on antimycin A-induced versus salicylic acid-induced resistance to *Tobacco mosaic* virus. *Plant Physiology*, 132(3), pp.1518-1528.
- Hahl, T., van Moorsel, S.J., Schmid, M.W., Zuppinger-Dingley, D., Schmid, B. and Wagg, C., 2020. Plant responses to diversity-driven selection and associated rhizosphere microbial communities. *Functional Ecology*, 34(3), pp.707-722.
- Haney, C.H., Samuel, B.S., Bush, J. and Ausubel, F.M., 2015. Associations with rhizosphere bacteria can confer an adaptive advantage to plants. *Nature plants*, *1*(6), pp.1-9.
- Hannula, S.E., Kielak, A.M., Steinauer, K., Huberty, M., Jongen, R., Jonathan, R., Heinen, R. and Bezemer, T.M., 2019. Time after time: temporal variation in the effects of grass and forb species on soil bacterial and fungal communities. *MBio*, 10(6).
- Harrison, K.A. and Bardgett, R.D., 2010. Influence of plant species and soil conditions on plant-soil feedback in mixed grassland communities. *Journal of Ecology*, 98(2), pp.384-395.
- Hein, J.W., Wolfe, G.V. and Blee, K.A., 2008. Comparison of rhizosphere bacterial communities in *Arabidopsis thaliana* mutants for systemic acquired resistance. *Microbial ecology*, 55(2), pp.333-343.
- Hickey, R.J., Abdo, Z., Zhou, X., Nemeth, K., Hansmann, M., Osborn III, T.W., Wang, F. and Forney, L.J., 2013. Effects of tampons and menses on the composition and diversity of vaginal microbial communities over time. *BJOG: An International Journal of Obstetrics & Gynaecology*, 120(6), pp.695-706.
- Hodge, A. and Fitter, A.H., 2013. Microbial mediation of plant competition and community structure. *Functional Ecology*, 27(4), pp.865-875.
- Hu, Y., Zhang, Z., Huang, L., Qi, Q., Liu, L., Zhao, Y., Wang, Z., Zhou, H., Lv, X., Mao, Z. and Yang, Y., 2019. Shifts in soil microbial community functional gene structure across a 61-year desert revegetation chronosequence. *Geoderma*, 347, pp.126-134.
- Jamal, M., Bukhari, S.M., Andleeb, S., Ali, M., Raza, S., Nawaz, M.A., Hussain, T., Rahman, S.U. and Shah, S.S., 2019. *Bacteriophages*: an overview of the control strategies against multiple bacterial infections in different fields. *Journal of basic microbiology*, 59(2), pp.123-133.
- Joosten, L., Mulder, P.P., Klinkhamer, P.G. and van Veen, J.A., 2009. Soil-borne microorganisms and soil-type affect pyrrolizidine alkaloids in *Jacobaea vulgaris*. *Plant and Soil*, 325(1-2), p.133.

- Kardol, P., De Deyn, G.B., Laliberté, E., Mariotte, P. and Hawkes, C.V., 2013. Biotic plant-soil feedbacks across temporal scales. *Journal of Ecology*, 101(2), pp.309-315.
- Kirk, H., Vrieling, K., Van Der Meijden, E. and Klinkhamer, P.G., 2010. Species by environment interactions affect pyrrolizidine alkaloid expression in *Senecio jacobaea*, *Senecio aquaticus*, and their hybrids. *Journal of chemical ecology*, 36(4), pp.378-387. '
- Klironomos, J.N., 2002. Feedback with soil biota contributes to plant rarity and invasiveness in communities. *Nature*, 417(6884), pp.67-70.
- Koo, Y.M., Heo, A.Y. and Choi, H.W., 2020. Salicylic acid as a safe plant protector and growth regulator. *The Plant Pathology Journal*, *36*(1), p.1.
- Kortright, K.E., Chan, B.K., Koff, J.L. and Turner, P.E., 2019. Phage therapy: a renewed approach to combat antibiotic-resistant bacteria. *Cell host & microbe*, 25(2), pp.219-232.
- Kostenko, O., van de Voorde, T.F., Mulder, P.P., van der Putten, W.H. and Martijn Bezemer, T., 2012. Legacy effects of aboveground-belowground interactions. *Ecology Letters*, *15*(8), pp.813-821.
- Kowalchuk, G.A., Hol, W.G. and Van Veen, J.A., 2006. Rhizosphere fungal communities are influenced by *Senecio jacobaea* pyrrolizidine alkaloid content and composition. *Soil Biology and Biochemistry*, 38(9), pp.2852-2859.
- Kulmatiski, A., Beard, K.H., Stevens, J.R. and Cobbold, S.M., 2008. Plant-soil feedbacks: a metaanalytical review. *Ecology letters*, 11(9), pp.980-992.
- Kumar, D., 2020. Plant immune response strategies against pathogens. *Plant Archives*, 20(1), pp.1169-1174.
- Kundu, S., Chakraborty, D. and Pal, A., 2011. Proteomic analysis of salicylic acid induced resistance to *Mungbean yellow mosaic India* virus in Vigna mungo. *Journal of proteomics*, 74(3), pp.337-349.
- Lauber, C.L., Ramirez, K.S., Aanderud, Z., Lennon, J. and Fierer, N., 2013. Temporal variability in soil microbial communities across land-use types. *The ISME journal*, 7(8), pp.1641-1650.
- Lebeis, S.L., Paredes, S.H., Lundberg, D.S., Breakfield, N., Gehring, J., McDonald, M., Malfatti, S., Del Rio, T.G., Jones, C.D., Tringe, S.G. and Dangl, J.L., 2015. Salicylic acid modulates colonization of the root microbiome by specific bacterial taxa. *Science*, 349(6250), pp.860-864.
- Li, T., Huang, Y., Xu, Z.S., Wang, F. and Xiong, A.S., 2019. Salicylic acid-induced differential resistance to the Tomato yellow leaf curl virus among resistant and susceptible tomato cultivars. *BMC plant biology*, 19(1), pp.1-14.
- Liu, Y.R., Delgado-Baquerizo, M., Bi, L., Zhu, J. and He, J.Z., 2018. Consistent responses of soil microbial taxonomic and functional attributes to mercury pollution across China. *Microbiome*, 6(1), pp.1-12.
- Liu, Y.R., Delgado-Baquerizo, M., Yang, Z., Feng, J., Zhu, J. and Huang, Q., 2020. Microbial taxonomic and functional attributes consistently predict soil CO2 emissions across contrasting croplands. *Science of The Total Environment*, 702, p.134885.
- Maurhofer, M., Reimmann, C., Schmidli-Sacherer, P., Heeb, S., Haas, D. and Défago, G., 1998. Salicylic acid biosynthetic genes expressed in *Pseudomonas fluorescens* strain P3 improve the induction of systemic resistance in tobacco against tobacco necrosis virus. *Phytopathology*, 88(7), pp.678-684.
- Massalha, H., Korenblum, E., Tholl, D. and Aharoni, A., 2017. Small molecules below-ground: the role of specialized metabolites in the rhizosphere. *The plant journal*, *90*(4), pp.788-807.
- Mayers, C.N., Lee, K.C., Moore, C.A., Wong, S.M. and Carr, J.P., 2005. Salicylic acid-induced resistance to *Cucumber mosaic* virus in squash and *Arabidopsis thaliana*: contrasting mechanisms of induction and antiviral action. *Molecular plant-microbe interactions*, 18(5), pp.428-434.

- Miki, T., 2012. Microbe-mediated plant-soil feedback and its roles in a changing world. *Ecological Research*, 27(3), pp.509-520.
- Murphy, A.M., Chivasa, S., Singh, D.P. and Carr, J.P., 1999. Salicylic acid-induced resistance to viruses and other pathogens: a parting of the ways? *Trends in plant science*, *4*(4), pp.155-160.
- Nahar, K., Kyndt, T., De Vleesschauwer, D., Höfte, M. and Gheysen, G., 2011. The jasmonate pathway is a key player in systemically induced defense against root knot nematodes in rice. *Plant physiology*, 157(1), pp.305-316.
- Nijjer, S., Rogers, W.E. and Siemann, E., 2007. Negative plant-soil feedbacks may limit persistence of an invasive tree due to rapid accumulation of soil pathogens. *Proceedings of the Royal Society B: Biological Sciences*, 274(1625), pp.2621-2627.
- Nishad, R., Ahmed, T., Rahman, V.J. and Kareem, A., 2020. Modulation of plant defense system in response to microbial interactions. *Frontiers in Microbiology*, 11, p.1298.
- Padda, K.P., Puri, A. and Chanway, C.P., 2018. Isolation and identification of endophytic diazotrophs from lodgepole pine trees growing at unreclaimed gravel mining pits in central interior British Columbia, Canada. *Canadian Journal of Forest Research*, 48(12), pp.1601-1606.
- Packer, A. and Clay, K., 2000. Soil pathogens and spatial patterns of seedling mortality in a temperate tree. *Nature*, 404(6775), pp.278-281.
- Puri, A., Padda, K.P. and Chanway, C.P., 2018. Evidence of endophytic diazotrophic bacteria in lodgepole pine and hybrid white spruce trees growing in soils with different nutrient statuses in the West Chilcotin region of British Columbia, Canada. *Forest Ecology and Management*, 430, pp.558-565.
- Puri, A., Padda, K.P. and Chanway, C.P., 2020. Can naturally-occurring endophytic nitrogen-fixing bacteria of hybrid white spruce sustain boreal forest tree growth on extremely nutrient-poor soils?. Soil Biology and Biochemistry, 140, p.107642.
- Rasmann, S., Hiltpold, I. and Ali, J., 2012. The role of root-produced volatile secondary metabolites in mediating soil interactions. In *Advances in selected plant physiology aspects*. IntechOpen.
- Rehman, S., Ali, Z., Khan, M., Bostan, N. and Naseem, S., 2019. The dawn of phage therapy. *Reviews in medical virology*, 29(4), p.e2041.
- Reymond, P. and Farmer, E.E., 1998. Jasmonate and salicylate as global signals for defense gene expression. *Current opinion in plant biology*, *1*(5), pp.404-411.
- Shen, C., Shi, Y., Ni, Y., Deng, Y., Van Nostrand, J.D., He, Z., Zhou, J. and Chu, H., 2016. Dramatic increases of soil microbial functional gene diversity at the treeline ecotone of Changbai Mountain. *Frontiers in microbiology*, 7, p.1184.
- Smith, L.M. and Reynolds, H.L., 2012. Positive plant-soil feedback may drive dominance of a woodland invader, *Euonymus fortunei*. *Plant Ecology*, 213(5), pp.853-860.
- Smith, M.E., Delean, S., Cavagnaro, T.R. and Facelli, J.M., 2018. Evidence for species-specific plant responses to soil microbial communities from remnant and degraded land provides promise for restoration. *Austral Ecology*, 43(3), pp.301-308.
- Sun, X.G. and Tang, M., 2013. Effect of arbuscular mycorrhizal fungi inoculation on root traits and root volatile organic compound emissions of *Sorghum* bicolor. *South African Journal of Botany*, 88, pp.373-379.
- Svircev, A., Roach, D. and Castle, A., 2018. Framing the future with bacteriophages in agriculture. *Viruses*, 10(5), p.218.
- Trivedi, P., Leach, J.E., Tringe, S.G., Sa, T. and Singh, B.K., 2020. Plant-microbiome interactions: from community assembly to plant health. *Nature Reviews Microbiology*, pp.1-15.

- Tscherko, D., Rustemeier, J., Richter, A., Wanek, W. and Kandeler, E., 2003. Functional diversity of the soil microflora in primary succession across two glacier forelands in the Central Alps. *European Journal of Soil Science*, 54(4), pp.685-696.
- Van Der Heijden, M.G., Bardgett, R.D. and Van Straalen, N.M., 2008. The unseen majority: soil microbes as drivers of plant diversity and productivity in terrestrial ecosystems. *Ecology letters*, 11(3), pp.296-310.
- Vos, I.A., Pieterse, C.M. and Van Wees, S.C., 2013. Costs and benefits of hormone-regulated plant defences. *Plant Pathology*, 62, pp.43-55.
- Wang, X., Sager, R., Cui, W., Zhang, C., Lu, H. and Lee, J.Y., 2013. Salicylic acid regulates plasmodesmata closure during innate immune responses in *Arabidopsis*. *The Plant Cell*, 25(6), pp.2315-2329.
- Wang, B., Guo, X.W., Li, K., Han, X., Xu, S.J., Liu, Z.D., Guo, Y.S. and Xie, H.G., 2015. Effects of salicylic acid on grape plants and the soil microbial community. *Allelopathy Journal*, 36(1).
- Wang, M., Ruan, W., Kostenko, O., Carvalho, S., Hannula, S.E., Mulder, P.P., Bu, F., van der Putten, W.H. and Bezemer, T.M., 2019. Removal of soil biota alters soil feedback effects on plant growth and defense chemistry. *New Phytologist*, 221(3), pp.1478-1491.
- Watt, M., Silk, W.K. and Passioura, J.B., 2006. Rates of root and organism growth, soil conditions, and temporal and spatial development of the rhizosphere. *Annals of botany*, 97(5), pp.839-855.
- Yang, F., Zhang, Y., Huang, Q., Yin, G., Pennerman, K.K., Yu, J., Liu, Z., Li, D. and Guo, A., 2015. Analysis of key genes of jasmonic acid mediated signal pathway for defense against insect damages by comparative transcriptome sequencing. *Scientific reports*, 5, p.16500.
- Yuan, W., Jiang, T., Du, K., Chen, H., Cao, Y., Xie, J., Li, M., Carr, J.P., Wu, B., Fan, Z. and Zhou, T., 2019. Maize phenylalanine ammonia-lyases contribute to resistance to *Sugarcane mosaic* virus infection, most likely through positive regulation of salicylic acid accumulation. *Molecular plant pathology*, 20(10), pp.1365-1378.
- Zhu, X., Mao, L. and Chen, B., 2019. Driving forces linking microbial community structure and functions to enhanced carbon stability in biochar-amended soil. *Environment international*, 133, p.105211.

### Summary

Many plant species grow better in sterilized soil than in soil that contains a live microbial community. One hypothesis to explain this phenomenon is that the overall net pathogenic effect of soil microbial communities reduces plant performance. Induced plant defenses triggered by the application of the plant hormones jasmonic acid (JA) and salicylic acid (SA) may help to mitigate this pathogenic effect. However, little is known about how the activation of SA-induced resistance impacts the microbial composition and the expression of functional genes in the rhizosphere soil.

We manipulated and induced the plant defense system through foliar application of phytohormones (JA or SA), and examined whether the negative effect of live soil on plant growth was reduced. The growth of four plant species (Jacobaea vulgaris, *Cirsium vulgare, Trifolium repens* and *Daucus carota*) was affected differently in live soil and by the hormone treatments. Foliar application of SA increased plant growth in live soil for the species, J. vulgaris and C. vulgare, which were the two species that both produced less biomass in live soil than in sterilized soil, SA application slightly reduced plant growth in live soils for the species T. repens and D. carota that were not affected in live soil. Application of JA reduced plant growth in live and sterile soil for all species. For J. vulgaris the treatments were repeated for three more generations. In each generation, the live soil consisted of a mixture of 10% of soil collected from pots from the previous generation mixed with 90% sterilized soil. In all four generations, plant biomass was measured. The reduction in growth in live soil was consistent in each generation, and in each generation, this negative effect was mitigated by the application of SA to plants. Hence, we found no evidence for an increase in the negative plant-soil feedback over generations, but also no selection effect of SA application over time.

RNA extracted from the rhizosphere soil from each generation was subsequently sequenced. Soil microbial composition at genus level was studied and the expression of functional genes of live soils where plants grown under SA treatments and control were compared. Application of SA to *J. vulgaris* leaves altered the composition of bacterial communities in the rhizosphere soil but only in the second, third and fourth

growth cycle. However, the SA effects on bacterial community composition were small, while there was a substantial temporal effect on rhizosphere bacterial composition. As there were no genera of bacteria that responded to SA application in the first generation this suggests that there are no immediate responses of bacteria in the rhizosphere to SA application to plants.

Subsequently, the effects of the application of SA to *J. vulgaris* on the gene expression and functions of the soil-borne microbial community were examined for each of the four plant generations. Gene expression and functions of the soil-borne microbial community responded to the exogenous application of SA but these effects differed per generation. The number of differentially expressed genes tended to increase over generations, but remarkably there was no overlap for these annotated genes among the four generations. Moreover, we found that foliar application of SA upregulated GO terms of biological processes that were related to viral RNA genome replication, to interactions with host cells, to organelles of the host cells and to RNA polymerase activities. There were six GO terms of which the expression decreased in the second, third and fourth generation, and these were associated with processing nitrogen and macromolecules.

Finally, in a series of experiments, we examined for J. vulgaris, how plant responses to live soil changed over time, by repeatedly harvesting plants over time. In all experiments, plant growth was worse in live soil than in sterilized soil and this effect on plant biomass was consistent over time. However, relative growth rates of plants in the sterilized soil and live soil only differed for young plants and a reverse pattern was even observed during the latest stage where relative growth rates were higher for plants in live soil. This shows that while the soil treatment may result in plant biomass being consistently lower, this could have been caused solely by initial effects of the treatment on plant growth. Hence, to better understand plant-soil interactions, it is important to examine not only plant biomass but also plant growth rates. In a third growth experiment, we also examined the effect of the timing of soil inoculation prior to planting on the relative growth rate of J. vulgaris plants with four different timing treatments. Biomass was reduced in all inoculated soils and there was a negative relationship between time since inoculation and plant biomass. Again, in all inoculated soils the negative effect of the soil microbial community on plant growth 190

disappeared two weeks after planting. Overall, these results suggest that young plants or seedlings are most sensitive to soil pathogens.

In conclusion, our research shows that aboveground activation of defenses in the plant affects soil microbial communities and as soil microbes can greatly influence plant performance, this implies that induction of plant defenses, can lead to complex abovebelowground feedbacks.

# Nederlandse samenvatting

Veel plantensoorten groeien beter in gesteriliseerde grond dan in grond met een levende microbiële gemeenschap. Een hypothese om dit fenomeen te verklaren is dat het algehele netto pathogene effect van microbiële gemeenschappen in de bodem de groei van planten vermindert. Dit pathogene effect kan in theorie verminderd worden door de afweer van planten te induceren met de plantenhormonen jasmonzuur (JA) en salicylzuur (SA). Er is echter weinig bekend over hoe de activering van SA-geïnduceerde resistentie de microbiële samenstelling en de expressie van functionele genen van bacteriën in de rhizosfeerbodem beïnvloedt.

Ik manipuleerde en induceerde de afweer van planten door deze te behandelen met plantenhormonen (JA of SA), en onderzocht of het negatieve effect van levende grond op plantengroei inderdaad verminderd was. De levende grond en de hormoonbehandelingen hadden een verschillend effect op de groei van vier plantensoorten (Jacobaea vulgaris, Cirsium vulgare, Trifolium repens en Daucus carota). Het behandelen van bladeren met SA verhoogde de plantengroei in levende grond voor J. vulgaris en C. vulgare ten opzichte van de controle in levende grond. Ook produceerden deze twee soorten minder biomassa in levende grond dan in gesteriliseerde grond. De SA behandeling verminderde de plantengroei in levende grond enigszins voor de soorten T. repens en D. carota terwijl hun groei niet verminderd was ten opzichte van de groei op gesteriliseerde grond. De behandeling met JA verminderde plantengroei in levende en steriele grond voor alle soorten. Alleen voor J. vulgaris werden de behandelingen voortgezet voor nog drie generaties. In elke generatie bestond de levende grond uit een mengsel van 10% grond, verzameld uit potten van de vorige generatie van dezelfde behandeling, gemengd met 90% gesteriliseerde grond. In alle vier de generaties werd de biomassa gemeten. De afname van de groei in levende grond ten opzichte van steriele grond was

consistent in elke generatie en in elke generatie werd dit negatieve effect verminderd door behandeling met SA. Ik vond geen bewijs voor een toename van de negatieve plant-bodem-terugkoppeling over generaties, maar ook geen selectie-effect van SA behandeling in opeenvolgende generaties.

Uit de rhizosfeerbodem van alle behandelingen en generaties van *J. vulgaris* werd RNA geëxtraheerd en gesequenced. De microbiële samenstelling van de bodem werd bestudeerd en de expressie van functionele genen in de SAbehandeling en controle werden vergeleken. Behandeling van *J. vulgaris* bladeren met SA veranderde de samenstelling van bacteriële gemeenschappen in de rhizosfeerbodem, echter alleen in de tweede, derde en vierde generatie. Het effect van de SA behandeling op de samenstelling van de bacteriële gemeenschap was echter klein, terwijl de bacteriële gemeenschap sterk verschilde tussen generaties. Aangezien er in de eerste generatie geen bacteriegenera waren die reageerden op de SA behandeling van bladeren van *J. vulgaris*, suggereert dit dat er geen onmiddellijke reacties zijn van bacteriën in de rhizosfeer op SA-toediening op planten.

Vervolgens werden de effecten van de SA behandeling van bladeren van *J. vulgaris* op de genexpressie en functies van de microbiële gemeenschap in de rhizosfeer onderzocht voor elk van de vier generaties. De exogene toepassing van SA beïnvloedde genexpressie en functies van de microbiële gemeenschap, maar deze effecten verschilden per generatie. Het aantal differentieel tot expressie gebrachte geannoteerde genen nam over generaties toe, maar opmerkelijk genoeg was er geen overlap voor deze genen tussen de vier generaties. Bovendien ontdekte ik dat toediening van SA op het blad de biologische processen die verband hielden met virale RNA-genoomreplicatie, met interacties met gastheercellen, met organellen van de gastheercellen en met RNA-polymerase-activiteiten beïnvloedde. Er waren zes GO-termen waarvan de expressie afnam in de tweede, derde en vierde generatie, en deze waren gerelateerd aan het verwerken van stikstof en macromoleculen. Ten slotte heb ik in een reeks experimenten onderzocht hoe de relatieve groeisnelheid van J. vulgaris planten op levende grond in de loop van de tijd veranderde. In alle experimenten was de absolute plantengroei in levende grond slechter dan in gesteriliseerde grond en dit effect op de biomassa van planten was consistent in de tijd. De relatieve groeisnelheden van planten in de gesteriliseerde grond en levende grond verschilden alleen voor jonge planten en een omgekeerd patroon werd zelfs waargenomen tijdens de latere groeifase waarin de relatieve groeisnelheden hoger waren voor planten in levende grond. Hoewel een levende bodem ertoe kan leiden dat de plantenbiomassa constant lager is wordt dit dus uitsluitend veroorzaakt door een lagere relatieve groeisnelheid op levende bodems in de eerste weken. Om de interacties tussen plant en bodem beter te begrijpen, is het daarom belangrijk om niet alleen de biomassa van planten te onderzoeken, maar ook de relatieve groeisnelheden. In een derde groei-experiment onderzocht ik het effect van de timing van bodeminoculatie voorafgaand aan het planten van zaailingen in de bodem, op de relatieve groeisnelheid van J. vulgaris planten. In dit experiment waren er vier verschillende momenten van inoculatie van de bodem voor beplanting. Op alle geïnoculeerde bodems was de biomassa lager dan op steriele bodems en er was een negatief verband tussen de tijd sinds inoculatie en biomassa. In alle geïnoculeerde bodems verdween het negatieve effect van de microbiële bodemgemeenschap op de plantengroei echter al twee weken na het planten. Deze resultaten suggereren dat alleen jonge planten of zaailingen gevoelig zijn voor bodempathogenen.

Concluderend laat mijn onderzoek zien dat bovengrondse activering van afweermechanismen in de plant de microbiële gemeenschappen in de bodem beïnvloedt. Aangezien bodemmicroben de groei en chemie van planten sterk kunnen beïnvloeden, impliceert dit dat inductie van afweermechanismen van planten kan leiden tot complexe bovengrondse-ondergrondse terugkoppelingen.

### **Curriculum Vitae**

Jing Zhang was born in Jinan, Shandoing Province in China on 22nd, May, 1991 (28th April, 1990 on official documents). She finished her bachelor in Geographic Information Systems (GIS) in 2013 at Sichuan Agricultural University in China, then she obtained her MSc degree in Phytopathology in 2015 at the same university. In September 2015, she started her PhD project on "The impact of defense hormones on the interaction between plants and soil microbial community" under the supervision of Prof. Dr. Peter Klinkhamer, Prof. Dr. Martijn Bezemer and Dr. Klaas Vrieling in the group of Plant Sciences, at Leiden university, in the Netherlands.

# Acknowledgements

This dissertation of me is the output of the effort and support of a community of people to whom I feel extremely grateful.

I thank my promoters and supervisors Professor Peter Klinkhamer, Professor Martijn Bezemer and Dr. Klaas Vrieling. Thank you all for accepting me as your PhD student and helping me to overcome scientific challenges along the journey. I know without your help I will not go this far.

Many thanks to people from the group of Plant Ecology, Karin, Yangan, Tom de Jong, Sanae, Onno, Aneta, Gang, Marie, Martine, Xiaojie, Tiantian, Sofia, Xiangyu, Martina, Suzanne, Rick, Xianqin and Menno. It was great to know you all during the whole or part of the journey of my PhD life in PECO group, I appreciate the help you offered with my research.

I also want to thank people outside of our group who gave me valuable help for my research: Daniel, Robin, Niels, Maria, Michael, Bing, Paul, Chao, Joost, Yin, Alexander, Quanxiao, Ward and Jan.

I feel grateful to the people in IBL with whom I shared activities. Many thanks go to Luis, Gerardo, Lina, Martine, Hycelone, and Diana; I had fun with all of you during our trips to Germany, Luxemburg and Leiden liberation day. Thank also to Zheren, Shixiong, Sanne and Hein.

I thank my friends outside my research life. Weiya, you have been someone so important in my life and your way of thinking inspired me so much for enjoying life, as you said we are more than friends, we are family. Dami, I am so lucky to have you as a friend, you are a great listener and I always love it to share my happiness and sorrows with you. Sanae, I feel grateful that we become such close friends, you are a very sweet friend, and you are there for me whenever I need help. I am lucky to have you in my life. Xingxing, with you I believe friends can be so selfless and wish I go back soon and live in the life we described. Also, many thanks to Saskia, Judith, Connor, Casper, Marine, Jasmine, Alex, Merissa, Yufei, Xiaoqin, Quanchi, Qiang, Tingxian, Zhang Jiamei Laoshi, Xiaoyi, Yixin, Xuhan, Shujiang and Professor Zhu.

I love my parents, I thank them for adopting me, raising me up, offering me the best they can and setting me free to reach what I want for myself. Many thanks to my brother and sister-in-law for taking care of our parents when I am not there. Although, we do not communicate much, but we all know we love each other and thanks for the best gift ever in the world from my parents--- you give me a family.

## **Publications**

Zhan, J., K, Vrieling., P.G.L. Klinkhamer., T.M. Bezemer., 2021. Activation of hormone-associated plant defense pathways alters the effects of soil microbial communities on plant performance. *Under revision in Basic and applied ecology*.

Zhan, J., K, Vrieling., P.G.L. Klinkhamer., T.M. Bezemer., 2021. Activation of SA-associated plant defense pathway alters the composition of soil bacterial communities. *Submitted to Plant and soil*.

Zhan, J., K, Vrieling., P.G.L. Klinkhamer., T.M. Bezemer., 2021. Negative effects of soil microorganisms on plant growth only extend to the first weeks. *Under revision in Journal of plant ecology*.

<u>Zhang</u>, J. and Zhu, T.H., 2018. First report of *Colletotrichum boninense* causing leaf anthracnose on *Eucalyptus robusta* (Smith) in the upper reaches of Yangtze River. *Plant Disease*, *102*(7), pp.1446-1447.

Qiao, T.M., <u>Zhang, J.</u>, Li, S.J., Han, S. and Zhu, T.H., 2016. Development of nested PCR, multiplex PCR, and loop-mediated isothermal amplification assays for rapid detection of *Cylindrocladium scoparium* on *Eucalyptus*. *The plant pathology journal*, *32*(5), p.414.

Zheng, L., Peng, Y., <u>Zhang, J.</u>, Ma, W.J., Li, S.J. and Zhu, T.H., 2015. First report of *Fusarium solani* causing root rot of *Juglans sigllata* Dode in China. *Plant Disease*, 99(1), pp.159-159.

### Patents

Zhu TH, <u>Zhang J</u>, Zhu HMY, Li SJ, Qiao TM, Han S, Wang LM, Zhang LN. A detection kit for *Cylindrocladium scoparium* on *Eucalyptus* based on LAMP method. Patent number in China: CN104164486A

Zhu TH, Zhu HMY, Li SJ, <u>Zhang J</u>, Liu Y, Han S, Qiao TM, Zhang LN. A strain of *Bacillus amyloliquefaciens* and its liquid formulations application on controlling of *Cenangium ferruginosum*, Patent number in China: CN103555618A

Zhu TH, Yu Q, Li SJ, <u>Zhang J</u>, Han S, Qiao TM, Zhang LN, Zheng L. A Thiram-resistant mutant strain of *Streptomyces sampsonii* MV-2 and its liquid formulations. Patent number in China: CN103396961A

Zhu TH, Zhu HMY, Li SJ, <u>Zhang J</u>, Han S, Qiao TM, Zhang LN. A method of prevention and control of forest pest in fields. Patent number in China: CN103493811A