



Universiteit
Leiden
The Netherlands

Modulation of the inflammatory response following myocardial infarction

Pluijmert, N.J.

Citation

Pluijmert, N. J. (2021, June 3). *Modulation of the inflammatory response following myocardial infarction*. Retrieved from <https://hdl.handle.net/1887/3182529>

Version: Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/3182529>

Note: To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <https://hdl.handle.net/1887/3182529> holds various files of this Leiden University dissertation.

Author: Pluijmert, N.J.

Title: Modulation of the inflammatory response following myocardial infarction

Issue Date: 2021-06-03

2

Post-ischemic myocardial inflammatory response: a complex and dynamic process susceptible to immunomodulatory therapies (review)

Niek J. Pluijmert¹
Douwe E. Atsma¹
Paul H.A. Quax^{2,3}

¹ Department of Cardiology, Leiden University Medical Center, Leiden, The Netherlands

² Department of Surgery, Leiden University Medical Center, Leiden, The Netherlands

³ Einthoven Laboratory for Experimental Vascular Medicine, Leiden University Medical Center, Leiden, The Netherlands

Front Cardiovasc Med 2021;8:647785

ABSTRACT

Following acute occlusion of a coronary artery causing myocardial ischemia and implementing first-line treatment involving rapid reperfusion, a dynamic and balanced inflammatory response is initiated to repair and remove damaged cells. Paradoxically, restoration of myocardial blood flow exacerbates cell damage as a result of myocardial ischemia-reperfusion (MI-R) injury, which eventually provokes accelerated apoptosis. In the end the infarct size still corresponds to the subsequent risk of developing heart failure. Therefore, true understanding of the mechanisms regarding MI-R injury, and its contribution to cell damage and cell death, are of the utmost importance in the search for successful therapeutic interventions to finally prevent the onset of heart failure. This review focusses on the role of innate immunity, chemokines, cytokines, and inflammatory cells in all three overlapping phases following experimental, mainly murine, MI-R injury known as the inflammatory, reparative, and maturation phase. It provides a complete state-of-the-art overview including most current research of all post-ischemic processes and phases and additionally summarizes the use of immunomodulatory therapies translated into clinical practice.

INTRODUCTION

Ischemic myocardial injury causes decreased oxygen tension within the cell, subsequent degradation and loss of oxidative phosphorylation, and decreased generation of high-energy phosphates resulting in loss of membrane integrity¹. Irreversible cardiomyocyte injury, demonstrated by sarcolemmal disruption and presence of small amorphous densities in the mitochondria, develops after 20-40 minutes of sustained severe ischemia². Cardiomyocyte cell death causes release of cardiomyocyte-specific proteins, such as myoglobin, cardiac troponin (cTn) T and I, creatine kinase MB (CK-MB) and creatine phosphokinase (CPK), which are used clinically as markers for early detection³, especially by use of high-sensitivity cTn assays nowadays⁴. The dominant mechanism of cardiomyocyte death is coagulation necrosis, peaking after 12 hours up to 4 days, with cell swelling and rupture of cell membranes resulting in extrusion of intracellular contents after passing the point of no return even though reperfusion is applied at a certain moment. This triggers an extensive inflammatory reaction activating reparative pathways ultimately contributing to mature scar formation⁵. The other mechanism is apoptosis, peaking after 6 to 8 hours, concerning programmed cell death, in particular induced by reperfusion and also affecting non-infarcted areas of the LV wall and interventricular septum⁶.

Myocardial infarct repair comprises three overlapping phases: the inflammatory, reparative, and maturation phase. The inflammatory phase contains activation of chemokine and cytokine cascades resulting in recruitment of local infiltration of leukocytes into the infarct area. Dead cells and matrix debris are cleared by neutrophils and macrophages. During the reparative or proliferative phase, activated mononuclear cell and macrophage subpopulations release cytokines and growth factors that recruit and activate predominantly myofibroblasts and vascular cells, while pro-inflammatory mediators are suppressed. Activated myofibroblasts abundantly produce extracellular matrix proteins and an extensive microvascular network is formed with preservation of the structural integrity. In the maturation phase, reparative fibroblasts and vascular cells become apoptotic and a cross-linked collagen-based scar is formed⁷. This review addresses the post-ischemic inflammatory response in its entirety focusing on the role of innate immunity, chemokines, cytokines, and inflammatory cells in all three overlapping inflammatory, reparative, and maturation phases following experimental myocardial ischemia-reperfusion (MI-R) injury. The complexity of this process makes it amenable to pharmacologic interventions of which some has been successfully translated towards clinical practice as discussed in the final section.

MYOCARDIAL ISCHEMIA-REPERFUSION INJURY

As mentioned previously, early reperfusion by primary percutaneous coronary intervention (pPCI) is the recommended therapy in acute myocardial infarction (MI) since reperfusion restores oxygen and nutrient supply accentuating the post-ischemic inflammatory response

and accelerating wound healing, making it a prerequisite for cardiomyocyte salvage⁸. Even late reperfusion showed to be beneficial in humans, where it results in considerable myocardial salvage⁹, as well as in animals, exhibiting permanent reduction of infarct expansion and ventricular remodeling¹⁰. Paradoxically, restoration of myocardial blood flow comes at a price, as it initiates myocardial reperfusion injury by a series of events, which eventually provoke accelerated apoptosis¹¹.

There are four types of MI-R injury recognized. The first two are reversible and consist of reperfusion-induced arrhythmias, in particular ventricular arrhythmias, which are usually self-terminating or effectively treated¹², and myocardial stunning, referring to reversible post-ischemic contractile dysfunction resulting from the detrimental effects of oxidative stress and intracellular calcium overload on the myocardial contractile apparatus¹³. The other two are irreversible and are known as microvascular obstruction with reduced blood flow to the infarct zone despite patency of the infarcted-related artery^{14,15}, and lethal reperfusion injury, which may account for up to 50% of the final infarct size¹¹.

The innate immune system is triggered following tissue injury resulting in the release of soluble inflammatory mediators soon after reperfusion, inducing an extensive inflammatory response as described above. Recruitment of inflammatory cells, oxidative stress, and endothelial barrier dysfunction compose myocardial healing in the early phase. Cytokines activate and recruit neutrophils to the injured infarct area, which cause direct injury to the endothelial cells via production of reactive oxygen species (ROS)¹⁶, inflammatory cytokines, and adhesion molecules that facilitate binding of leucocytes and platelets. Other mediators of myocardial reperfusion injury are opening of the mitochondrial permeability transition pore, cardiomyocyte calcium overload with hypercontracture, and intracellular pH changes resulting in a wave front of reperfusion injury with a strict therapeutic window¹⁷. Reperfusion also induces and aggravates apoptosis⁶. Through binding and ingestion of dying cells, myeloid cells can markedly influence immune responses. MI-R induced apoptosis results in changes of cellular structures including loss of the asymmetric distribution of plasma membrane phospholipids¹⁸.

Innate immunity (Figure 1)

Cell death induced by necrosis causes a release of intracellular contents and triggers an inflammatory response by activating the innate immune system. Expression of endogenous ligands upon reperfusion are judged as “danger signals” These danger-associated molecular patterns (DAMPs) are recognized by both signaling and endocytic pattern recognition receptors (PRRs), such as Toll-like receptors (TLRs), expressed by cells of the innate immune system. Activation of inflammatory pathways by activation of, e.g., TLR-mediated pathways, chemoattractants, the complement cascade, and ROS as a result of MI induces the expression of a large panel of pro-inflammatory genes driven by the activation of mitogen-activated protein kinases (MAPKs) and especially nuclear factor (NF)- κ B⁷.




Figure 1: Innate immunity following myocardial ischemia-reperfusion injury. Ischemic induced myocardial cell death activates innate immunity. Expression of endogenous ligands upon reperfusion are known as danger-associated molecular patterns (DAMPs) including high-mobility group box-1 (HMGB1), heat shock proteins (HSPs), oxidation-specific epitopes (OSEs), and hyaluronic acid (HA). These DAMPs are recognized by pattern recognition receptors (PRRs) like TLRs, NLRs, RAGE, complement cascade, reactive oxygen species (ROS), and chemoattractants. Finally, activation of the nuclear factor (NF- κ B) and mitogen-activated protein kinase (MAPK) pathways induce the expression of pro-inflammatory chemokines, cytokines, and adhesion molecules regulating a complex post-ischemic inflammatory response.

Danger-associated molecular patterns

The innate immune system can be triggered by endogenous, non-pathogenous signals, referred to as DAMPs¹⁹. These endogenous ligands released in response to MI include high-mobility group box-1 (HMGB1), heat shock proteins (HSPs) and S100-proteins, nuclear and mitochondrial DNAs, RNAs, adenosine triphosphate (ATP), low molecular weight hyaluronic acid, and fibronectin fragments²⁰. Oxidation-specific epitopes (OSEs) can act as endogenous DAMPs as well²¹.

HMGB1, the best characterized danger signal, is a key initiator of inflammatory injury following myocardial ischemia controlled by action mechanisms involving TLRs^{22,23} and RAGEs, the receptor for advanced glycation end products²². HMGB1 mediates MI-R injury by activation of inflammatory pathways²⁴, but also confers favorable effects by enhancing angiogenesis, reducing infarct size and improving cardiac function following MI²⁵. Also, HSPs showed both beneficial and detrimental effects following myocardial ischemia. For instance, HSP20 and HSP70 protect cardiomyocytes against MI-R injury in rodents, with improved recovery of cardiac contractile performance and decreased infarct size²⁶. However, HSP60 has strongly been associated with the development of ischemic heart failure²⁷, and HSP60-mediated activation of TLR4 induced myocardial apoptosis and cytokine expression following MI-R injury²⁸. Also, low molecular weight hyaluronic acid and fibronectin are capable of activating the innate immune system via TLRs^{29,30}.

Toll-like receptors

TLRs are highly conserved transmembrane receptors and comprise the major group of PRRs, and recognize a variety of DAMPs and pathogen-associated molecular patterns (PAMPs). The TLR family consists of 13 different receptors, classified into the plasma membrane or endosomal localized TLR subfamilies, triggering the innate immune system upon binding of their ligands

leading to the dimerization of the cytosolic Toll/interleukin-1 receptor (TIR) domain, ultimately activating nuclear factor (NF)- κ B and MAPK pathways to upregulate pro-inflammatory mediators^{31,32}. Myocardial infarction induced local TLR4 expression³³, and upregulated TLR4 in cardiomyocytes exacerbated heart failure³⁴. Furthermore, TLR4 deficiency resulted in decreased MI-R injury³⁵, restricted cardiomyocyte apoptosis²³, and attenuated adverse cardiac remodeling resulting in improved LV function³⁶. In addition, administration of a chemical TLR4 inhibitor reduced the recruitment of Ly-6C^{hi} monocytes accompanied by impaired NF- κ B activation and cytokine expression, which resulted in a decreased infarct size³⁷. TLR2 deficiency resulted in reduced LV dilation and improved LV function due to decreased fibrosis in the non-infarcted area³⁸, and antagonizing TLR2 reduced infarct size after MI-R injury³⁹. Although less extensive, other TLRs have been analyzed as well following MI-R injury. Signaling of TLR3-TIR domain-containing adaptor inducing IFN- β -mediated transcription factor (Trif) represents an injurious pathway, since TLR3-Trif deletion reduced infarct size and improved LV function probably by mediating myocardial apoptosis⁴⁰. In addition, administration of the TLR9 ligand activated the PI3K/Akt signaling pathway conferring cardioprotective effects by induction of TLR9 tyrosine phosphorylation and association with the p85 subunit of PI3K⁴¹. Taken together, TLRs are promising therapeutic targets, and several TLR (ant)agonists already have been developed and investigated following MI-R injury^{42,43}.

Nucleotide-binding oligomerization domain-like receptors

NLRs are a group of intracellular PRRs with over 20 members. NLRs consist of a pyrin (PYD) or caspase recruitment (CARD) domain at the N-terminus combined with a central nucleotide binding (NACHT) followed by a C-terminal leucine-rich repeat (LRR) domain. The inflammasome comprises an activated NLR protein, the adaptor apoptosis speck-like protein containing a caspase-recruitment (ASC) domain, and procaspase-1⁴⁴. After a MI, major components of the NLRP3 inflammasome are upregulated and/or activated in leukocytes, fibroblasts, and endothelial cells, as well as in border zone cardiomyocytes, resulting in increased interleukin (IL)-1 β and IL-18, its cytokine end products. Targeted gene disruption⁴⁵ and antibody neutralization⁴⁶ have been shown to reduce infarct size after MI-R injury, whereas NLRP3 deficiency⁴⁷ and selective NLRP3 inhibition⁴⁸ exhibited additional preservation of cardiac function.

Receptor for advanced glycation end products

RAGE also serves as a multiligand PRR, which triggers a number of cytosolic signaling pathways, including NF- κ B and MAPK-dependent inflammatory genes⁴⁹. RAGE deficiency protected against MI-R injury resulting in improved cardiac function and remodeling, while RAGE-mediated signaling is essential for circulating cells to migrate to the myocardium and exert detrimental effects⁵⁰. Accordingly, co-treatment of soluble RAGE and siRNA RAGE exhibited synergistic cardioprotective effects following MI-R injury⁵¹, and modulation of the AGE-RAGE/MAPK pathways resulted in improved cardiac function by attenuating MI-R injury⁵².

Complement cascade

Involvement of the complement system in myocardial ischemia was first demonstrated almost five decades ago⁵³, and activation has been suggested to extend ischemic injury⁵⁴. The complement cascade operates on the transition from innate to adaptive immunity and is activated through three different mechanisms defined as the classical, alternative, and lectin pathways. All three lead to cleavage of C3 with subsequent activation of C5 and formation of the membrane attack complex (MAC)⁵⁵. It consists of numerous plasma and cell membrane proteins, which are activated following binding of antibodies to PAMPs and release of DAMPs in response to tissue injury. Also, C-reactive protein and IgM play an important role since they are co-localized with activated complement in human infarcted myocardium⁵⁶. Activation causes opsonization of pathogens, phagocytosis, augmentation of inflammation, and direct cell lysis, which is regulated by many regulatory proteins⁵⁷.

All three pathways are involved in the pathogenesis of MI-R injury, by activation-mediated neutrophil and monocyte recruitment in the ischemic myocardium especially in the first hours of reperfusion⁵⁸. The lectin pathway, initiated by recognition of the circulating pathogen recognition molecule mannose-binding lectin (MBL), seemed to be the major contributor. Complement inhibition by antibodies against MBL reduced neutrophil infiltration and attenuated pro-inflammatory gene expression⁵⁹. In addition, MBL deficiency with concomitant active alternative and classical pathways protected against MI-R injury with resultant preservation of cardiac function⁶⁰.

Blocking the complement cascade may reduce myocardial injury, which has been accomplished by consumptive depletion with cobra venom factor⁶¹, antibody-induced inhibition⁶², genetic deficiency⁶³, or ablated plasma activation⁶⁴ of individual complement components such as C5, or by infusion of modified native components that block activation like soluble complement receptor type 1 (sCR1)⁶⁵ and C1-esterase inhibitor (C1-INH)⁶⁶. Despite of these promising results in animal models, clinical studies concerning C5 inhibition in humans have been performed with disappointing results⁶⁷.

Activation of NF- κ B

Induction of pro-inflammatory mediators in ischemic myocardium is controlled by activation of the NF- κ B system, a central transcriptional effector of inflammatory signaling, triggering transcription of various genes including inflammatory cytokines, CXC and CC chemokines, and adhesion molecules upon MI⁶⁸. The family of transcription factors consists of at least five subunits, RelA (p65), RelB, c-Rel, p50, and p52⁶⁹. NF- κ B signaling in the heart has been controversial and seems to regulate three genetic programs depending on the timing and cellular activation, known as acute cytoprotection following hypoxia/ischemia, chronic cytotoxicity generated by a prolonged inflammatory response, and hypertrophy⁶⁸.

Pharmacological blockade of inhibitor of NF- κ B (I κ B α)⁷⁰ and inhibitor of NF- κ B kinase subunit β (IKK β)⁷¹, cardiomyocyte-specific NF- κ B deletion⁷², and blocking of the inflammatory

gene activation⁷³ decreased infarct size, reduced inflammatory responses, and improved cardiac function following MI-R. However, several studies reported conflicting results endorsing the various cellular processes and molecular pathways affected by the NF- κ B system, which are also strictly dependent on the cellular context and timing of activation since prolonged NF- κ B activation promoted heart failure by provoking chronic inflammation⁶⁸.

Chemokines (Figure 2)

Following MI, a variety of pro-inflammatory chemokines and cytokines are upregulated as a result of activation of the innate immune system. Chemokines are divided into subfamilies on the basis of the number and sequential relationship of their conserved cysteine residues (CC, CXC, CX3C, and XC) and play a critical role in basal and inflammatory leukocyte trafficking⁷⁴. Functionally, they can be divided in homeostatic chemokines, which are constitutively expressed in certain tissues and execute basal leukocyte trafficking, and inducible or inflammatory chemokines, which are markedly upregulated following tissue injury and actively induce leukocyte recruitment⁷⁵. Expression of chemokines is managed by activation of TLR-mediated pathways, the complement cascade, ROS generation, and the NF- κ B system following myocardial ischemia, while reperfusion accentuates chemokine expression⁷⁶.

CC chemokines

CC chemokines are the largest and most distinct subfamily and function as potent mononuclear cell chemoattractants. The monocyte chemoattractant protein (MCP)-1/chemokine ligand (CCL)2 is rapidly upregulated in the ischemic myocardium and operates as a potent chemoattractant to mononuclear cells following reperfusion⁷⁷. In addition, MCP-1 deficiency attenuated post-ischemic LV remodeling as a result of a prolonged inflammatory phase with decreased and delayed macrophage infiltration⁷⁸. However, in the absence of reperfusion, cardiac overexpression of MCP-1 induced macrophage infiltration, neovascularization, myocardial IL-6 secretion, and myofibroblast accumulation, which prevented LV dysfunction and adverse remodeling after MI⁷⁹. Furthermore, administration of a CCL2 competitor reduced monocyte recruitment, which attenuated MI-R injury⁸⁰, and pharmacological blockage of the CCL5-CXCR4 interaction impaired the inflammatory response resulting in a reduced infarct size and preserved cardiac function⁸¹. Similar results were observed with anti-CCL5 antibody therapy following permanent ischemia as a result of impaired infiltration of mononuclear cells⁸². Absence of the chemokine receptor CCR1 resulted in reduced functional impairment and structural remodeling due to an abolished early inflammatory recruitment of neutrophils accompanying improved tissue healing⁸³. More recently, CCR9 deficiency was demonstrated to reduce mortality as a result of a decreased infarct size and limited adverse LV remodeling associated with an attenuated inflammatory response⁸⁴.




Figure 2: Post-ischemic inflammatory response. Myocardial infarction especially followed by reperfusion results in a complex post-ischemic inflammatory response encompassing a multifactorial and dynamic process. Apoptosis and necrosis activate the innate immune system resulting in the upregulation of pro-inflammatory chemokines and cytokines. As a result, inflammatory cells are attracted to the ischemic myocardium known as the inflammatory phase. In addition, the production of anti-inflammatory cytokines cause a transition towards the reparative phase accompanied by reparative monocyte and macrophage subtypes. Finally, (myo)fibroblasts and angiogenic cells ensure the formation and maturation of myocardial scar tissue.

CXC chemokines

CXC chemokines that contain an ELR motif are critically involved in chemotactic recruitment of neutrophils, and those lacking the ELR motif regulate the recruitment and activation of lymphocytes. From the ELR containing CXC chemokines, CXCL8/IL-8 is upregulated in experimental MI-R and participates in neutrophil-mediated myocardial injury^{85,86}. IL-8 induces

chemotaxis and granulocyte recruitment, enhances $\beta 2$ integrin-dependent cellular adhesion⁸⁷, and inhibition with a neutralizing monoclonal antibody reduced the degree of necrosis after MI-R independent of neutrophil infiltration⁸⁸. Other ELR-containing CXC chemokines as growth-regulated oncogene (GRO)- α /KC/CXCL1, macrophage inflammatory protein (MIP)-2/CXCL2, and lipopolysaccharide-induced chemokine (LIX)/CXCL5, induced neutrophil chemotaxis and activation, and are expressed by infiltrated inflammatory cells and ischemic myocardium following MI-R injury⁸⁹. In addition, deficiency of the chemokine receptor CXCR2 showed predominant cardioprotective effects with reduced infarct size and recruitment of inflammatory cells after MI-R injury⁹⁰. Also, pharmacological inhibition by a chemokine-binding protein of CXC chemokines, such as CXCL1 and CXCL2, reduced infarct size by preventing neutrophil recruitment and ROS production⁹¹.

Of the ELR-negative CXC chemokines, interferon-inducible protein (IP)-10/CXCL10 peaks in the early reperfusion phase and regulates the cellular composition attenuating adverse remodeling, since IP-10 deficiency resulted in a hypercellular early reparative response with intense infiltration of myofibroblasts suggesting the exertion of anti-fibrotic actions following myocardial ischemia⁹². Finally, stromal cell-derived factor (SDF)-1/CXCL12 may recruit progenitor cells with angiogenic potential, contributing to neovascularization of the scar⁹³, and is upregulated after MI regulating therapeutic stem-cell homing to ischemic myocardium, which resulted in preserved cardiac function⁹⁴. Improved cardiac function after myocardial ischemia was attributed to increased angiogenesis and decreased scar formation as a result of SDF-1 α treatment⁹⁵, and the protective effect was blocked by the administration of a selective antagonist following MI-R injury⁹⁶. However, conflicting observations have been reported following MI-R with detrimental effects of CXCR4-CXCL12 overexpression possibly as a result of enhanced inflammatory cell recruitment, and activation of cell death or apoptosis⁹⁷. These conflicting results again reflect the complexity of the reparative process with multifunctional, pleiotropic, and context- and timing-dependent actions of chemokine signaling.

Cytokines (Figure 2)

Following acute MI, various inflammatory cytokines are secreted by circulating inflammatory cells and cardiac resident cells, amplifying the pro-inflammatory response to acute ischemia-reperfusion injury by moderating the local recruitment of inflammatory cells. Pro-inflammatory cytokines tumor necrosis factor (TNF)- α , IL-1, and IL-6 have consistently been found to be induced and released following MI^{98,99}, whereas IL-10 and transforming growth factor (TGF)- β exerted anti-inflammatory effects^{5,98}.

Pro-inflammatory cytokines

Induction and release of pro-inflammatory cytokines play an important role in mediating chemokine upregulation in the injured myocardium. TNF- α stimulates expression of other pro-inflammatory cytokines, chemokines, and adhesion molecules by leukocytes and endothe-

lial cells, and regulates extracellular matrix metabolism by decreasing collagen synthesis and activating matrix metalloproteinase (MMP) activity¹⁰⁰. Targeted TNF- α overexpression caused adverse cardiac remodeling provoked by progressive cardiomyocyte apoptosis¹⁰¹. Following MI-R injury, TNF- α deficiency exhibited attenuated chemokine expression and NF- κ B activation in the infarcted heart resulting in reduced infarct size and improved cardiac function¹⁰². Additional remote ischemic preconditioning prior to reperfusion attenuated the inflammatory response causing decreased levels of TNF- α and IL-1 β accompanied with an improved LV function¹⁰³. Pharmacological treatment with morphine following MI-R injury was shown to decrease circulating TNF- α levels and reduce infarct size with concomitant improvement of LV function¹⁰⁴. Furthermore, G protein-coupled receptor kinase 2 (GRK2) deficiency resulted in decreased TNF- α expression and fibrosis resulting in reduced infarct size and preserved cardiac function¹⁰⁵. In experimental MI, TNF- α deficiency protected against myocardial rupture and chronic LV dysfunction by inhibiting abundant inflammation, matrix and collagen degradation, and apoptosis¹⁰⁶. In contrast, protective effects of TNF- α signaling are described as well, since neutralizing TNF- α with adenoviral TNFR1 exhibited detrimental effects by promoting ventricular rupture and exacerbating ventricular dysfunction and remodeling following MI¹⁰⁷. Complexity of cytokine expression regarding therapeutic interventions remains a challenge and influences its effects, which might partially depend on blockage of specific receptors since TNF- α effects showed to be toxic via TNFR1 and protective via TNFR2^{108,109}.

Neutralization of IL-1 by genetic overexpression of IL-1 receptor antagonist protected the myocardium from MI-R injury by attenuating the inflammatory response associated with decreased apoptosis¹¹⁰, and reduced inflammatory markers in patients with acute coronary syndrome¹¹¹, but exerted no functional improvement¹¹². Renal sympathetic denervation resulted in a reduced circulating production of pro-inflammatory cytokines IL-1 as well as IL-6 and TNF- α , which was associated with decreased circulating inflammatory cells, reduced infarct size, and improved LV ejection fraction following MI-R injury¹¹³. Administration of a caspase-1 inhibitor combined with a P2Y₁₂ antagonist decreased circulating IL-1 β , reducing infarct size and preserving LV-function¹¹⁴. In addition, treatment with an IL-1 receptor (IL-1R) antagonist as well as IL-1RI deficiency exhibited decreased neutrophil and macrophage infiltration of the injured myocardium with reduced chemokine and cytokine expression subsequently resulting in reduced cardiomyocyte apoptosis and an attenuated fibrotic response preserving adverse remodeling^{115,116}.

Synthesis of IL-6 is induced rapidly in the ischemic-reperfused myocardium¹¹⁷, and IL-6 has been associated with acute coronary syndrome¹¹⁸. In mendelian randomization studies, IL-6R inhibition resulted in reduced cardiovascular events suggesting a novel therapeutic approach to prevent coronary heart disease¹¹⁹. In addition, administration of tocilizumab, an IL-6R antagonist, in patients suffering a non-ST-elevation MI attenuated the inflammatory response with reduced levels of high-sensitivity C-reactive protein and cTnT¹²⁰. However, pre-clinical studies reported conflicting results, as IL-6 deficiency did not affect infarct size, LV function or remodel-

ing, and survival after unperfused MI, which was explained by a compensatory role of other mediators¹²¹, but, on the other hand, reduced infarct size following MI-R injury¹²². Furthermore, administration of the IL-6R antibody MR16-1 after MI, suppressed myocardial inflammation causing impaired LV-remodeling and improved LV contractile function¹²³, but worsened LV remodeling following MI-R injury¹²⁴. Regarding pharmacological treatment, administration of bisoprolol conferred cardioprotective effects following MI-R injury by suppressing both IL-6 as TNF- α secretion and attenuating the unfolded protein response, thereby reducing infarct size and improving post-ischemic cardiac function¹²⁵. Linagliptin, a dipeptidylpeptidase (DPP)-4 inhibitor, attenuated the increase of IL-6 as well as IL-1 β and TNF- α following MI-R injury, which was associated with a limited infarct size and improved ejection fraction¹²⁶.

Overall, attenuating the effects of pro-inflammatory cytokines seemed cardioprotective, however, conflicting results suggest them to exert diverse biological effects and emphasize the complex and pleiotropic cytokine actions in biological processes.

Anti-inflammatory cytokines

Inhibitory mediators, such as IL-10, TGF- β , or lipid-derived substances may be the predominant mechanism for deactivation of chemokine signals. They exert pleiotropic effects in regulating the immune response and are leading in mediating the post-ischemic reparative response¹²⁷.

IL-10 is predominantly expressed by activated T lymphocytes and stimulated monocytes, and possesses potent anti-inflammatory properties. Induction of IL-10 coincides with the suppression of IL-6 formation in macrophages and monocytes, and further inhibits production of IL-1 α , IL-1 β , TNF- α , and IL-8, suppressing the inflammatory response. Furthermore, it stabilizes the matrix by decreasing metalloproteinase biosynthesis¹²⁸. IL-10 is upregulated in reperfused myocardium, and neutralizing IL-10 activity caused reduced expression of tissue inhibitor of metalloproteinases (TIMP)-1, suggesting IL-10 to contribute to stabilization of the matrix¹²⁹. Following MI-R injury, IL-10 deficiency resulted in increased fibrosis and apoptosis¹³⁰ and an enhanced inflammatory response, associated with increased infarct size and mortality¹³¹. In addition, administration of recombinant IL-10 following MI suppressed the inflammatory response and contributed to improved LV function and remodeling by inhibiting fibrosis and enhancing capillary density¹³². Pharmacological treatment with colchicine before¹³³ or a selective β_2 adrenergic receptor agonist during¹³⁴ MI-R injury increased systemic IL-10 levels exhibiting a cardioprotective effect with a reduced infarct size. However, conflicting evidence regarding the role of IL-10 in the resolution of the inflammatory response following MI-R exists, as IL-10 deficiency also showed timely repression of pro-inflammatory cytokines and chemokine mRNA synthesis with a similar course of neutrophil infiltrate resolution, indicating the involvement of multiple overlapping modulating mechanisms¹³⁵.

Cell proliferation, differentiation, and apoptosis as well as modulation of the immune response are all regulated by TGF- β . Because of its anti-inflammatory and fibrogenic properties, it affects cardiac remodeling by mediating transition from inflammation to fibrosis¹³⁶. Upon

TGF- β stimulation, myofibroblast transdifferentiation is induced, and the activity of proteases that degrade the extracellular matrix (ECM) is suppressed¹³⁷. Following myocardial ischemia TGF- β is increased, of which TGF- β 1 and TGF- β 2 isoforms are upregulated within the first few days in contrast to the delayed and prolonged expression of the TGF- β 3 isoform showing a positive correlation with parameters of the ECM metabolism¹³⁸. TGF- β is predominantly located in the infarct border zone activated by the induction of thrombospondin (TSP)-1 following MI-R injury¹³⁹. The signaling pathway is regulated through Smad proteins¹⁴⁰ of which Smad3 signaling seemed to be critical in cardiac remodeling¹⁴¹.

Post-ischemic treatment with TGF- β prevented severe cardiac injury, presumably by attenuating the deleterious effects of pro-inflammatory cytokines¹⁴². Loss of the anti-inflammatory mechanism of growth differentiation factor (GDF)-15, a TGF- β related cytokine and inhibitor of leukocyte integrin activation, resulted in fatal cardiac rupture after MI¹⁴³. Moreover, prevention of myeloid-epithelial-reproductive tyrosine kinase (MerTK) cleavage following MI-R injury resulted in secretion of TGF- β and IL-10, which was associated with a reduced infarct size and improved systolic function¹⁴⁴. However, reducing TGF- β activity by inhibiting the TGF- β type I receptor, attenuated systolic dysfunction and LV remodeling¹⁴⁵, and gene therapy against TGF- β mitigated LV remodeling as well¹⁴⁶. Also, inhibition of TGF- β expression due to combination therapy of angiotensin-converting enzyme (ACE) inhibitor and angiotensin receptor blocker attenuated cardiac fibrosis and remodeling¹⁴⁷. Furthermore, targeted disruption of TGF- β signaling by conditional deficiency of the TGF- β receptor 1 or 2 resulted in a marked decline of neutrophil recruitment and upregulation of protective cardiokines protecting from LV wall rupture¹⁴⁸. Finally, Smad3 deficiency following MI-R injury resulted in reduced interstitial fibrosis and attenuated cardiac remodeling due to abrogation of the fibrogenic TGF- β responses¹⁴⁹. Discrepancies might be explained by a specific time window of action since activation of TGF- β signaling seemed protective in the early post-ischemic phase, but caused adverse LV remodeling and dysfunction when expression persists over time¹⁵⁰.

The inhibitory cytokines exert pleiotropic effects regarding their inflammatory and reparative properties, which can be either stimulatory or inhibitory, depending on the state of cellular differentiation, microenvironmental cues, and the cellular tissue origin. The related complexity has hampered the translation from attractive hypothesis towards novel therapeutic strategies exhibiting functional improvement in experimental MI models.

Inflammatory cells (Figure 2)

Acute MI results in a complex inflammatory response associated with induction of endothelial adhesion molecules and enhanced permeability of the microvasculature. Upregulated chemokines and cytokines cause extravasation of activated blood-derived cells into the infarcted area, which play an important role in regulating the reparative response and are possible targets for cardioprotective therapies¹⁵¹. Single-cell RNA-sequencing following unreperfused MI of the non-cardiomyocyte fraction revealed over 30 cell subtype populations, which highlighted post-

ischemic non-linear dynamics in myeloid and fibroblast lineages crucial for understanding of cardiac homeostasis, inflammation, fibrosis, repair, and regeneration¹⁵².

At first, platelets are recruited into injured myocardium representing an important linkage between tissue injury and repair¹⁵³. They target monocytes, macrophages, and T cells by releasing mitogenic factors, recruit leukocytes by production of cytokines and chemokines, and trigger complement activation^{154,155}. Following MI-R injury, platelet-derived serotonin induced neutrophil degranulation with release of myeloperoxidase, which was abolished after additional genetic or pharmacological depletion preserving cardiac function¹⁵⁶. In addition, targeting activated platelets with a single-chain antibody-CD39 (Targ-CD39) resulted in attenuated post-ischemic inflammation, reduced infarct size, and increased neovascularization resulting in a remarkable restoration of ejection fraction and strain rate¹⁵⁷.

Briefly, MI triggers a biphasic reaction of inflammation and remodeling, with immediate influx of neutrophils and CCR2⁺Ly-6C^{hi} pro-inflammatory M1 monocytes/macrophages in the immediate phase, which phagocytose debris and secrete pro-inflammatory cytokines IL-1 β , IL-6, and TNF- α . After around 3 days transition to the reparative phase is achieved in which accumulated Ly-6C-F4/80^{hi} M2 macrophages secrete anti-inflammatory cytokines IL-10 and TGF- β ¹⁵⁸.

Inflammatory and reparative phases

Neutrophils

Prior to MI, neutrophils as well as monocytes already circulate in the blood in a steady state. In response to experimental MI, neutrophils, which are efficient phagocytes that engulf and degrade microorganisms using oxidative and non-oxidative mechanisms, are migrated into the injured myocardium as the first line of defense to phagocytose death myocardial cells¹⁵⁹. In the clinical setting, elevated neutrophil as well as leukocyte counts following primary PCI in ST-segment elevation MI are directly related to infarct size and LV ejection fraction and are independent predictors of cardiovascular outcomes¹⁶⁰. Neutrophils infiltrate the injured myocardium within hours, peaking at 1-3 days, and primarily tend to target the border zone showing accentuated accumulation at reperfusion¹⁶¹. They are recruited to the injured myocardium by a high concentration of chemotactic factors, such as MIP-2 α , IL-8(/CXCL8), CXCL1(/GRO- α /KC), complement 5a, and leukotriene B4. Migration from the circulation into the injured myocardium depends on binding to P-selectin, E-selectin, intercellular adhesion molecule (ICAM)-1, and vascular cell adhesion molecules¹⁶². Until recently, it was believed that cardiac neutrophils demonstrate a pro-inflammatory N1 phenotype, being predominant in the infarct area correlating with wall thinning, and an anti-inflammatory N2 phenotype, increasing over the course of MI¹⁶³. Excessive infiltration and/or delayed removal may exacerbate injury by intensifying the inflammatory response. Novel insights show a more complex concept of temporal cardiac neutrophil heterogeneity in experimental MI. Single-cell RNA-sequencing combined with surface epitope detection demonstrated more complex neutrophil dynamics with two major subsets. The Siglec-F^{lo} neutrophils, resembling circulating blood neutrophils, and the SiglecF^{hi} neutrophil subset,

exclusively found in the heart characterized by specific effector functions as phagocytosis and production of ROS¹⁶⁴.

Targeting neutrophils to reduce accumulation into the injured myocardium has been shown to reduce infarct size and mostly improved cardiac function following neutralization of the receptor activator of NF- κ B ligand¹⁶⁵, plasminogen activator inhibitor-1 deficiency¹⁶⁶, brahma-related gene 1 (BRG1) deficiency¹⁶⁷, pharmacological inhibition, as well as deficiency of transient receptor potential melastatin 2 (TRPM2)¹⁶⁸, or proteasome-mediated I κ B α inhibition¹⁶⁹. Furthermore, inhibition of neutrophil activity exposed beneficial effects with a reduced infarct size as a result of pretreatment with a dual cyclooxygenase-lipoxygenase blocking agent before MI-R injury¹⁷⁰. Improved LV function and attenuated adverse remodeling was shown following administration of an inhibitor of the triggering receptor expressed on myeloid cells-1 (TREM-1)¹⁷¹, a S100A8/A9 blocker¹⁷², or myeloperoxidase inhibitor¹⁷³ early after initiating reperfusion. In addition, neutralization of L-selectin and P-selectin with monoclonal antibodies attenuated neutrophil accumulation and reduced myocardial necrosis after MI-R injury^{174,175}. However, combined P-selectin and ICAM-1 deficiency demonstrated impaired neutrophil trafficking but did not affect infarct size in response to MI-R¹⁷⁶. Moreover, inhibition or loss of the CD11/CD18-integrin receptor, which allows neutrophil binding and transmigration, resulted in diminished neutrophil accumulation accompanied by a reduced infarct size and preserved cardiac function after experimental MI-R injury¹⁷⁷⁻¹⁷⁹. Disappointingly, clinical studies targeting CD11/CD18 subunits did not report any cardioprotective effect following acute MI, endorsing its challenging aspects^{180,181}. Finally, complete neutrophil depletion showed to be detrimental, since neutrophils are crucially involved by polarizing macrophages towards a reparative phenotype, emphasizing the importance of carefully balanced novel anti-inflammatory therapeutic strategies¹⁸².

Mononuclear cells

Following experimental MI-R injury, mononuclear cells (monocytes and lymphocytes) are rapidly recruited into the ischemic myocardium, visualized by real-time *in vivo* imaging¹⁸³, due to various monocyte chemoattractants driven by MCP-1⁷⁸. Monocytes and macrophages have a critical role in post-ischemic cardiac repair, since they promote both injury and repair, and express wide heterogeneity and dynamics during the immediate inflammatory response. As a result of cardiomyocyte injury, monocytes and monocyte-derived macrophages infiltrate the heart and largely replace tissue-resident macrophages. Cardiac tissue-resident macrophages originate from CX3CR1+ progenitors¹⁸⁴, are self-renewing and long-lived¹⁸⁵, and display an anti-inflammatory F4/80^{hi}Ly-6C^{lo} or M2 phenotype¹⁸⁶. Combination of single-cell RNA-sequencing with genetic fate mapping of the healthy adult myocardium revealed four transcriptionally distinct cardiac macrophage subsets, which are, on the one hand, maintained independently of blood monocytes, or on the other hand, partially or fully replaced by monocytes. Following myocardial injury, resident cardiac macrophages, suppressed by peripherally CCR2+ monocyte-derived macrophages, accounted for only 2-5% of the cardiac macrophages within the infarct area¹⁸⁷. Resident cardiac

macrophages undergo continuous and distinct transcriptomic changes¹⁸⁶, and their depletion promoted adverse remodeling and impaired cardiac function¹⁸⁷.

Of the monocytes, the pro-inflammatory (Ly-6C^{hi}) monocytes migrate to the site of injured myocardium at first, peaking 3 days after MI, which express TNF- α and IL-1 β , produce proteolytic enzymes, and secrete MMPs. They exhibit rapid kinetics with a relatively short circulating life-span¹⁸⁸, and differentiate, in the presence of IFN- γ or lipopolysaccharide (LPS), into activated pro-inflammatory M1 (CCR2+) macrophages secreting large amounts of pro-inflammatory mediators¹⁸⁹. In addition, nuclear receptor subfamily 4, group a, member 1 (Nr4a1) seems critical for Ly-6C^{hi} monocytes to exert either inflammatory or reparative effects following unperfused MI. Although differentiated cardiac macrophages depend on Nr4a1 to limit inflammation, the apoptotic or proliferation processes seem unaffected¹⁹⁰. Second, after about 7 days, the reparative (Ly-6C^{lo}) monocytes, also suggested to directly correspond with F4/80^{hi}Ly-6C^{lo} macrophages originated from Ly-6C^{hi} monocytes^{190,191}, become predominant, which differentiate, in the presence of IL-4 and IL-10, towards a reparative M2 (CCR2-) phenotype, promoting the healing response, and contributing to angiogenesis and scar maturation¹⁸⁹. Tissue-resident CCR2+ and CCR2- macrophages differentially regulated cardiac mobilization and recruitment of peripheral monocytes where depletion of tissue-resident CCR2+ macrophages substantially reduced the recruitment of recipient monocytes and neutrophils accompanied by an improved LV systolic function following murine MI-R injury¹⁸⁵. In the clinical setting, higher levels of pro-inflammatory monocytes were associated with severe myocardial injury and poor functional outcome following acute MI^{192,193}. Furthermore, the macrophage migration inhibitory factor (MIF) seemed to exert pleiotropic effects following MI-R injury. By activating the cardioprotective AMP-activated protein kinase (AMPK) pathway, it seemed beneficial, on the one hand¹⁹⁴, but MIF deficiency, on the other hand, turned out to protect the heart by suppressing the inflammatory response as well¹⁹⁵. The compartmentalized and opposing effects are largely mediated by CXCR2, where CXCR2-bearing inflammatory cells revealed to be detrimental by increasing monocyte infiltration¹⁹⁶. Efferocytosis of apoptotic cardiomyocytes by macrophages suppressed expression of pro-inflammatory mediators, which may drive resolution of inflammation¹⁹⁷.

Attenuating the pro-inflammatory phase by targeting Ly-6C^{hi} monocytes or M1 macrophages has been shown cardioprotective following reduction of interferon regulatory factor 5 (IRF5) expression¹⁹⁸, injection of phosphatidylserine (PS)-presenting liposomes¹⁹⁹ or irbesartan-nanoparticles²⁰⁰, as well as administration of annexin A5 (AnxA5)²⁰¹ or phosphorylcholine monoclonal immunoglobulin G antibodies²⁰², and CCR-2 silencing²⁰³. In addition, expediting the differentiation from reparative Ly-6C^{lo} monocytes towards M2 macrophages by administration of pioglitazone-nanoparticles²⁰⁴ or topiramate²⁰⁵, T_{reg} cell activation²⁰⁶, and silencing the collapsin response mediator protein-2 (CRMP2)²⁰⁷ limited adverse remodeling. Moreover, exacerbation of the post-ischemic inflammatory response by administering 2-arachidonolglycerol resulted in increased neutrophil and monocyte counts associated with an increased infarct size and worsened cardiac function²⁰⁸. Inhibition of the reparative phase finally, by selective depletion of

M2 macrophages as a result of deficiency of the kinase tribbles homolog 1 (TRIB1) caused an increased risk of cardiac rupture following MI, emphasizing their role in regulation of fibroblast activation²⁰⁹.

Other leukocytes, like lymphocytes, dendritic cells, and mast cells invading the injured myocardium includes smaller numbers, but may have an important role in regulating infarct healing and the myeloid response. CD4+ T lymphocytes were shown to become activated after MI and improved beneficial remodeling and survival²¹⁰. Lymphopenia, primarily due to loss of T cells in patients with an acute MI, was correlated with substantial microvascular obstruction²¹¹. Furthermore, T_{reg} cells are known to attenuate the innate immune response and inducing T_{reg} cells by injection of tolerogenic dendritic cells (tDCs) elicited an inflammatory-to-reparative macrophage shift yielding improved remodeling associated with preservation of LV function and increased survival²¹². However, CD4+ T cell deficiency also showed to reduce infarct size and preserve ejection fraction following MI-R injury²¹³, suggesting different roles of T cells regarding infarct remodeling compared with MI-R injury. B lymphocytes peak 5 days after acute MI and trigger monocyte mobilization, which deteriorated LV function²¹⁴. Following MI, granulocyte-macrophage colony-stimulating factor (GM-CSF)-producing B cells, dendritic cells, and T cells expand in pericardial adipose tissue, at which both B cell depletion and GM-CSF blockade, as well as removal of pericardial adipose tissue reduced fibrosis and preserved cardiac function²¹⁵.

Dendritic cells are antigen-presenting cells contributing to innate immunity and providing a link to the adaptive immune system. Following MI-R injury, they migrate to and accumulate in the injured myocardium, and deficiency resulted in increased pro-inflammatory cell and cytokine recruitment with worsened adverse remodeling suggesting immunoprotective effects²¹⁶. In addition, exosomes derived from dendritic cells activated CD4+ T lymphocytes, which improved cardiac function following MI²¹⁷. On the other hand, restriction of dendritic cell activation and migration by inhibiting HMGB1 reduced infarct size as well²¹⁸.

Resident cardiac mast cells degranulate quickly after MI releasing histamine and pro-inflammatory mediators as TNF- α . Mast cells are increased in reperfused myocardium and are associated with upregulation of stem cell factor (SCF), which may expedite recruitment and homing of primitive bone marrow-derived cells contributing to the reparative phase²¹⁹. The most abundant protease in mast cell granules is tryptase, which induced synthesis of chemokines and cytokines to mediate leukocyte accumulation and angiogenesis²²⁰ and was identified to regulate PKA activity modulating cardiomyocyte contractility after acute MI²²¹. Following MI-R injury, dual inhibition of mast cell and neutrophil-derived proteases reduced myocyte apoptosis and preserved cardiac function²²². Furthermore, inhibition of renin release from mast cells prevented activation of the renin-angiotensin system (RAS), reduced infarct size, and alleviated ventricular arrhythmias^{223,224}.

Reparative to maturation phase

Fibroblasts

Between days 3 and 7 following MI, activated cardiac fibroblasts proliferate, expand, and transdifferentiate into synthetic myofibroblasts precluding the reparative phase of cardiac repair due to the replacement of death cardiomyocytes by a matured fibrotic scar^{225,226}. Interstitial fibroblasts surviving the ischemic event, or recruited bone marrow progenitor or epicardial-derived cells may undergo myofibroblast transdifferentiation²²⁷ critically regulated by TGF- β . These myofibroblasts are marked by expression of α -smooth muscle actin (ASMA) and initiate the synthesis of ECM proteins and formation of neovessels²²⁸. In the early post-ischemic period, fibroblasts may even modulate survival pathways in cardiomyocytes, affecting their susceptibility to myocardial ischemia²²⁹. Moreover, myofibroblasts acquired anti-inflammatory properties following milk fat globule-epidermal growth factor (MFG-E)8-mediated engulfment of apoptotic cells²³⁰.

Following MI, generation of active TGF- β in the injured myocardium triggers Smad activation in both fibroblasts and cardiomyocytes. Fibroblast-specific Smad3 seems essential for scar organization, since deficiency accentuated adverse remodeling with perturbed alignment of myofibroblast arrays in the infarct. In contrast, cardiomyocyte-specific Smad3 seems detrimental, as deficiency was associated with attenuated cardiomyocyte apoptosis without affecting infarct size²³¹. In addition, impaired TGF- β and Smad3 activation in fibroblasts depleted of the primary cilium resulted in decreased ECM protein levels and contractile function²³². Attenuating post-ischemic inflammation by administration of anti-inflammatory IL-10 improved cardiac remodeling and function by stimulation of fibroblast activation influenced by M2 macrophage polarization²³³. Furthermore, increased cardiac hyaluronan synthesis contributed to post-ischemic infarct healing by promoting the myofibroblast response and supporting macrophage survival²³⁴.

Neovascularization

Neovascularization is initiated by angiogenic growth factors followed by the recruitment of a muscular coat and formation of neoarterioles during scar maturation, which may stabilize infarct vasculature by inhibition of endothelial cell proliferation and vascular sprouting²³⁵. Vascular endothelial growth factor (VEGF)-induced endothelial cell migration and proliferation cause hyperpermeable neovessel formation. Numerous factors, such as fibroblast growth factors (FGFs), TGF- β , MCP-1, IL-8, and IP-10 modulate angiogenesis, indicating its complexity. The post-ischemic increase in angiopoietin (Ang)-2 expression and reciprocal decrease in Ang-1 expression, suggested a predominant role for Ang-2 in the angiogenic response to MI²³⁶. Furthermore, platelet-derived growth factor receptor (PDGFR) activation is involved in vasculature maturation, which promotes resolution of inflammation and stabilization of the myocardial scar²³⁷. Following MI, cardiomyocyte-specific deficiency of HSPB1 resulted in decreased angiogenesis and collagen deposition with aggravated adverse remodeling and cardiac dysfunction as a result of enhanced leukocyte infiltration and expression of inflammatory cytokines²³⁸.

Resolution of the post-ischemic inflammatory response

The acute transient inflammatory response finally promotes to a stage of tissue repair and scar formation^{98,225}. Inhibition of chemokine and cytokine synthesis after they peaked is crucial for the repair process resulting in resolution of the inflammatory response²³⁹. Although identifying the precise mechanisms controlling the switch from formation of pro-inflammatory to anti-inflammatory reparative pathways is an ongoing process, components of resolution include apoptosis, autophagy, necrosis, and formation of anti-inflammatory cytokines²⁴⁰. Clearance of the neutrophilic infiltrate and removal of debris, inhibition of chemokine and cytokine synthesis, removal of the fibrin-based provisional matrix, and activation of fibroblasts and collagen deposition finally contribute to the myocardial reparative process and mature scar formation. This complex process of infiltration, differentiation, activation, and interaction of various cell types eventually provides accomplishment of complete infarct wound healing, which takes about 2 weeks in mice compared with 6 months in humans²⁴¹.

IMMUNOMODULATORY THERAPIES IN CLINICAL PRACTICE

Although therapeutic interventions following myocardial infarction have been dramatically improved over the years, morbidity and mortality remain high. This could largely be explained by the practically lacking capability of the heart to regenerate²⁴², since differentiated cardiomyocytes predominantly lost their capacity to proliferate. In addition, myocardial wound healing occurs under continuous rhythmic contraction of the non-infarcted myocardium, thus under continuous mechanical stress²⁴¹. Treatment of patients to interfere in myocardial remodeling by healing of ischemic myocardium and preventing scar formation with concomitant preservation of cardiac function would be the ultimate goal. In the last years, cardiac regeneration has experienced a watershed moment, and it is believed that even in adult mammals, latent regeneration can be awakened. Combining stem cell-based methods with gene regulatory elements seems promising, however, correct electromechanical incorporation of transplanted cardiomyocytes remains a challenge as yet²⁴³. In addition, combination of selected therapies might be promising exerting synergistic effects, for example, by inhibition of the early pro-inflammatory response with subsequent activation of the anti-inflammatory or so called reparative response. At the same time, effects regarding the non-inflammatory components, such as cardiomyocytes, platelet activation, microvascular obstruction, and coronary endothelial dysfunction should also be taken into account²⁴⁴.

Already in 1973, the use of the anti-inflammatory agent corticosteroid for the first time showed limitation of myocardial infarct size in dogs²⁴⁵. However, a subsequent clinical trial resulted in deleterious effects²⁴⁶, possibly caused by delayed wound healing due to an impaired inflammatory response²⁴⁷. This emphasizes the necessity of wound healing and illustrates the difficulty of counteracting maladaptive inflammatory pathways without interfering in benign

wound healing endorsing inflammatory mediators to be notoriously pleiotropic. Current research focusses on targeted anti-inflammatory or immunomodulatory treatment, cardiac cell therapy and regeneration, mechanical therapeutic interventions, pharmacological therapies, and microRNA (miRNA)-targeted treatment. Below, anti-inflammatory and immunomodulatory therapies will be briefly highlighted. In addition, the selection of an appropriate animal model to ensure optimal translation of novel cardioprotective therapies from bench-to bedside is of the utmost importance^{248,249}, ideally allowing for an atherosclerotic phenotype and myocardial ischemia-reperfusion injury²⁵⁰.

Although the immune response includes an essential part of the reparative myocardial wound healing process, no specific immunomodulatory treatments are available for patients on a large scale. Finished or discontinued clinical trials failed to translate promising results from experimental MI studies in animals towards clinical practice in human patients, although the post-ischemic inflammatory response remains amenable for therapeutic intervention where success depends on timing and topography emphasizing the necessity of real-time visualization of the extent and time course of cellular damage²⁵¹. In addition, right selection of subpopulations regarding distinct pathophysiologic perturbations and underlying mechanisms needs to be taken into account. Glucocorticoids were the first to be thought beneficial in post-ischemic remodeling. On the one hand, they attenuate the responses to danger signals during the acute inflammatory phase and reduce leukocyte infiltration, on the other hand, they are known to promote clearance of apoptotic cells and direct macrophages towards an anti-inflammatory phenotype. However, clinical studies showed conflicting results, and considering their broad actions on cell types involved in ischemic injury and effects on several molecular cascades, they are unattractive as a therapeutic option because of a wide variety of adverse events²⁵².

IL-1 signaling is suggested to play a crucial role in post-ischemic adverse remodeling and cardiac dysfunction, which is attenuated following IL-1 blockade with anakinra in animals¹¹⁵, and represents a promising therapeutic approach following acute MI in patients^{111,112}. Moreover, targeting the IL-1 β innate immunity pathway with canakinumab reduced recurrent cardiovascular events in high-risk patients²⁵³. Furthermore, administration of an IL-6R antagonist tocilizumab in patients suffering an acute coronary syndrome attenuated the inflammatory response¹²⁰. TNF- α blockade by etanercept reduced systemic inflammation but increased platelet activation following acute MI and was therefore concluded to be unfavorable²⁵⁴. Targeted interventions regarding anti-inflammatory effects of IL-10 and TGF- β have not made it to a serious clinical trial yet. Trials with humanized recombinant antibody therapy directed against integrins¹⁸⁰ or the complement system^{67,255} failed to show positive effects. On the other hand, intracoronary nitrite following primary PCI for acute MI reduced neutrophil numbers and activation associated with a reduction in infarct size and microvascular obstruction²⁵⁶. Immunomodulation with cyclosporine, which inhibits the opening of mitochondrial permeability-transition pores and exerts potent immunosuppressive effects, attenuated lethal myocardial injury²⁵⁷, but did not pre-

serve cardiac function following acute MI²⁵⁸. Finally, anti-inflammatory therapy with colchicine following MI²⁵⁹ as well as in patients with chronic coronary disease²⁶⁰ led to reduced recurrent ischemic cardiovascular events.

CONCLUSION

Acute myocardial infarction especially followed by reperfusion results in a complex post-ischemic inflammatory response encompassing a multifactorial and dynamic process consisting of overlapping inflammatory, reparative, and maturation phases. Therefore, MI-R injury is exceptionally amenable and, above all, suitable for immunomodulatory therapies. However, successful development and translation of future therapeutic strategies should focus on the sensitive balance between anti-inflammatory and pro-reparative effects and should be tested in translational research models exposed to clinical relevant pathophysiologic mechanisms and comorbidity.

REFERENCES

1. Lochner A, de Villiers M. Phosphatidylcholine biosynthesis in myocardial ischaemia. *J Mol Cell Cardiol* 1989;21(2):151-63.
2. Jennings RB, Murry CE, Steenbergen C, Jr., Reimer KA. Development of cell injury in sustained acute ischemia. *Circulation* 1990;82(3 Suppl):II2-12.
3. Rajappa M, Sharma A. Biomarkers of cardiac injury: an update. *Angiology* 2005;56(6):677-91.
4. Mueller C. Biomarkers and acute coronary syndromes: an update. *Eur Heart J* 2014;35(9):552-6.
5. Frangogiannis NG. Regulation of the inflammatory response in cardiac repair. *Circ Res* 2012;110(1):159-73.
6. Eefting F, Rensing B, Wigman J, et al. Role of apoptosis in reperfusion injury. *Cardiovasc Res* 2004;61(3):414-26.
7. Frangogiannis NG. The inflammatory response in myocardial injury, repair, and remodelling. *Nat Rev Cardiol* 2014;11(5):255-65.
8. Ibanez B, James S, Agewall S, et al. 2017 ESC Guidelines for the management of acute myocardial infarction in patients presenting with ST-segment elevation: The Task Force for the management of acute myocardial infarction in patients presenting with ST-segment elevation of the European Society of Cardiology (ESC). *Eur Heart J* 2018;39(2):119-77.
9. Schomig A, Ndrepepa G, Kastrati A. Late myocardial salvage: time to recognize its reality in the reperfusion therapy of acute myocardial infarction. *Eur Heart J* 2006;27(16):1900-7.
10. Boyle MP, Weisman HF. Limitation of infarct expansion and ventricular remodeling by late reperfusion. Study of time course and mechanism in a rat model. *Circulation* 1993;88(6):2872-83.
11. Yellon DM, Hausenloy DJ. Myocardial reperfusion injury. *N Engl J Med* 2007;357(11):1121-35.
12. Manning AS, Hearse DJ. Reperfusion-induced arrhythmias: mechanisms and prevention. *J Mol Cell Cardiol* 1984;16(6):497-518.
13. Kloner RA, Bolli R, Marban E, et al. Medical and cellular implications of stunning, hibernation, and preconditioning: an NHLBI workshop. *Circulation* 1998;97(18):1848-67.
14. Ito H. No-reflow phenomenon and prognosis in patients with acute myocardial infarction. *Nat Clin Pract Cardiovasc Med* 2006;3(9):499-506.
15. Hausenloy DJ, Chilian W, Crea F, et al. The coronary circulation in acute myocardial ischaemia/reperfusion injury: a target for cardioprotection. *Cardiovasc Res* 2019;115(7):1143-55.
16. Griendling KK, FitzGerald GA. Oxidative stress and cardiovascular injury: Part I: basic mechanisms and in vivo monitoring of ROS. *Circulation* 2003;108(16):1912-6.
17. Frohlich GM, Meier P, White SK, et al. Myocardial reperfusion injury: looking beyond primary PCI. *Eur Heart J* 2013;34(23):1714-22.
18. Savill J, Dransfield I, Gregory C, Haslett C. A blast from the past: clearance of apoptotic cells regulates immune responses. *Nat Rev Immunol* 2002;2(12):965-75.
19. Matzinger P. The danger model: a renewed sense of self. *Science* 2002;296(5566):301-5.
20. Zuurbier CJ, Abbate A, Cabrera-Fuentes HA, et al. Innate immunity as a target for acute cardioprotection. *Cardiovasc Res* 2019;115(7):1131-42.
21. Miller YI, Choi SH, Wiesner P, et al. Oxidation-specific epitopes are danger-associated molecular patterns recognized by pattern recognition receptors of innate immunity. *Circ Res* 2011;108(2):235-48.
22. Tian J, Avalos AM, Mao SY, et al. Toll-like receptor 9-dependent activation by DNA-containing immune complexes is mediated by HMGB1 and RAGE. *Nat Immunol* 2007;8(5):487-96.
23. Ding HS, Yang J, Chen P, et al. The HMGB1-TLR4 axis contributes to myocardial ischemia/reperfusion injury via regulation of cardiomyocyte apoptosis. *Gene* 2013;527(1):389-93.
24. Andrassy M, Volz HC, Igwe JC, et al. High-mobility group box-1 in ischemia-reperfusion injury of the heart. *Circulation* 2008;117(25):3216-26.
25. Kitahara T, Takeishi Y, Harada M, et al. High-mobility group box 1 restores cardiac function after myocardial infarction in transgenic mice. *Cardiovasc Res* 2008;80(1):40-6.
26. Fan GC, Ren X, Qian J, et al. Novel cardioprotective role of a small heat-shock protein, Hsp20, against ischemia/reperfusion injury. *Circulation* 2005;111(14):1792-9.

27. Tanonaka K, Yoshida H, Toga W, et al. Myocardial heat shock proteins during the development of heart failure. *Biochem Biophys Res Commun* 2001;283(2):520-5.
28. Kim SC, Stice JP, Chen L, et al. Extracellular heat shock protein 60, cardiac myocytes, and apoptosis. *Circ Res* 2009;105(12):1186-95.
29. Jiang D, Liang J, Fan J, et al. Regulation of lung injury and repair by Toll-like receptors and hyaluronan. *Nat Med* 2005;11(11):1173-9.
30. Dobaczewski M, Gonzalez-Quesada C, Frangogiannis NG. The extracellular matrix as a modulator of the inflammatory and reparative response following myocardial infarction. *J Mol Cell Cardiol* 2010;48(3):504-11.
31. Akira S, Takeda K. Toll-like receptor signalling. *Nat Rev Immunol* 2004;4(7):499-511.
32. Kawai T, Akira S. The role of pattern-recognition receptors in innate immunity: update on Toll-like receptors. *Nat Immunol* 2010;11(5):373-84.
33. Frantz S, Kobzik L, Kim YD, et al. Toll4 (TLR4) expression in cardiac myocytes in normal and failing myocardium. *J Clin Invest* 1999;104(3):271-80.
34. Liu L, Wang Y, Cao ZY, et al. Up-regulated TLR4 in cardiomyocytes exacerbates heart failure after long-term myocardial infarction. *J Cell Mol Med* 2015;19(12):2728-40.
35. Oyama J, Blais C, Jr., Liu X, et al. Reduced myocardial ischemia-reperfusion injury in toll-like receptor 4-deficient mice. *Circulation* 2004;109(6):784-9.
36. Riad A, Jager S, Sobirey M, et al. Toll-like receptor-4 modulates survival by induction of left ventricular remodeling after myocardial infarction in mice. *J Immunol* 2008;180(10):6954-61.
37. Fujiwara M, Matoba T, Koga JI, et al. Nanoparticle incorporating Toll-like receptor 4 inhibitor attenuates myocardial ischaemia-reperfusion injury by inhibiting monocyte-mediated inflammation in mice. *Cardiovasc Res* 2019;115(7):1244-55.
38. Shishido T, Nozaki N, Yamaguchi S, et al. Toll-like receptor-2 modulates ventricular remodeling after myocardial infarction. *Circulation* 2003;108(23):2905-10.
39. Arslan F, Smeets MB, O'Neill LA, et al. Myocardial ischemia/reperfusion injury is mediated by leukocytic toll-like receptor-2 and reduced by systemic administration of a novel anti-toll-like receptor-2 antibody. *Circulation* 2010;121(1):80-90.
40. Chen C, Feng Y, Zou L, et al. Role of extracellular RNA and TLR3-Trif signaling in myocardial ischemia-reperfusion injury. *J Am Heart Assoc* 2014;3(1):e000683.
41. Cao Z, Ren D, Ha T, et al. CpG-ODN, the TLR9 agonist, attenuates myocardial ischemia/reperfusion injury: involving activation of PI3K/Akt signaling. *Biochim Biophys Acta* 2013;1832(1):96-104.
42. Arslan F, Houtgraaf JH, Keogh B, et al. Treatment with OPN-305, a humanized anti-Toll-Like receptor-2 antibody, reduces myocardial ischemia/reperfusion injury in pigs. *Circ Cardiovasc Interv* 2012;5(2):279-87.
43. Connolly DJ, O'Neill LA. New developments in Toll-like receptor targeted therapeutics. *Curr Opin Pharmacol* 2012;12(4):510-8.
44. Schroder K, Tschopp J. The inflammasomes. *Cell* 2010;140(6):821-32.
45. Kawaguchi M, Takahashi M, Hata T, et al. Inflammasome activation of cardiac fibroblasts is essential for myocardial ischemia/reperfusion injury. *Circulation* 2011;123(6):594-604.
46. Venkatchalam K, Prabhu SD, Reddy VS, et al. Neutralization of interleukin-18 ameliorates ischemia/reperfusion-induced myocardial injury. *J Biol Chem* 2009;284(12):7853-65.
47. Sandanger O, Ranheim T, Vinge LE, et al. The NLRP3 inflammasome is up-regulated in cardiac fibroblasts and mediates myocardial ischaemia-reperfusion injury. *Cardiovasc Res* 2013;99(1):164-74.
48. van Hout GP, Bosch L, Ellenbroek GH, et al. The selective NLRP3-inflammasome inhibitor MCC950 reduces infarct size and preserves cardiac function in a pig model of myocardial infarction. *Eur Heart J* 2017;38(11):828-36.
49. Xie J, Mendez JD, Mendez-Valenzuela V, Aguilar-Hernandez MM. Cellular signalling of the receptor for advanced glycation end products (RAGE). *Cell Signal* 2013;25(11):2185-97.
50. Volz HC, Laohachewin D, Seidel C, et al. S100A8/A9 aggravates post-ischemic heart failure through activation of RAGE-dependent NF-kappaB signaling. *Basic Res Cardiol* 2012;107(2):250.
51. Ku SH, Hong J, Moon HH, et al. Deoxycholic acid-modified polyethylenimine based nanocarriers for RAGE siRNA therapy in acute myocardial infarction. *Arch Pharm Res* 2015;38(7):1317-24.

52. Suchal K, Malik S, Khan SI, et al. Protective effect of mangiferin on myocardial ischemia-reperfusion injury in streptozotocin-induced diabetic rats: role of AGE-RAGE/MAPK pathways. *Sci Rep* 2017;7:42027.
53. Hill JH, Ward PA. The phlogistic role of C3 leukotactic fragments in myocardial infarcts of rats. *J Exp Med* 1971;133(4):885-900.
54. Yasojima K, Schwab C, McGeer EG, McGeer PL. Human heart generates complement proteins that are upregulated and activated after myocardial infarction. *Circ Res* 1998;83(8):860-9.
55. Dunkelberger JR, Song WC. Complement and its role in innate and adaptive immune responses. *Cell Res* 2010;20(1):34-50.
56. Krijnen PA, Ciurana C, Cramer T, et al. IgM colocalises with complement and C reactive protein in infarcted human myocardium. *J Clin Pathol* 2005;58(4):382-8.
57. Ehrnthaller C, Ignatius A, Gebhard F, Huber-Lang M. New insights of an old defense system: structure, function, and clinical relevance of the complement system. *Mol Med* 2011;17(3-4):317-29.
58. Birdsall HH, Green DM, Trial J, et al. Complement C5a, TGF-beta 1, and MCP-1, in sequence, induce migration of monocytes into ischemic canine myocardium within the first one to five hours after reperfusion. *Circulation* 1997;95(3):684-92.
59. Jordan JE, Montalto MC, Stahl GL. Inhibition of mannose-binding lectin reduces postischemic myocardial reperfusion injury. *Circulation* 2001;104(12):1413-8.
60. Walsh MC, Bourcier T, Takahashi K, et al. Mannose-binding lectin is a regulator of inflammation that accompanies myocardial ischemia and reperfusion injury. *J Immunol* 2005;175(1):541-6.
61. Gorsuch WB, Guikema BJ, Fritzing DC, et al. Humanized cobra venom factor decreases myocardial ischemia-reperfusion injury. *Mol Immunol* 2009;47(2-3):506-10.
62. Vakeva AP, Agah A, Rollins SA, et al. Myocardial infarction and apoptosis after myocardial ischemia and reperfusion: role of the terminal complement components and inhibition by anti-C5 therapy. *Circulation* 1998;97(22):2259-67.
63. De Hoog VC, Timmers L, Van Duijvenvoorde A, et al. Leucocyte expression of complement C5a receptors exacerbates infarct size after myocardial reperfusion injury. *Cardiovasc Res* 2014;103(4):521-9.
64. Pischke SE, Gustavsen A, Orrem HL, et al. Complement factor 5 blockade reduces porcine myocardial infarction size and improves immediate cardiac function. *Basic Res Cardiol* 2017;112(3):20.
65. Weisman HF, Bartow T, Leppo MK, et al. Soluble human complement receptor type 1: in vivo inhibitor of complement suppressing post-ischemic myocardial inflammation and necrosis. *Science* 1990;249(4965):146-51.
66. Horstick G, Heimann A, Gotze O, et al. Intracoronary application of C1 esterase inhibitor improves cardiac function and reduces myocardial necrosis in an experimental model of ischemia and reperfusion. *Circulation* 1997;95(3):701-8.
67. Investigators AA, Armstrong PW, Granger CB, et al. Pexelizumab for acute ST-elevation myocardial infarction in patients undergoing primary percutaneous coronary intervention: a randomized controlled trial. *JAMA* 2007;297(1):43-51.
68. Gordon JW, Shaw JA, Kirshenbaum LA. Multiple facets of NF-kappaB in the heart: to be or not to NF-kappaB. *Circ Res* 2011;108(9):1122-32.
69. Hayden MS, Ghosh S. Shared principles in NF-kappaB signaling. *Cell* 2008;132(3):344-62.
70. Onai Y, Suzuki J, Kakuta T, et al. Inhibition of IkappaB phosphorylation in cardiomyocytes attenuates myocardial ischemia/reperfusion injury. *Cardiovasc Res* 2004;63(1):51-9.
71. Moss NC, Stansfield WE, Willis MS, et al. IKKbeta inhibition attenuates myocardial injury and dysfunction following acute ischemia-reperfusion injury. *Am J Physiol Heart Circ Physiol* 2007;293(4):H2248-53.
72. Zhang XQ, Tang R, Li L, et al. Cardiomyocyte-specific p65 NF-kappaB deletion protects the injured heart by preservation of calcium handling. *Am J Physiol Heart Circ Physiol* 2013;305(7):H1089-97.
73. Morishita R, Sugimoto T, Aoki M, et al. In vivo transfection of cis element "decoy" against nuclear factor-kappaB binding site prevents myocardial infarction. *Nat Med* 1997;3(8):894-9.
74. Luster AD. Chemokines--chemotactic cytokines that mediate inflammation. *N Engl J Med* 1998;338(7):436-45.
75. Zlotnik A, Yoshie O. Chemokines: a new classification system and their role in immunity. *Immunity* 2000;12(2):121-7.
76. Frangogiannis NG. Chemokines in ischemia and reperfusion. *Thromb Haemost* 2007;97(5):738-47.

77. Kumar AG, Ballantyne CM, Michael LH, et al. Induction of monocyte chemoattractant protein-1 in the small veins of the ischemic and reperfused canine myocardium. *Circulation* 1997;95(3):693-700.
78. Dewald O, Zymek P, Winkelmann K, et al. CCL2/Monocyte Chemoattractant Protein-1 regulates inflammatory responses critical to healing myocardial infarcts. *Circ Res* 2005;96(8):881-9.
79. Morimoto H, Takahashi M, Izawa A, et al. Cardiac overexpression of monocyte chemoattractant protein-1 in transgenic mice prevents cardiac dysfunction and remodeling after myocardial infarction. *Circ Res* 2006;99(8):891-9.
80. Liehn EA, Piccinini AM, Koenen RR, et al. A new monocyte chemotactic protein-1/chemokine CC motif ligand-2 competitor limiting neointima formation and myocardial ischemia/reperfusion injury in mice. *J Am Coll Cardiol* 2010;56(22):1847-57.
81. Vajen T, Koenen RR, Werner I, et al. Blocking CCL5-CXCL4 heteromerization preserves heart function after myocardial infarction by attenuating leukocyte recruitment and NETosis. *Sci Rep* 2018;8(1):10647.
82. Montecucco F, Braunersreuther V, Lenglet S, et al. CC chemokine CCL5 plays a central role impacting infarct size and post-infarction heart failure in mice. *Eur Heart J* 2012;33(15):1964-74.
83. Liehn EA, Merx MW, Postea O, et al. Ccr1 deficiency reduces inflammatory remodelling and preserves left ventricular function after myocardial infarction. *J Cell Mol Med* 2008;12(2):496-506.
84. Huang Y, Wang D, Wang X, et al. Abrogation of CC chemokine receptor 9 ameliorates ventricular remodeling in mice after myocardial infarction. *Sci Rep* 2016;6:32660.
85. Ivey CL, Williams FM, Collins PD, et al. Neutrophil chemoattractants generated in two phases during reperfusion of ischemic myocardium in the rabbit. Evidence for a role for C5a and interleukin-8. *J Clin Invest* 1995;95(6):2720-8.
86. Kukielka GL, Smith CW, LaRosa GJ, et al. Interleukin-8 gene induction in the myocardium after ischemia and reperfusion in vivo. *J Clin Invest* 1995;95(1):89-103.
87. Takami M, Terry V, Petruzzelli L. Signaling pathways involved in IL-8-dependent activation of adhesion through Mac-1. *J Immunol* 2002;168(9):4559-66.
88. Boyle EM, Jr., Kovacich JC, Hebert CA, et al. Inhibition of interleukin-8 blocks myocardial ischemia-reperfusion injury. *J Thorac Cardiovasc Surg* 1998;116(1):114-21.
89. Chandrasekar B, Smith JB, Freeman GL. Ischemia-reperfusion of rat myocardium activates nuclear factor-KappaB and induces neutrophil infiltration via lipopolysaccharide-induced CXC chemokine. *Circulation* 2001;103(18):2296-302.
90. Tarzami ST, Miao W, Mani K, et al. Opposing effects mediated by the chemokine receptor CXCR2 on myocardial ischemia-reperfusion injury: recruitment of potentially damaging neutrophils and direct myocardial protection. *Circulation* 2003;108(19):2387-92.
91. Montecucco F, Lenglet S, Braunersreuther V, et al. Single administration of the CXC chemokine-binding protein Evasin-3 during ischemia prevents myocardial reperfusion injury in mice. *Arterioscler Thromb Vasc Biol* 2010;30(7):1371-7.
92. Bujak M, Dobaczewski M, Gonzalez-Quesada C, et al. Induction of the CXC chemokine interferon-gamma-inducible protein 10 regulates the reparative response following myocardial infarction. *Circ Res* 2009;105(10):973-83.
93. Liehn EA, Tuchscheerer N, Kanzler I, et al. Double-edged role of the CXCL12/CXCR4 axis in experimental myocardial infarction. *J Am Coll Cardiol* 2011;58(23):2415-23.
94. Askari AT, Unzek S, Popovic ZB, et al. Effect of stromal-cell-derived factor 1 on stem-cell homing and tissue regeneration in ischaemic cardiomyopathy. *Lancet* 2003;362(9385):697-703.
95. Saxena A, Fish JE, White MD, et al. Stromal cell-derived factor-1alpha is cardioprotective after myocardial infarction. *Circulation* 2008;117(17):2224-31.
96. Hu X, Dai S, Wu WJ, et al. Stromal cell derived factor-1 alpha confers protection against myocardial ischemia/reperfusion injury: role of the cardiac stromal cell derived factor-1 alpha CXCR4 axis. *Circulation* 2007;116(6):654-63.
97. Chen J, Chemaly E, Liang L, et al. Effects of CXCR4 gene transfer on cardiac function after ischemia-reperfusion injury. *Am J Pathol* 2010;176(4):1705-15.
98. Dewald O, Ren G, Duerr GD, et al. Of mice and dogs: species-specific differences in the inflammatory response following myocardial infarction. *Am J Pathol* 2004;164(2):665-77.

99. Nian M, Lee P, Khaper N, Liu P. Inflammatory cytokines and postmyocardial infarction remodeling. *Circ Res* 2004;94(12):1543-53.
100. Siwik DA, Chang DL, Colucci WS. Interleukin-1beta and tumor necrosis factor-alpha decrease collagen synthesis and increase matrix metalloproteinase activity in cardiac fibroblasts in vitro. *Circ Res* 2000;86(12):1259-65.
101. Engel D, Peshock R, Armstrong RC, et al. Cardiac myocyte apoptosis provokes adverse cardiac remodeling in transgenic mice with targeted TNF overexpression. *Am J Physiol Heart Circ Physiol* 2004;287(3):H1303-11.
102. Maekawa N, Wada H, Kanda T, et al. Improved myocardial ischemia/reperfusion injury in mice lacking tumor necrosis factor-alpha. *J Am Coll Cardiol* 2002;39(7):1229-35.
103. Pilz PM, Hamza O, Gidlof O, et al. Remote ischemic preconditioning attenuates adverse cardiac remodeling and preserves left ventricular function in a rat model of reperfused myocardial infarction. *Int J Cardiol* 2019;285:72-9.
104. Rajani SF, Imani A, Faghihi M, et al. Post-infarct morphine treatment mitigates left ventricular remodeling and dysfunction in a rat model of ischemia-reperfusion. *Eur J Pharmacol* 2019;847:61-71.
105. Woodall MC, Woodall BP, Gao E, et al. Cardiac Fibroblast GRK2 Deletion Enhances Contractility and Remodeling Following Ischemia/Reperfusion Injury. *Circ Res* 2016;119(10):1116-27.
106. Sun M, Dawood F, Wen WH, et al. Excessive tumor necrosis factor activation after infarction contributes to susceptibility of myocardial rupture and left ventricular dysfunction. *Circulation* 2004;110(20):3221-8.
107. Monden Y, Kubota T, Tsutsumi T, et al. Soluble TNF receptors prevent apoptosis in infiltrating cells and promote ventricular rupture and remodeling after myocardial infarction. *Cardiovasc Res* 2007;73(4):794-805.
108. Monden Y, Kubota T, Inoue T, et al. Tumor necrosis factor-alpha is toxic via receptor 1 and protective via receptor 2 in a murine model of myocardial infarction. *Am J Physiol Heart Circ Physiol* 2007;293(1):H743-53.
109. Hamid T, Gu Y, Ortines RV, et al. Divergent tumor necrosis factor receptor-related remodeling responses in heart failure: role of nuclear factor-kappaB and inflammatory activation. *Circulation* 2009;119(10):1386-97.
110. Suzuki K, Murtuza B, Smolenski RT, et al. Overexpression of interleukin-1 receptor antagonist provides cardioprotection against ischemia-reperfusion injury associated with reduction in apoptosis. *Circulation* 2001;104(12 Suppl 1):I308-I3.
111. Morton AC, Rothman AM, Greenwood JP, et al. The effect of interleukin-1 receptor antagonist therapy on markers of inflammation in non-ST elevation acute coronary syndromes: the MRC-ILA Heart Study. *Eur Heart J* 2015;36(6):377-84.
112. Abbate A, Van Tassel BW, Biondi-Zoccai G, et al. Effects of interleukin-1 blockade with anakinra on adverse cardiac remodeling and heart failure after acute myocardial infarction [from the Virginia Commonwealth University-Anakinra Remodeling Trial (2) (VCU-ART2) pilot study]. *Am J Cardiol* 2013;111(10):1394-400.
113. Sun X, Wei Z, Li Y, et al. Renal denervation restrains the inflammatory response in myocardial ischemia-reperfusion injury. *Basic Res Cardiol* 2020;115(2):15.
114. Audia JP, Yang XM, Crockett ES, et al. Caspase-1 inhibition by VX-765 administered at reperfusion in P2Y12 receptor antagonist-treated rats provides long-term reduction in myocardial infarct size and preservation of ventricular function. *Basic Res Cardiol* 2018;113(5):32.
115. Abbate A, Salloum FN, Vecile E, et al. Anakinra, a recombinant human interleukin-1 receptor antagonist, inhibits apoptosis in experimental acute myocardial infarction. *Circulation* 2008;117(20):2670-83.
116. Bujak M, Dobaczewski M, Chatila K, et al. Interleukin-1 receptor type 1 signaling critically regulates infarct healing and cardiac remodeling. *Am J Pathol* 2008;173(1):57-67.
117. Gwechenberger M, Mendoza LH, Youker KA, et al. Cardiac myocytes produce interleukin-6 in culture and in viable border zone of reperfused infarctions. *Circulation* 1999;99(4):546-51.
118. Empana JP, Jouven X, Canoui-Poitrine F, et al. C-reactive protein, interleukin 6, fibrinogen and risk of sudden death in European middle-aged men: the PRIME study. *Arterioscler Thromb Vasc Biol* 2010;30(10):2047-52.
119. Interleukin-6 Receptor Mendelian Randomisation Analysis C, Swerdlow DI, Holmes MV, et al. The interleukin-6 receptor as a target for prevention of coronary heart disease: a mendelian randomisation analysis. *Lancet* 2012;379(9822):1214-24.
120. Kleaveland O, Kunszt G, Bratlie M, et al. Effect of a single dose of the interleukin-6 receptor antagonist tocilizumab on inflammation and troponin T release in patients with non-ST-elevation myocardial infarction: a double-blind, randomized, placebo-controlled phase 2 trial. *Eur Heart J* 2016;37(30):2406-13.

121. Fuchs M, Hilfiker A, Kaminski K, et al. Role of interleukin-6 for LV remodeling and survival after experimental myocardial infarction. *FASEB J* 2003;17(14):2118-20.
122. Jong WM, Ten Cate H, Linnenbank AC, et al. Reduced acute myocardial ischemia-reperfusion injury in IL-6-deficient mice employing a closed-chest model. *Inflamm Res* 2016;65(6):489-99.
123. Kobara M, Noda K, Kitamura M, et al. Antibody against interleukin-6 receptor attenuates left ventricular remodeling after myocardial infarction in mice. *Cardiovasc Res* 2010;87(3):424-30.
124. Hartman MH, Vreeswijk-Baudoin I, Groot HE, et al. Inhibition of Interleukin-6 Receptor in a Murine Model of Myocardial Ischemia-Reperfusion. *PLoS One* 2016;11(12):e0167195.
125. Zhang C, He S, Li Y, et al. Bisoprolol protects myocardium cells against ischemia/reperfusion injury by attenuating unfolded protein response in rats. *Sci Rep* 2017;7(1):11859.
126. Birnbaum Y, Tran D, Bajaj M, Ye Y. DPP-4 inhibition by linagliptin prevents cardiac dysfunction and inflammation by targeting the Nlrp3/ASC inflammasome. *Basic Res Cardiol* 2019;114(5):35.
127. Li MO, Flavell RA. Contextual regulation of inflammation: a duet by transforming growth factor-beta and interleukin-10. *Immunity* 2008;28(4):468-76.
128. Lacraz S, Nicod LP, Chicheportiche R, et al. IL-10 inhibits metalloproteinase and stimulates TIMP-1 production in human mononuclear phagocytes. *J Clin Invest* 1995;96(5):2304-10.
129. Frangogiannis NG, Mendoza LH, Lindsey ML, et al. IL-10 is induced in the reperfused myocardium and may modulate the reaction to injury. *J Immunol* 2000;165(5):2798-808.
130. Bagchi AK, Akolkar G, Mandal S, et al. Toll-like receptor 2 dominance over Toll-like receptor 4 in stressful conditions for its detrimental role in the heart. *Am J Physiol Heart Circ Physiol* 2017;312(6):H1238-H47.
131. Yang Z, Zingarelli B, Szabo C. Crucial role of endogenous interleukin-10 production in myocardial ischemia/reperfusion injury. *Circulation* 2000;101(9):1019-26.
132. Krishnamurthy P, Rajasingh J, Lambers E, et al. IL-10 inhibits inflammation and attenuates left ventricular remodeling after myocardial infarction via activation of STAT3 and suppression of HuR. *Circ Res* 2009;104(2):e9-18.
133. Bakhta O, Blanchard S, Guihot AL, et al. Cardioprotective Role of Colchicine Against Inflammatory Injury in a Rat Model of Acute Myocardial Infarction. *J Cardiovasc Pharmacol Ther* 2018;23(5):446-55.
134. Tian Y, Miao B, Charles EJ, et al. Stimulation of the Beta2 Adrenergic Receptor at Reperfusion Limits Myocardial Reperfusion Injury via an Interleukin-10-Dependent Anti-Inflammatory Pathway in the Spleen. *Circ J* 2018;82(11):2829-36.
135. Zymek P, Nah DY, Bujak M, et al. Interleukin-10 is not a critical regulator of infarct healing and left ventricular remodeling. *Cardiovasc Res* 2007;74(2):313-22.
136. Bujak M, Frangogiannis NG. The role of TGF-beta signaling in myocardial infarction and cardiac remodeling. *Cardiovasc Res* 2007;74(2):184-95.
137. Mauviel A. Transforming growth factor-beta: a key mediator of fibrosis. *Methods Mol Med* 2005;117:69-80.
138. Deten A, Holzl A, Leicht M, et al. Changes in extracellular matrix and in transforming growth factor beta isoforms after coronary artery ligation in rats. *J Mol Cell Cardiol* 2001;33(6):1191-207.
139. Frangogiannis NG, Ren G, Dewald O, et al. Critical role of endogenous thrombospondin-1 in preventing expansion of healing myocardial infarcts. *Circulation* 2005;111(22):2935-42.
140. Shi Y, Massague J. Mechanisms of TGF-beta signaling from cell membrane to the nucleus. *Cell* 2003;113(6):685-700.
141. Dobaczewski M, Bujak M, Li N, et al. Smad3 signaling critically regulates fibroblast phenotype and function in healing myocardial infarction. *Circ Res* 2010;107(3):418-28.
142. Lefer AM, Tsao P, Aoki N, Palladino MA, Jr. Mediation of cardioprotection by transforming growth factor-beta. *Science* 1990;249(4964):61-4.
143. Kempf T, Zarbock A, Widera C, et al. GDF-15 is an inhibitor of leukocyte integrin activation required for survival after myocardial infarction in mice. *Nat Med* 2011;17(5):581-8.
144. DeBerge M, Yeap XY, Dehn S, et al. MerTK Cleavage on Resident Cardiac Macrophages Compromises Repair After Myocardial Ischemia Reperfusion Injury. *Circ Res* 2017;121(8):930-40.
145. Tan SM, Zhang Y, Connelly KA, et al. Targeted inhibition of activin receptor-like kinase 5 signaling attenuates cardiac dysfunction following myocardial infarction. *Am J Physiol Heart Circ Physiol* 2010;298(5):H1415-25.

146. Okada H, Takemura G, Kosai K, et al. Postinfarction gene therapy against transforming growth factor-beta signal modulates infarct tissue dynamics and attenuates left ventricular remodeling and heart failure. *Circulation* 2005;111(19):2430-7.
147. Yu CM, Tipoe GL, Wing-Hon Lai K, Lau CP. Effects of combination of angiotensin-converting enzyme inhibitor and angiotensin receptor antagonist on inflammatory cellular infiltration and myocardial interstitial fibrosis after acute myocardial infarction. *J Am Coll Cardiol* 2001;38(4):1207-15.
148. Rainer PP, Hao S, Vanhoutte D, et al. Cardiomyocyte-specific transforming growth factor beta suppression blocks neutrophil infiltration, augments multiple cytoprotective cascades, and reduces early mortality after myocardial infarction. *Circ Res* 2014;114(8):1246-57.
149. Bujak M, Ren G, Kweon HJ, et al. Essential role of Smad3 in infarct healing and in the pathogenesis of cardiac remodeling. *Circulation* 2007;116(19):2127-38.
150. Ikeuchi M, Tsutsui H, Shiomi T, et al. Inhibition of TGF-beta signaling exacerbates early cardiac dysfunction but prevents late remodeling after infarction. *Cardiovasc Res* 2004;64(3):526-35.
151. Andreadou I, Cabrera-Fuentes HA, Devaux Y, et al. Immune cells as targets for cardioprotection: new players and novel therapeutic opportunities. *Cardiovasc Res* 2019;115(7):1117-30.
152. Farbehi N, Patrick R, Dorison A, et al. Single-cell expression profiling reveals dynamic flux of cardiac stromal, vascular and immune cells in health and injury. *Elife* 2019;8.
153. von Hundelshausen P, Weber C. Platelets as immune cells: bridging inflammation and cardiovascular disease. *Circ Res* 2007;100(1):27-40.
154. Gawaz M. Role of platelets in coronary thrombosis and reperfusion of ischemic myocardium. *Cardiovasc Res* 2004;61(3):498-511.
155. Schanze N, Bode C, Duerschmied D. Platelet Contributions to Myocardial Ischemia/Reperfusion Injury. *Front Immunol* 2019;10:1260.
156. Mauler M, Herr N, Schoenichen C, et al. Platelet Serotonin Aggravates Myocardial Ischemia/Reperfusion Injury via Neutrophil Degranulation. *Circulation* 2019;139(7):918-31.
157. Ziegler M, Hohmann JD, Searle AK, et al. A single-chain antibody-CD39 fusion protein targeting activated platelets protects from cardiac ischaemia/reperfusion injury. *Eur Heart J* 2018;39(2):111-6.
158. Swirski FK, Nahrendorf M. Cardioimmunology: the immune system in cardiac homeostasis and disease. *Nat Rev Immunol* 2018;18(12):733-44.
159. Pham CT. Neutrophil serine proteases: specific regulators of inflammation. *Nat Rev Immunol* 2006;6(7):541-50.
160. Chia S, Nagurney JL, Brown DE, et al. Association of leukocyte and neutrophil counts with infarct size, left ventricular function and outcomes after percutaneous coronary intervention for ST-elevation myocardial infarction. *Am J Cardiol* 2009;103(3):333-7.
161. Vinten-Johansen J. Involvement of neutrophils in the pathogenesis of lethal myocardial reperfusion injury. *Cardiovasc Res* 2004;61(3):481-97.
162. Ebnet K, Vestweber D. Molecular mechanisms that control leukocyte extravasation: the selectins and the chemokines. *Histochem Cell Biol* 1999;112(1):1-23.
163. Ma Y, Yabluchanskiy A, Iyer RP, et al. Temporal neutrophil polarization following myocardial infarction. *Cardiovasc Res* 2016;110(1):51-61.
164. Vafadarnejad E, Rizzo G, Krampert L, et al. Dynamics of Cardiac Neutrophil Diversity in Murine Myocardial Infarction. *Circ Res* 2020;127(9):e232-e49.
165. Carbone F, Crowe LA, Roth A, et al. Treatment with anti-RANKL antibody reduces infarct size and attenuates dysfunction impacting on neutrophil-mediated injury. *J Mol Cell Cardiol* 2016;94:82-94.
166. Praetner M, Zuchtriegel G, Holzer M, et al. Plasminogen Activator Inhibitor-1 Promotes Neutrophil Infiltration and Tissue Injury on Ischemia-Reperfusion. *Arterioscler Thromb Vasc Biol* 2018;38(4):829-42.
167. Zhang X, Liu S, Weng X, et al. Brg1 deficiency in vascular endothelial cells blocks neutrophil recruitment and ameliorates cardiac ischemia-reperfusion injury in mice. *Int J Cardiol* 2018;269:250-8.
168. Hiroi T, Wajima T, Negoro T, et al. Neutrophil TRPM2 channels are implicated in the exacerbation of myocardial ischaemia/reperfusion injury. *Cardiovasc Res* 2013;97(2):271-81.

169. Bao J, Sato K, Li M, et al. PR-39 and PR-11 peptides inhibit ischemia-reperfusion injury by blocking proteasome-mediated I kappa B alpha degradation. *Am J Physiol Heart Circ Physiol* 2001;281(6):H2612-8.
170. Amsterdam EA, Pan HL, Rendig SV, et al. Limitation of myocardial infarct size in pigs with a dual lipoxygenase-cyclooxygenase blocking agent by inhibition of neutrophil activity without reduction of neutrophil migration. *J Am Coll Cardiol* 1993;22(6):1738-44.
171. Boufenzar A, Lemarie J, Simon T, et al. TREM-1 Mediates Inflammatory Injury and Cardiac Remodeling Following Myocardial Infarction. *Circ Res* 2015;116(11):1772-82.
172. Marinkovic G, Grauen Larsen H, Yndigeegn T, et al. Inhibition of pro-inflammatory myeloid cell responses by short-term S100A9 blockade improves cardiac function after myocardial infarction. *Eur Heart J* 2019;40(32):2713-23.
173. Ali M, Pulli B, Courties G, et al. Myeloperoxidase Inhibition Improves Ventricular Function and Remodeling After Experimental Myocardial Infarction. *JACC Basic Transl Sci* 2016;1(7):633-43.
174. Ma XL, Weyrich AS, Lefer DJ, et al. Monoclonal antibody to L-selectin attenuates neutrophil accumulation and protects ischemic reperfused cat myocardium. *Circulation* 1993;88(2):649-58.
175. Weyrich AS, Ma XY, Lefer DJ, et al. In vivo neutralization of P-selectin protects feline heart and endothelium in myocardial ischemia and reperfusion injury. *J Clin Invest* 1993;91(6):2620-9.
176. Briaud SA, Ding ZM, Michael LH, et al. Leukocyte trafficking and myocardial reperfusion injury in ICAM-1/P-selectin-knockout mice. *Am J Physiol Heart Circ Physiol* 2001;280(1):H60-7.
177. Aversano T, Zhou W, Nedelman M, et al. A chimeric IgG4 monoclonal antibody directed against CD18 reduces infarct size in a primate model of myocardial ischemia and reperfusion. *J Am Coll Cardiol* 1995;25(3):781-8.
178. Arai M, Lefer DJ, So T, et al. An anti-CD18 antibody limits infarct size and preserves left ventricular function in dogs with ischemia and 48-hour reperfusion. *J Am Coll Cardiol* 1996;27(5):1278-85.
179. Palazzo AJ, Jones SP, Girod WG, et al. Myocardial ischemia-reperfusion injury in CD18- and ICAM-1-deficient mice. *Am J Physiol* 1998;275(6):H2300-7.
180. Baran KW, Nguyen M, McKendall GR, et al. Double-blind, randomized trial of an anti-CD18 antibody in conjunction with recombinant tissue plasminogen activator for acute myocardial infarction: limitation of myocardial infarction following thrombolysis in acute myocardial infarction (LIMIT AMI) study. *Circulation* 2001;104(23):2778-83.
181. Faxon DP, Gibbons RJ, Chronos NA, et al. The effect of blockade of the CD11/CD18 integrin receptor on infarct size in patients with acute myocardial infarction treated with direct angioplasty: the results of the HALT-MI study. *J Am Coll Cardiol* 2002;40(7):1199-204.
182. Horckmans M, Ring L, Duchene J, et al. Neutrophils orchestrate post-myocardial infarction healing by polarizing macrophages towards a reparative phenotype. *Eur Heart J* 2017;38(3):187-97.
183. Jung K, Kim P, Leuschner F, et al. Endoscopic time-lapse imaging of immune cells in infarcted mouse hearts. *Circ Res* 2013;112(6):891-9.
184. Ensan S, Li A, Besla R, et al. Self-renewing resident arterial macrophages arise from embryonic CX3CR1(+) precursors and circulating monocytes immediately after birth. *Nat Immunol* 2016;17(2):159-68.
185. Bajpai G, Bredemeyer A, Li W, et al. Tissue Resident CCR2- and CCR2+ Cardiac Macrophages Differentially Orchestrate Monocyte Recruitment and Fate Specification Following Myocardial Injury. *Circ Res* 2019;124(2):263-78.
186. Mouton AJ, DeLeon-Pennell KY, Rivera Gonzalez OJ, et al. Mapping macrophage polarization over the myocardial infarction time continuum. *Basic Res Cardiol* 2018;113(4):26.
187. Dick SA, Macklin JA, Nejat S, et al. Self-renewing resident cardiac macrophages limit adverse remodeling following myocardial infarction. *Nat Immunol* 2019;20(1):29-39.
188. Leuschner F, Rauch PJ, Ueno T, et al. Rapid monocyte kinetics in acute myocardial infarction are sustained by extramedullary monocytopenesis. *J Exp Med* 2012;209(1):123-37.
189. Nahrendorf M, Swirski FK. Monocyte and macrophage heterogeneity in the heart. *Circ Res* 2013;112(12):1624-33.
190. Hilgendorf I, Gerhardt LM, Tan TC, et al. Ly-6Chigh monocytes depend on Nr4a1 to balance both inflammatory and reparative phases in the infarcted myocardium. *Circ Res* 2014;114(10):1611-22.
191. Rizzo G, Vafadarnejad E, Arampatzi P, et al. Single-cell transcriptomic profiling maps monocyte/macrophage transitions after myocardial infarction in mice. *bioRxiv* 2020.
192. Tsujioka H, Imanishi T, Ikejima H, et al. Impact of heterogeneity of human peripheral blood monocyte subsets on myocardial salvage in patients with primary acute myocardial infarction. *J Am Coll Cardiol* 2009;54(2):130-8.

193. van der Laan AM, Hirsch A, Robbers LF, et al. A proinflammatory monocyte response is associated with myocardial injury and impaired functional outcome in patients with ST-segment elevation myocardial infarction: monocytes and myocardial infarction. *Am Heart J* 2012;163(1):57-65 e2.
194. Miller EJ, Li J, Leng L, et al. Macrophage migration inhibitory factor stimulates AMP-activated protein kinase in the ischaemic heart. *Nature* 2008;451(7178):578-82.
195. Gao XM, Liu Y, White D, et al. Deletion of macrophage migration inhibitory factor protects the heart from severe ischemia-reperfusion injury: a predominant role of anti-inflammation. *J Mol Cell Cardiol* 2011;50(6):991-9.
196. Liehn EA, Kanzler I, Konschalla S, et al. Compartmentalized protective and detrimental effects of endogenous macrophage migration-inhibitory factor mediated by CXCR2 in a mouse model of myocardial ischemia/reperfusion. *Arterioscler Thromb Vasc Biol* 2013;33(9):2180-6.
197. Wan E, Yeap XY, Dehn S, et al. Enhanced efferocytosis of apoptotic cardiomyocytes through myeloid-epithelial-reproductive tyrosine kinase links acute inflammation resolution to cardiac repair after infarction. *Circ Res* 2013;113(8):1004-12.
198. Courties G, Heidt T, Sebas M, et al. In vivo silencing of the transcription factor IRF5 reprograms the macrophage phenotype and improves infarct healing. *J Am Coll Cardiol* 2014;63(15):1556-66.
199. Harel-Adar T, Ben Mordechai T, Amsalem Y, et al. Modulation of cardiac macrophages by phosphatidylserine-presenting liposomes improves infarct repair. *Proc Natl Acad Sci U S A* 2011;108(5):1827-32.
200. Nakano Y, Matoba T, Tokutome M, et al. Nanoparticle-Mediated Delivery of Irbesartan Induces Cardioprotection from Myocardial Ischemia-Reperfusion Injury by Antagonizing Monocyte-Mediated Inflammation. *Sci Rep* 2016;6:29601.
201. de Jong RCM, Pluijmert NJ, de Vries MR, et al. Annexin A5 reduces infarct size and improves cardiac function after myocardial ischemia-reperfusion injury by suppression of the cardiac inflammatory response. *Sci Rep* 2018;8(1):6753.
202. Pluijmert NJ, de Jong RCM, de Vries MR, et al. Phosphorylcholine Antibodies Preserve Cardiac Function and Reduce Infarct Size by Attenuating the Post-Ischemic Inflammatory Response. *JACC Basic Transl Sci* 2020;5(12):1228-39.
203. Leuschner F, Dutta P, Gorbатов R, et al. Therapeutic siRNA silencing in inflammatory monocytes in mice. *Nat Biotechnol* 2011;29(11):1005-10.
204. Tokutome M, Matoba T, Nakano Y, et al. Peroxisome proliferator-activated receptor-gamma targeting nanomedicine promotes cardiac healing after acute myocardial infarction by skewing monocyte/macrophage polarization in preclinical animal models. *Cardiovasc Res* 2019;115(2):419-31.
205. Wang Z, Huang S, Sheng Y, et al. Topiramate modulates post-infarction inflammation primarily by targeting monocytes or macrophages. *Cardiovasc Res* 2017;113(5):475-87.
206. Weirather J, Hofmann UD, Beyersdorf N, et al. Foxp3+ CD4+ T cells improve healing after myocardial infarction by modulating monocyte/macrophage differentiation. *Circ Res* 2014;115(1):55-67.
207. Zhou LS, Zhao GL, Liu Q, et al. Silencing collapsin response mediator protein-2 reprograms macrophage phenotype and improves infarct healing in experimental myocardial infarction model. *J Inflamm (Lond)* 2015;12:11.
208. Schloss MJ, Horckmans M, Guillaumat-Prats R, et al. 2-Arachidonoylglycerol mobilizes myeloid cells and worsens heart function after acute myocardial infarction. *Cardiovasc Res* 2019;115(3):602-13.
209. Shiraishi M, Shintani Y, Shintani Y, et al. Alternatively activated macrophages determine repair of the infarcted adult murine heart. *J Clin Invest* 2016;126(6):2151-66.
210. Hofmann U, Beyersdorf N, Weirather J, et al. Activation of CD4+ T lymphocytes improves wound healing and survival after experimental myocardial infarction in mice. *Circulation* 2012;125(13):1652-63.
211. Boag SE, Das R, Shmeleva EV, et al. T lymphocytes and fractalkine contribute to myocardial ischemia/reperfusion injury in patients. *J Clin Invest* 2015;125(8):3063-76.
212. Choo EH, Lee JH, Park EH, et al. Infarcted Myocardium-Primed Dendritic Cells Improve Remodeling and Cardiac Function After Myocardial Infarction by Modulating the Regulatory T Cell and Macrophage Polarization. *Circulation* 2017;135(15):1444-57.
213. Mathes D, Weirather J, Nordbeck P, et al. CD4(+) Foxp3(+) T-cells contribute to myocardial ischemia-reperfusion injury. *J Mol Cell Cardiol* 2016;101:99-105.

214. Zouggari Y, Ait-Oufella H, Bonnin P, et al. B lymphocytes trigger monocyte mobilization and impair heart function after acute myocardial infarction. *Nat Med* 2013;19(10):1273-80.
215. Horckmans M, Bianchini M, Santovito D, et al. Pericardial Adipose Tissue Regulates Granulopoiesis, Fibrosis, and Cardiac Function After Myocardial Infarction. *Circulation* 2018;137(9):948-60.
216. Anzai A, Anzai T, Nagai S, et al. Regulatory role of dendritic cells in postinfarction healing and left ventricular remodeling. *Circulation* 2012;125(10):1234-45.
217. Liu H, Gao W, Yuan J, et al. Exosomes derived from dendritic cells improve cardiac function via activation of CD4(+) T lymphocytes after myocardial infarction. *J Mol Cell Cardiol* 2016;91:123-33.
218. Xue J, Ge H, Lin Z, et al. The role of dendritic cells regulated by HMGB1/TLR4 signalling pathway in myocardial ischaemia reperfusion injury. *J Cell Mol Med* 2019;23(4):2849-62.
219. Frangogiannis NG, Lindsey ML, Michael LH, et al. Resident cardiac mast cells degranulate and release preformed TNF-alpha, initiating the cytokine cascade in experimental canine myocardial ischemia/reperfusion. *Circulation* 1998;98(7):699-710.
220. Somasundaram P, Ren G, Nagar H, et al. Mast cell tryptase may modulate endothelial cell phenotype in healing myocardial infarcts. *J Pathol* 2005;205(1):102-11.
221. Ngkelo A, Richart A, Kirk JA, et al. Mast cells regulate myofilament calcium sensitization and heart function after myocardial infarction. *J Exp Med* 2016;213(7):1353-74.
222. Hooshdaran B, Kolpakov MA, Guo X, et al. Dual inhibition of cathepsin G and chymase reduces myocyte death and improves cardiac remodeling after myocardial ischemia reperfusion injury. *Basic Res Cardiol* 2017;112(6):62.
223. Aldi S, Marino A, Tomita K, et al. E-NTPDase1/CD39 modulates renin release from heart mast cells during ischemia/reperfusion: a novel cardioprotective role. *FASEB J* 2015;29(1):61-9.
224. Marino A, Sakamoto T, Robador PA, et al. S1P receptor 1-Mediated Anti-Renin-Angiotensin System Cardioprotection: Pivotal Role of Mast Cell Aldehyde Dehydrogenase Type 2. *J Pharmacol Exp Ther* 2017;362(2):230-42.
225. Prabhu SD, Frangogiannis NG. The Biological Basis for Cardiac Repair After Myocardial Infarction: From Inflammation to Fibrosis. *Circ Res* 2016;119(1):91-112.
226. Frangogiannis NG. Cardiac fibrosis: Cell biological mechanisms, molecular pathways and therapeutic opportunities. *Mol Aspects Med* 2019;65:70-99.
227. Ruiz-Villalba A, Simon AM, Pogontke C, et al. Interacting resident epicardium-derived fibroblasts and recruited bone marrow cells form myocardial infarction scar. *J Am Coll Cardiol* 2015;65(19):2057-66.
228. Willems IE, Havenith MG, De Mey JG, Daemen MJ. The alpha-smooth muscle actin-positive cells in healing human myocardial scars. *Am J Pathol* 1994;145(4):868-75.
229. Frangogiannis NG. The Functional Pluralism of Fibroblasts in the Infarcted Myocardium. *Circ Res* 2016;119(10):1049-51.
230. Nakaya M, Watari K, Tajima M, et al. Cardiac myofibroblast engulfment of dead cells facilitates recovery after myocardial infarction. *J Clin Invest* 2017;127(1):383-401.
231. Kong P, Shinde AV, Su Y, et al. Opposing Actions of Fibroblast and Cardiomyocyte Smad3 Signaling in the Infarcted Myocardium. *Circulation* 2018;137(7):707-24.
232. Villalobos E, Criollo A, Schiattarella GG, et al. Fibroblast Primary Cilia Are Required for Cardiac Fibrosis. *Circulation* 2019;139(20):2342-57.
233. Jung M, Ma Y, Iyer RP, et al. IL-10 improves cardiac remodeling after myocardial infarction by stimulating M2 macrophage polarization and fibroblast activation. *Basic Res Cardiol* 2017;112(3):33.
234. Petz A, Grandoch M, Gorski DJ, et al. Cardiac Hyaluronan Synthesis Is Critically Involved in the Cardiac Macrophage Response and Promotes Healing After Ischemia Reperfusion Injury. *Circ Res* 2019;124(10):1433-47.
235. Dobaczewski M, Akrivakis S, Nasser K, et al. Vascular mural cells in healing canine myocardial infarcts. *J Histochem Cytochem* 2004;52(8):1019-29.
236. Sandhu R, Teichert-Kuliszewska K, Nag S, et al. Reciprocal regulation of angiotensin-1 and angiotensin-2 following myocardial infarction in the rat. *Cardiovasc Res* 2004;64(1):115-24.
237. Zymek P, Bujak M, Chatila K, et al. The role of platelet-derived growth factor signaling in healing myocardial infarcts. *J Am Coll Cardiol* 2006;48(11):2315-23.

238. Wang Y, Liu J, Kong Q, et al. Cardiomyocyte-specific deficiency of HSPB1 worsens cardiac dysfunction by activating NFkappaB-mediated leucocyte recruitment after myocardial infarction. *Cardiovasc Res* 2019;115(1):154-67.
239. Nathan C. Points of control in inflammation. *Nature* 2002;420(6917):846-52.
240. Christia P, Frangogiannis NG. Targeting inflammatory pathways in myocardial infarction. *Eur J Clin Invest* 2013;43(9):986-95.
241. Cleutjens JP, Blankesteijn WM, Daemen MJ, Smits JF. The infarcted myocardium: simply dead tissue, or a lively target for therapeutic interventions. *Cardiovasc Res* 1999;44(2):232-41.
242. Bergmann O, Bhardwaj RD, Bernard S, et al. Evidence for cardiomyocyte renewal in humans. *Science* 2009;324(5923):98-102.
243. Tzahor E, Poss KD. Cardiac regeneration strategies: Staying young at heart. *Science* 2017;356(6342):1035-9.
244. Davidson SM, Ferdinandy P, Andreadou I, et al. Multitarget Strategies to Reduce Myocardial Ischemia/Reperfusion Injury: JACC Review Topic of the Week. *J Am Coll Cardiol* 2019;73(1):89-99.
245. Libby P, Maroko PR, Bloor CM, et al. Reduction of experimental myocardial infarct size by corticosteroid administration. *J Clin Invest* 1973;52(3):599-607.
246. Roberts R, DeMello V, Sobel BE. Deleterious effects of methylprednisolone in patients with myocardial infarction. *Circulation* 1976;53(3 Suppl):I204-6.
247. Kloner RA, Fishbein MC, Lew H, et al. Mummification of the infarcted myocardium by high dose corticosteroids. *Circulation* 1978;57(1):56-63.
248. Hausenloy DJ, Garcia-Dorado D, Botker HE, et al. Novel targets and future strategies for acute cardioprotection: Position Paper of the European Society of Cardiology Working Group on Cellular Biology of the Heart. *Cardiovasc Res* 2017;113(6):564-85.
249. Lindsey ML, Bolli R, Cauty JM, Jr., et al. Guidelines for experimental models of myocardial ischemia and infarction. *Am J Physiol Heart Circ Physiol* 2018;314(4):H812-H38.
250. Pluijmert NJ, Bart CI, Bax WH, et al. Effects on cardiac function, remodeling and inflammation following myocardial ischemia-reperfusion injury or unreperfused myocardial infarction in hypercholesterolemic APOE*3-Leiden mice. *Sci Rep* 2020;10(1):16601.
251. Merz SF, Korste S, Bornemann L, et al. Contemporaneous 3D characterization of acute and chronic myocardial I/R injury and response. *Nat Commun* 2019;10(1):2312.
252. Huang S, Frangogiannis NG. Anti-inflammatory therapies in myocardial infarction: failures, hopes and challenges. *Br J Pharmacol* 2018;175(9):1377-400.
253. Ridker PM, Everett BM, Thuren T, et al. Antiinflammatory Therapy with Canakinumab for Atherosclerotic Disease. *N Engl J Med* 2017;377(12):1119-31.
254. Padfield GJ, Din JN, Koushiappi E, et al. Cardiovascular effects of tumour necrosis factor alpha antagonism in patients with acute myocardial infarction: a first in human study. *Heart* 2013;99(18):1330-5.
255. de Zwaan C, Kleine AH, Diris JH, et al. Continuous 48-h C1-inhibitor treatment, following reperfusion therapy, in patients with acute myocardial infarction. *Eur Heart J* 2002;23(21):1670-7.
256. Jones DA, Khambata RS, Andiapien M, et al. Intracoronary nitrite suppresses the inflammatory response following primary percutaneous coronary intervention. *Heart* 2017;103(7):508-16.
257. Piot C, Croisille P, Staat P, et al. Effect of cyclosporine on reperfusion injury in acute myocardial infarction. *N Engl J Med* 2008;359(5):473-81.
258. Cung TT, Morel O, Cayla G, et al. Cyclosporine before PCI in Patients with Acute Myocardial Infarction. *N Engl J Med* 2015;373(11):1021-31.
259. Tardif JC, Kouz S, Waters DD, et al. Efficacy and Safety of Low-Dose Colchicine after Myocardial Infarction. *N Engl J Med* 2019;381(26):2497-505.
260. Nidorf SM, Fiolet ATL, Mosterd A, et al. Colchicine in Patients with Chronic Coronary Disease. *N Engl J Med* 2020;383(19):1838-47.