



Universiteit  
Leiden  
The Netherlands

# **The role of glomerular filtration and active tubular secretion in predicting renal clearance of drugs in children using population pharmacokinetic and physiology-based pharmacokinetic modeling approaches: unspinning the yarn**

Cristea, S.

## **Citation**

Cristea, S. (2021, June 16). *The role of glomerular filtration and active tubular secretion in predicting renal clearance of drugs in children using population pharmacokinetic and physiology-based pharmacokinetic modeling approaches: unspinning the yarn*. Retrieved from <https://hdl.handle.net/1887/3188573>

Version: Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/3188573>

**Note:** To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <https://hdl.handle.net/1887/3188573> holds various files of this Leiden University dissertation.

**Author:** Cristea, S.

**Title:** The role of glomerular filtration and active tubular secretion in predicting renal clearance of drugs in children using population pharmacokinetic and physiology-based pharmacokinetic modeling approaches: unspinning the yarn

**Issue Date:** 2021-06-16

**The role of glomerular filtration and active tubular secretion in predicting renal clearance of drugs in children using population pharmacokinetic and physiology-based pharmacokinetic modeling approaches**

Unspinning the yarn

1. Model-driven dose optimization supports an effective and safe treatment from the start of therapy in special populations. *This thesis*
2. PBPK principles are a prerequisite to establish guidelines for dosing renally excreted drugs in children for drugs with various properties. *This thesis*
3. Knowing the unbound drug fraction in adults is essential to accurately scale pediatric drug doses by the maturation function of glomerular filtration rate for drugs mainly eliminated by glomerular filtration. *This thesis*
4. Ignoring the ontogeny of renal transporters for children younger than 2 years leads to inaccurate renal clearance predictions for drugs that are mainly eliminated by active tubular secretion. *This thesis*
5. Transporter ontogeny functions that cannot be measured *in vivo* can be derived from pediatric PK profiles by combining mechanistic insight of PBPK models with population PK estimation methods. *This thesis*
6. Trough concentrations are overemphasized as surrogates for safe and effective antibiotic exposure and treatment. *Modified from Neely, 2013*
7. For drugs that are being subject to active tubular secretion the proportionality between unbound drug fraction and renal clearance changes with extraction ratio as a consequence of the blood flow limitation. *Modified from Levy, 1980*
8. The audience for nifty modeling methods and techniques is not as large as the audience that would benefit from understanding and applying model-informed precision dosing in the clinic. *Modified from Henning, 2020*
9. Following up on the work of others can be more of a challenge than a head-start.
10. Great colleagues who cook well are an asset to any office.
11. Never look into your modelling results as in the mirror of Erised. *Modified from Harry Potter*
12. The right data are indispensable for a good model. The right analysis methods and the right graphics are indispensable for a good modeler. *Modified from Vandemeulebroecke 2019*