



Patterns of conflict-related trauma exposure and their relation to psychopathology: A person-centered analysis in a population-based sample from eastern DRC



Lars Dumke^{a,*}, Roos van der Haer^b, Carlo Koos^c, Tobias Hecker^a

^a Department of Psychology, Bielefeld University, Bielefeld, Germany

^b Institute of Political Science, University of Leiden, Leiden, Netherlands

^c Chr. Michelsen Institute, Bergen, Norway

ARTICLE INFO

Keywords:

Trauma exposure
PTSD
Latent class analysis
Democratic Republic of Congo
Violence
Trauma patterns

ABSTRACT

Populations in war-torn regions are exposed to a wide array of traumatic events that can cause an enormous psychological burden. Individual characteristics influence the likelihood of being exposed to certain events, pointing to systematic interindividual differences in trauma exposure. However, there is a dearth of studies examining potential patterns of trauma exposure in war regions. In this cross-sectional epidemiological study, we applied a person-centered approach to identify patterns in the exposure to conflict-related traumatic events and determine their impact on commonly reported mental health problems in a population-based sample ($N = 1000$) from the eastern Democratic Republic of Congo. We implemented multi-stage random cluster sampling to randomly select adults from 100 villages. Of 1000 adults ($M_{\text{age}} = 43.19$ years) included in the study, 50% were female. Results showed high prevalence of PTSD (17.0%), depression (27.8%), anxiety (25.4%) and suicidality (15.1%) following exposure to conflict-related traumatic events since 2002. Latent Class Analysis identified three distinct classes of trauma exposure: Class 1 “*low-trauma-exposure*” (51.4%, $n = 514$) was characterized by the lowest probabilities of trauma exposure. Class 2 “*non-physical-trauma*” (39.1%, $n = 391$) consisted of individuals with a high probability for exposure to non-physical trauma types only. Class 3 “*interpersonal-trauma*” (9.5%, $n = 95$) had the overall highest probability of exposure to traumatic events and was the only class affected by interpersonal-trauma types. Class membership was related to gender, age and place of living. Vulnerability to mental health problems increased from *low-trauma-exposure* to *non-physical-trauma* to *interpersonal-trauma* class. Our findings indicate that the exposure to traumatic events in conflict-affected populations underlies distinct patterns, with interpersonal trauma as a distinguishing marker. Vulnerability to psychopathology varies with trauma patterns, revealing patterns that include both non-physical and interpersonal traumata as most detrimental for mental health. Identification of underlying trauma patterns and their effects may improve mental health care in war-affected populations.

1. Introduction

Mental health research has highlighted the dire psychological burden for populations affected by armed conflict (De Jong and Komproe, 2002; Steel et al., 2009). Population-based epidemiological studies have revealed the full scale of detrimental outcomes. Across conflict-affected regions prevalence for psychopathology ranged from 11% to 50% for PTSD, 5%–40% for depression, 10%–40% for anxiety disorder and 11%–26% for suicidality (De Jong et al., 2003; Farhood and Dimassi, 2012; Johnson et al., 2008, 2010; Lopes Cardozo et al., 2000; Pham et al., 2004,

2010; Stammel et al., 2013). Exposure to conflict-related traumatic events proved to be a major risk factor (Charlson et al., 2012; Pham et al., 2004; Steel et al., 2009). The psychological consequences of these events are typically evaluated by means of variable-oriented approaches. These approaches analyze, among others, the effect of cumulative trauma exposure or of individual events. For instance, cumulative trauma exposure has been associated with increased vulnerability for PTSD and depression, also referred to as *dose-effect* relation (Farhood and Dimassi, 2012; Lopes Cardozo et al., 2000; Neuner et al., 2004; Pham et al., 2004). Similarly, individual types of traumatic events vary in their potential to

* Corresponding author. Department of Psychology, Clinical Psychology and Psychotherapy, Bielefeld University, P.O. Box 100131, 33501, Bielefeld, Germany.
E-mail address: lars.dumke@uni-bielefeld.de (L. Dumke).

contribute to the development of mental health disorders (Johnson et al., 2008; Lopes Cardozo et al., 2000; Pham et al., 2004).

On closer inspection, both variable-oriented approaches have methodological limitations and may not reflect the reality of conflict-affected regions (O'Donnell et al., 2017). When investigating cumulative trauma exposure, aggregation of events leads to a loss of information about the actual experiences and in turn the simplistic assumption of identical effects of individual traumatic events (Hagan et al., 2016). Furthermore, analysis of cumulative trauma exposure does not reflect existing research suggesting systematic interindividual variations in the number and types of conflict-related trauma along socio-demographic lines, e.g. women facing higher risks to experience conflict-related sexual violence (Johnson et al., 2008, 2010; Pham et al., 2004). At the same time, examining the effects of individual trauma types also has limitations. For example, persons classified as exposed to a specific traumatic event may in fact have heterogeneous traumatic histories, ranging from single incident to long-term exposure that may include other types of trauma. Consequently, the prevailing variable-oriented approaches may not be able to adequately evaluate the complexity of trauma exposure and its effects in conflict-affected populations. A person-oriented approach, grouping individuals into classes of exposure, may be able to overcome some of the aforementioned limitations. It may capture more of the heterogeneity in trauma exposure and also provide the opportunity to gain insights into systematic interindividual patterns of traumatic histories and their consequences on the mental health of the general population of conflict-affected regions (O'Donnell et al., 2017; Sengoelge et al., 2019). Building in-depth knowledge from population-based studies about patterns of trauma exposure and their relation to the development of psychopathology is crucial for identifying specific vulnerable populations and tailoring effective psychological support (De Jong and Komproe, 2002; Gibbs et al., 2020; Lopes Cardozo et al., 2000). This is even more essential in unstable regions with recurring incidence of armed violence, where the availability of, and access to, psychological support is extremely limited.

The objectives of the present study were twofold. First, to obtain recent and representative data on the prevalence and association of conflict-related traumatic events and mental health problems among the general population of South Kivu, DRC, a region immensely affected by various armed groups. We focused on the commonly reported mental health problems after exposure to conflict, namely PTSD, depression, anxiety disorder and suicidality. We hypothesized strong relations between experienced trauma types and psychopathology. Second, to adequately unravel the complex relation between trauma exposure and psychopathology in conflict-affected populations, the aim of the current study was to identify common patterns of conflict-related trauma exposure in the general population. We hypothesized that different patterns of trauma exposure have distinct consequences for mental health. We expected to observe sociodemographic differences, particularly between gender and age groups, in the prevalence of specific trauma types, patterns of trauma exposure and psychopathology.

2. Method

2.1. Design and sampling

This research is based on a cross-sectional study conducted with a population-based sample (N = 1000) from South Kivu region in the eastern DRC from March to May 2017. South Kivu was chosen as the survey site, due to its large proportion of the population of eastern DRC and its central role in the Congo Wars (Johnson et al., 2010). A multi-level randomized cluster sampling approach (see Fig. 1) based on the administrative hierarchy of political subdivisions in the DRC was employed (De Juan and Koos, 2019). South Kivu region is in descending order subdivided into territories, sectors, groupings and villages. Initially, the study was to take place in all territories and sectors; however due to lack of comparability and acute security threats, two of eight




Fig. 1. Design and sampling procedures.

territories and four of 14 sectors of the remaining six territories were excluded from the sampling. In Shabunda territory inaccessibility due to ongoing armed conflict and lack of infrastructure prevented data assessment. Idjwi territory, an isolated island in Lake Kivu, was mostly unaffected by armed conflict and excluded from the sampling as it is therefore not comparable to the political and cultural structures in the other territories (Autesserre, 2017). 33 groupings were randomly chosen from the selected ten sectors. The number of selected groupings within each selected sector was calculated using Probability Proportional to Size sampling. From each selected grouping, three villages were selected at random (in one grouping four villages were sampled), resulting in the selection of 100 villages. Due to the unpredictable security situation, random sampling included three replacement groupings per sector and three replacement villages per grouping. For security reasons, ten percent of initially sampled villages had to be replaced with replacement villages. In each village ten households were randomly selected. Additionally, a list of five randomly selected replacement households per village was created beforehand, in case a household was not available. The initial

response rate on household level was 73%. The remaining households were substituted by replacement households. Within each household, the head of the household or their spouse was randomly selected. If none of them was available, any person ≥ 18 years was randomly chosen. Female enumerators interviewed women and male enumerators interviewed men. Data assessment took place with the assistance of a local research organization (Research Initiatives for Social Development, Bukavu, DRC). The multi-level randomized cluster sampling ensured representativity of the sample to the general population of South Kivu.

2.2. Participants and procedures

The team of experienced local researchers received a four-day workshop to familiarize with the specific methods, instruments and ethical guidelines used in this study. Introduction as well as questionnaire administration procedures were standardized, and consent of all participants was obtained orally prior to interviews. To evaluate the feasibility of questionnaire implementation, the questionnaires were pilot tested before survey administration. All measures were translated into French and Swahili and carried out verbally in one-to-one settings. In total 1000 participants between the age of 18 and 100 years ($M = 43.19$, $SD = 16.83$) completed the assessment. The sociodemographic characteristics can be found in [Table 1](#). Ethical approval was granted by the Ethics Committee of the University of Konstanz, Germany.

2.3. Measures

Traumatic Events. Based on literature review of previous research from war-affected countries ([Neuner et al., 2004](#); [Farhood et al., 2006](#); [Ibrahim and Hassan, 2017](#)), including research from the DRC ([Pham et al., 2010](#); [Johnson et al., 2010](#); [Ainamani et al., 2020](#)), a list of six different trauma types was compiled (see [Table 2](#)). The selection includes trauma types with high and low prevalence and followed recommendations to reflect context-specific evidence ([Netland, 2005](#)). Individuals were asked to indicate direct experience of any of the traumatic events on this list. Events were only recorded if they occurred after 2002 and if they were related to violence by armed groups. Furthermore, an additional variable describing the witnessing of traumatic events was calculated by asking participants whether they witnessed any of the six trauma types

happening to another person.

Refugee Health Screener (RHS-15). The RHS-15 ([Hollifield et al., 2013](#)) includes four items based on three core PTSD symptoms clusters: Re-experiencing (two items), arousal (one item), and negative cognition and mood (one item). Items assessed symptoms occurring in the previous month and were rated on a five-point Likert scale from 0 (not at all) to 4 (extremely). When an item was rated 2 (moderately) or higher, the associated symptom cluster was considered to be present. Respondents screened positive for PTSD if all three symptom clusters were fulfilled and the DSM-5 A-criterion was met. In the current study, the DSM-5 A-criterion was met if the respondent reported direct exposure to or witnessing any of the events on the traumatic events list. The RHS-15 has previously been used in various conflict-affected populations, e.g. in samples from Iraq, Myanmar and Syria ([Fellmeth et al., 2018](#); [Hollifield et al., 2013](#); [Kaltenbach et al., 2017](#)). The Cronbach's alpha in the current study was 0.83.

Patient Health Questionnaire (PHQ-4). The PHQ-4 is an ultra-short screening tool for depression and anxiety ([Kroenke et al., 2009](#)). It is a well validated measure and has previously been used in similar cultural contexts, large-scale household surveys and with individuals affected by conflict ([Kocalevent et al., 2014](#); [Kuringe et al., 2019](#); [Löwe et al., 2010](#)). The questionnaire consists of two subscales (depression, anxiety) with two items each. Items are rated on a four-point Likert scale from 0 (not at all) to 3 (nearly every day). Scores ≥ 3 on the respective subscale indicated the heightened risk of a depressive or anxiety disorder. In our sample, the Cronbach's alpha was .73 and .72 for the depression and the anxiety subscale, respectively. In addition, participants were asked whether they have ever felt suicidal in the last twelve months ("Have you felt suicidal in the last 12 months?").

2.4. Data analysis

Statistical testing was conducted using IBM SPSS 26. Simple logistic regression analyses were used to calculate unadjusted Odds Ratios (OR) regarding the likelihood of psychopathology in relation to exposure to specific traumatic event types. Latent Class Analysis was conducted to identify classes of individuals with similar patterns of trauma exposure. Using the statistical software JMP®, [Version 15](#) latent classes were constructed based on participants answers on seven binary variables

Table 1
Socio-demographics and prevalence rates of mental health disorders by trauma exposure class.

	Overall sample N = 1000 N (%)	Low-trauma-exposure class	Non-physical- trauma class	Interpersonal- trauma class	χ^2 (df)	p value
Gender						
Female	500 (50.0%)	278 (54.1%)	188 (48.1%)	34 (35.8%)	11.68 (2)	0.003
Male	500 (50.0%)	236 (45.9%)	203 (51.9%)	61 (64.2%)		
Age in years						
18-29	260 (26.0%)	143 (27.8%)	98 (25.1%)	19 (20.0%)	10.90 (4)	0.028
30-59	529 (52.9%)	250 (48.6%)	217 (55.5%)	62 (65.3%)		
60-100	211 (21.1%)	121 (23.5%)	76 (19.4%)	14 (14.7%)		
Education						
No school	585 (58.5%)	312 (60.7%)	222 (56.8%)	51 (53.7%)	4.31 (4)	0.366
Primary School	324 (32.4%)	163 (31.7%)	129 (33.0%)	32 (33.7%)		
Secondary School	91 (9.1%)	39 (7.6%)	40 (10.2%)	12 (12.6%)		
Household income (per month)						
Less than 24.35 \$	759 (75.9%)	399 (77.6%)	288 (73.7%)	72 (75.9%)	5.30 (4)	0.258
24.35-48.69 \$	162 (16.2%)	84 (16.3%)	64 (16.4%)	14 (14.7%)		
More than 48.70 \$	79 (7.9%)	31 (6.0%)	39 (10.0%)	9 (9.5%)		
Distance household to village center						
Less than 2 km	617 (61.7%)	322 (62.6%)	234 (59.8%)	61 (64.2%)	13.32 (4)	0.010
2-5 km	236 (23.6%)	132 (25.7%)	91 (23.3%)	13 (13.7%)		
More than 6 km	147 (14.7%)	60 (11.7%)	66 (16.9%)	21 (22.1%)		
Psychological Outcomes						
PTSD	170 (17.0%)	53 (10.3%)	89 (22.8%)	28 (29.5%)	35.97 (2)	0.000
Depression	278 (27.8%)	128 (24.9%)	120 (30.7%)	30 (31.6%)	4.45 (2)	0.108
Anxiety	254 (25.4%)	110 (21.4%)	107 (27.4%)	37 (39.9%)	14.38 (2)	0.001
Suicidality	151 (15.1%)	67 (13.0%)	53 (13.6%)	31 (32.6%)	25.21 (2)	0.000

Table 2
Lifetime prevalence of various traumatic experiences.

Traumatic Experiences	Total (N = 1000)		Women (N = 500)		Men (N = 500)		Sex Difference χ^2	18–29 yrs (N = 260)		30–59 yrs (N = 529)		60–100 yrs (N = 211)		Age Difference χ^2
	n	%	n	%	n	%		n	%	n	%	n	%	
Experience of any Event	696	69.6	341	68.2	348	71.0	0.93	174	66.9	389	73.5	133	63.0	9.05**
Witness of any traumatic event	678	67.8	289	57.8	389	77.8	45.80***	166	63.8	377	71.3	135	64.0	6.18*
Displacement	607	60.7	296	59.2	311	62.2	0.94	155	59.6	338	63.9	114	54.0	6.33*
Burning or looting of home or property	452	45.2	222	44.4	230	46.0	0.26	110	42.3	258	48.8	84	39.8	6.08*
Murder of relative	173	17.3	79	15.8	94	18.8	1.57	46	17.7	100	18.9	27	12.8	3.97
Serious injury by armed attack	58	5.8	22	4.4	36	7.2	3.59	10	3.8	37	7.0	11	5.2	3.33
Abduction	44	4.4	8	1.6	36	7.2	18.64***	10	3.8	29	5.5	5	2.4	3.73
Experienced Rape	13	1.3	13	2.6	0.0	0.0	13.17***	5	2.0	7	1.3	1	0.5	1.91

Note. * $p < .05$. ** $p < .01$. *** $p < .001$.

measuring the exposure to traumatic events. We identified the most appropriate number of classes by considering commonly used information criteria as well as the theoretical adequacy and utility of the class solution. Information criteria used for model selection were the Bayesian Information Criterion (BIC), Sample-size Adjusted Bayesian Information Criterion (SABIC), Akaike Information Criterion (AIC), and the Consistent Akaike Information Criterion (CAIC). Running the analysis with a set of random start values lead to identical solutions, suggesting that the model has converged on the global maximum of the likelihood function. Individuals were categorized into the classes based on their largest posterior probability estimate across classes. This *maximum-probability assignment* does not take uncertainty in class assignment into account, as it treats the most likely class as the true class to which an individual belongs (Bray et al., 2015). The potentially occurring classification error can lead to bias in the estimates of subsequent analyses, e.g. regression models (Bray et al., 2015). In studies with large sample size, i.e., $N > 500$, and well distinguished classes, i.e., mean posterior probability of class membership for each class exceeding 0.7–0.9, regression models with latent classes are little affected by classification error (Elliott et al., 2020; Nagin, 2005). In the current study, the average posterior probabilities for individuals assigned to each class ranged from 0.76 to 0.95, indicating the majority of orders were likely to be in a given class and not warranting to control for classification uncertainty (Nagin, 2005). Multiple logistic regression models were conducted for each psychopathological outcome, using trauma exposure class as primary prediction variable, and sociodemographic information as control variables. Adjusted ORs were reported to describe the likelihood of psychopathology depending on class membership and control variables. Necessary assumptions for logistic regression analysis, including linearity of logits, absence of multicollinearity and lack of influential outliers, were met for all models.

3. Results

3.1. Prevalence of conflict-related traumatic events

In total 69.6% ($n = 696$) of respondents directly experienced at least one traumatic event involving armed groups since 2002. Furthermore, 67.8% ($n = 678$) of respondents witnessed at least one traumatic event type happening to other people. Table 2 shows the prevalence for the different traumatic event types by gender and age group. Women reported being a victim of rape more often, while the frequency of abduction and witnessing of events was higher for men. Age group differences were found for burning or looting of home or property, displacement, and witnessing of events. The exposure to traumatic event types was most prevalent in the age group of 30–59 years old.

3.2. Prevalence of psychopathology and the association with specific traumatic events

In total, 17.0% ($n = 170$) of the sample displayed post-traumatic stress symptoms at a level indicating potentially clinically relevant

symptoms of PTSD. Results further showed that 27.8% ($n = 278$) and 25.4% ($n = 254$) of participants screened positive for depression and anxiety, respectively. Furthermore, 15.1% ($n = 151$) of participants reported to feel suicidal. The conditional prevalence and unadjusted ORs for psychopathology across the individual trauma types are documented in the supplementary files (see eTable 1). The unadjusted ORs indicated that six of the seven traumatic event types, including the witnessing of events, were associated with an increased risk of psychopathology. Only the experience of abduction did not show an increased risk of mental health problems in the sample. Being a victim of rape by armed personnel showed the strongest association with each PTSD, depression, and anxiety and suicidality.

3.3. Classifying trauma exposure

Latent class models with two to six classes were analyzed to classify participants by their exposure to traumatic events (eTable 2). The three-class solution was selected for further analyses based on the information criteria (eFig. 1) and theoretical considerations. The estimated and observed conditional probabilities for each traumatic event are listed in eTable 3. The class profiles based on the observed conditional probabilities for the three-class solution are presented in Fig. 2. Accordingly, class 1 (51.4%, $n = 514$) included individuals with the lowest probabilities of exposure to traumatic events (0%–46%). This class was labeled *low-trauma-exposure class*. Individuals in class 2 (39.1%, $n = 391$) were primarily characterized by high probabilities of exposure to non-physical violence types, i.e., exposure to burning/looting of home or property (93%), displacement (89%), murder of close persons (31%), and witnessing traumatic events (89%). The probability of exposure to interpersonal traumatic event types was, similar to class 1, almost zero. Class 2 was therefore labeled *non-physical-trauma class*. In class 3 (9.5%, $n = 95$) probabilities were high for non-physical trauma types and additionally the highest for interpersonal trauma types, i.e., being injured (60%), abducted (46%), and raped (14%) by armed groups. This class was thus labeled *interpersonal-trauma class*.

Table 1 presents socio-demographic information and prevalence of mental health disorders for the three classes. The *interpersonal-trauma class* had a significantly higher proportion of men, consisted to a greater extent of middle-aged (30–59 years) respondents and lived further away from the village center. Furthermore, the prevalence of clinically significant symptoms of PTSD, anxiety and suicidality increased from *low-trauma-exposure* to *non-physical-trauma* to *interpersonal-trauma class*. No significant differences were found for depression.

3.4. Predicting psychopathology with trauma-exposure class

To examine the relation between class membership and psychopathology, four separate logistic regressions were conducted with the prospective diagnosis of PTSD, depression, anxiety and suicidality as outcome variables. Each regression model included participants' age, education, household income and distance to village center as control




Fig. 2. Observed probabilities of conflict-related trauma exposure by class membership in a three-class solution.

variables. Our results showed a constant increase in the risks for psychopathology from *low-trauma-exposure* to *non-physical-trauma* to *interpersonal-trauma* class (see Table 3). Using the *low-trauma-exposure* class as reference category, membership in the *non-physical-trauma* class increased risks for PTSD, depression and anxiety. Similarly, individuals in the *interpersonal-trauma* class showed significantly increased risks for PTSD, depression, anxiety, but also suicidality compared to the *low-trauma-exposure* class. Furthermore, comparing individuals from the *non-physical-trauma* and the *interpersonal-trauma* class revealed higher risks for PTSD, anxiety, and suicidality in the latter class. Being female significantly increased the risks for all mental health problems. Higher education and household income significantly decreased the risk for suicidality.

4. Discussion

This study investigated the prevalence of traumatic experiences and mental health problems in a representative sample of the general population from conflict-affected South Kivu, providing recent data from a continuously unstable post-conflict setting. Consistent with previous population-based epidemiological studies from post-conflict settings, our findings showed high prevalence of traumatic experiences in a war-affected population (De Jong et al., 2001; Pham et al., 2004). The tremendous trauma exposure represents a high psychological burden for the population, which reflects in elevated screening rates of mental health problems (De Jong et al., 2003; Johnson et al., 2008; Lopes Carozo et al., 2000).

Acknowledging the complexity of trauma exposure in conflict-affected populations, we implemented a person-oriented analysis

approach to adequately disentangle the relation between trauma exposure and mental health outcomes. This resulted in the identification of classes of individuals with shared trauma histories and the determination of classes most vulnerable to mental health problems. We found three distinct classes of individuals: those who experienced low trauma exposure (*low-trauma-exposure* class), those affected by non-physical trauma types only (*non-physical-trauma* class) and those subjected to non-physical and additionally interpersonal trauma types (*interpersonal-trauma* class). The markers distinguishing between classes were the number of experienced trauma types, but also the type of trauma exposure itself, i.e., non-physical trauma and interpersonal trauma. Especially the *interpersonal-trauma* class featured an alarmingly high likelihood of being exposed to multiple traumata, including non-physical events, and as the only class the occurrence of interpersonal trauma types. Our findings are concurrent with person-centered research, including studies with war-affected samples, which found similar distinct trauma patterns based on the presence of interpersonal traumata (Gibbs et al., 2020; O'Donnell et al., 2017; Sengoelge et al., 2019). However, these studies have either examined samples not living in war-affected areas anymore (e.g., refugees (Sengoelge et al., 2019)) or only subgroups of the population (e.g., women (Gibbs et al., 2020)). Compared to studies with refugees, our identification of trauma patterns is specifically based on the experiences of events within a war-affected region, i.e., our identified patterns are free of an influence of (post-) migration associated traumatic experiences and thus more specific to populations living in war-affected areas. Moreover, our representative sample including both women and men, extends existing evidence of studies conducted within war-affected regions for further population subgroups and allows for the examination of gender effects in trauma patterns. Regarding the composition of the

Table 3
Adjusted Odds Ratios of Variables associated with PTSD, Depression, Anxiety and Suicidality.

	PTSD OR (95% CI)	Depression OR (95% CI)	Anxiety OR (95% CI)	Suicidality OR (95% CI)
Trauma Class				
non-physical	3.10*** (2.10–4.57)	1.50* (1.10–2.05)	1.54** (1.12–2.12)	1.18 (0.79–1.76)
vs. low- exposure				
interpersonal	5.80*** (3.25–10.32)	1.81* (1.09–3.00)	3.24*** (1.97–5.33)	4.43*** (2.57–7.66)
vs. low- exposure				
interpersonal	1.87* (1.09–3.22)	1.20 (0.72–2.00)	2.11** (1.29–3.46)	3.76*** (2.15–6.57)
vs. non- physical				
Gender (female)	3.94*** (2.55–6.10)	3.14*** (2.24–4.42)	3.26*** (2.28–4.65)	2.00** (1.31–3.07)
Age	1.00 (0.99–1.01)	1.00 (0.99–1.01)	1.00 (0.99–1.01)	1.00 (0.99–1.01)
Education				
None	Reference category			
Primary School	0.60* (0.38–0.94)	0.80 (0.56–1.14)	1.00 (0.70–1.45)	0.42** (0.26–0.70)
Secondary	0.49 (0.20–1.17)	0.60 (0.31–1.15)	0.67 (0.34–1.33)	0.34* (0.13–0.91)
School				
Household income (per month)				
Less than 24.35 \$	Reference category			
24.35–48.69 \$	1.43 (0.87–2.35)	0.80 (0.52–1.22)	0.90 (0.58–1.39)	0.54* (0.29–0.98)
More than 48.70 \$	1.13 (0.53–2.41)	1.01 (0.55–1.86)	1.12 (0.61–2.06)	0.09* (0.01–0.69)
Distance household to village center				
Less than 2 km	Reference category			
2–5 km	1.03 (0.68–1.60)	0.57* (0.40–0.82)	0.93 (0.65–1.33)	0.64 (0.40–1.01)
More than 6 km	0.65 (0.38–1.11)	0.59* (0.38–0.92)	0.77 (0.49–1.19)	0.70 (0.41–1.20)

Note. *p < .05. **p < .01. ***p < .001.

trauma classes, it is noteworthy that in the current study only the *interpersonal-trauma class* reported exposure to sexual violence and abduction. In previous studies with war-affected samples, the trauma classes mostly differed in their probability of exposure to traumatic events but not the presence of certain event types (Gibbs et al., 2020; Sengoelge et al., 2019). In the current study, the observation of a clear distinction of trauma patterns based on the presence of certain trauma types (i.e., interpersonal trauma) offers a more nuanced picture. Our findings indicate that sexual violence and abduction are rarely single events, but rather accompanied or preceded by other severe types of interpersonal trauma. Furthermore, the composition of the classes showed that men, middle-aged and those who stay far from village centers may be predisposed to higher exposure to violence by armed groups including interpersonal trauma.

Focusing on the psychopathological consequences of the trauma exposure patterns, we observed significant differences in the probabilities of developing mental health problems. The risk of mental health problems increased as the exposure to trauma types increased, rendering those exposed to both non-personal and interpersonal trauma types as most vulnerable to psychopathology. This replicates findings of an association between the number of trauma types and an increased symptomatology of mental health problems (i.e. *dose-effect*), but also suggests that the level of psychopathology varies with the type of exposure (Farhood and Dimassi, 2012; Lopes Cardozo et al., 2000). More specifically, our findings revealed that trauma histories involving the exposure to interpersonal trauma types, such as rape and injury by armed attack, are more detrimental for mental health outcomes than trauma histories encompassing non-physical trauma types only. These findings highlight the relevance of taking into account not only the number of trauma types, but also common trauma histories in which trauma types co-occur to build nuanced models of vulnerability to psychopathological outcomes.

Furthermore, we found gender differences in the vulnerability to mental health problems. Despite men being more likely to be grouped in the *interpersonal-trauma class*, women were more likely to develop symptoms of PTSD, depression, anxiety or suicidality. This inverse relation between gender and trauma exposure and gender and psychopathology mirrors the conclusion of studies describing lower risks to experience traumatic events, but a higher PTSD symptom severity for women (Sengoelge et al., 2019; Tolin and Foa, 2006). A higher vulnerability to mental health problems has also been observed among female refugees from eastern DRC (Ainamani et al., 2020). In a systematic analysis, such gender differences were attributed to differences in pre-, peri-, and posttraumatic risk factors, including type of traumatic experience, peritraumatic fear and helplessness, and negative posttraumatic cognitions (Christiansen and Hansen, 2015). Regarding the increased exposure of women to sexual violence and the associated powerful negative emotions and cognitions, these risk factors may be a plausible explanation for the gender differences found in the present study. Context-specific factors, such as high levels of stigmatization and daily stressors, e.g., inadequate medical care, may further contribute to vulnerability to mental health problems among female victims of sexual violence from Eastern DRC (Verelst et al., 2014a, 2014b).

4.1. Strengths and limitations

The implemented person-oriented approach in a population-based sample complements existing post-conflict research with several benefits and contributes to a more comprehensive picture of the lived reality in post-conflict settings (O'Donnell et al., 2017). Firstly, it offers more information about the heterogeneity of trauma exposure in conflict-affected populations than the analysis of cumulative trauma exposure. Secondly, it describes shared patterns of trauma experience among the population caused by the presence of armed groups, instead of focusing on the exposure to single traumatic events. And thirdly, it offers more differentiated explanations for differences in the prevalence of psychopathology beyond the study of cumulative and isolated traumatic events. Lastly, analytical inclusion of socio-demographic characteristics allows for the determination of population strata-specific vulnerabilities for trauma exposure and mental health problems. However, some important limitations must be noted: The data for this study were collected in six of eight territories of South Kivu. Insecurity at the time of data collection limited the access to ten percent of the sampled villages, which were replaced using the initial sampling procedure. It is likely, that residents of these insecure areas have experienced high levels of violence, which may not be reflected by the observations in the replacement villages. The instruments used to measure mental health problems and trauma exposure, although validated in non-western samples and for use in studies like this (Kaltenbach et al., 2017; Kocalevent et al., 2014; Löwe et al., 2010; Neuner et al., 2004), have been developed for screening purpose and do not substitute for diagnoses determined by clinicians. The use of mental health screening instruments as opposed to structured diagnostic interviews may lead to an overestimation of prevalence rates in the current study (Levis et al., 2020; Tsai, 2014). Furthermore, the validity of the assessment of suicidal ideation as well as the experience of rape may be limited by the use of single-item measures. Regarding the experience of rape, previous studies have reported higher prevalence rates in samples from the DRC using more detailed measures (Johnson et al., 2010). Utilizing standardized instruments and following detailed measurement guidelines for the assessment of suicidality and sexual violence can reduce potential bias and the risk of misclassifications in future studies (World Health Organization, 2019). As data for this study was assessed in self-report, it may generally not be free of potential bias, e.g., recall-bias or social-desirability. Nevertheless, pre-tests indicated participants' comprehension and disclosure to all questions, including the single items on rape and suicidal ideation, in the current cultural context. In addition, although we provided up-to-date information on the prevalence of traumatic events in the general population of South Kivu, our

analyses did not include monitoring for events before 2002. Despite dating back many years, these experiences may have influenced the predisposition to mental disorders in the current study. Furthermore, we only recorded a range of possible traumatic events, which does not make our assessment exhaustive and does not preclude exposure to other traumatic events. Moreover, other aspects of trauma exposure, such as severity, frequency, and duration, may provide incremental value in understanding patterns of trauma exposure, but were not considered in this study. However, the prevalence of traumatic events and the effects on mental health in the current study were largely consistent with previous reports (Johnson et al., 2010; Pham et al., 2010). Moreover, use of a multi-stage randomized cluster sampling procedure and large sample size have been found to minimize biases in mental health problems prevalence rates (Steel et al., 2009).

4.2. Implications

The high trauma burden in the population poses an enormous challenge to the porous health systems of war-affected countries (De Jong and Komproe, 2002). Our approach to analyze trauma histories highlights the diverse mental health needs of populations in post-conflict settings, thus placing the implementation of population needs-based approaches into the focus of research and policies. To adequately address the complex needs and overcome shortages in mental health-care workforce, this may include decentralized and community-oriented interventions, such as the training of community-based lay personnel (De Jong and Komproe, 2002; Ertl et al., 2011). Provision of special treatment services for those most at risk for psychopathology, such as treatment centers for violence-affected women, may further facilitate treatment access for vulnerable groups. Moreover, given the multifaceted impact of different trauma histories on psychopathology, increased attention to traumatic experiences is indicated for the clinical practice. Therefore, health-care providers in post-conflict regions should implement extensive screening procedures with a special emphasis on interpersonal traumatic events. The treatment of psychopathological symptoms, not only PTSD, in post-conflict settings can benefit from the inclusion of trauma-focused interventions (Robjant and Fazel, 2010).

5. Conclusions

This is one of the first studies to implement a person-centered approach for the analysis of trauma exposure in a representative sample living in a conflict-affected region. Our findings revealed distinct patterns of exposure to traumatic events in the population of eastern DRC, primarily differentiated by the number of trauma types and the presence of interpersonal trauma. Thus, this is the first study showing that populations in (post-) conflict regions are comprised of different subgroups regarding their trauma exposure. Analyses of the relation between trauma patterns and mental health provided nuanced insights into vulnerability to psychopathology, beyond the analysis of cumulative trauma exposure and isolated trauma types. The study has important clinical implications by identifying a gradual increase in the risk of suffering from mental disorders depending on the trauma history, with trauma histories that include both non-physical and interpersonal trauma as most detrimental for mental health outcomes.

Funding/support

This study was funded by the German Research Foundation (Deutsche Forschungsgemeinschaft, DFG; project number: 282722353).

Role of the funder/sponsor

The funding organization had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; and decision to

submit the manuscript for publication.

Additional contributions

We are grateful to all individuals who supported our work with their eagerness to participate and willingness to reveal intimate information.

CRediT authorship contribution statement

Lars Dumke: Conceptualization, Methodology, Software, Formal analysis, Writing – original draft, Writing – review & editing, Visualization. **Roos van der Haer:** Conceptualization, Validation, Writing – review & editing. **Carlo Koos:** Validation, Investigation, Resources, Writing – review & editing, Funding acquisition, Project administration. **Tobias Hecker:** Conceptualization, Methodology, Formal analysis, Resources, Writing – original draft, Writing – review & editing, Supervision.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.ssmmh.2021.100005>.

References

- Ainamani, H. E., Elbert, T., Olema, D. K., & Hecker, T. (2020). Gender differences in response to war-related trauma and posttraumatic stress disorder - a study among the Congolese refugees in Uganda. *BMC Psychiatry*, 20(1), 17. <https://doi.org/10.1186/s12888-019-2420-0>
- Autesserre, S. (2017). *Resolving the political crisis in the democratic republic of the Congo*. Presented at the.
- Bray, B. C., Lanza, S. T., & Tan, X. (2015). Eliminating bias in classify-analyze approaches for latent class Analysis. *Structural Equation Modeling*, 22(1), 1–11. <https://doi.org/10.1080/10705511.2014.935265>
- Charlson, F. J., Steel, Z., de genhardt, L., et al. (2012). Predicting the impact of the 2011 conflict in Libya on population mental health: PTSD and depression prevalence and mental health service requirements. *PLoS One*, 7(7). <https://doi.org/10.1371/journal.pone.0040593>
- Christiansen, D. M., & Hansen, M. (2015). Accounting for sex differences in PTSD: A multi-variable mediation model. *European Journal of Psychotraumatology*, 6, 1–10. <https://doi.org/10.3402/ejpt.v6.26068>
- De Jong, J. T. V. M., & Komproe, I. H. (2002). Closing the gap between psychiatric epidemiology and mental health in post-conflict situations. *Lancet*, 359(9320), 1793–1794. [https://doi.org/10.1016/S0140-6736\(02\)08729-9](https://doi.org/10.1016/S0140-6736(02)08729-9)
- De Jong, J. T. V. M., Komproe, I. H., & Van Ommeren, M. (2003). Common mental disorders in postconflict settings. *Lancet*, 361(9375), 2128–2130. [https://doi.org/10.1016/S0140-6736\(03\)13692-6](https://doi.org/10.1016/S0140-6736(03)13692-6)
- De Jong, J. T. V. M., Komproe, I. H., Van Ommeren, M., et al. (2001). Lifetime events and posttraumatic stress disorder in 4 postconflict settings. *Journal of the American Medical Association*, 286(5), 555–562. <https://doi.org/10.1001/jama.286.5.555>
- De Juan, A., & Koos, C. (2019). The historical roots of cooperative behavior—evidence from eastern Congo. *World Development*, 116, 100–112. <https://doi.org/10.1016/j.worlddev.2018.11.011>
- Elliott, M. R., Zhao, Z., Mukherjee, B., Kanaya, A., & Needham, B. L. (2020). Methods to account for uncertainty in latent class Assignments when using latent classes as predictors in regression models, with application to acculturation Strategy measures. *Epidemiology*, 31(2), 194–204. <https://doi.org/10.1097/EDE.0000000000001139>
- Ertl, V., Pfeiffer, A., Schauer, E., Elbert, T., & Neuner, F. (2011). Community-implemented trauma therapy for former child soldiers in northern Uganda: A randomized controlled trial. *The Journal of the American Medical Association*, 306(5), 503–512. <https://doi.org/10.1001/jama.2011.1060>
- Farhood, L., & Dimassi, H. (2012). Prevalence and predictors for post-traumatic stress disorder, depression and general health in a population from six villages in South Lebanon. *Social Psychiatry and Psychiatric Epidemiology*, 47(4), 639–649. <https://doi.org/10.1007/s00127-011-0368-6>
- Farhood, L., Dimassi, H., & Lehtinen, T. (2006). Exposure to war-related traumatic events, prevalence of PTSD, and general psychiatric morbidity in a civilian population from southern Lebanon. *Journal of Transcultural Nursing*, 17(4), 333–340. <https://doi.org/10.1177/1043659606291549>
- Fellmeth, G., Plugge, E., Fazel, M., et al. (2018). Validation of the Refugee Health Screener-15 for the assessment of perinatal depression among Karen and Burmese

- women on the Thai-Myanmar border. A. Bazzano (Ed.). *PLoS One*, 13(5), Article e0197403. <https://doi.org/10.1371/journal.pone.0197403>
- Gibbs, A., Abdelatif, N., Said, N., & Jewkes, R. (2020). Associations between exposures to occupation-related events, depression and intimate partner violence among women in the occupied Palestinian Territories. *Global Public Health*. <https://doi.org/10.1080/17441692.2020.1849349>. Published online.
- Hagan, M. J., Sulik, M. J., & Lieberman, A. F. (2016). Traumatic life events and psychopathology in a high risk, ethnically diverse sample of young children: A person-centered approach. *Journal of Abnormal Child Psychology*, 44(5), 833–844. <https://doi.org/10.1007/s10802-015-0078-8>
- Hollifield, M., Verbillis-Kolp, S., Farmer, B., et al. (2013). The refugee health screener-15 (RHS-15): Development and validation of an instrument for anxiety, depression, and PTSD in refugees. *General Hospital Psychiatry*, 35(2), 202–209. <https://doi.org/10.1016/j.genhosppsych.2012.12.002>
- Ibrahim, H., & Hassan, C. Q. (2017). Post-traumatic stress disorder symptoms resulting from torture and other traumatic events among Syrian Kurdish refugees in Kurdistan region, Iraq. *Frontiers in Psychology*, 8(FEB), 241. <https://doi.org/10.3389/fpsyg.2017.00241>
- JMP®. Version 15.
- Johnson, K., Asher, J., Rosborough, S., et al. (2008). Association of combatant status and sexual violence with health and mental health outcomes in postconflict Liberia. *The Journal of the American Medical Association*, 300(6), 676–690. <https://doi.org/10.1001/jama.300.6.676>
- Johnson, K., Scott, J., Rughita, B., et al. (2010). Association of sexual violence and human rights violations with physical and mental health in territories of the Eastern Democratic Republic of the Congo. *The Journal of the American Medical Association*, 304(5), 553–562. <https://doi.org/10.1001/jama.2010.1086>
- Kaltenbach, E., Härdtner, E., Hermentau, K., Schauer, M., & Elbert, T. (2017). Efficient identification of mental health problems in refugees in Germany: The refugee health screener. *European Journal of Psychotraumatology*, 8(sup2), 1389205. <https://doi.org/10.1080/20008198.2017.1389205>
- Kocalevent, R. D., Finck, C., Jimenez-Leal, W., Sautier, L., & Hinz, A. (2014). Standardization of the Colombian version of the PHQ-4 in the general population. *BMC Psychiatry*, 14(1). <https://doi.org/10.1186/1471-244X-14-205>
- Kroenke, K., Spitzer, R. L., Williams, J. B. W., & Lowe, B. (2009). An ultra-brief screening scale for anxiety and depression: The PHQ-4. *Psychosomatics*, 50(6), 613–621. <https://doi.org/10.1176/appi.psy.50.6.613>
- Kuringe, E., Materu, J., Nyato, D., et al. (2019). Prevalence and correlates of depression and anxiety symptoms among out-of-school adolescent girls and young women in Tanzania: A cross-sectional study. *PLoS One*, 14(8). <https://doi.org/10.1371/journal.pone.0221053>
- Levis, B., Benedetti, A., Ioannidis, J. P. A., et al. (2020). Patient health questionnaire-9 scores do not accurately estimate depression prevalence: Individual participant data meta-analysis. *Journal of Clinical Epidemiology*, 122, 115–128. <https://doi.org/10.1016/j.jclinepi.2020.02.002>. e1.
- Lopes Cardozo, B., Vergara, A., Agani, F., & Gotway, C. A. (2000). Mental health, social functioning, and attitudes of Kosovar Albanians following the war in Kosovo. *Journal of the American Medical Association*, 284(5), 569–577. <https://doi.org/10.1001/jama.284.5.569>
- Löwe, B., Wahl, I., Rose, M., et al. (2010). A 4-item measure of depression and anxiety: Validation and standardization of the patient health questionnaire-4 (PHQ-4) in the general population. *Journal of Affective Disorders*, 122(1–2), 86–95. <https://doi.org/10.1016/j.jad.2009.06.019>
- Nagin, D. (2005). *Group-based modeling of development*. Harvard University Press. <https://doi.org/10.4159/9780674041318>
- Netland, M. (2005). Event-list construction and treatment of exposure data in research on political violence. *Journal of Traumatic Stress*, 18(5), 507–517. <https://doi.org/10.1002/jts.20059>
- Neuner, F., Schauer, M., Karunakara, U., Klaschik, C., Robert, C., & Elbert, T. (2004). Psychological trauma and evidence for enhanced vulnerability for posttraumatic stress disorder through previous trauma among West Nile refugees. *BMC Psychiatry*, 4, 34. <https://doi.org/10.1186/1471-244X-4-34>
- O'Donnell, M. L., Schaefer, I., Varker, T., et al. (2017). A systematic review of person-centered approaches to investigating patterns of trauma exposure. *Clinical Psychology Review*, 57, 208–225. <https://doi.org/10.1016/j.cpr.2017.08.009>
- Pham, P. N., Vinck, P., Kinkodi, D. K., & Weinstein, H. M. (2010). Sense of coherence and its association with exposure to traumatic events, posttraumatic stress disorder, and depression in eastern Democratic Republic of Congo. *Journal of Traumatic Stress*, 23(3). <https://doi.org/10.1002/jts.20527>. n/a-n/a.
- Pham, P. N., Weinstein, H. M., & Longman, T. (2004). Trauma and PTSD symptoms in Rwanda: Implications for attitudes toward justice and reconciliation. *Journal of the American Medical Association*, 292(5), 602–612. <https://doi.org/10.1001/jama.292.5.602>
- Robjant, K., & Fazel, M. (2010). The emerging evidence for narrative exposure therapy: A review. *Clinical Psychology Review*, 30(8), 1030–1039. <https://doi.org/10.1016/j.cpr.2010.07.004>
- Sengoelge, M., Johnson-Singh, C. M., Mittendorfer-Rutz, E., Vaez, M., & Saboonchi, F. (2019). Identifying subgroups of refugees from Syria resettled in Sweden based on multiple trauma exposures: A latent class analysis of trauma history and mental health outcomes. *Journal of Psychosomatic Research*, 125, 109814. <https://doi.org/10.1016/j.jpsychores.2019.109814>
- Stammel, N., Heeke, C., Bockers, E., et al. (2013). Prolonged grief disorder three decades post loss in survivors of the Khmer Rouge regime in Cambodia. *Journal of Affective Disorders*, 144(1–2), 87–93. <https://doi.org/10.1016/j.jad.2012.05.063>
- Steel, Z., Chey, T., Silove, D., Marnane, C., Bryant, R. A., & van Ommeren, M. (2009). Association of torture and other potentially traumatic events with mental health outcomes among populations exposed to mass conflict and displacement. *Journal of the American Medical Association*, 302(5), 537. <https://doi.org/10.1001/jama.2009.1132>
- Tolin, D. F., & Foa, E. B. (2006). Sex differences in trauma and posttraumatic stress disorder: A quantitative review of 25 years of research. *Psychological Bulletin*, 132(6), 959–992. <https://doi.org/10.1037/0033-2909.132.6.959>
- Tsai, A. C. (2014). Reliability and validity of depression assessment among persons with HIV in sub-Saharan Africa: Systematic review and meta-analysis. *Journal of Acquired Immune Deficiency Syndromes*, 66(5), 503–511. <https://doi.org/10.1097/QAI.0000000000000210>
- Verelst, A., De Schryver, M., Broekaert, E., & Derluyn, I. (2014b). Mental health of victims of sexual violence in eastern Congo: Associations with daily stressors, stigma, and labeling. *BMC Women's Health*, 14(1), 1–12. <https://doi.org/10.1186/1472-6874-14-106>
- Verelst, A., De Schryver, M., De Haene, L., Broekaert, E., & Derluyn, I. (2014a). The mediating role of stigmatization in the mental health of adolescent victims of sexual violence in Eastern Congo. *Child Abuse & Neglect*, 38(7), 1139–1146. <https://doi.org/10.1016/j.chiabu.2014.04.003>
- World Health Organization (WHO), United Nations Population Fund (UNFPA), United Nations High Commissioner for Refugees (UNHCR). (2019). *Clinical management of rape and intimate partner violence survivors: Developing protocols for use in humanitarian settings*.