



Universiteit  
Leiden  
The Netherlands

## Transdiagnostic treatment for eating disorders

Jong, M. de

### Citation

Jong, M. de. (2021, September 1). *Transdiagnostic treatment for eating disorders*. Retrieved from <https://hdl.handle.net/1887/3209226>

Version: Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/3209226>

**Note:** To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <https://hdl.handle.net/1887/3209226> holds various files of this Leiden University dissertation.

**Author:** Jong, M. de

**Title:** Transdiagnostic treatment for eating disorders

**Issue Date:** 2021-09-01

# Chapter 4

## **Enhanced cognitive behavioral therapy for patients with eating disorders: a systematic review**

Martie de Jong<sup>1,2</sup>, Maartje Schoorl<sup>3,4</sup>, and Hans W. Hoek<sup>1,5,6</sup>

1. Parnassia Psychiatric Institute, The Hague, The Netherlands
2. Eating Disorders Center, PsyQ Haaglanden, The Hague, The Netherlands
3. Psychotrauma Center, PsyQ Haaglanden, The Hague, The Netherlands
4. Department of Clinical Psychology, Leiden University, Leiden, The Netherlands
5. Department of Psychiatry, University Medical Center Groningen, University of Groningen, Groningen, The Netherlands
6. Department of Epidemiology, Mailman School of Public Health, Columbia University, New York, New York, United States

*Published in: Current Opinion in Psychiatry 2018;31(6):436-444.*  
*DOI: 10.1097/YCO.0000000000000452*

## Abstract

**Purpose of review:** The aim of this study was to provide an update of the most recent (since January 2014) enhanced cognitive behavioral therapy (CBT-E) effectiveness studies (randomized controlled trials and open trials) on bulimia nervosa, binge eating disorder, and transdiagnostic samples.

**Recent findings:** Of 451 screened studies, seven effectiveness studies (five randomized and two open trials) were included in this review: of these, three had a bulimia nervosa sample and four a transdiagnostic sample (all conducted in an outpatient setting). Substantial differences in posttreatment remission rates were found (range: 22.2% to 67.6%) due, in part, to differences in samples and operationalization of clinical significant change.

**Summary:** There is robust evidence that CBT-E is an effective treatment for patients with an eating disorder. However, more studies on differential effects and working mechanisms are required to establish the specificity of CBT-E.

## Keywords

cognitive behavioral therapy, eating disorders, effectiveness, transdiagnostic, treatment

## Introduction

Eating disorders are severe mental disorders, which often begin in adolescence (Mitchison et al., 2012), frequently have a chronic course (Steinhausen, 2009) and can have considerable impact on quality of life (Jenkins et al., 2011). Eating disorders make a substantial contribution to the global burden of disease, especially among young women (Erskine, Whiteford, & Pike, 2016). Although anorexia nervosa is a relatively rare disorder in many non-western countries, bulimia nervosa and binge eating disorder (BED) are common disorders worldwide (Hoek, 2016). Previous reviews showed that, among young women in Europe, Asia, Africa and Latin America, bulimia nervosa is reported by 1-2% and BED by 1-4% (Keski-Rahkonen & Mustelin, 2016; Kolar et al., 2016; Perez, Ohrt, & Hoek, 2016; Thomas, Lee, & Becker, 2016; van Hoeken, Burns, & Hoek, 2016). Recent studies show that eating disorders (especially bulimia nervosa and BED) are also common among older persons; according to the DSM-5 criteria, the prevalence of all eating disorders combined is around 3.5% in older (aged >40 years) women and around 1-2% in older men (Mangweth-Matzek & Hoek, 2017). Despite that increasing numbers of individuals with eating disorders are receiving treatment, European samples show that only about one-third are detected via health care (Keski-Rahkonen & Mustelin, 2016).

In terms of the DSM-IV, the most common eating disorder diagnosis in both clinical and community samples was 'Eating disorder not otherwise specified' (EDNOS). With the introduction of the DSM-5 and concurrent changes in the eating disorder section (including the introduction of BED as an official category, and lowering the threshold for anorexia nervosa and bulimia nervosa) the percentage of 'Other specified feeding or eating disorder' (OSFED; DSM-IV EDNOS) was significantly reduced, even though this diagnosis might still be the most common one in this population (Keel et al., 2011; Machado, Goncalves, & Hoek, 2013; Smink, van Hoeken, & Hoek, 2013).

According to a recent international comparison between nine evidence-based clinical guidelines for eating disorders, cognitive behavioral therapy (CBT) is widely used as the preferred treatment for bulimia nervosa and BED (Hilbert, Hoek, & Schmidt, 2017). The major guidelines for the treatment of eating disorders (Hay et al., 2014; National Institute for Health and Care Excellence (NICE), 2017; Yager et al., 2014) recommend CBT as the psychological treatment of first choice for bulimia nervosa and BED. CBTE(nhanced) is a specific form of CBT and is designed to be suitable for the full range of

eating disorder diagnoses (Fairburn, 2008). It is based on the transdiagnostic theory of the maintenance of eating disorders, in which it is assumed that most of the mechanisms involved in the persistence of eating disorders are common to all eating disorders, rather than being specific to each diagnostic group separately. It asserts that central to all eating disorders is a dysfunctional evaluation of self-worth that is overly based on shape and weight (Fairburn, Cooper, & Shafran, 2003). CBT-E uses strategies and procedures to address this overevaluation of shape and weight by focusing on targeting these mechanisms (known as the 'focused' version of CBT-E). The treatment protocol can be extended with interventions that target additional maintaining mechanisms, that is core low self-esteem, clinical perfectionism, and interpersonal problems (known as the 'broad' version of CBT-E). For the OSFED diagnoses, CBT-E has an advantage over other CBT protocols because of its transdiagnostic reach. CBT-E has been investigated in several samples in which CBT-E for bulimia nervosa, BED and EDNOS proved to be a successful treatment in the first studies after development of the CBT-E protocol (Byrne et al., 2011; Fairburn et al., 2009).

This review provides an update of the most recent (i.e. published since 2014) CBT-E effectiveness studies (randomized controlled trials (RCTs) and open trials) on bulimia nervosa, BED and transdiagnostic samples. Studies on the transdiagnostic samples include bulimia nervosa, BED, OSFED and, sometimes (i.e. in studies with lower BMI inclusion criteria), anorexia nervosa. However, excluded from the present review were studies with an anorexia nervosa sample alone, due to differences in treatment duration and other treatment variables (e.g. a focus on weight gain).

In this review the characteristics of the included studies are described, possible explanations for the variability in outcome are proposed, recommendations are made for future research, and the methodological quality of the RCTs is described. Due to the small number of included studies, no meta-analysis was performed.

## Materials and methods

### Search strategy and study selection

The primary search strategy was made in Medline, PsycInfo and EMBASE; the search covered the period from January 2014 up to March 2018. The following concepts were combined and searched for in the title and abstract:

1. Eating Disorder OR disordered eat\* OR binge eating disorder OR bulimia nervosa
2. Cognitive-behavioral OR CBT OR CBT-E

Articles had to meet the following criteria: i) a peer-reviewed study; ii) including a sample that meets the criteria for bulimia nervosa or BED, or a transdiagnostic sample with an eating disorder; and iii) an effectiveness study that includes (at least one condition of) manualized CBT-E.

After removing duplicates, 451 articles (published January 2014 to March 2018) were selected. The titles and abstracts of these articles were screened by the first author. The full-text versions of potential articles (n=35) were read to check for eligibility. The reference lists of the included articles and reviews were also examined for relevant studies.

Finally, seven articles met the inclusion criteria (Figure 4.1).

This review also includes an assessment of the methodological quality of the included RCTs. Tarrier and Wykes (Tarrier & Wykes, 2004) developed the Clinical Trials Assessment Test (CTAM), based on relevant features from the CONSORT guidelines (Moher et al., 2001), to assess the quality of trials of psychological treatments in mental health. This test contains 15 items grouped into six areas. Total scores range from 0 (no criterion is reached) to 100 (maximum score). The CTAM has good blind inter-rater agreement and adequate internal consistency (Tarrier & Wykes, 2004).

Ratings were done by the first author and one other independent rater. When required information was missing, the first author contacted the trial researchers for (possible) clarification.




Figure 4.1. Flow diagram of inclusion of studies for this review

## Results

If data were not reported, a calculation was made (when possible) based on the available data.

### Design

Of the seven included studies, five were RCTs (Fairburn et al., 2015; Poulsen et al., 2014; Thompson-Brenner et al., 2016; Wade, Byrne, & Allen, 2017; Wonderlich et al., 2014), and two were open trials (Dalle Grave et al., 2015;



Signorini et al., 2018). Of the two open trials, one specifically aimed to find evidence that CBT-E is generalizable to treatment conducted in a non-controlled clinical context (Signorini et al., 2018).

## **Recruitment and population**

All seven studies were conducted in an outpatient setting. Three studies included participants who were seeking help and had been referred (Dalle Grave et al., 2015; Fairburn et al., 2015; Signorini et al., 2018). Four studies also recruited participants through distribution of information in local papers, flyers, e-mails or (online) advertisements (Poulsen et al., 2014; Thompson-Brenner et al., 2016; Wade et al., 2017; Wonderlich et al., 2014). Four studies included a transdiagnostic sample (Dalle Grave et al., 2015; Fairburn et al., 2015; Signorini et al., 2018; Wade et al., 2017), two studies included participants with bulimia nervosa only (Poulsen et al., 2014; Wonderlich et al., 2014), and one study included participants with bulimia nervosa and comorbid (subthreshold) borderline personality disorder (Thompson-Brenner et al., 2016). Two transdiagnostic samples also included participants with anorexia nervosa (Signorini et al., 2018; Wade et al., 2017); this is explained by the use of a variable low-range cut-off for BMI, ranging substantially from 16 to 18.5. The Eating Disorder Examination (EDE) (Fairburn, Cooper, & O'Connor, 2008) is generally regarded as the gold standard in the assessment of an eating disorder. In five studies the diagnoses were assessed with the EDE (Fairburn et al., 2015; Poulsen et al., 2014; Thompson-Brenner et al., 2016; Wade et al., 2017; Wonderlich et al., 2014). In one study, the eating disorder was assessed by the treating therapists based on the DSM-IV criteria (Signorini et al., 2018), and in one study, no information was provided on how the eating disorder was diagnosed (Dalle Grave et al., 2015). Most studies included adults, although one study evaluated the effects of CBT-E in a cohort of non-underweight adolescents (Dalle Grave et al., 2015). There was a considerable difference in the number of participants per study (see Table 1).

## **Primary outcome measure and operationalization of clinical significant change**

In all studies the EDE (Fairburn et al., 2008), or its self-report version (EDE-Q) (Fairburn & Beglin, 2008), was used as the primary outcome measure. Four studies used the EDE (Fairburn et al., 2015; Poulsen et al., 2014; Thompson-Brenner et al., 2016; Wonderlich et al., 2014) and three the EDE-Q (Dalle Grave et al., 2015; Signorini et al., 2018; Wade et al., 2017). In all four studies using the EDE as primary outcome measure, the EDE was assessed by independent blinded assessors (Fairburn et al., 2015; Poulsen et al., 2014;

Thompson-Brenner et al., 2016; Wonderlich et al., 2014). However, studies used different definitions of clinical significant change to indicate relevant change (e.g. remission, good outcome, abstinence, minimal residual eating disorder psychopathology, etc.) and different operationalizations of these concepts. In the studies with a bulimia nervosa sample (Poulsen et al., 2014; Thompson-Brenner et al., 2016; Wonderlich et al., 2014), abstinence from binges and purging was the main definition for clinical significant change. In the transdiagnostic samples, an global EDE-(Q) score less than 1 SD above the community mean (sometimes combined with BMI  $\geq 18.5$ ) was defined as clinical significant change (Dalle Grave et al., 2015; Fairburn et al., 2015; Signorini et al., 2018; Wade et al., 2017) (Table 1). The two studies conducted in Australia (Signorini et al., 2018; Wade et al., 2017) used different EDE-Q norms; although both studies refer to Mond et al. (2006) for the norms used to indicate clinical significant change (less than 1 SD above the community mean, i.e.  $\leq 2.77$ ), the EDE-Q norms reported by Signorini et al. (2018) were  $\leq 2.46$  or less.

### **Cognitive behavioral therapy enhanced variant**

The seven included studies varied in: i) the setting in which therapy took place, ii) whether the focused or broad version of CBT-E was investigated, iii) the duration of therapy, and iv) whether extra sessions were planned involving significant others.

Four studies investigated the individual 20-session variant of the focused version of CBT-E (Fairburn et al., 2015; Poulsen et al., 2014; Thompson-Brenner et al., 2016; Wonderlich et al., 2014). In the study of Dalle Grave et al. (2015), parents were involved more closely, as participants were adolescents; the parental involvement consisted of five sessions of patients and parents together. Details about which version of CBT-E was investigated in this study were not reported. Wade et al. (2017) developed a treatment manual for group CBT-E based on the individual broad version of CBT-E including sessions to address the additional maintaining mechanisms (i.e. core low self-esteem, clinical perfectionism, and interpersonal problems). Eighteen group sessions of 2 h each were offered (with 5-10 min of individual work before each group session), and two additional individual sessions of 50 min each. In the study of Signorini et al. (2018), although CBT-E was investigated according to the manual (Fairburn, 2008), there was variability in the number of sessions (40 sessions for underweight participants, 20 sessions for nonunderweight participants) and also in the use of the focused or the broad version of CBT-E.

## Control group

Of the five RCTs, three compared CBT-E to another active condition (Fairburn et al., 2015; Poulsen et al., 2014; Wonderlich et al., 2014). In one study CBT-E was compared with psychoanalytic psychotherapy (Poulsen et al., 2014). In the study of Wonderlich et al. (2014), CBT-E was compared with a new psychotherapeutic treatment for bulimia nervosa, that is integrative cognitive-affective therapy (ICAT). In the study of Fairburn et al. (2015), CBT-E was compared with another evidence-based treatment for bulimia nervosa: interpersonal psychotherapy (IPT) (Fairburn et al., 2015). In two of these three studies, the therapy dosage was the same in both groups (Fairburn et al., 2015; Wonderlich et al., 2014), but in one study the duration of therapy differed greatly due to the nature of psychoanalytic psychotherapy (Poulsen et al., 2014), that is the psychoanalytic psychotherapy involved weekly sessions of 50 minutes each over 2 years (mean number of sessions 72.3). Thompson-Brenner et al. (2016) compared the focused and broad version of CBT-E in persons with comorbid bulimia nervosa and borderline personality disorder. In the RCT of Wade et al. (2017), the control group was a waiting list group; however, in that study, only the first 8 weeks were controlled for; after this period, the control group had a delayed treatment start.

## Therapist competence/treatment integrity

In four of the seven studies, the founder of CBT-E (Christopher Fairburn) or his colleague (Zafra Cooper) was closely involved in the training and supervision of the therapists (Dalle Grave et al., 2015; Fairburn et al., 2015; Poulsen et al., 2014; Thompson-Brenner et al., 2016). The remaining studies were supervised by experienced therapists (Signorini et al., 2018; Wade et al., 2017; Wonderlich et al., 2014). In six studies, the frequency of supervision was weekly or biweekly (Dalle Grave et al., 2015; Fairburn et al., 2015; Poulsen et al., 2014; Thompson-Brenner et al., 2016; Wade et al., 2017; Wonderlich et al., 2014). In the study of Signorini et al. (2018) the frequency of supervision was reported to be 'regular'. In three studies, the sessions were audio-recorded and a selection of these sessions was used and/or reviewed for purposes of supervision (Dalle Grave et al., 2015; Fairburn et al., 2015; Thompson-Brenner et al., 2016).

In three studies, treatment integrity was measured (Fairburn et al., 2015; Poulsen et al., 2014; Wonderlich et al., 2014). The quality of the delivery of the treatment condition was assessed by independent raters using diverse adherence scales. In these three studies, the raters scored adherence as 'good' (Wonderlich et al., 2014) or as 'high' (Fairburn et al., 2015; Poulsen et al., 2014).

## **Noncompleters**

The operationalization of 'completion' also differs between studies. In four studies 'completion' was operationalized as finishing the complete treatment (Dalle Grave et al., 2015; Fairburn et al., 2015; Poulsen et al., 2014; Wade et al., 2017). Wonderlich et al. (2014) defined completion as attending at least 16 sessions (of 21). In two studies (Signorini et al., 2018; Thompson-Brenner et al., 2016), it is not clear how completion was operationalized. Noncompletion rates ranged from 22.2% to 50%. In the open trial of Signorini et al. (2018), an attrition rate of 50% was reported whereas the other open trial (Dalle Grave et al., 2015) reported a substantially lower rate (25%) of noncompleters. In four of the RCTs the rate of noncompleters was similar, ranging from 22.2% (Poulsen et al., 2014) to 26.2% (Fairburn et al., 2015). In the RCT of Wade et al. (2017), 30% of the participants did not complete treatment.

## **Analysis**

All reported results are based on an intention-to-treat analysis.

### ***Randomized controlled trials***

Of the five RCTs, three reported significant differences between groups in favor of CBT-E (Fairburn et al., 2015; Poulsen et al., 2014; Wade et al., 2017). Wade et al. (2017) found that the first 8 weeks of group CBT-E were more effective in terms of reducing EDE-Q global scores compared with no treatment. In the study of Fairburn et al. (2015) the levels of eating disorder psychopathology decreased (global EDE score) in both conditions (CBT-E and IPT); however, the changes were significantly greater among CBT-E participants. The percentage of CBT-E participants in remission was almost twice as high as that in participants who received IPT (65.5% vs 33.3%). In the study of Poulsen et al. (2014), there was a large variation in treatment duration (5 months CBT-E vs. 24 months psychoanalytic psychotherapy). Significant differences were found between groups for binge eating and purging; 42% of the patients in CBT-E had ceased binge eating and purging (after 5 months) compared with 15% of the patients in psychoanalytic psychotherapy (after 24 months). By the end of both treatments, although there were substantial improvements in eating disorder psychopathology (global EDE scores), these changes took place more rapidly in CBT-E. In two out of five RCTs, no significant differences were found. In the study of Thomson-Brenner et al. (2016), two versions of CBT-E were compared (focused version vs. broad version). The groups did not differ in primary

outcome and the remission rate of the total sample was 42%. In addition, in the study of Wonderlich et al. (2014), comparing CBT-E with ICAT, no significant differences in treatment outcome were found between groups.

### ***Open trials***

In both open trials, there was a significant decrease in EDE-Q scores (Dalle Grave et al., 2015; Signorini et al., 2018) (Table 2). Dalle Grave et al. (2015) reported a remission rate of 67.6%; however, a substantial percentage of their patients (25%) met the criteria for remission before treatment started. Signorini et al. (2018) used two different definitions of remission and reported a remission rate of 42.2% and 35.4%, respectively. As mentioned, in the study of Wade et al. (2017), a control condition was included only in the first 8 weeks; after having received a full dosage of CBT-E, the remission rate for all patients (whether in the experimental or control group) was 66.7% (Table 1).

Differences in outcome, in RCTs and open trials, are explained in part by differences in the definition of clinical significant change and in the level of the EDE-Q community mean (Table 1).

### **Follow-up**

Of the seven included studies, five had a follow-up assessment period varying from: 3 months (Wade et al., 2017), 4 months (Wonderlich et al., 2014), 20 weeks (Signorini et al., 2018), 6 months (Thompson-Brenner et al., 2016) to 60 weeks (Fairburn et al., 2015). Generally, in most studies, the posttreatment results were maintained during follow-up. In the study of Wade et al. (2017), during follow-up, the percentage 'good outcome' decreased from 66.7% to 46.2%. In the study of Fairburn et al. (2015), the proportion of participants meeting the criteria for remission during follow-up increased in the IPT condition (33.3% to 49.0%), but the rate remained higher (69.4%) in CBT-E.

### **Assessing quality and variability in psychological treatment trials: the Clinical Trial Assessment Measure (CTAM)**

We used the CTAM (Tarrier & Wykes, 2004) to assess the methodological quality of the included RCTs (see Materials and methods). Three of the five RCTs had a similarly high CTAM score of 89 (Fairburn et al., 2015; Poulsen et al., 2014; Wonderlich et al., 2014), indicating good methodological quality. One of the RCTs described the process of assessor blinding (Thompson-Brenner et al., 2016), but none of them verified the blinding of assessors at

**Table 4.1.** CBT-E studies (published Jan 2014-March 2018): study characteristics and operationalization of clinical significant change

First author	Country	Design	N	Sample	BMI	Measure	Condition
Poulsen et al., 2014	Denmark	RCT	70	BN	-	EDE	CBT-Ef psychoanalytic psychotherapy
Wonderlich et al., 2014	USA	RCT	80	BN	≥18	EDE	CBT-Ef ICAT
Fairburn et al., 2015	UK	RCT	130	BN; 40.8% BED; 6.2% EDNOS; 53.1%	17.5-40	EDE	CBT-Ef IPT
Thompson-Brenner et al., 2016	USA	RCT	50	BN & BPS	-	EDE	CBT-Ef CBT-Eb
Wade et al., 2017	Australia	RCT <sup>b</sup>	40	AN; 20% BN; 57.5% BED; 5% OSFED; 17.5%	17.5-30	EDE-Q	Group CBT-Eb waiting list
Dalle Grave et al., 2015 <sup>a</sup>	Italy	Open trial	68	BN; 29.4% BED; 20.6% EDNOS; 50%	≥18.5	EDE-Q	CBT-E <sup>c</sup>
Signorini et al., 2018	Australia	Open trial	114	AN; 20.8% BN; 36.8% EDNOS; 42.5%	≥16	EDE-Q	CBT-Ef/Eb

AN = anorexia nervosa; BED = binge eating disorder; BMI = body mass index; BN = bulimia nervosa; BPS = borderline personality disorder; CBT-Eb = cognitive behavioral therapy enhanced broad version; CBT-Ef = cognitive behavioral therapy enhanced focused version; EDE = Eating Disorder Examination; EDE-Q = Eating Disorder Examination Questionnaire; EDNOS = eating disorder not otherwise specified; ICAT = integrative cognitive-affective therapy; IPT = interpersonal psychotherapy; ns = not significant; OSFED = other specified feeding or eating disorder; RCT = randomized controlled trial; WT = waiting list

<sup>a</sup> Sample: adolescents

<sup>b</sup> First eight weeks controlled

<sup>c</sup> Version not defined

<sup>d</sup> Criterion: abstinence of bingeing/purging in the past 4 weeks

<sup>e</sup> Global EDE(-Q) score less than 1 SD above the community mean and BMI ≥18.5

<sup>f</sup> Global EDE(-Q) score less than 1 SD above the community mean

Operationalization of clinical significant change				
Global EDE(-Q) less than 1 SD above community mean	BMI $\geq 18.5$	Binging and/or purging <sup>d</sup>	Clinical significant change posttreatment	Result
no	no	yes	42% 15%	CBT-E > psychoanalytic psychotherapy
no	no	yes	22.5% 37.5%	ns
yes (i.e. $\leq 1.74$ )	no	no	65.5% 33.3%	CBT-E > IPT
no	no	yes	44% 40%	ns
yes (i.e. $\leq 2.77$ )	yes	no	66.7% <sup>e</sup>	CBT-E > WT <sup>b</sup>
yes (i.e. $\leq 2.77$ )	no	no	67.6%	-
yes (i.e. $\leq 2.46$ )	yes	no	42.2% <sup>f</sup> 35.4% <sup>e</sup>	-

the end of the study. In the study of Thompson-Brenner et al. (2016), due to the small sample size and lack of measurement of treatment quality, the CTAM score was 7 points lower. Compared with the other four RCTs, the trial of Wade et al. (2017) had a lower CTAM score; this latter study had a small sample size, no independent randomization, no description of randomization, no active control condition, and no assessment of treatment quality.

A full description and ratings of the CTAM are available on request from the first author.

**Table 4.2.** Changes in EDE-Q global score in open trials: intention-to-treat analysis

First author	N	Pre-treatment Mean Global EDE(-Q) (SD)	Post-treatment Mean Global EDE(-Q) (SD)	Follow-up Mean Global EDE(-Q) (SD)
Dalle Grave et al., 2015	68	3.6 (1.5)	1.8 (1.8) <sup>a</sup>	–
Signorini et al., 2018	108	4.03 (1.29)	3.09 (1.76) <sup>a</sup>	3.10 (1.76)
Wade et al., 2017	39	4.37 (1.19)	2.36 (1.31) <sup>a</sup>	2.67 (1.44)

EDE = Eating Disorder Examination; EDE-Q = Eating Disorder Examination Questionnaire;  
SD = standard deviation

<sup>a</sup> significant at p <0.05

## Discussion

The findings of this systematic review of seven effectiveness studies (five RCTs and two open trials) replicate and extend findings from two earlier studies (Byrne et al., 2011; Fairburn et al., 2009), demonstrating that CBT-E is an effective treatment for bulimia nervosa, BED and transdiagnostic samples of adult patients with an eating disorder. Since 2014, several RCTs made a direct comparison between CBT-E and other active treatment conditions, such as interpersonal psychotherapy (IPT), psychoanalytic psychotherapy, and integrative cognitive-affective therapy (ICAT). Although IPT is also an established evidence-based treatment for bulimia nervosa and BED (Kass, Kolko, & Wilfley, 2013), the first direct comparison made between IPT and CBT-E in a transdiagnostic eating disorder sample, showed that CBT-E was more effective (Fairburn et al., 2015). In another comparison in a bulimia nervosa sample, 20 weeks of CBT-E was compared with two years of psychoanalytic psychotherapy (Poulsen et al., 2014). At the end of treatment, the considerable difference in remission rates of binge eating and purging in favour of CBT-E (in combination with the substantial differences



in treatment duration), demonstrates that CBT-E for bulimia nervosa is highly cost-effective compared with psychoanalytic psychotherapy. One study was the first to show that ICAT (a new psychotherapeutic treatment for bulimia nervosa) might be as effective as CBT-E (Wonderlich et al., 2014). Furthermore, group CBT-E seems to be an acceptable alternative to individual CBT-E (Wade et al., 2017). In a bulimia nervosa sample with comorbid borderline personality disorder, no difference in effect was found between the focused and the broad version of CBT-E (Thompson-Brenner et al., 2016). The study of Dalle Grave et al. (2015) showed that CBT-E might be a potential treatment approach for nonunderweight adolescents with an eating disorder. Although Family-Based Treatment (FBT) is the preferred treatment for adolescents with bulimia nervosa (Le Grange et al., 2015), CBT-E might be a possible alternative when, for example, FBT is not sufficiently effective or not available. Finally, the study of Signorini et al. (2018) showed that CBT-E is generalizable to a noncontrolled clinical context. However, that study had a high attrition rate of up to 50%, possibly due to the high percentage of participants with anorexia nervosa (20.8%) in their sample. In an earlier open trial (Byrne et al., 2011) with a transdiagnostic sample including anorexia nervosa, the attrition rate was also high (40%).

In this review, substantial differences were found in posttreatment remission rates (22.2% to 67.6%); when interpreting these differences, several issues need to be considered. First, studies are difficult to compare due to variation in the included samples, differences in the definition of clinical significant change, and differences in the methodological quality of the studies. For example, in the study of Dalle Grave et al. (2015), the high proportion that met the criteria for remission at baseline (25%) biases the relatively high posttreatment remission rate (67.6%). Also, the difference in 'good outcome' between the studies of Wade et al. (2017) and Signorini et al. (2018), both carried out in Australia, can be explained, in part, by the different EDE-Q community mean used for the definition of clinical significant change. Signorini et al. (2018) found a posttreatment remission rate of 42.2% (EDE-Q score  $\leq 2.46$ ), whereas Wade et al. (2017) reported 66.7% (EDE-Q score  $\leq 2.77$ ).

Moreover, differences between the studies are not always easy to explain. For example, the substantial difference in outcome of CBT-E between the study of Poulsen et al. (2014), with a posttreatment abstinence rate of 42% compared with the 22.5% reported by Wonderlich et al. (2014) is puzzling, as both studies are similar regarding their samples, operationalization of

clinical significant change (abstinence of binge eating/purging), and both are of good quality. One difference between these studies is that, in the study of Poulsen et al. (2014), the founder of CBT-E was closely involved in the training and supervision of the therapists. Another is how completion was operationalized. Poulsen et al. (2014) defined completion as finishing the complete treatment, whereas Wonderlich et al. (2014) defined completion as attending at least 16 sessions.

A strong point of the present study is that it is the first review on CBT-E to assess the methodological quality of the included RCTs. Moreover, the results of this assessment indicate that, overall, the quality of the studies was high.

Taken together, the effectiveness studies of CBT-E for bulimia nervosa, BED and transdiagnostic samples (published since January 2014), of which four RCTs with high methodological quality, provide additional and robust evidence that CBT-E is indeed an effective treatment for patients with eating disorders.

This systematic review excluded CBT-E trials which studied patients with anorexia nervosa alone; however, the two open studies with transdiagnostic samples also included patients with anorexia nervosa (Signorini et al., 2018; Wade et al., 2017). Although these latter studies show positive effects of CBT-E in these samples, the anorexia nervosa subgroups were not analysed separately. Also, although CBT-E has been described as promising for the treatment of anorexia nervosa (Fairburn et al., 2013), the results are not consistent (Calugi, El Ghoch, & Dalle Grave, 2017; Dalle Grave et al., 2016; Egger et al., 2016; Frostad et al., 2018). In an open trial, preliminary support was found for the use of CBT-E for anorexia nervosa (Fairburn et al., 2013). In a subsequent implementation study of CBT-E for outpatients with anorexia nervosa, half of the patients did not complete CBT-E whereas the remaining patients achieved a significant increase in BMI at 1-year follow-up (Frostad et al., 2018). In an open study among inpatients with anorexia nervosa, Calugi et al. (2017) found that CBT-E was well accepted and might be a viable and promising treatment, even for those with severe and enduring anorexia nervosa. Overall treatment results of CBT-E for anorexia nervosa were poorer than results of CBT-E for other eating disorders; however, this finding needs to be interpreted in the broader context of treatment studies on anorexia nervosa with overall poor posttreatment outcome (Waller, 2016).

Some recommendations can be made for future research. A trial with a direct comparison between CBT-E and another CBT protocol might help unravel the differential effects of CBT-E, and studies on the working mechanisms of CBT-E could strengthen its theoretical foundation. On the basis of our results, we also recommend that researchers facilitate comparability between CBT-E studies. Agreement should be reached concerning, for example, what outcome variable to use to establish clinical significant change, what level of competence is needed for a CBT-E therapist, what tool should be used to measure treatment integrity, and what specifically constitutes 'not completed' therapy.

This review has some limitations. First, the literature search and identification of relevant studies was done by one researcher (first author), implying that studies might have been missed and/or study characteristics or results may have been misinterpreted. Second, for practical reasons, only studies in the English language were included. Finally, the literature search was restricted to Medline, PsycINFO and Embase; although we tried to address this limitation by examining the reference lists of earlier meta-analyses and of the articles in this review, eligible articles may unintentionally have been missed.

## **Conclusion**

There is robust evidence that CBT-E is an effective treatment for adult patients with an eating disorder, especially for bulimia nervosa, BED and OSFED. Future research on the working mechanisms and differential effects of CBT-E compared with other CBT protocols might reveal the theoretical foundations and specificity of CBT-E. To establish good comparability between studies, we recommend that agreement be made between researchers, in particular regarding the operationalization of clinical significant change and the use of standard definitions.

## **KEY POINTS**

- There is robust evidence that CBT-E is an effective treatment for adult patients with an eating disorder, especially for bulimia nervosa, BED and OSFED.
- The substantial range in remission rates between studies is partly due to differences in study samples and the definition used for clinical significant change.
- Although IPT is an evidence-based treatment for bulimia nervosa and BED, the first direct comparison between IPT and CBT-E showed CBT-E to be more effective.
- CBT-E is a far more (cost-)effective treatment for bulimia nervosa than psychoanalytic treatment on the main parameters of bulimia nervosa, that is binge eating and purging.
- Future research should focus on the working mechanisms and differential effects of CBT-E compared with other CBT protocols to establish the specificity of CBTE.

## **Acknowledgements**

The authors thank professor Philip Spinhoven, PhD and professor Kees Korrelboom, PhD, for critically reading draft versions of this article; Willemke Marinus for her help with the literature search; Iris van der Meer, MSc, for assistance in examining included articles and reviews for relevant studies; Marie-Louise Kullberg, MSc, for CTAM ratings; and Judith Offringa for her editorial assistance.

## **Financial support and sponsorship**

None.

## **Conflicts of interest**

There are no conflicts of interest.

## References and recommended reading

Papers of particular interest, published within the annual period of review, have been highlighted as:

- of special interest
- of outstanding interest

Byrne, S. M., Fursland, A., Allen, K. L., & Watson, H. (2011). The effectiveness of enhanced cognitive behavioural therapy for eating disorders: an open trial. *Behaviour Research and Therapy*, 49(4), 219-226. <https://doi.org/10.1016/j.brat.2011.01.006>

▪ Calugi, S., El Ghoch, M., & Dalle Grave, R. (2017). Intensive enhanced cognitive behavioural therapy for severe and enduring anorexia nervosa: a longitudinal outcome study. *Behaviour Research and Therapy*, 89, 41-48. <https://doi.org/10.1016/j.brat.2016.11.006>

In this open trial, the effect of inpatient CBT-E for anorexia nervosa was evaluated. It showed that CBT-E was well accepted and might be a viable and promising treatment, even for those with severe and enduring anorexia nervosa.

Dalle Grave, R., Calugi, S., Sartirana, M., & Fairburn, C. G. (2015). Transdiagnostic cognitive behaviour therapy for adolescents with an eating disorder who are not underweight. *Behaviour Research and Therapy*, 73, 79-82. <https://doi.org/10.1016/j.brat.2015.07.014>

Dalle Grave, R., El Ghoch, M., Sartirana, M., & Calugi, S. (2016). Cognitive behavioral therapy for anorexia nervosa: an update. *Current Psychiatry Reports*, 18(1), 2. <https://doi.org/10.1007/s11920-015-0643-4>

Egger, N., Wild, B., Zipfel, S., Junne, F., Konnopka, A., Schmidt, U., et al. (2016). Cost-effectiveness of focal psychodynamic therapy and enhanced cognitive-behavioural therapy in out-patients with anorexia nervosa. *Psychological Medicine*, 46(16), 3291-3301. <https://doi.org/10.1017/S0033291716002002>

Erskine, H. E., Whiteford, H. A., & Pike, K. M. (2016). The global burden of eating disorders. *Current Opinion in Psychiatry*, 29(6), 346-353. <https://doi.org/10.1097/YCO.0000000000000276>

Fairburn, C. G. (2008). *Cognitive behavior therapy and eating disorders*. New York, NY: Guilford Press.

▪▪ Fairburn, C. G., Bailey-Straebl, S., Basden, S., Doll, H. A., Jones, R., Murphy, R., et al. (2015). A transdiagnostic comparison of enhanced cognitive behaviour therapy (CBT-E) and interpersonal psychotherapy in the treatment of eating disorders. *Behaviour Research and Therapy*, 70, 64-71. <https://doi.org/10.1016/j.brat.2015.04.010>  
This is the first RCT that made a direct comparison between CBT-E and another evidence-based treatment for eating disorders, IPT; CBT-E showed to be more effective.

Fairburn, C. G., & Beglin, S. J. (2008). Eating Disorder Examination Questionnaire (EDE-Q 6.0). In C. G. Fairburn (Ed.), *Cognitive behavior therapy and eating disorders* (pp. 309-313). New York, NY: Guilford Press.

Fairburn, C. G., Cooper, Z., Doll, H. A., O'Connor, M. E., Bohn, K., Hawker, D. M., et al. (2009). Transdiagnostic cognitive-behavioral therapy for patients with eating disorders: a two-site trial with 60-week follow-up. *American Journal of Psychiatry*, 166(3), 311-319. <https://doi.org/10.1176/appi.ajp.2008.08040608>

Fairburn, C. G., Cooper, Z., Doll, H. A., O'Connor, M. E., Palmer, R. L., & Dalle Grave, R. (2013). Enhanced cognitive behaviour therapy for adults with anorexia nervosa: a UK-Italy study. *Behaviour Research and Therapy*, 51(1), R2-8. <https://doi.org/10.1016/j.brat.2012.09.010>

Fairburn, C. G., Cooper, Z., & O'Connor, M. E. (2008). Eating Disorder Examination (16.0D). In C. G. Fairburn (Ed.), *Cognitive behaviour therapy and eating disorders* (pp. 265-308). New York: Guilford Press.

Fairburn, C. G., Cooper, Z., & Shafran, R. (2003). Cognitive behaviour therapy for eating disorders: a "transdiagnostic" theory and treatment. *Behaviour Research and Therapy*, 41(5), 509-528. [https://doi.org/10.1016/s0005-7967\(02\)00088-8](https://doi.org/10.1016/s0005-7967(02)00088-8)

- Frostad, S., Danielsen, Y. S., Rekkedal, G. A., Jevne, C., Dalle Grave, R., Ro, O., & Kessler, U. (2018). Implementation of enhanced cognitive behaviour therapy (CBT-E) for adults with anorexia nervosa in an outpatient eating-disorder unit at a public hospital. *Journal of Eating Disorders*, 6, 12. <https://doi.org/10.1186/s40337-018-0198-y>  
In this open trial, the implementation of CBT-E for anorexia nervosa was evaluated; although the drop-out rate was high, the remaining patients achieved a significant increase in BMI at 1year follow-up.

Hay, P., Chinn, D., Forbes, D., Madden, S., Newton, R., Sugenor, L., et al. (2014). Royal Australian and New Zealand College of Psychiatrists clinical practice guidelines for the treatment of eating disorders. *Australian & New Zealand Journal of Psychiatry*, 48(11), 977-1008. <https://doi.org/10.1177/0004867414555814>

- Hilbert, A., Hoek, H. W., & Schmidt, R. (2017). Evidence-based clinical guidelines for eating disorders: international comparison. *Current Opinion in Psychiatry*, 30(6), 423-437. <https://doi.org/10.1097/YCO.0000000000000360>

In this systematic review, nine evidence-based clinical treatment guidelines for eating disorders were compared. It provides an overview of commonalities and differences between these current clinical guidelines.

Hoek, H. W. (2016). Review of the worldwide epidemiology of eating disorders. *Current Opinion in Psychiatry*, 29(6), 336-339. <https://doi.org/10.1097/YCO.0000000000000282>

Jenkins, P. E., Hoste, R. R., Meyer, C., & Blissett, J. M. (2011). Eating disorders and quality of life: a review of the literature. *Clinical Psychology Review*, 31(1), 113-121. <https://doi.org/10.1016/j.cpr.2010.08.003>

- Kass, A. E., Kolko, R. P., & Wilfley, D. E. (2013). Psychological treatments for eating disorders. *Current Opinion in Psychiatry*, 26(6), 549-555. <https://doi.org/10.1097/YCO.0b013e328365a30e>
- Keel, P. K., Brown, T. A., Holm-Denoma, J., & Bodell, L. P. (2011). Comparison of DSM-IV versus proposed DSM-5 diagnostic criteria for eating disorders: reduction of eating disorder not otherwise specified and validity. *International Journal of Eating Disorders*, 44(6), 553-560. <https://doi.org/10.1002/eat.20892>
- Keski-Rahkonen, A., & Mustelin, L. (2016). Epidemiology of eating disorders in Europe: prevalence, incidence, comorbidity, course, consequences, and risk factors. *Current Opinion in Psychiatry*, 29(6), 340-345. <https://doi.org/10.1097/YCO.0000000000000278>
- Kolar, D. R., Rodriguez, D. L., Chams, M. M., & Hoek, H. W. (2016). Epidemiology of eating disorders in Latin America: a systematic review and meta-analysis. *Current Opinion in Psychiatry*, 29(6), 363-371. <https://doi.org/10.1097/YCO.0000000000000279>
- Le Grange, D., Lock, J., Agras, W. S., Bryson, S. W., & Jo, B. (2015). Randomized clinical trial of family-based treatment and cognitive-behavioral therapy for adolescent bulimia nervosa. *Journal of the American Academy of Child and Adolescent Psychiatry*, 54(11), 886-894 e882. <https://doi.org/10.1016/j.jaac.2015.08.008>
- Machado, P. P., Goncalves, S., & Hoek, H. W. (2013). DSM-5 reduces the proportion of EDNOS cases: evidence from community samples. *International Journal of Eating Disorders*, 46(1), 60-65. <https://doi.org/10.1002/eat.22040>
- Mangweth-Matzek, B., & Hoek, H. W. (2017). Epidemiology and treatment of eating disorders in men and women of middle and older age. *Current Opinion in Psychiatry*, 30(6), 446-451. <https://doi.org/10.1097/YCO.0000000000000356>
- Mitchison, D., Hay, P., Slewa-Younan, S., & Mond, J. (2012). Time trends in population prevalence of eating disorder behaviors and their relationship to quality of life. *PLoS One*, 7(11), e48450. <https://doi.org/10.1371/journal.pone.0048450>
- Moher, D., Schulz, K. F., Altman, D., & Group, C. (2001). The CONSORT statement: revised recommendations for improving the quality of reports of parallel-group randomized trials. *JAMA*, 285(15), 1987-1991. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/11308435>
- Mond, J. M., Hay, P. J., Rodgers, B., & Owen, C. (2006). Eating Disorder Examination Questionnaire (EDE-Q): norms for young adult women. *Behaviour Research and Therapy*, 44(1), 53-62. <https://doi.org/10.1016/j.brat.2004.12.003>
- National Institute for Health and Care Excellence (NICE). (2017). Eating disorders: recognition and treatment. In *NICE guideline [NG69]* (pp. 41). London, UK: NICE.
- Perez, M., Ohrt, T. K., & Hoek, H. W. (2016). Prevalence and treatment of eating disorders among Hispanics/Latino Americans in the United States. *Current Opinion in Psychiatry*, 29(6), 378-382. <https://doi.org/10.1097/YCO.0000000000000277>

- ■ Poulsen, S., Lunn, S., Daniel, S. I., Folke, S., Mathiesen, B. B., Katznelson, H., & Fairburn, C. G. (2014). A randomized controlled trial of psychoanalytic psychotherapy or cognitive-behavioral therapy for bulimia nervosa. *American Journal of Psychiatry*, 171(1), 109-116. <https://doi.org/10.1176/appi.ajp.2013.12121511>

In this RCT, 20 weeks of CBT-E for BN was compared with 2 years of psychoanalytic psychotherapy; CBT-E appeared a far more (cost-)effective treatment for BN than psychoanalytic treatment on the main parameters of bulimia nervosa, that is binge eating and purging.

- Signorini, R., Sheffield, J., Rhodes, N., Fleming, C., & Ward, W. (2018). The effectiveness of enhanced cognitive behavioural therapy (CBT-E): a naturalistic study within an out-patient eating disorder service. *Behavioural and Cognitive Psychotherapy*, 46(1), 21-34. <https://doi.org/10.1017/S1352465817000352>

In this open trial, which includes a transdiagnostic eating disorder sample, the implementation of CBT-E was evaluated and showed that CBT-E is generalizable to a noncontrolled clinical context.

Smink, F. R., van Hoeken, D., & Hoek, H. W. (2013). Epidemiology, course, and outcome of eating disorders. *Current Opinion in Psychiatry*, 26(6), 543-548. <https://doi.org/10.1097/YCO.0b013e328365a24f>

Steinhausen, H. C. (2009). Outcome of eating disorders. *Child and Adolescent Psychiatric Clinics of North America*, 18(1), 225-242. <https://doi.org/10.1016/j.chc.2008.07.013>

Tarrier, N., & Wykes, T. (2004). Is there evidence that cognitive behaviour therapy is an effective treatment for schizophrenia? A cautious or cautionary tale? *Behaviour Research and Therapy*, 42(12), 1377-1401. <https://doi.org/10.1016/j.brat.2004.06.020>

Thomas, J. J., Lee, S., & Becker, A. E. (2016). Updates in the epidemiology of eating disorders in Asia and the Pacific. *Current Opinion in Psychiatry*, 29(6), 354-362. <https://doi.org/10.1097/YCO.0000000000000288>

- Thompson-Brenner, H., Shingleton, R. M., Thompson, D. R., Satir, D. A., Richards, L. K., Pratt, E. M., & Barlow, D. H. (2016). Focused vs. broad enhanced cognitive behavioral therapy for bulimia nervosa with comorbid borderline personality: a randomized controlled trial. *International Journal of Eating Disorders*, 49(1), 36-49. <https://doi.org/10.1002/eat.22468>

In this RCT with a comorbid bulimia nervosa and borderline personality disorder sample, no difference in effect was found between the focused and the broad version of CBT-E.

van Hoeken, D., Burns, J. K., & Hoek, H. W. (2016). Epidemiology of eating disorders in Africa. *Current Opinion in Psychiatry*, 29(6), 372-377. <https://doi.org/10.1097/YCO.0000000000000274>

Wade, S., Byrne, S., & Allen, K. (2017). Enhanced cognitive behavioral therapy for eating disorders adapted for a group setting. *International Journal of Eating Disorders*, 50(8), 863-872. <https://doi.org/10.1002/eat.22723>



Waller, G. (2016). Recent advances in psychological therapies for eating disorders. *F1000Research*, 5. <https://doi.org/10.12688/f1000research.7618.1>

- Wonderlich, S. A., Peterson, C. B., Crosby, R. D., Smith, T. L., Klein, M. H., Mitchell, J. E., & Crow, S. J. (2014). A randomized controlled comparison of integrative cognitive-affective therapy (ICAT) and enhanced cognitive-behavioral therapy (CBT-E) for bulimia nervosa. *Psychological Medicine*, 44(3), 543-553. <https://doi.org/10.1017/S0033291713001098>

First study that showed that ICAT (a new psychotherapeutic treatment for bulimia nervosa) might be as effective as CBT-E.

Yager, J., Devlin, M. J., Halmi, K. A., B., H. D., Mitchell III, J. E., Powers, P., & Zerbe, K. J. (2014). Guideline watch (August 2012): Practice guideline for the treatment of patients with eating disorders, 3rd Edition. *Focus*, 12(4), 416-431. <https://doi.org/10.1176/appi.focus.120404>

