



Universiteit
Leiden
The Netherlands

Physiological measurements of the effect of cord clamping strategies

Brouwer, E.

Citation

Brouwer, E. (2021, September 22). *Physiological measurements of the effect of cord clamping strategies*. Retrieved from <https://hdl.handle.net/1887/3213482>

Version: Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/3213482>


Note: To cite this publication please use the final published version (if applicable).



PART TWO

Placental transfusion and the effect of
spontaneous breathing





1

CHAPTER 1

Effect of spontaneous breathing on umbilical venous blood flow and placental transfusion during delayed cord clamping in preterm lambs

E Brouwer
AB te Pas
GR Polglase
E McGillick
KJ Crossley
K Rodgers
D Blank
S Yamaoka
AW Gill
M Kluckow
SB Hooper

Arch Dis Child Fetal Neonatal Ed 2020;105:F26-F32

ABSTRACT

INTRODUCTION

During delayed umbilical cord clamping, the factors underpinning placental transfusion remain unknown. We hypothesised that reductions in thoracic pressure during inspiration would enhance placental transfusion in spontaneously breathing preterm lambs.

OBJECTIVE

Investigate the effect of spontaneous breathing on umbilical venous flow and body weight in preterm lambs.

METHODS

Pregnant sheep were instrumented at 132-133 days gestational age to measure fetal common umbilical venous, pulmonary and cerebral blood flows as well as arterial and intra-pleural (IP) pressures. At delivery, doxapram and caffeine were administered to promote breathing. Lamb body weights were measured continuously and breathing was assessed by IP pressure changes.

RESULTS

In 6 lambs, 491 out of 1117 breaths were analysed for change in body weight. Weight increased in 46.6% and decreased in 47.5% of breaths. An overall mean increase of 0.02 ± 2.5 g per breath was calculated, and no net placental transfusion was observed prior to cord clamping (median difference in body weight 52.3 [-54.9-166.1]g, $p=0.418$). Umbilical venous (UV) flow transiently decreased with each inspiration, and in some cases ceased, before UV flow normalised during expiration. The reduction in UV flow was positively correlated with the standardised reduction in IP pressure, increasing by 109 mL/min for every SD reduction in IP pressure. Thus, the reduction in UV flow was closely related to inspiratory depth.

CONCLUSION

Spontaneous breathing had no net effect on body weight in preterm lambs at birth. UV blood flow decreased as inspiratory effort increased, possibly due to constriction of the inferior vena cava caused by diaphragmatic contraction, as previously observed in human fetuses.

INTRODUCTION AND RATIONALE

Placental to infant blood transfusion during delayed umbilical cord clamping (DCC) after birth⁽¹⁾ is evidenced by an increase in birth weight and a decrease in placental blood volume⁽²⁻⁴⁾ and is thought to occur at 2-3 mL/kg/min over the first 3 min after delivery.⁽⁵⁾ Animal studies have also shown that DCC until after ventilation onset prevents the reduction in cardiac output caused by immediate cord clamping (ICC).⁽⁶⁾ This finding has led to the concept that cord clamping should be based on the infant's physiology (i.e. whether it is breathing) rather than on 'time after birth', which can be unrelated to the infant's physiological state.⁽⁷⁻¹⁰⁾ This concept is termed 'physiological based cord clamping (PBCC)' and highlights the importance of pulmonary ventilation in mediating the benefits of DCC.⁽¹¹⁻¹³⁾

In term newborns, DCC increases haemoglobin and ferritin levels, reduces rates of iron deficiency and anaemia in the first year of life.⁽¹⁴⁾ In preterm infants, small randomised trials have shown that DCC is associated with less intraventricular haemorrhage, necrotizing enterocolitis, sepsis and need for blood transfusions, which are common complications associated with preterm birth.⁽¹⁵⁻¹⁷⁾ As a result, the WHO now recommends DCC for >60 s in infants not requiring resuscitation.

While placental transfusion is an accepted feature of DCC, little is known about the physiology causing the net movement of blood from the placenta into the infant during DCC. Most commentaries suggest that placental transfusion results from gravity (i.e. when the infant is placed below the mother) or from uterine contractions squeezing blood out of the placenta into the infant.^(1, 18) However, recent clinical trials in humans and animal experiments have failed to demonstrate an effect of gravity or uterine contractions on placental transfusion.^(4, 19) In a unique observational study in normal term infants, Doppler ultrasound was used to investigate blood flow patterns in both the umbilical artery and vein during DCC.⁽²⁰⁾ This study showed that umbilical venous (UV) flow was heavily influenced by breathing and crying. During inspiration, umbilical venous flow appeared to increase, whereas between breaths it appeared to cease. Similarly, umbilical artery flow was influenced by breathing and crying and at times was bidirectional, which was thought to result from uterine contractions.⁽²⁰⁾

Animal experiments investigating the mechanisms of placental transfusion have consistently failed to detect a shift in blood volume from the placenta into the newborn.^(9, 19) Possible explanations for this relate to the animal models used, which have mostly involved lambs delivered by caesarean section, followed by mechanical ventilation.^(9, 19) In this situation, intermittent increases in intrathoracic pressure combined with end-expiratory pressures may restrict net placenta-to-infant blood transfusion. In contrast, all human studies showing placental

transfusion have been in vaginally delivered spontaneously breathing infants.⁽²¹⁾ In this study, we hypothesised that the sub atmospheric (negative) intrathoracic pressure created during inspiration would enhance UV flow into the newborn, leading to a net increase in weight.

METHODS

All experimental procedures were conducted in accordance with the National Health and Medical Research Council code of practice for the care and use of animals for scientific purposes.

Fetal surgery

Sterile fetal surgery was performed on six pregnant ewes (Merino X Border Leicester) at 131-132 days (term is 147 days; 131-134 is considered late preterm) of gestational age, 3 days prior to delivery.⁽²²⁾ Briefly, anaesthesia was induced using an intravenous bolus of sodium thiopentane (20 mL at 50 mg/mL) and maintained following intubation by inhalation of isoflurane (1%-3% Isoflow, Abbot, Australia) in air/oxygen. Antibiotics (cefazoline, 1 g/5 mL) were administered before surgery. The fetal head, neck and abdomen were exteriorised via hysterotomy and jugular vein and carotid artery catheters inserted. Ultrasonic flow probes (Transonic Systems, Ithaca, New York, USA) were placed around the left pulmonary artery, common umbilical vein and right carotid artery. In lambs, the two umbilical veins within the cord join at the umbilicus to form the intra-abdominal common umbilical vein before it enters the fetal liver and divides into the ductus venosus or joins the hepatic portal system. A sterile, saline-filled intra-pleural (IP) balloon catheter was also inserted into the IP space to measure changes in IP pressure during breathing. Following instrumentation, the fetus was returned to the amniotic sac and both catheters and flow-probe leads were exteriorised via the ewe's right flank for continuous monitoring until delivery, 3 days later. During the post-operative period, ewes received analgesia via a fentanyl patch (75 µg/hour) and daily antibiotics (2.5 mL cefazoline 1 g/5mL).

Experimental protocol

Prior to delivery, electronic recordings of blood flows in the left pulmonary artery, umbilical vein and right carotid artery as well as pressures in the carotid artery and IP space commenced using a data acquisition system (Powerlab ADI, Sydney, Australia). As general anaesthesia is known to inhibit spontaneous breathing activity, lambs were delivered under spinal anaesthesia at 133-134 days of

gestations. Ewes were sedated with 10 mL propofol 1% (10 mg/mL) to administer spinal anaesthesia (lignocain 2%, 0.1 mL/kg) via the sacral lumbar space. Following successful induction of spinal anaesthesia, mild sedation of the ewe continued using midazolam (1.0 mg/kg/hour at 20-25 mL/hour). The fetal head and neck were exteriorised via caesarean section and the lambs were intubated using an endotracheal tube (4.5 mm cuffed endotracheal tube). Lambs were intubated as the application of facemask without leak is problematic and, in addition intubation ensured opening of the larynx. A transcutaneous oxygen saturation probe (Masimo, Radical 7, California, USA) was placed around the right foreleg, a near infrared spectroscopy (NIRS) optode (Casmed Foresight, CAS Medical Systems, Branford, Connecticut, USA) was placed over the left frontal cortex, and a rectal temperature probe was inserted and used to correct blood gas measurements. The lamb was dried and placed on electronic scales (Bilanciari D70, Campogalliano, Italy) immediately beside the ewe, taking care to avoid stretching or mechanically interfering with the umbilical cord; the lamb's weight was recorded continuously. Lambs were given naloxone (400 µg intravenous) and anexate (0.04 mg bolus intravenous) to counter the inhibitory effects of fentanyl (postoperative analgesia of the ewe) and midazolam (maternal sedation), respectively, on respiration. They were also given caffeine (20 mg/kg loading dose and 10 mg/kg/hour) and doxapram (5 mg/kg loading dose and 2.5 mg/kg/hour) to stimulate breathing.

If the lamb was breathing spontaneously, it was given continuous positive airway pressure (5 cmH₂O) via endotracheal tube (4.5 mm cuffed endotracheal tube) and disturbed as little as possible to avoid affecting the body weight recordings and other physiological measurements. If the lamb did not initiate spontaneous breathing following delivery and physical stimulation, a sustained inflation (SI) was administered (peak inspiratory pressure (PIP) of 35 cmH₂O for 30 s). Following the SI, if spontaneous breathing did not commence or if the lambs became apnoeic, brief periods of intermittent positive pressure ventilation (PIP of 35 cmH₂O with a positive end expiratory pressure (PEEP) of 5 cmH₂O) was applied to avoid hypoxia. During the experiment, no supplemental oxygen was administered to the lamb as it was still receiving oxygenation from the ewe via the umbilical cord. All physiological parameters were recorded continuously throughout the experiment. Umbilical cord clamping occurred once the lamb was assessed to have established a stable breathing pattern and the pulmonary blood flow (PBF) waveform had adopted the neonatal phenotype (figure 1); this took up to 19:15 [15:55-20:25] mins postdelivery to achieve in our experiments.




Figure 1 | Pulmonary blood flow (PBF) wave form before and after lung aeration

Following cord clamping, the lamb was placed under a radiant heater and was ventilated using a Babylog 8000+ ventilator (Dräger, Lübeck, Germany) with a PEEP of 5 cmH₂O for an additional 30 minutes. All physiological recordings commenced prior to, or at delivery (NIRS, oxygen saturation (SpO₂), weight) and arterial blood gasses were taken at regular intervals following delivery to assess lamb well-being. All blood samples and liquid volumes taken or given to the lamb while they were being weighed were recorded to adjust the weight measurements.

The ewe was euthanised following cord clamping, using an overdose of pentobarbitone (100 mg/kg intravenous). Lambs were euthanised after completion of the ventilation period using an overdose of pentobarbitone administered intravenously. Lamb body and organ weights were recorded post mortem.

Analytical methods

Recordings of physiological measurements made prior to, during, and after delivery were analysed using LabChart 8 (ADInstruments). A breath-by-breath analysis, which involved analysing every individual spontaneous breath in all animals, was performed to observe the effect of breathing on weight and UV flow. All breathing events were annotated manually by the researcher (EB) to determine change in lamb weight, IP pressure and UV blood flow with each spontaneous breath. To make IP pressure comparable across animals, measurements were standardised per individual (mean was subtracted and values were divided by the SD). The basal IP pressure is a relative measure that varies between animals, whereas the change in IP pressure from baseline is a precise measurement. Standardised values per animal allow interpretation of the data in terms of quantiles, for example, a value of 2 corresponds to the 97.5% quantile of IP measurements within this animal. Analysis of carotid arterial (CA) pressure, CA blood flow and PBF, involved averaging values over five consecutive heartbeats for each time point.

Statistical analysis

Statistical analysis was performed using IBM SPSS Statistics V.24.0. Data on CA pressure, CA blood flow, PBF, UV flow and temperature are presented as median and IQR. Paired t-test was used to compare weight measurements at delivery with weight measurements prior to cord clamping. A linear regression analysis was used to determine the effect of breathing on UV flow and weight in separate analyses. Animal ID were added as fixed factor to each model to account for clustering of measurements within animals. A p-value ≤ 0.05 was considered to be statistically significant. A sensitivity analysis was performed for the relationships between the reduction in IP pressure and both UV flow and weight by excluding the animal with the largest number of measurements. These analyses (data not shown) were performed to exclude the possibility that measurements from a single animal dominated the results. As results were very similar to the analysis with the complete data, the outcomes were not influenced by the number of measurements from any one animal.

RESULTS

Fetal characteristics and blood gas status at the beginning of experiments are displayed in table 1. One out of the six lambs was male.

Table 1 | Fetal characteristics

N=6	Fetal	5 min prior to cord clamping	5 min after cord clamping	10 min after cord clamping
Gestational age [days]	134 [134-134]	-	-	-
pH	7.39 [7.37-7.42]	7.27 [7.27-]	7.13 [7.12-7.22]	7.12 [7.07-7.24]
pCO ₂	45.6 [44.6-49.0]	54.4 [44.9-]	70.2 [64.4-73.3]	72.1 [52.5-85.8]
pO ₂	23.6 [21.2-27.1]	35.1 [32.9-]	42.4 [22.7-51.7]	74.0 [47.0-96.1]
Saturation [%]	69.4 [58.9-72.3]	82.0 [74.9-]	84.4 [50.5-90.9]	96.8 [74.5-98.1]

Characteristics displayed as median [IQR]

Effect of spontaneous breaths on lamb weight

The breath-by-breath analysis evaluated 1117 breaths, which resulted in an average reduction of IP pressure of 8.6 ± 0.3 mmHg (range 2.2 to 72.2 mmHg); this equates to a mean reduction in pressure of 11.2 ± 0.6 cmH₂O (range 2.9 to 93.8 cmH₂O). Of the 1117 breaths analysed, 626 breaths were excluded from weight analysis due to major changes caused by artefacts such as body movement. Of the 491 breaths analysed for weight, 46.6% of breaths were associated with an

increase in weight, 5.9% were associated with no change in weight and 47.5% of breaths were associated with a decrease in weight. Overall, the mean increase in lamb weight per breath was $0.02 \pm 2.5\text{g}$ (figure 2A). The increase in weight was positively correlated with the standardised reduction in IP pressure. The regression coefficient was 0.839 meaning that for an IP pressure that is 1 SD above the mean, weight increased by 0.839 g. Median [IQR] lamb body weight was 3969 [3635-4131] g at delivery, 3924 [3668-4080] g at 10 minutes before cord clamping and 4038 [3610-4201] g at time of cord clamping. No net placental transfusion was measured or observed during spontaneous breathing prior to cord clamping as measured by body weight change ($p=0.418$).




Figure 2 | Relationship between intra-pleural pressure and body weight change and common umbilical venous (UV) blood flow per breath

Common Umbilical Venous flow

In 44% of the breaths analysed, blood flow in the common UV transiently (0.31 ± 0.19 s in duration) decreased with inspiration (figure 3 and 4); the delay between peak reduction in IP pressure and peak reduction in UV flow varied between 120 and 300ms. The mean reduction in common UV flow per breath was 183.4 ± 4.5 mL/min (range 79.4 to 627.8 mL/min). Reductions in UV flow that decreased below zero were also commonly observed, indicating that flow in the common UV had reversed (indicated by negative blood flow value) and transiently flowed back towards the placenta (figure 4). These reversals in flows were most often associated with large inspiratory efforts that were immediately followed by transient increases in IP pressure. Following these reversals in flow, common UV flow transiently increased above baseline values following the restoration in IP pressure. Reductions in UV flow that often resulted in a reversal of common UV flow, also occurred during transient increases in IP pressure that were not preceded by inspiration (figure 4); this was relatively common in all lambs.




Figure 3 | Physiological recordings of intra-pleural (IP) pressure, body weight and common umbilical venous (UV) blood flow

The reduction in common UV flow was found to be positively correlated with the standardised reduction in IP pressure. The regression coefficient was 109.0 and as such, a 1 SD reduction in IP pressure below the mean basal value will reduce UV flow by 109.0 mL/min (figure 2B). Thus, as the depth of the inspiration increased, the associated reduction in common UV flow increased, causing flow to cease or even reverse with some breaths (figures 3 and 4).




Figure 4 | Physiological recordings of intra-pleural (IP) pressure and common umbilical venous (UV) blood flow

Haemodynamics

Carotid arterial pressure and flow

Following delivery and ventilation onset, mean CA pressure and blood flow remained relatively constant for the 10 min leading up to cord clamping. Immediately following cord clamping, CA pressure (by ≈ 0.5 mmHg) slightly increased. While CA pressure remained relatively constant, CA flow gradually increased, reaching a maximum of 86 mL/min at 8 min following cord clamping (table 2).

Pulmonary artery blood flow

At 10 mins prior to cord clamping, PBF was quite variable (table 2) between lambs, with some lambs having already aerated their lungs and transitioned their pulmonary circulation (as indicated by PBF wave form; figure 1), whereas others still had a predominantly fetal PBF waveform phenotype. Immediately following cord clamping, PBF increased and tended to increase further over the subsequent 10 min following umbilical cord clamping.

Table 2 | Averaged measurements in the 10 min prior to and 10 min after cord clamping

Time	CA pressure (mmHg)	CA flow (mL/min)	PA flow (mL/min)	Body Temperature
-10 min	48.73 [40.37 - 56.11]	60.35 [43.10 - 83.63]	137.82 [62.22 - 254.00]	39.5 [39.0 - 39.8]
-5 min	50.32 [46.94 - 58.39]	43.32 [27.25 - 61.75]	240.53 [138.00 - 298.67]	38.6 [37.7 - 39.5]
-2 min	48.02 [41.16 - 55.40]	42.33 [32.94 - 57.20]	312.64 [179.14 - 403.90]	38.2 [37.4 - 38.9]
Cord clamped	54.45 [49.03 - 61.00]	49.27 [40.37 - 62.33]	354.40 [224.28 - 457.53]	38.2 [37.8 - 38.8]
1 min	54.98 [50.06 - 62.14]	52.15 [46.46 - 73.45]	255.87 [218.88 - 422.99]	38.2 [37.8 - 38.7]
2 min	58.96 [47.83 - 65.98]	58.14 [50.32 - 86.52]	314.78 [218.00 - 462.60]	38.1 [37.2 - 38.4]
3 min	55.51 [50.16 - 64.59]	62.65 [47.64 - 92.08]	281.72 [224.34 - 446.64]	38.1 [37.8 - 38.4]
4 min	58.83 [49.43 - 70.06]	68.41 [61.64 - 105.60]	313.45 [239.18 - 541.80]	38.0 [37.2 - 38.3]
5 min	59.97 [48.41 - 69.04]	73.61 [61.91 - 92.85]	319.15 [217.48 - 449.91]	38.0 [37.5 - 38.6]
6 min	55.28 [48.74 - 70.17]	74.36 [53.45 - 93.81]	260.98 [200.38 - 320.13]	37.9 [37.0 - 38.6]
7 min	58.27 [49.60 - 74.29]	82.93 [69.51 - 126.51]	389.07 [268.02 - 463.93]	36.7 [35.7 - 38.5]
8 min	55.30 [47.03 - 79.75]	85.66 [64.12 - 106.34]	343.40 [280.24 - 437.31]	37.8 [37.4 - 38.6]
9 min	71.65 [48.77 - 75.96]	71.93 [59.60 - 102.05]	356.79 [286.36 - 600.84]	38.1 [37.2 - 38.7]
10 min	64.26 [46.97 - 71.41]	78.96 [48.34 - 97.13]	355.93 [275.70 - 567.71]	38.0 [37.3 - 38.7]

CA pressure, CA flow and PA flow are displayed as median [IQR], 10 min before and 10 min after cord clamping. CA pressure increases after cord clamping with ≈ 4 mmHg at 4 minutes, after which it remains stable. CA flow increases up to 8 minutes with ≈ 36 mL/min. PA flow increases with a maximum of 35 mL/min after cord clamping. Temperature decreases with 0.2°C.

CA, carotid arterial; PA, pulmonary artery.

DISCUSSION

International resuscitation guidelines currently recommend DCC for 60 s in infants who do not need resuscitation, largely based on the concept of placental to infant blood transfusion.^(23,24) While previous studies have shown a beneficial effect of DCC on neonatal outcomes,^(14, 21) the physiological mechanisms underpinning the net shift of blood from the placenta into the infant during DCC remains unknown. We consider that a better understanding of this physiology and the factors influencing placental transfusion could help optimise the practice of DCC in the delivery room. We hypothesised that the reduction in intrathoracic pressure associated with inspiration would provide a pressure gradient favouring blood flow from the placenta into the infant and restrict flow in the opposite direction. However, contrary to our hypothesis, we found that UV flow into newborn lambs transiently reduced, ceased or even reversed during individual breaths, depending upon the respiratory pattern, and that the reduction in flow was positively correlated with the depth of the inspiration. It is not surprising, therefore, we also did not observe a relationship between spontaneous breathing and weight gain of the lamb.

Evidence for placental transfusion in humans is well established⁽⁵⁾ and includes an increase in infant weight during DCC, an increase in neonatal blood volume and a decrease in blood volume remaining within the placenta after DCC.^(1, 2, 5, 25, 26) However, our previous animal studies have been unable to provide any evidence of placental transfusion, which we considered was due to the absence of spontaneous breathing; instead all lambs were mechanically ventilated.^(9, 19) As such, in this study we aimed to investigate whether sub atmospheric pressures generated during inspiration resulted in placental transfusion. To measure placental transfusion, lamb body weights and UV flows were measured continuously throughout a period of DCC. However, the body weight measurements proved to be challenging as any slight movement or touching of the lamb caused large rapid changes in measured weight that were artefactual and unrelated to body weight changes; these sections of physiological recordings were manually detected and excluded from analysis. To counter the inhibitory effects of maternal sedation on spontaneous breathing, all lambs received drugs to reverse these effects and therefore body movements were common during the experiment, resulting in some unreliable weight measurements (figure 3). Nevertheless, even when the lambs were not moving and the weights were stable, we could not identify a relationship between breathing (IP pressure) and weight change over an entire breathing episode (figure 2A). Nevertheless, the linear regression analysis detected a relationship between the standardised IP reduction and a very small increase in weight. Considering that the UV flow was markedly reduced with each breath (see below), it is difficult to explain how this might occur. However, the effect of individual breaths on weight change are transient, reversed at the end of the breath and may involve a transient discrepancy between UV flow into and umbilical arterial (UA) flow out of the lamb.

Although we found that UV flow was influenced by spontaneous breathing and the associated decrease in IP pressure, the relationship was opposite to what we hypothesised. We hypothesised that the decrease in intrathoracic pressure associated with inspiration would provide a pressure gradient to increase UV flow into the lamb. However, we found that UV blood flow transiently decreased during inspiration, commonly decreasing to zero at end-inspiration. Furthermore, we found a strong positive relationship between the degree of decrease in UV flow and the size of the reduction in intrathoracic pressure generated by inspiration (figure 2B). While we were initially surprised by this finding, it is consistent with previously described findings measured by ultrasound in human fetuses during periods of fetal breathing movements (FBM).⁽²⁷⁾ This study showed that during inspiration, contraction of the diaphragm causes narrowing and closure of the inferior vena cava (IVC) as it passes through the diaphragm, causing IVC flow to cease. Blood returning from the UV, either via the ductus venosus or hepatic veins, also has to pass through the diaphragm via the same orifice as the IVC. As such, it is likely

that the ductus venosus constricts along with the IVC as the diaphragm contracts, causing flow in the UV to cease during inspiration. Due to the large compliance of the low pressure venous system, it is highly likely that these reductions in flow are slightly delayed in relation to the timing of inspiration. We found the delay to be 120-300ms in the intra-abdominal common UV, but this delay is likely to be larger when measured further upstream, closer to the placenta. As a result, when measured upstream in the cord, depending on the breathing rate, the reductions in UV flow may be completely out of synchrony with breathing. This finding explains why increases in UV flow appeared to be related to inspiration, whereas cessations in UV flow appeared to be related to expiration, in our previous study in human infants; in that study UV flows were measured upstream in the cord.⁽²⁰⁾ Nevertheless, the findings convincingly demonstrate that large sub atmospheric reductions in intrathoracic pressure cannot account for placental transfusion.

As noted in figure 3, both increases in IP pressure by itself, or following a large inspiratory effort, can result in a transient reversal of blood flow in the common UV, pushing blood back out of the lamb toward the placenta. These increases in IP pressure were undoubtedly due to abdominal muscle contractions that must also have resulted in large increases in intra-abdominal pressure. In the physiological recording displayed in figure 3, each inspiratory effort is immediately followed by a forced expiratory effort, as indicated by a large increase in IP pressure (up to 40 mmHg), which must have been driven by abdominal muscle contractions. In addition, a reversal in UV flow was also observed when transient increases in IP pressure were not preceded by inspiration. The precise reasons for these are unclear, but abdominal muscle contractions associated with grunting or body movements are possible explanations. Nevertheless, whether or not the increases in intrathoracic (and abdominal) pressure are associated with breathing, they are associated with a transient reversal in UV flow. This finding is not surprising given that the IP pressure increase (up to 40 mmHg; figure 3) is considerably larger than UV pressure in the cord. However, once the IP pressures begin to stabilise, UV blood flow transiently increases above baseline, which likely reflects a rebound increase in flow caused by blood pooling upstream in the cord and placenta (figure 3).

Ventilating the lung before clamping the umbilical cord, also known as PBCC, is known to promote haemodynamic stability immediately after birth.^(6,28) In particular, it avoids the large rapid increase in blood pressure caused by increasing systemic vascular resistance when the cord is clamped and the reduction in cardiac output caused by the loss of umbilical venous return and ventricular preload.⁽¹⁰⁾ Avoiding these significant cardiovascular events is likely to be a major contributing factor to the reduction in neonatal morbidities, such as intraventricular haemorrhage, associated with DCC.⁽⁶⁾ We found that CA pressure increased slightly following umbilical cord clamping (by ≈ 0.5 mmHg) but this increase is substantially less than

the increase that occurs with cord clamping before ventilation/breathing onset (≈ 20 mmHg).⁽⁶⁾ While the CA pressure tended to increase over the subsequent 9 min (by ≈ 15 mmHg), this increase appeared more to do with the lamb's activity levels than to cord clamping. Similarly, CA blood flow increased over the same time frame (by 30 mL/min), which indicates that cardiac output was sustained throughout the cord clamping period. PBF doubled in the minutes leading up to cord clamping, which was expected as it is well established that the increase in PBF after birth is triggered by lung aeration resulting from breathing onset.^(29, 30) Although PBF tended to increase in response to cord clamping, before cord clamping the majority of right ventricular output was mainly directed through the lungs rather than across the ductus arteriosus. This is clearly indicated by the PBF wave form (figure 1), whereby after transition, forward flow into the lung occurs throughout the cardiac cycle. Forward flow into the lung during diastole results from a reversal of flow through the ductus arteriosus due to the reduction in PVR causing a reversal in the pressure gradient between systemic and pulmonary circulations.⁽²²⁾ As clamping of the umbilical cord increases systemic arterial pressure, the increase in PBF following cord clamping likely results from increased left-to-right shunting of blood through the ductus arteriosus during diastole. These results are consistent with our previous findings in ventilated lambs.⁽⁶⁾ Taken together, these results clearly demonstrate a haemodynamically stable transition when lung aeration precedes cord clamping, irrespective of whether the lambs were ventilated or breathed spontaneously.

Temperature measurements stayed stable throughout the experiments, with the median temperature varying between 36.7 °C and 39.5 °C (39.0°C is the normal core body temperature in fetal sheep). Before cord clamping, the external application of heat required to maintain lamb body temperature was minimal and restricted to one hot water bottle and a wrap. However, after cord clamping an overhead radiant heater, two hot water bottles and a wrap were needed to maintain a stable temperature. While anecdotal, this indicates that when the cord is intact, umbilical blood flow may provide an effective source of heat that helps to maintain body temperature. While some heat loss may occur over the length of the umbilical cord, presumably this was compensated for by the continuous flow of blood flowing through the umbilical vessels at the ewe's body temperature.

CONCLUSION

In summary, contrary to our hypothesis, we found that during inspiration, UV flow was either reduced or ceased depending upon the size of the inspiration. This finding was likely caused by narrowing or closure of the IVC during contraction of the diaphragm during inspiration as previously described in human fetuses during FBM *in utero*.⁽²⁷⁾ The reduction in UV flow was positively correlated with the depth of the inspiration. Increases in IP pressure, likely caused by abdominal contractions, also reduced UV flow and caused it to transiently reverse, resulting in blood flowing out of the lamb and pooling in the cord or placenta. This study confirms the previous results of Bhatt *et al*, showing haemodynamic stability when ventilation onset precedes cord clamping and demonstrates that the mode of ventilation (spontaneous breathing vs mechanical ventilation) does not impact on this stability or on placental transfusion.⁽⁶⁾ Instead, the haemodynamic stability is reliant on the increase in PBF. Clearly further research is required to understand the mechanisms responsible for placental transfusion.

REFERENCES

1. Yao AC, Hirvensalo M, Lind J. Placental transfusion-rate and uterine contraction. *Lancet* (London, England). 1968;1(7539):380-3.
2. Law GR, Cattle B, Farrar D, Scott EM, Gilthorpe MS. Placental blood transfusion in newborn babies reaches a plateau after 140 s: Further analysis of longitudinal survey of weight change. *SAGE open medicine*. 2013;1:2050312113503321.
3. Mercer JS, Erickson-Owens DA, Collins J, Barcelos MO, Parker AB, Padbury JF. Effects of delayed cord clamping on residual placental blood volume, hemoglobin and bilirubin levels in term infants: a randomized controlled trial. *Journal of perinatology : official journal of the California Perinatal Association*. 2017;37(3):260-4.
4. Vain NE, Satragno DS, Gorenstein AN, Gordillo JE, Berazategui JP, Alda MG, et al. Effect of gravity on volume of placental transfusion: a multicentre, randomised, non-inferiority trial. *Lancet* (London, England). 2014;384(9939):235-40.
5. Yao AC, Moinian M, Lind J. Distribution of blood between infant and placenta after birth. *Lancet* (London, England). 1969;2(7626):871-3.
6. Bhatt S, Alison BJ, Wallace EM, Crossley KJ, Gill AW, Kluckow M, et al. Delaying cord clamping until ventilation onset improves cardiovascular function at birth in preterm lambs. *The Journal of physiology*. 2013;591(8):2113-26.
7. Polglase GR, Dawson JA, Kluckow M, Gill AW, Davis PG, Te Pas AB, et al. Ventilation onset prior to umbilical cord clamping (physiological-based cord clamping) improves systemic and cerebral oxygenation in preterm lambs. *PloS one*. 2015;10(2):e0117504.
8. Polglase GR, Blank DA, Barton SK, Miller SL, Stojanovska V, Kluckow M, et al. Physiologically based cord clamping stabilises cardiac output and reduces cerebrovascular injury in asphyxiated near-term lambs. *Archives of disease in childhood Fetal and neonatal edition*. 2018;103(6):F530-f8.
9. Blank DA, Polglase GR, Kluckow M, Gill AW, Crossley KJ, Moxham A, et al. Haemodynamic effects of umbilical cord milking in premature sheep during the neonatal transition. *Archives of disease in childhood Fetal and neonatal edition*. 2018;103(6):F539-f46.
10. Hooper SB, Polglase GR, Roehr CC. Cardiopulmonary changes with aeration of the newborn lung. *Paediatric respiratory reviews*. 2015;16(3):147-50.
11. Knol R, Brouwer E, Vernooij ASN, Klumper F, DeKoning P, Hooper SB, et al. Clinical aspects of incorporating cord clamping into stabilisation of preterm infants. *Archives of disease in childhood Fetal and neonatal edition*. 2018;103(5):F493-f7.
12. Hooper SB, Polglase GR, te Pas AB. A physiological approach to the timing of umbilical cord clamping at birth. *Archives of disease in childhood Fetal and neonatal edition*. 2015;100(4):F355-60.
13. Hooper SB, Binder-Heschl C, Polglase GR, Gill AW, Kluckow M, Wallace EM, et al. The timing of umbilical cord clamping at birth: physiological considerations. *Maternal health, neonatology and perinatology*. 2016;2:4.
14. McDonald SJ, Middleton P, Dowswell T, Morris PS. Effect of timing of umbilical cord clamping of term infants on maternal and neonatal outcomes. *The Cochrane database of systematic reviews*. 2013;2013(7):Cd004074.
15. Rabe H, Reynolds G, Diaz-Rossello J. A systematic review and meta-analysis of a brief delay in clamping the umbilical cord of preterm infants. *Neonatology*. 2008;93(2):138-44.

16. Fogarty M, Osborn DA, Askie L, Seidler AL, Hunter K, Lui K, et al. Delayed vs early umbilical cord clamping for preterm infants: a systematic review and meta-analysis. *American journal of obstetrics and gynecology*. 2018;218(1):1-18.
17. Tarnow-Mordi W, Morris J, Kirby A, Robledo K, Askie L, Brown R, et al. Delayed versus Immediate Cord Clamping in Preterm Infants. *The New England journal of medicine*. 2017;377(25):2445-55.
18. Yao AC, Lind J. Effect of gravity on placental transfusion. *Lancet (London, England)*. 1969;2(7619):505-8.
19. Hooper SB, Crossley KJ, Zahra VA, van Vonderen J, Moxham A, Gill AW, et al. Effect of body position and ventilation on umbilical artery and venous blood flows during delayed umbilical cord clamping in preterm lambs. *Archives of disease in childhood Fetal and neonatal edition*. 2017;102(4):F312-f9.
20. Boere I, Roest AA, Wallace E, Ten Harkel AD, Haak MC, Morley CJ, et al. Umbilical blood flow patterns directly after birth before delayed cord clamping. *Archives of disease in childhood Fetal and neonatal edition*. 2015;100(2):F121-5.
21. Rabe H, Diaz-Rossello JL, Duley L, Dowswell T. Effect of timing of umbilical cord clamping and other strategies to influence placental transfusion at preterm birth on maternal and infant outcomes. *The Cochrane database of systematic reviews*. 2012(8):Cd003248.
22. Crossley KJ, Allison BJ, Polglase GR, Morley CJ, Davis PG, Hooper SB. Dynamic changes in the direction of blood flow through the ductus arteriosus at birth. *The Journal of physiology*. 2009;587(Pt 19):4695-704.
23. Wyckoff MH, Aziz K, Escobedo MB, Kapadia VS, Kattwinkel J, Perlman JM, et al. Part 13: Neonatal Resuscitation: 2015 American Heart Association Guidelines Update for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care. *Circulation*. 2015;132(18 Suppl 2):S543-60.
24. WHO Guidelines Approved by the Guidelines Review Committee. Guideline: Delayed Umbilical Cord Clamping for Improved Maternal and Infant Health and Nutrition Outcomes. Geneva: World Health Organization. Copyright © World Health Organization 2014.; 2014.
25. Jahazi A, Kordi M, Mirbehbahani NB, Mazloom SR. The effect of early and late umbilical cord clamping on neonatal hematocrit. *Journal of perinatology : official journal of the California Perinatal Association*. 2008;28(8):523-5.
26. Linderkamp O, Nelle M, Kraus M, Zilow EP. The effect of early and late cord-clamping on blood viscosity and other hemorheological parameters in full-term neonates. *Acta paediatrica (Oslo, Norway : 1992)*. 1992;81(10):745-50.
27. Nyberg MK, Johnsen SL, Rasmussen S, Kiserud T. Hemodynamics of fetal breathing movements: the inferior vena cava. *Ultrasound in obstetrics & gynecology : the official journal of the International Society of Ultrasound in Obstetrics and Gynecology*. 2011;38(6):658-64.
28. Brouwer E, Knol R, Vernooij ASN, van den Akker T, Vlasman PE, Klumper F, et al. Physiological-based cord clamping in preterm infants using a new purpose-built resuscitation table: a feasibility study. *Archives of disease in childhood Fetal and neonatal edition*. 2019;104(4):F396-f402.
29. Lang JA, Pearson JT, Binder-Heschl C, Wallace MJ, Siew ML, Kitchen MJ, et al. Increase in pulmonary blood flow at birth: role of oxygen and lung aeration. *The Journal of physiology*. 2016;594(5):1389-98.
30. Lang JA, Pearson JT, te Pas AB, Wallace MJ, Siew ML, Kitchen MJ, et al. Ventilation/perfusion mismatch during lung aeration at birth. *Journal of applied physiology (Bethesda, Md : 1985)*. 2014;117(5):535-43.