Universiteit
Leiden
The Netherlands

Molecular and functional characterization of the scavenger receptor CD36 in zebrafish and common carp

Fink, I.R.; Benard, E.L.; Hermsen, T.; Meijer, A.H.; Forlenza, M.; Wiegertjes, G.F.

Citation

Fink, I. R., Benard, E. L., Hermsen, T., Meijer, A. H., Forlenza, M., & Wiegertjes, G. F. (2015). Molecular and functional characterization of the scavenger receptor CD36 in zebrafish and common carp. *Molecular Immunology*, 63(2), 381-393.
doi:10.1016/j.molimm.2014.09.010

Version: Publisher's Version

License: [Licensed under Article 25fa Copyright Act/Law \(Amendment Taverne\)](#)

Downloaded from: <https://hdl.handle.net/1887/3197041>

Note: To cite this publication please use the final published version (if applicable).

Molecular and functional characterization of the scavenger receptor CD36 in zebrafish and common carp

Inge R. Fink^a, Erica L. Benard^b, Trudi Hermsen^a, Annemarie H. Meijer^b,
Maria Forlenza^a, Geert F. Wiegertjes^{a,*}

^a Cell Biology and Immunology Group, Wageningen Institute of Animal Sciences, Wageningen University, PO Box 338, 6700 AH Wageningen, The Netherlands

^b Institute of Biology, Leiden University, Einsteinweg 55, 2333 CC Leiden, The Netherlands

ARTICLE INFO

Article history:

Received 20 June 2014

Received in revised form

11 September 2014

Accepted 17 September 2014

Available online 11 October 2014

Keywords:

CD36

Scavenger receptor

Zebrafish

Carp

Immune function

ABSTRACT

CD36 is a scavenger receptor which has been studied closely in mammals where it is expressed by many different cell types and plays a role in highly diverse processes, both homeostatic and pathologic. It is among other things important in the innate immune system, in angiogenesis, and in clearance of apoptotic cells, and it is also involved in lipid metabolism and atherosclerosis. Recently, in the cephalochordate amphioxus a primitive CD36 family member was described, which was present before the divergence of CD36 from other scavenger receptor B family members, SCARB1 and SCARB2. Not much is known on the Cd36 molecule in teleost fish. We therefore studied Cd36 in both zebrafish and common carp, two closely related cyprinid fish species. Whereas a single *cd36* gene is present in zebrafish, carp has two *cd36* genes, and all show conserved synteny compared to mammalian CD36. The gene expression of carp *cd36* is high in brain, ovary and testis but absent in immune organs. Although in mammals CD36 expression in erythrocytes, monocytes and macrophages is high, gene expression studies in leukocyte subtypes of adult carp and zebrafish larvae, including thrombocytes and macrophages provided no indication for any substantial expression of *cd36* in immune cell types. Surprisingly, analysis of the *cd36* promoter region does show the presence of several binding sites for transcription factors known to regulate immune responses. Overexpression of carp *cd36* locates the receptor on the cell surface of mammalian cell lines consistent with the predicted topology of cyprinid Cd36 with a large extracellular domain, two trans-membrane domains, and short cytoplasmic tails at both ends. Gene expression of *cd36* is down-regulated during infection of zebrafish with *Mycobacterium marinum*, whereas knockdown of *cd36* in zebrafish larvae led to higher bacterial burden upon such infection. We discuss the putative role for Cd36 in immune responses of fish in the context of other members of the scavenger receptor class B family.

© 2014 Elsevier Ltd. All rights reserved.

1. Introduction

Scavenger receptors are soluble or membrane-associated molecules which have in common the ability to bind chemically modified lipoproteins or other poly-anionic ligands (Plüddemann et al., 2007). Each molecule additionally binds to other ligands, and scavenger receptors are thus functionally diverse. They have

been grouped together into eight classes (A–H) where members within a class display structural homology. However, between the eight classes the scavenger receptors are structurally very different. CD36 is a member of the scavenger receptor class B family, which also contains scavenger receptor class B member 1 and 2 (SCARB1 and SCARB2) (Calvo and Vega, 1993; Calvo et al., 1995). SCARB1 is important in lipid metabolism as a receptor mediating cholesterol transfer to and from high density lipoprotein HDL (Brundert et al., 2005; Chadwick and Sahoo, 2013). SCARB2 (also referred to as lysosomal integral membrane protein-2, LIMP-2) is required for the normal biogenesis and maintenance of lysosomes and endosomes and as such plays a role in various human diseases (Gonzalez et al., 2014). CD36 is an integral membrane glycoprotein with two trans-membrane domains, a large extracellular loop containing multiple glycosylation sites, and two short intracellular tails. In mammals, CD36 is expressed by many different cell types such as platelets,

Abbreviations: CD36, cluster of differentiation 36; dpf, days post fertilization; dpi, days post injection; hpf, hours post fertilization; LDL, low density lipoprotein; LTA, lipoteichoic acid; PBL, peripheral blood leukocytes; RT-PCR, reverse transcription polymerase chain reaction; RT-qPCR, real-time quantitative polymerase chain reaction; SCARB, scavenger receptor class B; TLR, Toll-like receptor.

* Corresponding author. Tel.: +31 317 482732.

E-mail address: geert.wiegertjes@wur.nl (G.F. Wiegertjes).

monocytes and macrophages, endothelial cells and hepatocytes, and CD36 seems to play a role in highly diverse processes, both homeostatic and pathologic (Febbraio et al., 2001). Mammalian CD36 was originally identified as a macrophage receptor for oxidized LDL (Endemann et al., 1993), but its ligand-binding ability is very extensive and encompasses endogenous as well as exogenous ligands (reviewed in Silverstein and Febbraio, 2009). Among other molecules, CD36 recognizes microbial diacylglycerides such as macrophage-activating lipopeptide 2 from *Mycoplasma pneumoniae* (MALP-2) and lipoteichoic acid from *Staphylococcus aureus* and acts as a co-receptor for these molecules together with Toll-like receptor (TLR) 2/6 heterodimers (Hoebe et al., 2005; Stuart et al., 2005). As a receptor recognizing exogenous pathogen-associated molecular patterns (PAMPs), CD36 is acknowledged as a pattern recognition receptor (PRR) important in innate immunity.

CD36 orthologs have been described in many vertebrate species such as human, mouse (Endemann et al., 1993), rat (Abumrad et al., 1993), cow (Greenwalt et al., 1990) and chicken (Shu et al., 2011). Ancestral homologs are also found in invertebrate species such as *C. elegans* (Means et al., 2009), sea urchin and fruit fly, and a recent study reported several homologs in amphioxus, a basal chordate (Zhang et al., 2013). Presumably due to amphioxus-specific gene duplication events there are five *cd36* paralogs in this species. These genes are thought to represent the primitive form of scavenger receptor class B before the divergence of CD36, SCARB1 and SCARB2. One of the amphioxus genes was cloned and studied in more detail: its expression was up-regulated in the gills upon feeding and down-regulated during fasting. In addition, studies in Atlantic salmon and rainbow trout have focused on fatty acid transport proteins including Cd36 and described gene expression changes regulated by insulin, fasting and dietary composition (Sanchez-Gurmaches et al., 2011, 2012; Torstensen et al., 2011). This suggests a conservation of the role of SCARB family members in at least nutritional control and/or lipid metabolism. Contrary to the mammalian species, knowledge of the immune function of Cd36 in teleost fish is lacking.

In mammals, there is evidence of CD36 interaction with pre-formed Toll-like receptor TLR2–TLR6 heterodimers, where CD36 recognizes specific lipids or lipoproteins and brings the ligand to the TLRs. Ligand/receptor clusters localize within lipid rafts of the plasma membrane, after which intracellular signalling and phagocytosis is initiated (Triantafyllou et al., 2006). The intracellular signalling goes via MyD88 and a series of phosphorylation and ubiquitination events of cytosolic proteins, leading to activation of the transcription factor NF κ B and expression of immune related genes (Akira and Takeda, 2004). Of interest, out of eight accessory molecules classified in mammals as important mediators of ligand delivery and/or recognition and required for TLR function (Lee et al., 2012) only four molecules, including CD36, seem to have clear orthologs in fish genomes (Pietretti et al., 2013). Given our interest in the function of TLRs (Pietretti and Wiegertjes, 2014) among which Tlr2 (Ribeiro et al., 2010), we studied Cd36 in zebrafish and common carp, two closely related cyprinid fish species, taking advantage of the complementary tools available for these two species to achieve a more thorough understanding of the function of Cd36 in teleost fish.

We characterize for the first time in detail Cd36 of zebrafish and two Cd36 molecules of common carp finding a molecular structure consistent with mammalian CD36. Phylogenetic analysis as well as a synteny approach allowed us to confidently designate the cyprinid molecules as homologs of the mammalian CD36. Gene expression was studied in tissues and leukocyte subtypes, indicating a complete absence of *cd36* basal gene expression in immune organs and leukocytes of adult carp. However, knock-down of *cd36* in zebrafish embryos by a morpholino approach and subsequent *Mycobacterium marinum* infection indicated a role for

Cd36 in controlling bacterial burden. We discuss the possibility that the division of functions between the Cd36 family members Cd36, Scarb1 and Scarb2, all of which are present in teleosts, may not be exactly the same as in their mammalian counterparts.

2. Materials and methods

2.1. Animals

European common carp (*Cyprinus carpio carpio* L.) were reared in the central fish facility Carus, at Wageningen University, Wageningen, Netherlands. Fish were kept at 23 °C in recirculating UV-treated tap water and fed pelleted dry food (Sniff, Soest, Germany) daily. R3xR8 carp are the hybrid offspring of a cross between fish of Polish origin (R3 strain) and Hungarian origin (R8 strain) (Irnazarow, 1995). Carp were between 9 and 11 months old at the start of the experiments. All studies were performed with approval from the local animal welfare committee (DEC) of Wageningen University.

Zebrafish were handled in compliance with the local animal welfare regulations, maintained according to standard protocols (zfinfo.org), and culture was approved by the local animal welfare committee (DEC) of Leiden University. Embryos from the zebrafish AB/TL line were used for the infection experiments. Embryos were grown at 28.5 °C in egg water (60 μ g/mL Instant Ocean sea salts). For the duration of bacterial injections and stereo fluorescence imaging embryos were kept under anaesthesia in egg water containing 200 μ g/mL tricaine methane sulfonate (Sigma-Aldrich, St. Louis, MO, USA). Embryos used for stereo fluorescence imaging were kept in egg water containing 0.003% 1-phenyl-2-thiourea (Sigma-Aldrich) to prevent melanisation which can produce background fluorescence during imaging. For fluorescence activated cell sorting (FACS) isolation of macrophages and neutrophils from zebrafish larvae, the transgenic lines *Tg(mpeg1:mCherry-F)*^{UMSF001} (Bernut et al., 2014) and *Tg(mpx:egfp)*ⁱ¹¹⁴ (Renshaw et al., 2006) were used, respectively.

2.2. Organ isolation

Carp aged 9–11 months were euthanized with 0.3 g/L tricaine methane sulfonate (TMS, Crescent Research Chemicals, Phoenix, AZ, USA) buffered with 0.6 g/L NaHCO₃. Carp were bled from the caudal vein using a needle and syringe containing cRPMI medium (RPMI 1640 with 25 mM HEPES, (Lonza, Basel, Switzerland) adjusted to an osmolality of 280 mOsm/kg with sterile water) containing 50 U/mL heparin (Leo Pharma, Ballerup, Denmark), 50 U/mL penicillin G (Sigma-Aldrich), and 50 μ g/mL streptomycin sulphate (Sigma-Aldrich). For isolation of peripheral blood leukocytes (PBL), the heparinized blood was centrifuged at 100 $\times$ g for 5 min at 4 °C and then another 5 min at 300 $\times$ g. The buffy coat was collected, carefully layered on Ficoll-Paque PLUS (GE Healthcare, Little Chalfont, UK) and centrifuged at 800 $\times$ g for 25 min at 4 °C without brake. The leukocyte layer was collected and washed twice with cRPMI. The obtained PBL were stored at –80 °C until used for RNA isolation. After bleeding the fish, the organs of interest were aseptically removed and immediately frozen in liquid nitrogen and stored at –80 °C until used for RNA isolation.

Zebrafish embryos for RNA isolation were snap-frozen in liquid nitrogen and stored at –80 °C.

2.3. Isolation of leukocyte subtypes

Carp leukocyte subtypes were isolated by density gradient separation and/or magnetic cell sorting using specific antibodies as

described before for thrombocytes (Rombout et al., 1996), thymocytes (Stolte et al., 2008), granulocytes (Forlenza et al., 2008), B cells (Koumans-van Diepen et al., 1995; Secombes et al., 1983), and macrophages (Romano et al., 1998). In short, PBL or single-cell suspensions derived from carp organs were incubated with primary mouse monoclonal antibody: WCL-6 for thrombocytes (from blood), TCL-BE8 for neutrophils (from mid kidney), WCL-12 for B cells (from blood), and WCL-15 for monocytes/macrophages (from spleen). After incubation and washing, cells were stained with phycoerythrin (PE)-conjugated goat anti-mouse secondary antibody. After washing and counting of cells, magnetic beads (anti-PE MicroBeads, Miltenyi Biotec GmbH, Bergisch Gladbach, Germany) were added and allowed to bind, before washing and magnetic separation on LS Midi Columns using a MidiMACS Separator (Miltenyi Biotec).

For isolation of leukocytes from *Tg(mpeg1:mCherry-F)*^{UMSF001} and *Tg(mpx:egfp)*^{j114} zebrafish, larvae at 5–6 days post fertilization (dpf) were dissociated according to Covassin et al. (2006). In short, anaesthetized embryos were dechorionated using pronase treatment, rinsed in calcium-free Ringer solution, followed by digestion with 0.25% trypsin and 1 mM EDTA. The obtained cell suspension was centrifuged, rinsed with PBS and resuspended in Leibovitz medium L15 without phenol red, 1% foetal calf serum, 0.8 mM CaCl₂, penicillin 50 U/mL and streptomycin 0.05 mg/mL. The single cell suspension was subjected to FACS at room temperature using a FACS Aria (Becton Dickinson) with the BD FACSDiva software version 5.0.3 and a Coherent Sapphire solid state laser 488 nm with 13 mW power. The fluorescence-positive and negative cell fractions were collected in medium as above but with 10% foetal calf serum.

2.4. RNA isolation

Total RNA from carp organs and leukocytes was extracted using the RNeasy Mini kit according to the manufacturer's protocol (Qiagen, Venlo, Netherlands) including on-column DNase treatment with the RNase-free DNase set (Qiagen). Final elution was performed with 30 µL nuclease-free water. The integrity of the RNA was determined by agarose gel electrophoresis and the RNA quality and concentrations were assessed spectrophotometrically by measuring the absorbance at 260 nm and 280 nm (Nanodrop, Thermo Scientific, Waltham, MA, USA). RNA from *M. marinum*-infected zebrafish was isolated at 4 dpi as described in Stockhammer et al. (2009). RNA from zebrafish larval leukocytes was isolated directly after FACS sorting using the RNAqueous-Micro Kit (Ambion, Carlsbad, CA, USA). Zebrafish RNA quality was checked by lab-on-a-chip analysis with an Agilent Bioanalyzer 2100 using the RNA 6000 Pico kit (Agilent, Santa Clara, CA, USA). RNA was stored at –80 °C until use.

2.5. cDNA synthesis

Prior to cDNA synthesis, 500 ng–2 µg of carp total RNA was subjected to an additional DNase treatment by using DNase I Amplification Grade (Invitrogen, Carlsbad, CA, USA). Synthesis of cDNA was performed with Invitrogen's SuperScript III Reverse Transcriptase, according to the manufacturer's instructions. As control for genomic contamination, for each sample a reaction without SuperScript III Reverse Transcriptase was performed. cDNA samples were diluted 25 times in nuclease-free water before use as templates in real-time quantitative PCR experiments. For zebrafish samples, cDNA synthesis was performed as described previously (Stockhammer et al., 2009), using iScript reverse transcriptase (Bio-Rad Laboratories, Hercules, CA, USA). Zebrafish cDNA was diluted 10 times before further use.

2.6. Real-time quantitative PCR

Real-time quantitative PCR (RT-qPCR) on samples from carp was performed in a Rotor-Gene 6000 with a 72-well rotor (Corbett Research, Qiagen) with the Absolute QPCR SYBR Green Mix (Thermo Scientific) as detection chemistry. All primers were from Eurogentec (Liège, Belgium) and were used as forward and reverse primer mix at a 2.1 µM final concentration. Master-mix for each RT-qPCR run was prepared as follows: per reaction, 2 µL of primer stock was mixed with 7 µL of SYBR Green. To 9 µL of master-mix, 5 µL of diluted cDNA was added in a 0.1 mL tube. The following amplification program was used: one denaturation step of 15 min at 95 °C; followed by 40 cycles of a three-step amplification (15 s at 95 °C for denaturation, 20 s at 60 °C for annealing, 20 s at 72 °C for elongation). Then a short hold for 1 min at 60 °C, before a melting analysis starting from 60 °C and increasing to 90 °C in 0.5 °C steps with 5 s waiting. Fluorescence acquisition was performed at the end of each cycle and during the waiting steps of the melting analysis.

RT-qPCR data were analysed by Rotor-Gene 6000 Series Software 1.7. The melting temperature and profile of the melting curves were used to validate the specificity of the amplification. The gene expression was analysed using the average amplification efficiency for each primer pair and the take-off value of each sample (as derived from the Comparative Quantitation Analysis of the Rotor-Gene 6000 Series Software 1.7). The relative gene expression was calculated as the average amplification efficiency to the power of the take-off value. The gene expression of the house-keeping gene *40S ribosomal protein S11* was used to normalize the data.

RT-qPCR analysis of zebrafish samples was performed as described in Stockhammer et al. (2009). Reactions contained 10 µL of 2× iQ SYBR Green Supermix (Bio-Rad Laboratories), 5 pmol of each primer, and 5 µL diluted cDNA in a total of 20 µL. All reactions were performed with biological triplicates and technical duplicates. Data from zebrafish leukocytes were normalised to the expression of *eukaryotic translation initiation factor 4A, isoform 1B (eif4)*, whereas data from *M. marinum*-infected zebrafish embryos were normalized to *peptidylprolyl isomerase A-like (ppial)*. For analysis purposes an amplification efficiency of 2 was assumed.

The sequences of primers used in this study are given in Table 1.

2.7. Cloning full-length *cd36a* from common carp

Carp head kidney leukocytes were obtained essentially as described previously (Verbarg-van Kemenade et al., 1994). RNA was isolated as described above and then used as template in reverse-transcription PCR using the LongRange 2Step RT-PCR kit (Qiagen). Reverse transcription was done with an oligo-dT primer and the subsequent PCR was done with gene-specific primers (CycCD36.FW1 and CycCD36.RV3, see Table 1) whose sequence was based on preliminary information from the carp genome. Semi-nested PCR was then performed with the Expand High Fidelity PCR System (Roche Diagnostics, Basel, Switzerland) with primers CycCD36.FW1 and CycCD36.RV2 (see Table 1). The PCR product was subsequently cloned into pGEM-T Easy (Promega, Madison, WI, USA) and transformed into *E. coli* JM109 competent cells (Promega). Eight clones were sequenced using the ABI prism BigDye Terminator Cycle Sequencing Ready Reaction kit (Applied Biosystems, Foster City, CA, USA) and analysed using a 3730 DNA Analyzer. Sequence data were analysed with Sequencher version 4.10 (Gene Codes, Ann Arbor, MI, USA) and aligned with Clustal Omega (<http://www.ebi.ac.uk/Tools/msa/clustalo/>). A clone with the consensus sequence was used for further studies.

Table 1
List of primers and morpholinos used in this study. Restriction sites are indicated with lower-case letters, Kozak sequence is shown in bold, and sequence encoding an HA-tag is underlined.

Oligo name	Sequence 5' → 3'	Purpose
CycCD36_FW1	GCTGCTGCTCGCCATC	Cloning of carp <i>cd36a</i>
CycCD36_RV3	GTTTTACTGAAATGTGCTGACATAC	Cloning of carp <i>cd36a</i>
CycCD36_RV2	CCACAGTACGACCACATAATG	Semi-nested PCR for cloning of carp <i>cd36a</i>
CD36_pcDNA3_FW	ACGAggatcc AACATGGC ATACCCATACGATGTTCCAGATTACGCTATGACCTGCTGTGATCTGAAATG	Cloning of carp <i>cd36a</i>
CD36_pcDNA3_RV	CATGctcgagACAATGAAGATTTTTTTTGTATTGC	Cloning of carp <i>cd36a</i>
CD36_qFW1	CGGGACTTTAGCATTTGAC	RT-qPCR of carp <i>cd36a</i>
CD36_qRV1	AGTGAAGTTATCATTGAAGGTG	RT-qPCR of carp <i>cd36a</i>
CD36b_qFW2	GCTGATTGGTCGGATGGAGT	RT-qPCR of carp <i>cd36b</i>
CD36b_qRV2	GAAGTTTGGTATTTTCGGTTTCA	RT-qPCR of carp <i>cd36b</i>
CD36_qFW2	CACCTTCAATGATAACTTCACTT	RT-qPCR of carp <i>cd36a</i>
CD36_qRV2	AGCGTTGGATGAGCAAG	RT-qPCR of carp <i>cd36a</i>
CD36_qFW3	GACAGATGGTTCCTCCTTC	RT-qPCR of carp <i>cd36a</i> and <i>cd36b</i>
CD36_qRV3	TTCCCTTCAGGTCAACAG	RT-qPCR of carp <i>cd36a</i> and <i>cd36b</i>
CD36_qFW4	GGGCTCATCACAGGGAC	RT-qPCR of carp <i>cd36a</i>
CD36_qRV4	TTGGCGTGTCTACAGACG	RT-qPCR of carp <i>cd36a</i>
CD36_qFW5	GACCTGCTGTGATCTGAAATG	RT-qPCR of carp <i>cd36a</i> and <i>cd36b</i>
CD36_qRV5	GTCCCGTTTCCAACACTG	RT-qPCR of carp <i>cd36a</i> and <i>cd36b</i>
CD36_qFW6	GTGAACCATCGCTGAGT	RT-qPCR of carp <i>cd36a</i> and <i>cd36b</i>
CD36_qRV6	GTCAACAGTCCCTCATACTC	RT-qPCR of carp <i>cd36a</i> and <i>cd36b</i>
40S_FW	CCGTGGGTGACATCGTTACA	RT-qPCR of carp 40S ribosomal protein S11
40S_RV	TCAGGACATTGAACCTCACTGTCT	RT-qPCR of carp 40S ribosomal protein S11
zf.cd36_qFW2	CACAGGCAAAGATGACATT	RT-qPCR of zebrafish <i>cd36</i>
zf.cd36_qRV2	CGGGTGGAAAGAGGAAC	RT-qPCR of zebrafish <i>cd36</i>
eif4a1b_FW	TTCAGAACTCAGTACTAGCATAACA	RT-qPCR of zebrafish <i>eif4a1b</i>
eif4a1b_RV	GTGACATCCAACACCTCTGC	RT-qPCR of zebrafish <i>eif4a1b</i>
ppial_FW	ACACTGAAACACGGAGGCAAG	RT-qPCR of zebrafish <i>ppial</i>
ppial_REV	CATCCACAACCTTCCCGAACAC	RT-qPCR of zebrafish <i>ppial</i>
MO1-CD36-e2i2	CTATGAGGCCACAAATATTACCTGT	Zebrafish morpholino knockdown
MO2-CD36-e3i3	TTTGAAATACATACTGCGACAGCAA	Zebrafish morpholino knockdown
MO-FW	TGCTTGGCGAATCCTCATC	RT-PCR to verify zebrafish knockdown
MO-RV	TCGGATAAGCCAATCTGCCAG	RT-PCR to verify zebrafish knockdown

2.8. Bioinformatics

The nucleotide sequence of zebrafish *cd36* is available at Ensembl (ENSDARG00000032639). We obtained the coding sequence of carp *cd36a* after cloning, and the carp *cd36b* coding sequence is predicted based on genome sequence data (Bioproject PRJNA73579) (Henkel et al., 2012). In the present manuscript, to distinguish zebrafish from carp *cd36*, the genes are denoted as *Cyprinus carpio* (*cc*) or *Danio rerio* (*dr*) *cd36* with labelling *a* or *b* for the two carp *cd36*. Exon-intron structure was studied by multiple alignments and open reading frame predictions (FGENESH at <http://linux1.softberry.com/berry.phtml?topic=fgenesht&group=programs&subgroup=gfind>, and GENSCAN at <http://genes.mit.edu/GENSCAN.html>), and exon-intron structure was visualized using WormWeb (<http://www.wormweb.org/exonintron>). Nucleotide sequences were translated into protein sequence using the EMBL-EBI tool (<http://www.ebi.ac.uk/Tools/st/>) and these were aligned with Clustal Omega (<http://www.ebi.ac.uk/Tools/msa/clustalo/>). Molecular weights were calculated with ProtParam (<http://web.expasy.org/protparam/>). Transmembrane regions were predicted using TMHMM Server v. 2.0 (<http://www.cbs.dtu.dk/services/TMHMM/>) and N-glycosylation sites were predicted with NetNGlyc v. 1.0 (<http://www.cbs.dtu.dk/services/NetNGlyc/>). O-glycosylation sites were predicted using NetOGlyc v. 4.0 (<http://www.cbs.dtu.dk/services/NetOGlyc/>). Synteny analysis was performed on the basis of sequence information from Ensembl release 74 (<http://www.ensembl.org/>). The multiple sequence alignment for the phylogenetic tree was made with ClustalX 2.1 (Larkin et al., 2007) and the tree was constructed using the Neighbour Joining method with the number of bootstrap trials set to 10,000, also using ClustalX 2.1. The phylogenetic tree was subsequently visualized with MEGA6 and displayed with the moderately related zebrafish scavenger receptor Scarf1 as the root. Amino acid identity and similarity between Cd36 proteins was calculated

with the software MatGAT 2.0. The transcriptional start site for zebrafish *cd36* was determined by Cap Analysis Gene Expression (CAGE) analysis. MatInspector (<http://www.genomatix.de/>) was used to predict the presence of various vertebrate transcription factor binding sites in the promoter regions of human, mouse, chicken and zebrafish *cd36*.

2.9. *cd36a*-mCherry expression plasmid

The gene encoding the fluorescent protein mCherry was cloned into pcDNA3 using the BamHI and XbaI sites. An XhoI site and a short linker sequence of 15 nucleotides encoding the amino acids GGSGG was placed upstream of the mCherry sequence. Subsequently, the full-length carp *cd36a* sequence was modified from the above-mentioned pGEM-T Easy construct to include sequence to encode an HA-tag at the N-terminus, and to remove the stop codon, using primers CD36_pcDNA3_FW and CD36_pcDNA3_RV (see Table 1). The modified *cccd36a* was then cloned between the BamHI site and the newly created XhoI site in pcDNA3, thus creating a fusion of *cccd36a*-mCherry in pcDNA3. Isolation of highly pure plasmid DNA suitable for transfection was performed with the S.N.A.P. MidiPrep Kit (Invitrogen) according to the manufacturer's instructions.

2.10. Subcellular localization of carp *Cd36a*

Human embryonic kidney 293 (HEK) cells were cultured at 37 °C at 5% CO₂ in DMEM (Gibco, Carlsbad, CA, USA) supplemented with 10% foetal bovine serum (Gibco), 2 mM L-glutamine (Merck, Darmstadt, Germany), 50 U/mL penicillin G (Sigma-Aldrich), and 50 µg/mL streptomycin sulphate (Sigma-Aldrich). HEK cells were seeded directly on untreated glass cover slips, 4.5×10^4 cells each. The following day, cells were transfected with 2 µg *cccd36a*-mCherry-pcDNA3 or mCherry-pcDNA3 using jetPRIME (Polyplus

Transfection, Illkirch, France) according to the manufacturer's instructions at a 1:2 ratio of plasmid:transfection reagent.

The Epithelioma papulosum cyprini (EPC) cell line, which is an adherent cell type derived from fathead minnow, was cultured at 27 °C at 5% CO₂ in RPMI 1640 (Lonza) supplemented with 10% foetal bovine serum, 2 mM L-glutamine, 50 U/mL penicillin G, and 50 µg/mL streptomycin sulphate. EPC cells were seeded on untreated glass cover slips in six-well plates, 1 × 10⁶ cells/well, and allowed to adhere. EPC cells were transfected the following day with 2 µg *cccd36a*-mCherry-pcDNA3 or mCherry-pcDNA3 using FuGENE HD Transfection Reagent (Promega) according to the manufacturer's instructions at a 1:3.5 ratio of plasmid:transfection reagent.

Three days after transfecting HEK and EPC cells with carp *cd36a*-mCherry-pcDNA3 or mCherry-pcDNA3, the cover slips were carefully washed with HBSS, fixed for 20 min at room temperature with 4% paraformaldehyde, washed with HBSS and stained with wheat germ agglutinin-Alexa Fluor 488 conjugate (Molecular Probes, Invitrogen) for 15 min at room temperature, before final washing with HBSS and mounting with Vectashield (Vector Laboratories, Burlingame, CA, USA).

Cells were visualized with a Zeiss LSM-510 (Zeiss, Oberkochen, Germany) confocal laser scanning microscope with a Plan-Apochromat 63×/1.4 oil immersion objective. Green fluorescent signal (Alexa Fluor 488) was excited with a 488 nm argon laser and detected using a band-pass filter (505–550 nm). Red fluorescence (mCherry protein) was excited with a 543 nm helium-neon laser and detected using a long-pass filter (585 nm). Image processing was performed with ImageJ (<http://imagej.nih.gov/ij/>).

2.11. Morpholino knockdown

Morpholino oligonucleotides (Gene Tools, Philomath, OR, USA) were diluted to the desired concentration in 1× Danieau buffer (58 mM NaCl, 0.7 mM KCl, 0.4 mM MgSO₄, 0.6 mM Ca(NO₃)₂, 5.0 mM HEPES; pH 7.6) containing 1% phenol red (Sigma-Aldrich) and 1 nL was injected into the yolk at the 1–2 cell stage using a Femtojet injector (Eppendorf, Hamburg, Germany). For knockdown of *cd36*, two morpholinos were used, one targeting the exon 2–intron 2 splice junction (*cd36* Mo1: 5'CTATGAGGCCACAAATATTACCTGT3', 0.2 mM), and the other targeting the exon 3–intron 3 splice junction (*cd36* Mo2: 5'TTTGAAATACATACTGCGACAGCAA3', 0.6 mM). As a control, the standard control morpholino from Gene Tools was used at the same concentrations as the corresponding *cd36* morpholino. Knockdown of *cd36* with both morpholinos was verified at 1 dpf by RT-PCR with the SuperScript® One-Step RT-PCR System (Invitrogen). Whole embryo RNA isolation was performed as described in Stockhammer et al. (2009) and 50 ng of DNase treated (Invitrogen) RNA template was used for the RT-PCR reaction. RT-PCR primers for *cd36* knockdown verification were MO-FW and MO-RV, whose sequence is given in Table 1. The following adjustments were made to the PCR settings: 56 °C for 30 s for the annealing step of the PCR amplification with 35 cycles. To verify that *cd36* knockdown did not affect leukocyte development an L-plastin immunostaining was performed as described (Mathias et al., 2007).

2.12. *M. marinum* infection

Mycobacterium marinum infections were performed using the wildtype Mma20 strain expressing mCherry in a pSMT3 vector (van der Sar et al., 2004). Embryos were staged at 24 hours post fertilization (hpf) by morphological criteria and manually dechorionated. Bacteria were prepared and injected into the blood circulation at 28 hpf and plated during injections to confirm the number of injected colony forming units (cfu) (Benard et al., 2012). At 4 days post injection (dpi) the uninfected and infected embryos were

Fig. 1. Exon-intron structure of the *cd36* genes of human, zebrafish and carp. The exon-intron structure is shown for human (*hscd36* *Homo sapiens* ENST00000309881), zebrafish (*drcd36* *Danio rerio* ENSDART0000050753), and carp (*cccd36a* *Cyprinus carpio* Scaffold 349; *cccd36b* *Cyprinus carpio* Scaffolds 102854 and 15047). Exons are shown as boxes, introns as lines. Only coding exons are shown. Intron 2 of *cccd36b* is composed of sequence from Scaffolds 102,854 and 15,047, which could not be precisely joined in the current assembly of the carp genome; this is indicated by the dotted line. Scale bar represents 1000 bp.

imaged with a stereo fluorescence microscope, and RNA was isolated for subsequent gene expression analysis. Knockdown with morpholino 1 was repeated four times with 174 morphants and 135 control embryos and the infection phenotype was confirmed with knockdown using morpholino 2 with 35 morphants and 60 control embryos. Fluorescence images were taken with a Leica MZ16FA stereo fluorescence microscope equipped with a DFC420 C digital colour camera. Overlay images of bright field and fluorescence stereomicroscopy were made in Fiji. *M. marinum* pixel count values were calculated using dedicated pixel quantification software (Stoop et al., 2011; Nezhinsky et al., 2012). All data for *M. marinum* pixel count (mean ± SEM) were analysed using unpaired, two-tailed *t*-tests for comparisons between two groups (ns, no significant difference; * *p* < 0.05; ** *p* < 0.01; *** *p* < 0.001).

3. Results

3.1. Identification of cyprinid *cd36* genes

We identified a single *cd36* gene in zebrafish (ENS-DARG0000032639) and two *cd36* genes in common carp (Bioproject PRJNA73579). BLAST analyses in the zebrafish and carp genomes did not identify additional *cd36* genes. Furthermore, the presence of two genes in carp versus a single gene in zebrafish is very common, considering the additional whole genome duplication event that has taken place in carp compared to zebrafish (Henkel et al., 2012).

All three cyprinid *cd36* genes are composed of 12 exons and 11 introns, similar to the coding parts of the human *cd36* gene (Fig. 1). The full-length sequences (*drcd36*, *cccd36a* and *cccd36b*) have open reading frames ranging from 1398 to 1410 bp encoding for proteins of 465–469 aa with predicted molecular weights of 51.6–52.1 kDa. All three cyprinid molecules are predicted to be integral membrane glycoproteins with two transmembrane domains, a large extracellular loop and two short intracellular tails, one at the N- and one at the C-terminus (Fig. 2). The extracellular part is predicted to contain multiple N-glycosylation sites as is the case for mammalian Cd36 (see Fig. 2). Furthermore, there are several conserved cysteines of which six are located in the extracellular part and two in the N-terminal intracellular tail. By similarity to the human Cd36

There are three amino acids in the C-terminus of the human protein which are important for the function, namely Y463, C464, and C466 (human numbering, dark grey shading in Fig. 2). The two cysteines are palmitoylated similar to the cysteines in the N-terminal tail, and the tyrosine is potentially a site of phosphorylation. Both Y463 and C464 are important for phagocytosis of bacteria and for subsequent signalling via TLR2/6, leading to NFκB activation and cytokine production (Stuart et al., 2005). The cyprinid Cd36 molecules lack a tyrosine in the C-terminal tail, and they only have a single

cysteine (C456, cyprinid numbering, dark grey shading in Fig. 2). In conclusion, cyprinid Cd36 molecules are integral membrane glycoproteins with two transmembrane domains, a large extracellular loop and two short intracellular tails, similar to mammalian Cd36.

3.2. Phylogenetic analysis

An amino acid sequence alignment indicates a high degree of similarity, not only among the cyprinid Cd36 proteins (88.7–93%) but also between cyprinid and other fish species, chicken, xenopus and mammalian Cd36 proteins (all >70%; Table 2). Also percentages of identity are very high with human to fish amino acid identities ranging from 50.8 to 52.3%. This indicates that the molecule has been well conserved during evolution. The amphioxus Cd36 protein stands out as being most different from all the other proteins with identities around 35% only.

Phylogenetic analysis was performed on amino acid sequences of multiple Cd36 and Scarb molecules (Fig. 3). A moderately related scavenger receptor, zebrafish Scarf1, was used as an outgroup to root the phylogenetic tree. The overall topology of the phylogenetic tree shows that the Cd36, Scarb1 and Scarb2 molecules form separate clusters. In this study we have identified a putative Cd36 of Atlantic salmon, a sequence we obtained by performing a BLAST search in salmon whole genome shotgun sequences using the zebrafish Cd36 sequence. This putative Cd36 molecule conforms with the phylogenetic tree contrary to the rainbow trout Cd36, whose position in the tree suggests it should be renamed into Scarb2. The ancestral amphioxus Cd36 seems to be closer related to the Scarb1 family. Amongst the remaining Cd36 molecules, the tetrapods (mammalian, bird and amphibian molecules) form one cluster and all the fish molecules branch out in a separate cluster. Zebrafish Cd36 and the two carp Cd36 molecules cluster together as expected from the high identities found between these. Bootstrap values are high in the entire tree with only a single exception, namely the split between stickleback and platyfish/medaka Cd36 molecules. Overall, the phylogenetic tree highlights the close relationship of all the teleost Cd36 proteins, visibly separate from the teleost Scarb1 and Scarb2 molecules and from the tetrapod proteins.

3.3. Synteny analysis

Conservation of synteny was investigated by comparing the genomic regions immediately up- and down-stream of human, mouse, chicken and xenopus *cd36* with the genomic regions up- and down-stream of *cd36* in a number of fish species with an annotated genome, including zebrafish and common carp (Fig. 4). Human, mouse and chicken genomes contain a block of five genes: *magi2*, *gnai1*, *gmat3*, *cd36*, and *sema3c*. This block is fairly conserved in the other species with the exception of xenopus and fish genomes lacking the *gmat3* gene. For carp, although the scaffolds are limited in length, the analysis does confirm conservation of synteny with *sema3c* downstream of *cd36a* and *cd36b*, and a partial *gnai1* sequence upstream of *cd36a*. Overall, analysis of the genes adjacent to *cd36* in cyprinid fish indicates conserved synteny between teleost *cd36* and non-teleost *cd36*.

3.4. Promoter analysis

In addition to synteny analysis we also studied in detail the transcription factor binding sites in the promoter region of zebrafish *cd36* and compared this to human, mouse and chicken (Supplementary Fig. 1). The transcriptional start site has not been determined for either of the carp *cd36* genes, so promoter analysis was not performed for these. The putative transcriptional start site for zebrafish

Fig. 3. Phylogenetic tree of Cd36 and Scarb sequences. Cd36, Scarb1 and Scarb2 molecules from various species were included in the phylogenetic analysis. A moderately related scavenger receptor (zebrafish Scarf1) was used as an outgroup. Carp Cd36a and Cd36b, as well as zebrafish Cd36 are indicated in bold. Amino acid sequences used: Zebrafish Scarf1 (*Danio rerio* ENSDARP00000113737), rainbow trout Cd36/Scarb2 (*Oncorhynchus mykiss* AY606034), tetraodon Scarb2 (*Tetraodon nigroviridis* ENSTNIP00000018658), zebrafish Scarb2 #1 (*Danio rerio* ENSDARP00000110051), stickleback Scarb2 (*Gasterosteus aculeatus* ENSGACP00000018249), zebrafish Scarb2 #2 (*Danio rerio* ENSDARP00000076810), zebrafish Scarb2 #3 (*Danio rerio* ENSDARP00000110098), salmon SR-BI (*Salmo salar* DQ914655), tetraodon Scarb1 (*Tetraodon nigroviridis* ENSTNIP00000016587), zebrafish Scarb1 (*Danio rerio* ENSDARP00000106590), amphioxus Cd36 (*Branchiostoma floridae* JG1126670), human CD36 (*Homo sapiens* ENSP00000399421), mouse CD36 (*Mus musculus* ENSMUSP00000080974), rat Cd36 (*Rattus norvegicus* ENSRNOP00000058398), chicken Cd36 (*Gallus gallus* ENSGALP00000038958), xenopus Cd36 (*Xenopus tropicalis* ENSXETP00000017231), carp Cd36a (*Cyprinus carpio* KM030422), carp Cd36b (*Cyprinus carpio* BioProject PRJNA73579), zebrafish Cd36 (*Danio rerio* ENSDARP00000050752), spotted gar Cd36 (*Lepisosteus oculatus* ENSLOCP00000019733), salmon Cd36 (*Salmo salar*, predicted from AGKD01002586.1), cod Cd36 (*Gadus morhua* ENSG-MOP00000012522), tetraodon Cd36 (*Tetraodon nigroviridis* ENSTNIP00000014345), fugu Cd36 (*Takifugu rubripes* ENSTRUP000000041077), tilapia Cd36 (*Oreochromis niloticus* ENSONIP00000000108), platyfish Cd36 (*Xiphophorus maculatus* ENSXMAP00000004793), medaka Cd36 (*Oryzias latipes* ENSORLP00000020106), stickleback Cd36 (*Gasterosteus aculeatus* ENSGACP00000026370).

cd36 has been experimentally verified by CAGE analysis (personal communication, H.P. Spaik), while human, mouse and chicken transcriptional start sites and promoters are already annotated in MatInspector. The promoter analyses suggest regulation of *cd36* transcription by the common transcription factors Pu.1 (part of the ETS family of transcription factors) (Zakrzewska et al., 2010), p53 (Lane and Levine, 2010), NF-κB (Oeckinghaus and Ghosh, 2009), and

Table 2
Percentage identity and similarity between Cd36 of various species. Identities (upper right) and similarities (lower left) were based on the following amino acid sequences: human (*Homo sapiens* ENSP00000399421), mouse (*Mus musculus* ENSMUSP00000080974), chicken (*Gallus gallus* ENSGALP00000038958), xenopus (*Xenopus tropicalis* ENSXETP00000017231), fugu (*Takifugu rubripes* ENSTRUP00000041077), stickleback (*Gasterosteus aculeatus* ENSGACP00000026370), spotted gar (*Lepisosteus oculatus* ENSLOCP00000019733), zebrafish (*Danio rerio* ENSDARP00000050752), carp a (*Cyprinus carpio* KM030422), carp b (*Cyprinus carpio* Bioproject PRJNA73579), and amphioxus (*Branchiostoma floridae* JGI126670).

	Human	Mouse	Chicken	Xenopus	Fugu	Stickleback	Spotted gar	Zebrafish	Carp a	Carp b	Amphioxus
Human		84.1	61.4	54.9	52.3	51.1	51.6	51.5	51.2	50.8	35.6
Mouse	94.1		62.9	55.5	53.9	51.7	52.3	52.5	52.6	52.7	34.8
Chicken	81.8	81.4		56.2	49.8	50.9	51.7	52.9	51.5	50.1	36.9
Xenopus	75.4	74.6	76		50.8	52.2	51.4	51.2	49.3	47.1	36.8
Fugu	72.8	72.6	72.8	71.6		65.4	57.5	56.5	54.5	56.1	34.2
Stickleback	73.5	72.5	71.5	72.3	80.2		59.3	58.8	58.3	57.7	35.3
Spotted gar	74	73.4	73.2	70.9	74.2	76.9		56.3	56.1	57.5	37.2
Zebrafish	73.1	71.6	74.3	70.2	74.9	77.4	73.4		77.4	76.1	35.6
Carp a	72.9	72.2	72	70.9	73.5	75.5	74	88.7		87	37.3
Carp b	73.1	72.5	73.2	70.9	75.2	76.5	75.5	88.7	93		36.1
Amphioxus	59.2	57.7	60	59.6	58.5	58.8	59.4	59	60.8	60.8

Fig. 4. Comparison of genomic regions in the vicinity of *cd36*. Synteny analysis comparing the genomic organisation of *cd36* in various species. Five genes are included in this figure, each represented by a unique colour/shaded box. The arrow indicates the direction of transcription. Intergenic regions are shown as a black line. Carp *cd36b* is split over two scaffolds. The size of each depicted genomic area is indicated to the right, in megabasepairs (Mbp).

Stat1 (Reich, 2013). Although the exact locations of the transcription factor binding sites are not necessarily conserved and their functionality has not been experimentally verified, the presence of binding sites for all four transcription factors likely influences *cd36* gene expression levels in various tissues and cell types, including those with an immune function, and thus could be of importance for Cd36 function.

3.5. Gene expression

Constitutive gene expression of *cccd36a* and *cccd36b* was measured in organs of four healthy carp and normalized to the expression of the housekeeping gene. The results from 15 organs are shown in Fig. 5A. The gene expression of *cccd36a* is generally higher than that of *cccd36b*. Highest expression for both genes

Fig. 5. Constitutive gene expression of *cd36*. (A) Carp organs and (B) Carp cell types. The figure shows gene expression of *cccd36a* (black bars) and *cccd36b* (white bars) in various tissues and cell types, including neuronal, reproductive and immune organs, normalized to the expression of the 40S ribosomal protein *S11* housekeeping gene. For comparison, head kidney and PBL are shown next to the carp cell types in (B). Note the different scale on the y-axis in (A) and (B). The bars indicate average gene expression of $n = 4$ adult fish and standard deviations are indicated. PBL = peripheral blood leukocytes. (C) Expression of *dracd36* in macrophages (*mpeg1*-positive cells) and neutrophils (*mpx*-positive cells) from zebrafish larvae relative to *mpeg1*- and *mpx*-negative cells, respectively. (*mpeg1* and *mpx* gene expression data from Benard et al., 2014a). Data represent $n = 3$ biological replicates and are normalized to the *eif4* housekeeping gene.

is found in brain tissue (0.10 and 0.04 times the housekeeping gene, respectively), with the pituitary gland contributing slightly to the total brain gene expression. High-to-moderate gene expression was also detected in the reproductive organs, ovary (0.05 and 0.002) and testis (0.02 and 0.004), followed by low-to-moderate expression (≤ 0.005) levels in heart and liver. The remaining organs, including typical immune organs such as head kidney and spleen, have a very low to undetectable *cd36* gene expression level. The relatively high gene expression in brain, ovary and testis, and low to non-detectable expression in other tissues including the immune organs tested was confirmed by transcriptome analysis of the same organs from carp (data not shown). In addition, several primer pairs were used for RT-qPCR yielding similar results. The sequences of these primers are listed in Table 1, and their location is shown in Supplementary Fig. 2. Furthermore, *cd36* gene expression was undetectable by RT-qPCR using as template various immune cell types from carp purified by magnetically activated cell sorting (MACS), i.e. thrombocytes, thymocytes, neutrophils, B cells, and macrophages (Fig. 5B).

Complementary to the analysis of multiple organs and cell types of adult carp, we took advantage of the zebrafish model to investigate *cd36* expression in leukocytes. To this end we used transgenic

zebrafish lines with different fluorescently marked leukocyte populations, dissociated larvae at 6 dpf and isolated the fluorescent and non-fluorescent cell fractions by fluorescence activated cell sorting (FACS). RT-qPCR analysis of FACS isolated macrophages (*mpeg1*-promoter driven fluorescence) and neutrophils (*mpx*-promoter driven fluorescence) from zebrafish 6 dpf larvae revealed a very low *cd36* gene expression compared to the levels of macrophage- or neutrophil-specific markers (*mpeg1* and *mpx*, respectively), as shown in Fig. 5C.

3.6. Subcellular localization

In order to investigate the subcellular localization of cyprinid Cd36 we first cloned and sequenced the full-length *cccd36a* coding region. We then transfected the HEK cell line with a construct expressing carp Cd36a fused to the fluorescent protein mCherry. The cell membrane was visualized by staining with wheat germ agglutinin-Alexa 488 conjugate which exhibits green fluorescence. Confocal microscopy showed the presence of Cd36a-mCherry at the rim of the cells, co-localizing with the green fluorescent surface stain (Fig. 6A). In contrast, the fluorescent protein mCherry alone does not localize to the cell membrane but is evenly

Fig. 6. Subcellular localization of Cd36. (A) HEK cells stained with wheat germ agglutinin-Alexa Fluor 488 (green, left) and expressing carp Cd36a-mCherry (red, centre). An overlay is shown to the right. (B) HEK cells stained with wheat germ agglutinin-Alexa Fluor 488 (green, left) and expressing mCherry (red, centre). An overlay is shown to the right. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

distributed throughout the cell (Fig. 6B), meaning the Cd36 molecule is responsible for the redirection of the fluorescent protein to the cell membrane. Similar results were obtained with EPC cells transfected with the same Cd36 construct (data not shown). The surface localization of carp Cd36a is in agreement with the localization of the mammalian Cd36 molecule as well as the predicted structure of the carp molecule with two transmembrane regions, an extracellular part, and two cytoplasmic tails.

3.7. Role of Cd36 in *M. marinum* infection

To further study the function of teleost Cd36, we took advantage of the rapid gene knockdown technology available for zebrafish (Sumanas and Larson, 2002). We used antisense morpholino oligonucleotides to knockdown *cd36* in zebrafish embryos, which were subsequently subjected to infection with mCherry-labelled *M. marinum*. The bacterial fluorescence per embryo was quantified at 4 dpi by image analysis. At this stage of *M. marinum* infection, the embryos have developed granuloma-like structures consisting of infected and uninfected leukocytes and the bacterial fluorescence within these granulomas can be detected with stereo fluorescence microscopy (Benard et al., 2012). Two different splice blocking morpholinos were used targeting different exon-intron splice junctions of the *cd36* gene. The knockdown effect was verified by RT-PCR (Supplementary Fig. 3) and embryos were stained with antibody against the pan-leukocytic marker L-plastin, showing that *cd36* knockdown did not affect leukocyte development in duct of Cuvier or in the tail region (Supplementary Fig. 4). Both *cd36* morphant groups showed a significantly increased bacterial fluorescence compared to embryos injected with a control morpholino (quantification shown in Fig. 7A, and representative pictures shown in Fig. 7B). We further analysed gene expression of

cd36 in *M. marinum*-infected zebrafish embryos at 4 dpi (Fig. 7C). Compared to uninfected fish, *cd36* expression was down-regulated approximately 3-fold in infected individuals.

4. Discussion

Given our interest in the function of TLRs and the evidence, at least in mammals, of CD36 interaction with preformed TLR2-TLR6 heterodimers, we aimed to characterize the function of Cd36 in zebrafish and common carp, two closely related cyprinid fish species. Cyprinid Cd36 molecules are integral membrane glycoproteins with two transmembrane domains, a large extracellular loop and two short intracellular tails and are similar to mammalian Cd36 at the sequence level. Similar to mammalian Cd36 proteins the disulphide bonds are conserved in the cyprinid proteins, and so is the presence of multiple *N*-glycosylation sites. At present, immune functions for the scavenger receptor class B family member Cd36 have not been described for teleost fish.

Although the cyprinid Cd36 sequences seem relatively similar to mammalian Cd36, there also appear to be a number of intriguing differences. In contrast to the human protein that has no *O*-glycosylation sites, there is one *O*-glycosylation site predicted in drCd36, three sites in ccCd36a, and one site in ccCd36b. The significance of these potential *O*-glycosylation sites in the fish Cd36 proteins is not known. Most notably, there is a clear difference between the mammalian and fish Cd36 sequences in the C-terminal tail which, at least in the human protein, contains two key amino acids, namely Y463 and C464. The cysteine is palmitoylated and plays a role in the localization of Cd36 to lipid raft domains in the plasma membrane (Tao et al., 1996), whereas the tyrosine is a potential site of phosphorylation and could provide a docking site for SH2 domain-containing proteins such as the Src family

Fig. 7. Interplay between *cd36* and *Mycobacterium marinum* during infection of zebrafish embryos. *cd36* knockdown impairs the control of *M. marinum* infection. AB/TL embryos were injected with two different splice blocking morpholinos against *cd36*, subsequently injected with mCherry-expressing *M. marinum* at 28 hpf and imaged at 4 dpi. (A) Bacterial burden was quantified by determining the number of fluorescent bacterial pixels per embryo with dedicated software and is represented as a percentage of the average control pixel count. (B) Representative stereo fluorescence images of an infected embryo per group are depicted with their bright field channel shown as inset. (C) Gene expression of *cd36* is decreased during *M. marinum* infection of zebrafish embryos. Zebrafish embryos were infected as above and at 4 dpi gene expression was analysed for *cd36* and normalized to the *ppia* housekeeping gene. Data represent the average and standard deviation obtained from $n = 3$ biological replicates.

of tyrosine kinases and the actin regulator p130Cas (Stuart et al., 2005). Src kinases that reside in lipid rafts have previously been shown to initiate downstream signalling cascades upon interaction with Cd36, and the actin regulator p130Cas is presumably involved in Cd36-dependent cytoskeletal rearrangements that are required for phagocytosis (Stuart et al., 2007). Upon examination of the C-terminal tail of cyprinid Cd36 molecules it is clear that no tyrosine is present but a single cysteine is found. Possibly, the presence of the cysteine could suggest that cyprinid Cd36 may localize to lipid rafts in the plasma membrane, but the absence of the tyrosine could have a determining effect on signalling and phagocytosis.

In our study we could demonstrate the presence of Cd36 on the plasma membrane by confocal microscopy. We could not clearly demonstrate, however, a role for Cd36 in phagocytosis; morpholino knockdown of zebrafish *cd36* did not affect phagocytosis of *M. marinum* (Benard et al., 2014b) and transfection of cell lines with carp

cd36 did not confer the ability to phagocytose *S. aureus* or zymosan particles (unpublished data, I. R. Fink). This suggests that differences in key amino acids in the C-terminal tail of cyprinid Cd36 may affect function of fish Cd36 as compared to function of mammalian Cd36.

We further used cell lines transfected with carp *cd36* to study its putative role as a pattern recognition receptor, by stimulating the cells with various ligands and measuring downstream transcription factor activation. Since we could not see any effect of *cd36* in these studies, this could indicate that *cd36* might primarily function as a co-receptor and as such would always require the presence of another receptor to fulfil its role.

High gene expression levels of cyprinid *cd36* were found in brain, ovary and testis. In contrast, gene expression was undetectable in immune organs and undetectable in all leukocyte subtypes tested, including monocytes/macrophages of adult carp.

Also macrophages and neutrophils from zebrafish larvae had very low levels of *cd36* expression. In mammals, *cd36* is expressed in many different tissues and cell types, e.g. skeletal and smooth muscle, intestinal epithelia, adipocytes, endothelial cells, platelets, dendritic cells and monocytes/macrophages (Febbraio et al., 2001). We refute the option that the surprising absence of *cd36* gene expression in immune organs and cell types from (cyprinid) fish could be due to the design of our primers; design of several additional primer pairs located in different parts of *cd36* did not provide any indication of *cd36* expression in immune organs and immune cell types. We consider it unlikely that all the primer combinations we tested for carp *cd36* would have failed to amplify particular truncated splice variants uniquely expressed in immune organs and immune cell types. Although our primers would not detect the possible existence of tissue-specific splice variants arising from alternative promoter usage (Andersen et al., 2006), CAGE analysis indicates the presence of only a single transcription start site for zebrafish *cd36*, which suggests tissue-specific splice variants of *cd36* may not be commonly seen in fish. Furthermore, transcriptome analyses of a multitude of carp organs revealed an expression pattern for *cd36* which confirms our findings (unpublished data). In fact, we have found no indication for alternative splicing of cyprinid *cd36*. Possibly, differences in transcriptional regulation of *cd36* and/or absence of splice variants in cyprinid *cd36* molecules might help explain differences in function between mammalian and fish Cd36. Of interest, particular mutations in the human *cd36* gene can give rise to aberrant expression patterns and phenotypes. One example is a rather commonly occurring mutation (C268T), which leads to lack of *cd36* expression in monocytes and platelets, a condition called type I deficiency (Rac et al., 2007). Although this particular nucleotide is not mutated in the cyprinid genes and thus does not explain the observed absence of *cd36* gene expression in these cell types in fish, clearly mammalian CD36 is prone to mutations that affect expression in monocytes and platelets, a phenotype comparable to what we observe in the teleosts we studied. In conclusion, the complete absence of *cd36* gene expression in immune organs and leukocyte cell types of cyprinid fish suggests that maybe not all prototypical immune functions ascribed to the function of CD36 as a PRR important in innate immunity would be present in (cyprinid) fish. Clearly, the exact function of fish Cd36 remains a subject of investigation.

Presence of high levels of cyprinid *cd36* gene expression particularly in brain, testis and ovary, tissues with high lipid and/or cholesterol uptake for e.g. conversion to neurosteroid and sex hormones, could be indicative of a function for Cd36 mediating cholesterol transfer, a normal characteristic of the Scarb1 protein, in particular. Previous studies in salmon and rainbow trout (Trattner et al., 2008; Torstensen et al., 2011; Sanchez-Gurmaches et al., 2012, and others) on Cd36 have pointed to lipid metabolism as a function conserved between mammalian and teleost Cd36. Although it may well be true that fish members of the class B family of scavenger receptors, including Cd36 and Scarb molecules have overlapping functions, our phylogenetic analysis shows that the rainbow trout Cd36 sequence is more likely a Scarb2 homolog, which would suggest these data have to be interpreted with care. Although we identified a putative Cd36 sequence in salmon that also exhibits conserved synteny and does cluster with other fish Cd36 sequences in the phylogenetic tree, we could not easily identify a *cd36* sequence in rainbow trout. In fact all candidates in rainbow trout are more closely related to Scarb2 molecules than to Cd36. Several *scarb1* and *scarb2* paralogs are present in both the zebrafish and carp genome and interestingly, in contrast to *cd36*, transcriptome data reveal moderate to high expression of both *scarb1* and *scarb2* genes in macrophages (unpublished data, I. R. Fink). This finding reinforces the possibility that the division of functions between members of the class B family of

scavenger receptors could be different between mammals and fish.

The ancestral *cd36* gene in amphioxus could be up-regulated by *in vivo* challenge with *E. coli*, and by *in vitro* LTA stimulation of a flounder cell line transfected with an amphioxus *cd36*-pcDNA3 construct (Zhang et al., 2013). However, our *in vivo* study showed that *cd36* gene expression in *M. marinum*-infected zebrafish embryos was down-regulated, which is in line with published data on infected adult zebrafish (van der Sar et al., 2009).

Although a null mutant would be required to confirm our results, knockdown of *cd36* in zebrafish embryos did not affect general development of leukocytes or the migration of neutrophils or macrophages towards locally induced inflammation, nor phagocytosis of *M. marinum* but did result in higher bacterial load following infection with *M. marinum*, suggesting a possible but presently undefined role for fish Cd36 in the control of bacterial infection. Given the suggestion that the division of functions between members of the class B family of scavenger receptors could be different between mammals and fish, future studies could address immune functions of the different class B family members found in teleost fish and explore further their functional relationships.

Acknowledgements

We thank Fons Verbeek and Alexander Nezhdinsky (Leiden University, The Netherlands) for making an improved version of the pixel quantification software available prior to publication. The research leading to these results has received funding from the European Commission under the 7th Framework Programme for Research and Technological Development (FP7) of the European Union (grant agreement NEMO PITN-GA-2008-214505) to IRF and GFW. ELB and AHM were supported by the Smart Mix Program of the Netherlands Ministry of Economic Affairs and the Ministry of Education, Culture and Science.

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.molimm.2014.09.010>.

References

- Abumrad, N.A., El-Maghrabi, M.R., Amri, E.Z., Lopez, E., Grimaldi, P.A., 1993. Cloning of a rat adipocyte membrane protein implicated in binding or transport of long-chain fatty acids that is induced during preadipocyte differentiation. Homology with human CD36. *J. Biol. Chem.* 268, 17665–17668.
- Akira, S., Takeda, K., 2004. Toll-like receptor signalling. *Nat. Rev. Immunol.* 4, 499–511.
- Andersen, M., Lenhard, B., Whatling, C., Eriksson, P., Odeberg, J., 2006. Alternative promoter usage of the membrane glycoprotein CD36. *BMC Mol. Biol.* 7, 8.
- Benard, E.L., Racz, P.I., Rougeot, J., Nezhdinsky, A.E., Verbeek, F.J., Spaink, H.P., Meijer, A.H., 2014a. Macrophage-expressed perforins Mpeg1 and Mpeg1.2 have an anti-bacterial function in zebrafish. *J. Innate. Immun.*, <http://dx.doi.org/10.1159/000366103>.
- Benard, E.L., Roobol, S.J., Spaink, H.P., Meijer, A.H., 2014b. Phagocytosis of mycobacteria by zebrafish macrophages is dependent on the scavenger receptor Marco, a key control factor of pro-inflammatory signalling. *Dev. Comp. Immunol.* 47, 223–233.
- Benard, E.L., van der Sar, A.M., Ellett, F., Lieschke, G.J., Spaink, H.P., Meijer, A.H., 2012. Infection of zebrafish embryos with intracellular bacterial pathogens. *J. Visualized Exp.: JoVE* 61, e3781.
- Bernut, A., Herrmann, J.L., Kissa, K., Dubremetz, J.F., Gaillard, J.L., Lutfalla, G., Kremer, L., 2014. *Mycobacterium abscessus* cording prevents phagocytosis and promotes abscess formation. *Proc. Natl. Acad. Sci. U.S.A.* 111, E943–E952.
- Brundert, M., Ewert, A., Heeren, J., Behrendt, B., Ramakrishnan, R., Greten, H., Merkel, M., Rinninger, F., 2005. Scavenger receptor class B type I mediates the selective uptake of high-density lipoprotein-associated cholesteryl ester by the liver in mice. *Arterioscler. Thromb. Vasc. Biol.* 25, 143–148.
- Calvo, D., Dopazo, J., Vega, M.A., 1995. The CD36, CLA-1 (CD36L1), and LIMPII (CD36L2) gene family: cellular distribution, chromosomal location, and genetic evolution. *Genomics* 25, 100–106.

- Calvo, D., Vega, M.A., 1993. Identification, primary structure, and distribution of CLA-1, a novel member of the CD36/LIMPII gene family. *J. Biol. Chem.* 268, 18929–18935.
- Chadwick, A.C., Sahoo, D., 2013. Functional genomics of the human high-density lipoprotein receptor scavenger receptor BI: an old dog with new tricks. *Curr. Opin. Endocrinol. Diabetes Obes.* 20, 124–131.
- Covassin, L., Amigo, J.D., Suzuki, K., Teplyuk, V., Straubhaar, J., Lawson, N.D., 2006. Global analysis of hematopoietic and vascular endothelial gene expression by tissue specific microarray profiling in zebrafish. *Dev. Biol.* 299, 551–562.
- Endemann, G., Stanton, L.W., Madden, K.S., Bryant, C.M., White, R.T., Protter, A.A., 1993. CD36 is a receptor for oxidized low density lipoprotein. *J. Biol. Chem.* 268, 11811–11816.
- Febbraio, M., Hajjar, D.P., Silverstein, R.L., 2001. CD36: a class B scavenger receptor involved in angiogenesis, atherosclerosis, inflammation, and lipid metabolism. *J. Clin. Invest.* 108, 785–791.
- Forlenza, M., Scharfack, J.P., Kachamakova, N.M., Taverne-Thiele, A.J., Rombout, J.H., Wiegertjes, G.F., 2008. Differential contribution of neutrophilic granulocytes and macrophages to nitrosative stress in a host-parasite animal model. *Mol. Immunol.* 45, 3178–3189.
- Gonzalez, A., Valeiras, M., Sidransky, E., Tayebi, N., 2014. Lysosomal integral membrane protein-2: a new player in lysosome-related pathology. *Mol. Genet. Metab.* 111, 84–91.
- Greenwalt, D.E., Watt, K.W., Hasler, T., Howard, R.J., Patel, S., 1990. Structural, functional, and antigenic differences between bovine heart endothelial CD36 and human platelet CD36. *J. Biol. Chem.* 265, 16296–16299.
- Henkel, C.V., Dirks, R.P., Jansen, H.J., Forlenza, M., Wiegertjes, G.F., Howe, K., van den Thillart, G.E., Spaik, H.P., 2012. Comparison of the exomes of common carp (*Cyprinus carpio*) and zebrafish (*Danio rerio*). *Zebrafish* 9, 59–67.
- Hoebe, K., Georgel, P., Rutschmann, S., Du, X., Mudd, S., Crozat, K., Sovath, S., Shamel, L., Hartung, T., Zähringer, U., Beutler, B., 2005. CD36 is a sensor of diacylglycerides. *Nature* 433, 523–527.
- Imnazarow, I., 1995. Genetic variability of Polish and Hungarian carp lines. *Aquaculture* 129, 215–219.
- Koumans-van Diepen, J.C.E., Egberts, E., Peixoto, B.R., Taverne, N., Rombout, J.H.W.M., 1995. B cell and immunoglobulin heterogeneity in carp (*Cyprinus carpio* L.): an immuno(cyto)chemical study. *Dev. Comp. Immunol.* 19, 97–108.
- Lane, D., Levine, A., 2010. p53 Research: the past thirty years and the next thirty years. *Cold Spring Harbor Perspect. Biol.* 2, a000893.
- Larkin, M.A., Blackshields, G., Brown, N.P., Chenna, R., McGettigan, P.A., McWilliam, H., Valentin, F., Wallace, I.M., Wilm, A., Lopez, R., Thompson, J.D., Gibson, T.J., Higgins, D.G., 2007. Clustal W and Clustal X version 2.0. *Bioinformatics* 23, 2947–2948.
- Lee, C.C., Avalos, A.M., Ploegh, H.L., 2012. Accessory molecules for Toll-like receptors and their function. *Nat. Rev. Immunol.* 12, 168–179.
- Mathias, J.R., Dodd, M.E., Walters, K.B., Rhodes, J., Kanki, J.P., Look, A.T., Huttenlocher, A., 2007. Live imaging of chronic inflammation caused by mutation of zebrafish Hai1. *J. Cell Sci.* 120, 3372–3383.
- Means, T.K., Mylonakis, E., Tampakakis, E., Colvin, R.A., Seung, E., Puckett, L., Tai, M.F., Stewart, C.R., Pukkila-Worley, R., Hickman, S.E., Moore, K.J., Calderwood, S.B., Hachohen, N., Luster, A.D., El Khoury, J., 2009. Evolutionarily conserved recognition and innate immunity to fungal pathogens by the scavenger receptors SCARF1 and CD36. *J. Exp. Med.* 206, 637–653.
- Nezhinsky, A.E., Stoop, E., van der Sar, A., Verbeek, F.J., 2012. Numerical Analysis of Image Based High Throughput Zebrafish Infection Screens-Matching Meaning with Data. pp. 257–262.
- Oeckinghaus, A., Ghosh, S., 2009. The NF- κ B family of transcription factors and its regulation. *Cold Spring Harbor Perspect. Biol.* 1, a000034.
- Pietretti, D., Spaik, H.P., Falco, A., Forlenza, M., Wiegertjes, G.F., 2013. Accessory molecules for Toll-like receptors in teleost fish. Identification of TLR4 interactor with leucine-rich repeats (TRIL). *Mol. Immunol.* 56, 745–756.
- Pietretti, D., Wiegertjes, G.F., 2014. Ligand specificities of Toll-like receptors in fish: indications from infection studies. *Dev. Comp. Immunol.* 43, 205–222.
- Plüddemann, A., Neyer, C., Gordon, S., 2007. Macrophage scavenger receptors and host-derived ligands. *Methods* 43, 207–217.
- Rac, M.E., Safranow, K., Poncyljusz, W., 2007. Molecular basis of human CD36 gene mutations. *Mol. Med.* 13, 288–296.
- Reich, N.C., 2013. STATs get their move on. *Jak-Stat* 2, e27080.
- Renshaw, S.A., Loynes, C.A., Trushell, D.M.I., Elworthy, S., Ingham, P.W., Whyte, M.K.B., 2006. A transgenic zebrafish model of neutrophilic inflammation. *Blood* 108, 3976–3978.
- Ribeiro, C.M., Hermens, T., Taverne-Thiele, A.J., Savelkoul, H.F., Wiegertjes, G.F., 2010. Evolution of recognition of ligands from Gram-positive bacteria: similarities and differences in the TLR2-mediated response between mammalian vertebrates and teleost fish. *J. Immunol.* 184, 2355–2368.
- Romano, N., Picchietti, S., Taverne-Thiele, J.J., Taverne, N., Abelli, L., Mastrolia, L., Verburg-van Kemenade, B.M.L., Rombout, J.H.W.M., 1998. Distribution of macrophages during fish development: an immunohistochemical study in carp (*Cyprinus carpio*, L.). *Anat. Embryol.* 198, 31–41.
- Rombout, J.H.W.M., Koumans-van Diepen, J.C.E., Emmer, P.M., Taverne-Thiele, J.J., Taverne, N., 1996. Characterization of carp thrombocytes with specific monoclonal antibodies. *J. Fish Biol.* 49, 521–531.
- Sanchez-Gurmaches, J., Cruz-Garcia, L., Gutierrez, J., Navarro, I., 2012. mRNA expression of fatty acid transporters in rainbow trout: *in vivo* and *in vitro* regulation by insulin, fasting and inflammation and infection mediators. *Comp. Biochem. Physiol. A: Mol. Integr. Physiol.* 163, 177–188.
- Sanchez-Gurmaches, J., Østbye, T.-K., Navarro, I., Torgersen, J., Hevrøy, E.M., Ruyter, B., Torstensen, B.E., 2011. *In vivo* and *in vitro* insulin and fasting control of the transmembrane fatty acid transport proteins in Atlantic salmon (*Salmo salar*). *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 301, R947–R957.
- Secombes, C.J., van Groningen, J.J.M., Egberts, E., 1983. Separation of lymphocyte subpopulations in carp *Cyprinus carpio* L. by monoclonal antibodies: immunohistochemical studies. *Immunol.* 48, 165–175.
- Shu, G., Liao, W.Y., Feng, J.Y., Yu, K.F., Zhai, Y.F., Wang, S.B., Khondowe, P., Wang, X.Q., Jiang, Q.Y., 2011. Active immunization of fatty acid translocase specifically decreased visceral fat deposition in male broilers. *Poult. Sci.* 90, 2557–2564.
- Silverstein, R.L., Febbraio, M., 2009. CD36, a scavenger receptor involved in immunity, metabolism, angiogenesis, and behavior. *Sci. Signal.* 2, re3.
- Stockhammer, O.W., Zakrzewska, A., Hegedus, Z., Spaik, H.P., Meijer, A.H., 2009. Transcriptome profiling and functional analyses of the zebrafish embryonic innate immune response to *Salmonella* infection. *J. Immunol.* 182, 5641–5653.
- Stolte, E.H., Savelkoul, H.F., Wiegertjes, G., Flik, K., Lidy Verburg-van Kemenade, B.M., 2008. Differential expression of two interferon- γ genes in common carp (*Cyprinus carpio* L.). *Dev. Comp. Immunol.* 32, 1467–1481.
- Stoop, E.J.M., Schipper, T., Rosendahl Huber, S.K., Nezhinsky, A.E., Verbeek, F.J., Gurucha, S.S., Besra, G.S., Vandenbroucke-Grauls, C.M.J.E., Bitter, W., van der Sar, A.M., 2011. Zebrafish embryo screen for mycobacterial genes involved in the initiation of granuloma formation reveals a newly identified ESX-1 component. *Dis. Models Mech.* 4, 526–536.
- Stuart, L.M., Bell, S.A., Stewart, C.R., Silver, J.M., Richard, J., Goss, J.L., Tseng, A.A., Zhang, A., El Khoury, J.B., Moore, K.J., 2007. CD36 signals to the actin cytoskeleton and regulates microglial migration via a p130Cas complex. *J. Biol. Chem.* 282, 27392–27401.
- Stuart, L.M., Deng, J., Silver, J.M., Takahashi, K., Tseng, A.A., Hennessy, E.J., Ezekowitz, R.A., Moore, K.J., 2005. Response to *Staphylococcus aureus* requires CD36-mediated phagocytosis triggered by the COOH-terminal cytoplasmic domain. *J. Cell Biol.* 170, 477–485.
- Sumanas, S., Larson, J.D., 2002. Morpholino phosphorodiamidate oligonucleotides in zebrafish: a recipe for functional genomics? *Brief Funct. Genomic Proteomic* 1, 239–256.
- Tao, N., Wagner, S.J., Lublin, D.M., 1996. CD36 is palmitoylated on both N- and C-terminal cytoplasmic tails. *J. Biol. Chem.* 271, 22315–22320.
- Torstensen, B.E., Espe, M., Stubhaug, I., Lie, O., 2011. Dietary plant proteins and vegetable oil blends increase adiposity and plasma lipids in Atlantic salmon (*Salmo salar* L.). *Br. J. Nutr.* 106, 633–647.
- Trattner, S., Ruyter, B., Østbye, T.K., Gjoen, T., Zlabek, V., Kamal-Eldin, A., Pickova, J., 2008. Sesamin increases alpha-linolenic acid conversion to docosahexaenoic acid in Atlantic salmon (*Salmo salar* L.) hepatocytes: role of altered gene expression. *Lipids* 43, 999–1008.
- Triantafyllou, M., Gamper, F.G., Haston, R.M., Mouratis, M.A., Morath, S., Hartung, T., Triantafyllou, K., 2006. Membrane sorting of toll-like receptor (TLR)-2/6 and TLR2/1 heterodimers at the cell surface determines heterotypic associations with CD36 and intracellular targeting. *J. Biol. Chem.* 281, 31002–31011.
- van der Sar, A.M., Abdallah, A.M., Sparrius, M., Reinders, E., Vandenbroucke-Grauls, C.M., Bitter, W., 2004. *Mycobacterium marinum* strains can be divided into two distinct types based on genetic diversity and virulence. *Infect. Immun.* 72, 6306–6312.
- van der Sar, A.M., Spaik, H.P., Zakrzewska, A., Bitter, W., Meijer, A.H., 2009. Specificity of the zebrafish host transcriptome response to acute and chronic mycobacterial infection and the role of innate and adaptive immune components. *Mol. Immunol.* 46, 2317–2332.
- Verburg-van Kemenade, B.M.L., Groeneveld, A., van Rens, B.T.T.M., Rombout, J.H.W.M., 1994. Characterization of macrophages and neutrophilic granulocytes from the pronephros of carp (*Cyprinus carpio*). *J. Exp. Biol.* 187, 143–158.
- Zakrzewska, A., Cui, C., Stockhammer, O.W., Benard, E.L., Spaik, H.P., Meijer, A.H., 2010. Macrophage-specific gene functions in Spi1-directed innate immunity. *Blood* 116, e1–e11.
- Zhang, M., Xu, Y., Li, L., Wei, S., Zhang, S., Liu, Z., 2013. Identification, evolution and expression of a CD36 homolog in the basal chordate amphioxus *Branchiostoma japonicum*. *Fish Shellfish Immunol.* 34, 546–555.