



Universiteit
Leiden
The Netherlands

Robustness of trait connections across environmental gradients and growth forms

Flores-Moreno, H.; Fazayeli, F.; Banerjee, A.; Datta, A.; Kattge, J.; Butler, E.E.; ... ; Reich, P.B.

Citation

Flores-Moreno, H., Fazayeli, F., Banerjee, A., Datta, A., Kattge, J., Butler, E. E., ... Reich, P. B. (2019). Robustness of trait connections across environmental gradients and growth forms. *Global Ecology And Biogeography*, 28(12), 1806-1826. doi:10.1111/geb.12996

Version: Publisher's Version

License: [Licensed under Article 25fa Copyright Act/Law \(Amendment Taverne\)](#)

Downloaded from: <https://hdl.handle.net/1887/3200449>

Note: To cite this publication please use the final published version (if applicable).

RESEARCH PAPER

Robustness of trait connections across environmental gradients and growth forms

Habacuc Flores-Moreno^{1,2}  | Farideh Fazayeli³ | Arindam Banerjee³ |
 Abhirup Datta⁴ | Jens Kattge^{5,6} | Ethan E. Butler²  | Owen K. Atkin^{7,8} |
 Kirk Wythers² | Ming Chen² | Madhur Anand⁹ | Michael Bahn¹⁰ | Chaeho Byun¹¹ |
 J. Hans C. Cornelissen¹² | Joseph Craine¹³ | Andres Gonzalez-Melo¹⁴ |
 Wesley N. Hattingh¹⁵ | Steven Jansen¹⁶ | Nathan J. B. Kraft¹⁷ | Koen Kramer¹⁸ |
 Daniel C. Laughlin¹⁹ | Vanessa Minden²⁰ | Ülo Niinemets²¹  | Vladimir Onipchenko²² |
 Josep Peñuelas^{23,24} | Nadejda A. Soudzilovskaia²⁵ | Rhiannon L. Dalrymple²⁶  |
 Peter B. Reich^{2,27}

¹Department of Ecology, Evolution, and Behavior, University of Minnesota, St. Paul, Minnesota

²Department of Forest Resources, University of Minnesota, St. Paul, Minnesota

³Department of Computer Science and Engineering, University of Minnesota, Minneapolis, Minnesota

⁴Department of Biostatistics, Johns Hopkins University, Baltimore, Maryland

⁵Max Planck Institute for Biogeochemistry, Jena, Germany

⁶German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, Leipzig, Germany

⁷ARC Centre of Excellence in Plant Energy, Research School of Biology, The Australian National University, Canberra, Australian Capital Territory, Australia

⁸Division of Plant Sciences, Research School of Biology, the Australian National University, Canberra, Australian Capital Territory, Australia

⁹School of Environmental Sciences, University of Guelph, Guelph, Ontario, Canada

¹⁰Institute of Ecology, University of Innsbruck, Innsbruck, Austria

¹¹School of Biological Sciences, Seoul National University, Seoul, South Korea

¹²Systems Ecology, Department of Ecological Science, Vrije Universiteit, Amsterdam, The Netherlands

¹³Jonah Ventures, Manhattan, Kansas

¹⁴Facultad de Ciencias Naturales y Matematicas, Universidad del Rosario, Bogota, Colombia

¹⁵School of Animal, Plant and Environmental Sciences, University of the Witwatersrand, Johannesburg, South Africa

¹⁶Institute of Systematic Botany and Ecology, Ulm University, Ulm, Germany

¹⁷Department of Ecology and Evolutionary Biology, University of California, Los Angeles, California

¹⁸Wageningen University and Research, Wageningen Environmental Research, Wageningen, The Netherlands

¹⁹Department of Botany, University of Wyoming, Laramie, Wyoming

²⁰Department of Biology, Ecology and Biodiversity, Vrije Universiteit Brussel, Brussels, Belgium

²¹Institute of Agricultural and Environmental Sciences, Estonian University of Life Sciences, Tartu, Estonia

²²Department of Geobotany, Moscow State Lomonosov University, Moscow, Russia

²³CSIC, Global Ecology Unit CREAM-CSIC-UAB, Bellaterra, Catalonia, Spain

²⁴CREAF, Cerdanyola del Vallès, Catalonia, Spain

²⁵Conservation Biology Department, Institute of Environmental Sciences, CML, Leiden University, Leiden, The Netherlands

²⁶Ecology and Evolution Research Centre, School of Biological, Earth and Environmental Sciences, UNSW Sydney, Sydney, New South Wales, Australia

²⁷Hawkesbury Institute for the Environment, Western Sydney University, New South Wales, Australia

Habacuc Flores-Moreno and Farideh Fazayeli would like to be considered joint first authors, and for this article to be cited as Flores-Moreno, H and Fazayeli, F, et al.

Correspondence

Habacuc Flores-Moreno, Department of Ecology, Evolution, and Behavior, University of Minnesota, St. Paul, MN 55108.
Email: habacucfm@gmail.com

Funding information

National Science Foundation, Grant/Award Number: IIS-1563950; Advanced Research Projects Agency - Energy, Grant/Award Number: DE-SL0012677; H2020 European Research Council, Grant/Award Number: ERC-SyG-2013-610028 IMBALANCE-P; University of Minnesota, Grant/Award Number: CE140100008, 226299, 19-14-00038 and 22; Australian Research Council, Grant/Award Number: CE140100008; FP7; European Research Council; Russian Science Foundation, Grant/Award Number: # 19-14-00038

Editor: Andrew Kerkhoff

Abstract

Aim: Plant trait databases often contain traits that are correlated, but for whom direct (undirected statistical dependency) and indirect (mediated by other traits) connections may be confounded. The confounding of correlation and connection hinders our understanding of plant strategies, and how these vary among growth forms and climate zones. We identified the direct and indirect connections across plant traits relevant to competition, resource acquisition and reproductive strategies using a global database and explored whether connections within and between traits from different tissue types vary across climates and growth forms.

Location: Global.

Major taxa studied: Plants.

Time period: Present.

Methods: We used probabilistic graphical models and a database of 10 plant traits (leaf area, specific leaf area, mass- and area-based leaf nitrogen and phosphorous content, leaf life span, plant height, stem specific density and seed mass) with 16,281 records to describe direct and indirect connections across woody and non-woody plants across tropical, temperate, arid, cold and polar regions.

Results: Trait networks based on direct connections are sparser than those based on correlations. Land plants had high connectivity across traits within and between tissue types; leaf life span and stem specific density shared direct connections with all other traits. For both growth forms, two groups of traits form modules of more highly connected traits; one related to resource acquisition, the other to plant architecture and reproduction. Woody species had higher trait network modularity in polar compared to temperate and tropical climates, while non-woody species did not show significant differences in modularity across climate regions.

Main conclusions: Plant traits are highly connected both within and across tissue types, yet traits segregate into persistent modules of traits. Variation in the modularity of trait networks suggests that trait connectivity is shaped by prevailing environmental conditions and demonstrates that plants of different growth forms use alternative strategies to cope with local conditions.

KEYWORDS

leaf traits, plant functional traits, plant strategy integration, seed traits, stem traits, trait interdependence, trait networks

1 | INTRODUCTION

Plant traits are not independent from each other biologically or statistically. An accurate description of their interdependency could give us a clearer view of a myriad of processes, from the links between morphological traits and physiological function inside the leaves of plants (Osnas, Lichstein, Reich, & Pacala, 2013; Poorter, Lambers, & Evans, 2014; Westoby, Reich, & Wright, 2013), to ecosystem-scale processes such as global gross primary production (Wang et al., 2012). Evidence of generalities in the coordination of traits exists, but it is mostly limited to single tissue types or a few traits at broad spatial scales (Chave et al., 2009; Moles et al., 2009;

Wright et al., 2004), or across tissue types but only for certain regions and/or growth forms (Ackerly, 2004; Baraloto et al., 2010; Cheng, Chu, Chen, & Bai, 2015; Fortunel, Fine, & Baraloto, 2012; Freschet, Cornelissen, Logtestijn, & Aerts, 2010; Kramer-Walter et al., 2016; Liu et al., 2010). Díaz et al. (2016) found strong evidence for coordination among six traits measured on different tissue types at global scales, but they did not disentangle direct and indirect connections among traits. As of yet, it remains unknown how widespread coordination is among multiple traits across tissue types, and across different biogeographical scales and growth forms, once direct and indirect connections have been unravelled. By coordination, we mean variation in a trait that is linked to variation in another

trait or traits (i.e. non-random variation between traits). Crucially, in this paper, we try to disentangle the causal rather than correlative relationships that give rise to coordination or lack thereof among traits, and then assess whether this causal coordination among traits differs across biogeographical scales and growth forms.

Most previous studies describe the coordination across multiple traits using some form of correlative analyses (e.g. Ackerly, 2004; Freschet et al., 2010; Kramer-Walter et al., 2016). In this study, we instead focus on describing the causal coordination among traits by identifying their "connections". We use connection in a specific way: as short-hand for undirected statistical dependencies among traits (Shipley, 2016). A connection represents a direct linkage between two traits arising as a result of conditional statistical dependence among these traits (sometimes, and perhaps often, functional). That is, the observed correlation among these two traits cannot be fully explained by their connections to other traits. Meanwhile, indirect connections are due to conditional statistical independence among traits where no direct connection between the two traits exists, and any observed statistical correlation between them is indirectly mediated by both having connections to one or more other traits or by an environmental variable (see examples in Supporting Information Appendix S3). We note that we do not a priori believe that correlations that are due to covariances are unimportant physiologically or ecologically; in fact, they likely help to contribute to the small fraction of trait space that is occupied (Díaz et al., 2016) and thus of "trait strategies" that are successful in nature. However, distinguishing direct and indirect connections among traits is necessary for understanding the mechanistic roots of the trait correlations that define plant strategies, and can thus help us to clarify the causal link between traits and fitness components (Shipley, Lechowicz, Wright, & Reich, 2006), connections among traits and function (Li et al., 2015; Poorter et al., 2014), and the role that traits play in influencing higher-level processes and vegetation attributes (e.g. relative growth rate, net primary productivity, Kramer-Walter et al., 2016; Reich, Walters, & Ellsworth, 1992).

Highly connected traits are expected as a result of biophysical and/or selection processes that favour the efficient use and acquisition of resources within and across plant tissues (Reich, 2014). On the other hand, a certain degree of independence among traits can give plants more flexibility to adjust function to the environmental conditions currently experienced by the plant, and allow for more variation in the overall phenotype of the plant (Li et al., 2015; Messier, Lechowicz, McGill, Violle, & Enquist, 2017), as different traits/tissues have their own trade-offs. Li et al. (2015) found evidence for independence between the hydraulic and economic dimensions of leaves, which may give leaves more freedom to respond to existing environmental conditions. Meanwhile, Kramer-Walter et al. (2016) described coordination among root, stem and leaf tissue that also impacted plants relative growth rate (RGR), while also finding independent variation in specific root length (SRL), which may allow for higher phenotypic variation in roots compared to leaves. We should expect that interdependence among traits is mediated by evolutionary history, biophysical constraints, the environment that plants experience and whether selection favours the integration or

independence of traits in those environments (Fortunel et al., 2012; Li et al., 2015; Messier et al., 2017; Reich, 2014).

While a number of studies have investigated the interdependence among traits across tissue types (e.g. Ackerly, 2004; Fortunel et al., 2012; Kramer-Walter et al., 2016; Messier et al., 2017; Reich, 2014), only a handful have done so at macroecological scales. At a global scale, Díaz et al. (2016) showed plant species occupy two orthogonal trait dimensions across six traits; one dimension, related to the size of plants, was formed by seed mass and stem traits, and the other dimension was composed of leaf traits that characterize the construction costs of photosynthetic tissue. Leaf and stem traits of Neotropical trees have also been shown to vary independently of each other (Baraloto et al., 2010). Meanwhile, interdependence among leaf, stem and root traits of trees in New Zealand has been shown (Kramer-Walter et al., 2016), as well as among traits across tissue types in Mongolian grasslands (Cheng et al., 2015), and northern European herbaceous species (Kleyer et al., 2018). Thus, there seems to be evidence for high interdependence among traits across tissue types; however, this seems to be variable across environments and growth forms and to be highly dependent on the traits examined. As of yet, it remains unclear how the causal interdependence of traits varies globally, across biogeographical scales and among plant groups.

Water availability is likely to influence the amount of interdependence among different tissues types. Plant communities from wetter environments exhibit independence among leaf, root and stem tissue types (Baraloto et al., 2010; Fortunel et al., 2012; Jager, Richardson, Bellingham, Clearwater, & Laughlin, 2015), while in drier environments interdependence among these tissue types has been observed (Laughlin, Leppert, Moore, & Sieg, 2010; Li & Bao, 2015; de la Riva et al., 2016). Similarly, the independence of traits related to water acquisition from traits related to nutrient acquisition increases in environments with higher precipitation (Ackerly, 2004; Baraloto et al., 2010; Fortunel et al., 2012; Ishida et al., 2008), and with increasing water availability (de la Riva et al., 2016). Thus, higher connectivity among traits may be selected for in drier environments or in environments with higher seasonal variation in water availability, as matching tissue strategies may allow the plants to function efficiently (Reich, 2014) and take up resources faster when conditions allow. Following this, we hypothesized that arid and polar/cold climates – where water availability is limited throughout most of the year or is available during a short season – will favour more connectivity across traits.

Numerous studies have described multi-trait interdependence among woody species (e.g. Ackerly, 2004; Baraloto et al., 2010; de la Riva et al., 2016; Jager et al., 2015; Li & Bao, 2015), while fewer studies have done so across non-woody species (e.g. Cheng et al., 2015; Craine, Froehle, Tilman, Wedin, & Chapin, 2001; Craine & Lee, 2003; Santini et al., 2017). To our knowledge, only Díaz et al. (2016) have specifically contrasted the difference in coordination among different traits across tissue types among woody and non-woody species, finding no difference in trait interdependence between growth forms, but some differences in the strength of specific trait correlations. Despite these efforts, differences in trait interdependence in woody and non-woody species still remain unexplored.

Several recent studies at local to regional scales have made more comprehensive examinations of multi-trait integration for either herbaceous or woody plants. For example, Kleyer et al. (2018) and Messier et al. (2017) explored trait integration within a growth form, employing higher trait resolution than typically done previously (i.e. their studies incorporated all vegetative tissues and sometimes even reproductive tissue types and > 20 traits). These studies both suggested that traits integrate by function rather than by tissue type, and also identified traits with high interdependence levels (we refer to these as traits with high centrality). Both traits also found evidence for interdependence among traits across tissue types with no clear-cut segregation of traits into distinct trait dimensions. While neither study quantified the connectedness of the whole trait network (what we refer to as the edge density of the network), a quick calculation using their reported numbers of observed trait connections divided by the number of possible connections suggests that the trait network of non-woody species (65 present connections/210 possible connections; Kleyer et al., 2018) is similar but perhaps slightly more connected than woody species (41 present connections/190 possible connections; Messier et al., 2017). However, this calculation and other comparisons among these studies are complicated by dissimilarities in spatial scales and suites of targeted traits. There is very limited knowledge available that directly addresses the differences/similarities in trait interdependence between growth forms; however, from the limited evidence available, we hypothesized that non-woody species may display higher interdependence among traits.

Here we explore three hypotheses: (a) that trait connections (i.e. undirected statistical dependencies) and trait correlation are identical, (b) that high trait connectivity will be favoured in biomes where water availability is limited such as in arid and polar/cold climates, and (c) that trait connectivity will be higher in non-woody than woody species. To address our hypotheses we (i) compare trait correlations and trait connections for each pair of traits in our trait network; (ii) describe the global trait network based on the connections between 10 functional traits using 16,281 trait observations from sites around the world; (iii) use network metrics to compare the differences in the trait networks of woody and non-woody species; and to (iv) assess how the trait networks of woody and non-woody species change across five broad climate regions (tropical, temperate, arid, cold and polar). We evaluate the connections among traits through a test of undirected statistical dependence using a structure learning algorithm using the well-studied graphical Lasso (Friedman, Hastie, & Tibshirani, 2008). The key feature of this method is its capacity to identify direct from indirect connections with high probability by identifying relationships among traits once covariation with all other traits has been accounted for.

2 | METHODS

2.1 | Trait, taxonomic data and climate zones

Our attention is on 10 plant traits related to the functions of leaf tissues, stem tissues, and reproductive tissues of plants, and are thus relevant to resource economy and uptake, competitive ability

(or stress tolerance) and reproductive strategy. Seed mass (mg) reflects allocation of energy to few large versus many small offspring, and impacts early seedling survival (Moles & Westoby, 2006). Plant height (m) and stem specific density (mg dry mass/mm³ fresh volume; hereon SSD) are related to light competition, growth rate and long-term viability of stems (Chave et al., 2009; Moles et al., 2009). Specific leaf area (mm²/mg; SLA), leaf life span (month; LLS), leaf nitrogen (mg/g; N) mass and leaf phosphorus (mg/g; P) mass are related to nutrient economy and acquisition, and are key components of the leaf economic spectrum (LES; Reich et al., 1999; Wright et al., 2004). SLA represents the potential return rate measured in terms of light capture area per unit of mass investment (Reich, Walters, & Ellsworth, 1997; Wright et al., 2004). LLS represents the time needed to generate payback on this investment (Reich et al., 1992). Leaf N is associated with carboxylation capacity and is integral to the photosynthetic machinery (Reich et al., 1999; Wright et al., 2004). Leaf P is essential for bioenergetic molecules (e.g. ATP) and is linked to the formation of indispensable nucleic acids and lipid membranes (Reich et al., 1999; Wright et al., 2004). Therefore, variations in leaf P and N are crucial to respiration and photosynthetic capacity, as well as energy generation and storage. Leaf area (mm²) is related to the water and energy balance of a plant and is relevant to light interception. Finally, leaf N and P can also be expressed on an area basis (g/m²) reflecting light capture and transaction of energy on an area basis (Wright et al., 2004). Consequently, we also use leaf N per area and leaf P per area, in parallel to their mass-based counterparts.

We obtained spatially explicit trait data for our 10 traits and growth form (woody, non-woody) data from TRY (www.try.db.org; Kattge et al., 2011; a list of the original data sources is found in the Appendix). The TRY data subset used in this study includes 16,281 georeferenced trait observation records for 15,284 species, of which 9,053 and 6,231 species were identified as woody and non-woody plants, respectively. When we had multiple observations for the same trait for a species within the same climatic region, we used the geometric mean to calculate a single species-climate region combination, with the exception of plant height, for which we used the maximum value. We standardized the species names to *The Plant List* (v. 1.1; The Plant List, 2013), and obtained the higher order taxonomy for our species with *taxonlookup* (v. 1.0.1; Pennell, FitzJohn, & Cornwell, 2016). In our dataset LLS was the trait with the lowest number of records (c. 0.67% of records have information for this trait), while plant height was the trait with the highest number of records present (c. 35%; Supporting Information Table S1.1). Thus, we used a hierarchical Bayesian extension of probabilistic matrix factorization (BHPMF) to fill in the trait gaps in our dataset (Fazayeli, Banerjee, Kattge, Schrodte, & Reich, 2014; Schrodte et al., 2015; Shan et al., 2012). This algorithm harnesses the available trait and taxonomic information to fill in the gaps in the trait data, by approximating the trait matrices at higher taxonomic levels, which then serve as priors for the next level in the hierarchy (Fazayeli et al., 2014; Schrodte et al., 2015). This approach has the advantage of leveraging any phylogenetic signals in traits, although a phylogenetic signal is not a prerequisite for the model to work efficiently (Schrodte et al., 2015).

Previous studies have evaluated BHPMF's prediction accuracy, uncertainty, and confidence in predictions. These studies have compared BHPMF to other methods such as probabilistic matrix factorization and the more commonly used MEAN – gap-filling with species or genus level mean values – using plant traits datasets spanning 49–92% missing entries per trait (Fazayeli et al., 2014; Moreno-Martínez et al., 2018; Schrod et al., 2015). BHPMF's prediction accuracy outperforms all other methods with lower root mean square error (RMSE) values, and higher correlation coefficient of observed versus predicted values (Fazayeli et al., 2014; Schrod et al., 2015). BHPMF's average error of predictions is about half a standard deviation, and crucially its prediction accuracy is not related to the number of entries per trait (Schrod et al., 2015). BHPMF preserves trait-trait correlations well and reproduces extreme values more accurately than MEAN, thus it also captures the shape of the scatter of observed trait data more accurately than MEAN (Schrod et al., 2015). Further, uncertainty evaluation tests based on the relationship between a model's confidence versus accuracy have shown that BHPMF is accurate when confidence is high (Fazayeli et al., 2014; Schrod et al., 2015). By contrast, MEAN can introduce large error and bias in some cases as this method adds new data points without adding new information, making confidence limits incorrect (Fazayeli et al., 2014; Schrod et al., 2015).

A number of studies have used the BHPMF method in other trait analyses with robust results (see Blonder et al., 2018; Caplan, Meiners, Flores-Moreno, & McCormack, 2018; Díaz et al., 2016; Moreno-Martínez et al., 2018). Nonetheless, we also checked the robustness of the BHPMF algorithm by comparing the trait-trait relationship from 1,000 gap-filled datasets versus trait-trait correlations only from the original dataset (Supporting Information Figure S1.1, and determining the correlation between the accuracy (RMSE) and confidence (SD) of the algorithm using our dataset (Supporting Information Figure S1.2). The first test shows that BHPMF reproduces extreme values accurately capturing the shape of the scatter of observed trait data and in most cases the original trait-trait correlation lies inside the 95% confidence interval of the 1,000 gap-filled datasets suggesting the algorithm is robust in assessing correlations among trait pairs (Supporting Information Figure S1.1). The relationship between RMSE and SD suggests that the algorithm is accurate when it is confident (Supporting Information Figure S1.2). Further, we ensured the robustness of our results by re-running our analyses excluding LLS, which is the most gap-filled trait in our dataset (Supporting Information Appendix S1).

Using a map of the Köppen climate zones we assigned the georeferenced plant records to five different climates: tropical climate (with little to moderate water deficiency through seasons), which includes tropical rainforest, tropical seasonal forest, and savannas; arid climate (with low water availability through seasons), which includes deserts and steppes; temperate climate (with little or no water deficiency through seasons and seasonal variation in temperature), which includes temperate forest, temperate rainforest and Mediterranean vegetation; cold climate (with low water availability and high temperature seasonality), which includes only taiga; polar climate (with low water availability and extreme variation in

temperature), which includes tundra, alpine and circumpolar zones (Peel, Finlayson, & McMahon, 2007). While variation in temperature and/or water availability exist within these climate regions, more variation in environmental conditions occurs among them (Peel et al., 2007). We chose the climate types above for two reasons: (a) the climate types described above are based on long-term precipitation and heat records, which combined capture broad thermal and moisture differences among regions (Feddema, 2005), (b) the sparsity of plant trait data at global scales precludes the use of the 30 Köppen climate subtypes described by Peel et al. (2007).

2.2 | Estimation of trait correlations

We used Pearson correlations to calculate the relationship for each trait pair across all land plants. To estimate the statistical significance of the correlations, we generated 1,000 bootstrapped trait datasets and calculated the 95% confidence interval of the correlation for each trait pair. To determine the relationship between correlation and connections (the derivation of these is described below) we compared their absolute value for each trait pair, since we did not have a hypothesis about the sign of these metrics. Prior to analyses we \log_{10} transformed and z-transformed all continuous trait data.

2.3 | Estimation of trait connections

To determine the connections among traits we calculated the precision matrix and the confidence interval for each trait-trait interaction in the trait network. The precision matrix (i.e. inverse of the covariance matrix) establishes the undirected statistical dependency among a set of variables, in this case a set of traits. Undirected statistical dependencies can be understood in terms of partial correlations. Given a pair of traits (x and y) with correlation $\rho = \rho_1 + \rho_2$, ρ_1 would be how much of this correlation can be explained by another trait z (z being a single trait or set of traits), while ρ_2 would be how much of this correlation cannot be explained by z . Here, ρ_2 is the partial correlation and would correspond to an entry in the precision matrix for the undirected statistical dependency between traits x and y . If $\rho_2 = 0$, then any observed correlation between traits x and y can be fully explained by z (i.e. traits x and y do not provide information about each other once trait z is considered). If $\rho_2 \neq 0$, then there is a direct connection between traits x and y , even after accounting for trait z . Thus, the precision matrix of traits provides the undirected statistical conditional dependency structure among traits for a multivariate (in this case, assumed log-normal) set of traits, which describes the direct probabilistic interactions among traits (Supporting Information Appendix S2; Cai, Li, Liu, & Xie, 2012; Ebert-Uphoff & Deng, 2012). Importantly, the sparsity structure of the precision matrix matches the sparsity structure of ρ_2 .

We derived the precision matrix for all plants, and then for non-woody and woody plants separately. Next, we derived the precision matrix for non-woody and woody plants within each of the five climate zones defined above. We used a "glasso" algorithm (Graphical Lasso; Friedman et al., 2008) to estimate the precision matrix for each plant group. The glasso algorithm assumes that traits have a

multivariate Gaussian distribution and estimates the precision matrix by minimizing the negative log-likelihood among all plant trait measurements. Following Jankova and van de Geer (2015), we accounted for differences in sample size across precision matrices and their effect as: $\lambda = 2 \sqrt{(\log p/n)}$, where n refers to the sample size and p denotes the number of variables (i.e. traits; Supporting Information Appendix S2). Then for each precision matrix, we tested the significance of the trait–trait connections using a glasso-based confidence interval (Jankova & van de Geer, 2015). Finally, we removed the connections among traits whose confidence interval contained zero.

2.4 | Characterization of networks of trait connections using network metrics

To characterize the differences in the connections among traits – which were obtained through the estimation of precision matrices – within and across trait networks, we used three common metrics in the analyses of networks: “modularity” to establish the connectivity

among trait modules, “edge density” to quantify the connectivity of all traits across the whole trait network, and “degree” to quantify the connectivity of a focal trait to other traits in the network. We describe each of these metrics in detail below.

“Modularity” is the difference between the fraction of connections among traits that fall within a given module (i.e. a module is a subset of traits that interact more among themselves than with other surrounding traits) minus the same fraction in a null model where connections among traits are randomly distributed (Clauset, Newman, & Moore, 2004; Figure 1). We used modularity to measure the connectivity among trait modules. Modularity helps identify the tendency for traits within a trait network to cluster, in that networks with high modularity have a great number of connections among traits within modules but sparse connections among traits between modules (Figure 1). Higher modularity of traits confers an advantage under variable conditions as it lends robustness (Alon, 2003; Ravasz, Somera, Mongru, Oltvai, & Barabási, 2002), also providing opportunities for the network to adapt and evolve, as not all components




FIGURE 1 Network metrics used to assess the causal interdependence (i.e. connections) among traits. The metrics focus on the trait network, and individual traits’ properties. *Modularity*: describes the tendency of nodes within a network to cluster by quantifying how separated modules within a network are from each other. Modules are groups of traits (blue and red polygons) that interact more strongly among themselves than with other traits. In networks with low modularity traits interact strongly among modules (panel a, left), in networks with high modularity (panel a, right) traits within a module interact more strongly among themselves than with traits in other modules. *Edge density*: proportion of present connections among traits out of all possible connections in the network. In networks with high edge density (panel b, left), connections exist between all traits, networks with low edge density (panel b, right figure) have few connections among traits. *Degree*: measures how many connections an individual trait (e.g. Trait A) has with other traits in the network. A trait with high degree has connections with many (sometimes all) other traits (panel c, left) in the network, a trait with low degree (panel c, right) has no or few connections with other traits in the network

in a modular network are optimally linked (i.e. ability to respond to changing external conditions/internal organization while maintaining normal behaviour; Barabasi & Oltvai, 2004). We used a Spinglass algorithm to detect the module placement in our networks. To identify module placement this algorithm is based on the principle that nodes should connect edges of the same community, while nodes of different communities should be disconnected (Yang, Algesheimer, & Tessone, 2016). The community detection through spin glass may be understood as finding the ground state of an infinite range Potts spin glass by combining the information from both present and missing links (Reichardt & Bornholdt, 2006). The network community structure is interpreted as the spin (i.e. community) configuration that minimizes the energy of the spin glass, where the spin states are the community indices. The Spinglass algorithm accurately detects modules in networks of small size (number of nodes – here, traits – ≤ 233) and with small or large mixing parameters at the network level (i.e. the summation of external degree of each node over the summation of its total degree; Yang et al., 2016). Once we determined the placement of modules in the networks, we recorded the number of modules present, their trait composition and calculated their modularity. Modularity values of zero represent networks without compartmentalization of modules (i.e. networks where there is no subset of traits that interact more among themselves than with other surrounding traits), while non-zero values represent networks that have compartmentalization of modules (i.e. networks that tend to have dense connections among traits within modules but sparse connections to traits in other modules; Figure 1).

“Edge density” defines the proportion of present connections among traits out of all possible connections in a trait network (Figure 1). We used edge density to assess the connectedness across traits of the whole trait network, regardless of their modules. In the context of biological networks, variation in edge density is used as an indicator of the compromise between efficiency of connections versus the cost of connections (Alon, 2003; Lipson, Pollack, Suh, & Wainwright, 2002). For instance, a network with high edge density may allow for the efficient acquisition and mobilization of resources (as all traits are connected with all other traits), nevertheless this may be costly in terms of the establishment and maintenance of connections among traits. Edge density ranges from zero to one. A value of zero represents no connection across the traits in the network; a value of one implies that all traits in the network are connected to all other traits (Figure 1).

“Degree” is the number of connections between a focal trait and other traits, normalized by the total number of potential connections (Horvath, 2011; Figure 1). This is a measure of the connectivity of a focal trait to the rest of the trait network. Thus, we used degree – a type of centrality measure – to quantify the relative importance (the centrality) of a trait for a given trait network; values for degree range between zero when a focal trait has no connection to any other trait, to one when a focal trait is connected to all other traits in the network. Degree is widely used in biological networks to identify essential characteristics in biological entities such as genes, metabolites and proteins (Koschützki & Schreiber, 2008). For example,

metabolites with higher degree may belong to the oldest part of the metabolism, while proteins with higher degree have been identified as essential, with their removal being lethal to the organism (Koschützki & Schreiber, 2008).

2.5 | Significance of differences in trait network metrics

To estimate the statistical significance of the differences in network metrics above), we did the following: First, using 1,000 bootstrapped trait datasets, we calculated the precision matrices for each plant group and climate region (as described above), and calculated each network metric for each of the 1,000 resulting trait networks. Then, for a given network metric (i.e. modularity, degree, edge density) we calculated the 95% confidence interval of the difference across all group pairs. Following Besag, Green, Higdon, and Mengersen (1995), we corrected for multiple comparison by calculating simultaneous credible intervals based on order statistics of the difference in a given network metric among groups.

We ran all analyses in R 3.5.3 (R Core Team, 2019). For calculating the precision matrices we used the camel package (Li, Zhao, & Liu, 2013), and for calculating the network metrics we used the igraph package (Csardi & Nepusz, 2006).

3 | RESULTS

Contrary to our first hypothesis, we found that trait connections and trait correlation are not identical for all land plants (Figure 2). From the 28 unique trait–trait relationships possible in our trait networks, we found that 23 have both significant trait–trait correlations and trait connections, four have only significant trait correlations and one has only a significant trait connection (Figure 2c,d, Supporting Information Table S4.1). The disagreements between connections and correlations occur in trait–trait relationships with medium to low correlation values ($|r| \leq .30$; Figure 2d). There is a connection between leaf N mass–seed mass, but no correlation between these traits. On the other hand, there are four trait–trait relationships with significant correlations (leaf P mass–seed mass, leaf N mass–plant height, leaf area–leaf P mass and leaf area–leaf N mass), but without significant connection among these traits (Figure 2c). We found that strong correlations, such as those between SLA–LLS, seed mass–plant height, leaf N mass–leaf P mass ($|r| \geq .5$), were always present as connections (Supporting Information Table S4.1).

Among all terrestrial plants, there is a high proportion of connections among traits across all tissue types (edge density = .86, Figure 3a and Supporting Information Figure S4.1), and no modularity among traits (modularity = 0, Figure 3b). LLS and SSD (both degree = 1) are more central to the global trait networks in that they have significantly higher numbers of connections to other traits compared to plant height (degree = .86), leaf N (degree = .71) and leaf area (degree = .71). Leaf area (degree = .71) and leaf N mass (degree = .71) are the traits with the fewest connections to other traits (Figure 3c and




FIGURE 2 Comparison of trait correlations (r) and connections (Ω). (a) Trait network based on correlations (coloured connections are significant). (b) Trait network based on connections (coloured connections are significant). (c) Trait connections that differ from correlations. (d) Comparison of absolute values of trait connections and correlations; black points are significant trait–trait correlations and significant trait connections, red points are trait–trait relations that had either connections or correlations that were not significant

Supporting Information Table S4.2, S4.3. All other traits fall somewhere in between these two extremes, in descending order: SLA also had a high proportion of connections to other traits, but only significantly higher than plant height and leaf N; seed mass had higher proportion of connections to other traits but only compared to leaf N; and leaf P had a low proportion of connections compared to other traits; however, this difference was not significant (Figure 2c and Supporting Information Table S4.2, S4.3). Details of the global trait networks and multiple comparison corrected 95% confidence intervals for all results are in Supporting Information Appendix S3.

Contrary to our hypothesis that non-woody species will have higher connectivity among traits compared to woody species, we found that both growth forms have similar proportions of connections among traits in the trait network (edge density_{woody} = .71; edge density_{non-woody} = .61; Figure 3a and Supporting Information Figure S4.1) and both have two trait modules with comparably low levels of modularity, showing higher integration of traits across modules than within modules (modularity_{non-woody} = .10; modularity_{woody} = .06; Figure 3b). However, we found some fundamental differences in the composition of the trait modules and in

the identity of most central traits in the trait networks. In terms of the modules' composition, in non-woody species one module is composed of LLS, leaf N and SSD, while the other is composed of leaf P, SLA, plant height, leaf area and seed mass (Table 1). Meanwhile, in woody species one module is formed by leaf area, leaf P and SSD and the other one by seed mass, LLS, SLA, leaf N and plant height (Table 1). Both growth forms have comparable levels of modularity, showing higher integration of traits across modules than within modules (modularity_{non-woody} = .10; modularity_{woody} = .06; Figure 3b). For non-woody species leaf N, leaf area, LLS and leaf P (degree = .71) are the traits with the highest number of connections to other traits in the trait network (Figure 3d). Meanwhile, for woody species the traits with the highest proportion of connections to other traits are LLS (degree = .86), leaf area (degree = .86) and seed mass (degree = .86, Figure 3e). In both cases stem-related traits are the traits with the fewest connections to other traits in the network (degree \leq .57 Figure 3d,e). Details on trait networks of woody and non-woody species are presented in Supporting Information Appendix S3.

In both woody and non-woody species, there are connections among all leaf-tissue traits, except SLA–LLS in non-woody species.




FIGURE 3 Trait networks for all plants, woody and non-woody growth forms. (a) Edge density, (b) modularity and (c) trait degree. Whiskers denote 95% confidence interval. See Figure 1 for hypothetical graphical examples of these metrics

TABLE 1 Modules' composition for all plants, non-woody and woody species

Growth form	Modules
All plants	Leaf area-SLA-leaf N-LLS-leaf P-plant ht-SSD-seed mass
Non-woody	Leaf area-leaf P-plant ht SLA-leaf N-LLS-SSD-seed mass
Woody	Leaf area-LLS-plant ht-seed mass SLA-leaf N-leaf P-SSD

Note: plant ht = plant height; LLS = leaf life span; SLA = specific leaf area; SSD = stem specific density. Modules in our networks are detected using the Spinglass algorithm (see Methods). The modules in this table are the most common modules across the 1,000 networks obtained from the analyses of 1,000 bootstrapped trait matrices. The pipe character “|” separates individual modules. Traits across modules may be connected (see graphical representation of modularity in Figure 1b); however, they tend to be more connected with other traits within the modules than with traits outside the module.

Also, in both growth form groups there are connections between plant height-leaf area, plant height-LLS, seed mass-leaf area, seed mass-SLA, seed mass-leaf N and seed mass-plant height. Connections between LLS-SLA, SSD-leaf area, SSD-leaf P, plant height-leaf N, seed mass-LLS, and seed mass-SSD are only present in woody species. Connections between SSD-leaf N, SSD-LLS and plant height-leaf P are only present in non-woody species.

3.1 | Woody and non-woody species by climate region

Connections among traits of woody and non-woody species differ across climate regions (Supporting Information Figure S4.2). However, contrary to our expectations, the proportion of connectedness among traits did not increase with decreasing water availability. Woody species show a higher proportion of connections among traits (i.e. higher edge density) in tropical (edge density = .64),

temperate (edge density = .68), arid (edge density = .64) and cold environments (edge density = .57), compared to polar ones (edge density = .39; Figure 4a and Supporting Information Figure S4.2, Table S4.4, 4.5). Non-woody species show a higher proportion of connections between traits in cold (edge density = .61) compared to polar environments (edge density = .43; Figure 4c and Supporting Information Figure S4.2, Table S4.4, 4.5). However, non-woody species also have a higher proportion of connections in temperate (edge density = .57) and cold environments compared to tropical environments (edge density = .36; Figure 4c and Supporting Information Figure S4.2, Table S4.4, 4.5).

Woody species show significantly higher integration of traits within modules in polar (modularity = .25) compared to temperate and tropical environments (for both modularity = .08; Figure 4b and Supporting Information Table S4.6, 4.7). Non-woody species show non-significant differences in modularity across environments (Figure 4d and Supporting Information Table S4.6, 4.7). Across the different climate regions, woody species always have two modules, while non-woody species have two modules in all climates except polar and tropical where they have three (Table 2). Both growth forms have a module mainly composed of traits related to the LES. In woody species, this consists of leaf N and leaf P, and SLA is part of this module in all climates except tropical, while LLS is part of this module in all climates except cold climate. In non-woody species, the analogous module contains SLA, leaf N and leaf P in all climate regions. The second module consists of traits related to reproductive strategy and plant architecture for woody species, and mainly plant

architecture in non-woody species. In woody species, the core traits in this module are seed mass, plant height, leaf area and SSD. In non-woody species this module consists of plant height and leaf area; SSD is part of this module in all climates except polar, and seed mass is part of this module in all climates except tropical and arid. When a third module is present in non-woody species, it consists of LLS and either plant height in tropical, or SSD in polar climates (Table 2).

For both growth forms, LLS is the trait with the highest centrality (i.e. it has the most connections to other traits) in temperate ($\text{degree}_{\text{woody}} = .71$, $\text{degree}_{\text{non-woody}} = .71$) and arid climates ($\text{degree}_{\text{woody}} = .86$, $\text{degree}_{\text{non-woody}} = .57$), while seed mass has high centrality in tropical areas ($\text{degree}_{\text{woody}} = .86$, $\text{degree}_{\text{non-woody}} = .43$; Supporting Information Table S4.8–4.11). For non-woody species leaf N has a high centrality in all climate zones, except cold, while leaf area has a high centrality in all environments except temperate and polar ones (Supporting Information Table S4.10, 4.11). For woody species LLS has the highest centrality in all regions except tropical ones (Supporting Information Table S4.8, 4.9).

Across climate types and both growth forms, connections between leaf N–leaf P, leaf N–SLA, plant height–leaf area and seed mass–leaf area are always present. For non-woody species connections among seed mass–SLA and seed mass–leaf N are also robust across climate types. Meanwhile, for woody species connections between SLA–leaf area, leaf P–SLA, LLS–leaf N, SSD–leaf P, seed mass–plant height and seed mass–SSD are also found across climate regions. Details of trait network differences across climate regions are presented in Supporting Information Appendix S3.




FIGURE 4 Plant trait network edge density, modularity, and trait degree for woody (a, b and e, respectively) and non-woody species (c, d and f, respectively) across five climate regions using mass-based leaf N and P. Whiskers denote 95% confidence interval

Climate	Woody species modules	Non-woody species modules
Tropical	Leaf area–SLA–plant ht–SSD–seed mass leaf N–LLS–leaf P	Leaf area–SSD SLA–leaf N–leaf P–seed mass LLS–plant ht
Temperate	Leaf area–plant ht–SSD–seed mass SLA–leaf N–LLS–leaf P	Leaf area–LLS–plant ht–SSD–seed mass SLA–leaf N–leaf P
Arid	Leaf area–plant ht–SSD–seed mass SLA–leaf N–LLS–leaf P	Leaf area–LLS–plant ht–SSD SLA–leaf N–leaf P–seed mass
Cold	Leaf area–LLS–plant ht–SSD–seed mass SLA–leaf N–leaf P	Leaf area–LLS–plant ht–SSD–seed mass SLA–leaf N–leaf P
Polar	Leaf area–plant ht–SSD–seed mass SLA–leaf N–LLS–leaf P	Leaf area–plant ht–seed mass SLA–leaf N–leaf P LLS–SSD

Note: plant ht = plant height; LLS = leaf life span; SLA = specific leaf area; SSD = stem specific density. Modules in our networks are detected using the Spinglass algorithm (see Methods). The modules in this table are the most common modules across the 1,000 networks obtained from the 1,000 bootstrapped trait matrix. The pipe character “|” separates individual modules.

3.2 | Effect of excluding LLS and using area-based leaf N and P on the networks of trait connections

The patterns presented above are largely consistent when excluding LLS – the most gap-filled trait in our dataset (Supporting Information Appendix S1), or when we run the analyses using area-based leaf N and P traits (Supporting Information Appendix S4). The robustness of connections across climates and growth forms is similar in area-based results to that observed in mass-based analyses, with the addition of a connection between LLS–SLA, seed mass–leaf N and seed mass–plant height. In the results excluding LLS, we observe the addition of a connection between SSD–leaf areas. As in the mass-based results, woody species show higher modularity in polar regions when these traits are area-based (Supporting Information Table S5.5, S5.6), and when excluding LLS (Supporting Information Table S2.5, S2.6). As in the mass-based results, no one trait is essential to all networks (as measured by their degree) across all climates for both woody and non-woody growth forms when results are derived using area-based traits (Supporting Information Table S5.7–S5.10), or when we excluded LLS (Supporting Information Table S2.7–2.10). Across leaf nutrient mass-based and area-based results, and results excluding LLS, we are always able to distinguish two modules, one mainly related to LES traits and another one related to reproductive strategy and plant architecture.

4 | DISCUSSION

Here we have identified emergent characteristics of the trait connections for all land plants at a global scale, indicating trait connections both within and across plant leaf, stem and reproductive tissues. Further, we have identified variation in the trait connection network across growth forms (woody and non-woody) and, for the first time (to our knowledge), explicitly accounted for the impact of broad environmental gradients on the trait network. In doing so, our study builds on – and extends – previous attempts that describe the correlations across several traits and several tissue types at global

TABLE 2 Modules of non-woody and woody species across climate regions

scales (Díaz et al., 2016), and efforts that focused on certain vegetation types (Wright et al., 2007) and narrower environmental gradients (Cheng et al., 2015). The analyses indicate that strong integration across traits across different tissue types exists at the global scale, and across growth forms – however, integration among traits changes across environments. We find that groups of traits cluster forming modules, and that the composition of these modules mostly agrees with previously described trait dimensions. Our analyses also highlight individual traits that are highly connected, as well as robust trait–trait connections that are persistent across growth form and climate regions; however, these results also suggest that the importance of these traits and trait–trait connections are not always supported by current knowledge based on correlative evidence. We found that strong correlations are present as connections, while some weaker correlations are either direct connections or absent as connections. Collectively our analyses of causal connections among traits using a network approach suggest that while the plant phenotype is highly integrated, differences among growth forms and environmental conditions that plants experience modify the strength, clustering and centrality of traits and trait connections in the trait network.

4.1 | Connections among traits across all terrestrial plants

When we compared trait–trait relationships across all land plants using connections and correlations we found that 15% of all the present trait–trait correlations disappear as connections, and the only non-significant trait–trait correlation in the trait network emerged as a connection (leaf N mass–seed mass). These results suggest that trait networks are somewhat sparser than suggested by correlation analyses. That correlations disappear as connections suggests that the interactions among these traits are mediated through another trait(s) in our trait network once the influence of the network is taken into consideration. Importantly, correlations that are critical to plant growth, survival and reproduction – extensively described elsewhere – and which tend to have high correlation values, were always present as connections, whereas only trait–trait correlations with

medium to low values disappear when examined as connections. The correlation between leaf N mass–seed mass was non-existent, yet our analyses suggests there is a direct connection between these traits. This direct connection perhaps arises via the key role that both traits have in plant survival and reproduction or more specifically through the mobilization of vegetative nitrogen needed for the generation of seed storage protein (Shibles & Sundberg, 1998).

Variation in edge density in biological networks has been linked to a compromise between efficiency of connections versus the cost of connection (Alon, 2003; Lipson et al., 2002). At a global scale we found that land plants have a high proportion of connectedness (high trait network edge density) across traits and high integration across tissue types, although not all possible trait connections exist or are strong enough for biological significance. Contrary to our hypothesis, we found no significant differences in the proportion of connection among traits across growth forms, and an overall high proportion of connectedness across traits. This suggests that while there are crucial physiological and anatomical differences across woody and non-woody strategies, these differences do not translate to higher or lower integration of the whole plant phenotype. This supports the idea that matching tissue strategies (i.e. high integration among traits) should be advantageous (Reich, 2014) across growth forms, as this may allow for the efficient acquisition and sharing of resources across the whole plant. On the other hand, our analyses also show that the trait network proportion of connectedness varies across climate regions, suggesting that strong trait coordination and therefore efficient resource sharing comes with a cost and may not be advantageous under all environmental conditions. We, however, reject our hypothesis that traits will be more integrated in water-restricted environments, instead finding some evidence in woody species for lower trait integration in polar regions compared to all other climate regions, and a mixed pattern in non-woody species. Disentangling which abiotic conditions explain the most variation in trait integration, as well as the relationship between trait integration and these leading abiotic variables will be an important next step.

4.2 | Modularity varies across climates

Modules are groups of traits that interact more strongly among themselves, and tend to perform a common function (Alon, 2003); conceptually, modules are similar to trait dimensions/axes (Kleyer et al., 2018). We found there are at least two core modules that are present in both plant growth forms, and a third module present in trait networks of non-woody species. This finding is in agreement with studies describing the dimensionality of plant trait strategies, who usually define at least two independent axes of trait variation (e.g. Ackerly, 2004; Díaz et al., 2016); although up to four or even six independent axes have been described (Jager et al., 2015; Laughlin et al., 2010). Meanwhile, emergent analyses using network approaches have revealed highly integrated trait networks at local and regional scales, concluding a lack of support for the notion of independent trait dimensions (Kleyer et al., 2018; Messier et al., 2017). However, it is important to note that our present study and most

other studies that have focused on defining axes of trait variation have asked which traits interact strongly regardless of their tissue type, meanwhile Messier et al. (2017) focused on assessing the modularity of predefined trait dimensions, which is conceptually different. Differences in focus, scale and number as well as type of traits explored may somewhat explain the differences among studies.

SLA, leaf N and leaf P (mass- or area-based) always constituted a module in woody species across climate regions. These three physiological leaf-tissue traits are central to the LES (Wright et al., 2004). Seed mass, plant height and leaf area formed the core traits in a second persistent module (Table 2). These reproductive and architectural traits are relevant to plant size, and plant–water and –light relationships (Moles et al., 2005; Niklas, 1994; Westoby, Falster, Moles, Vesk, & Wright, 2002). Thus, in woody species we have one persistently connected module of traits within a tissue type, and one persistent module of connections of traits that come from different tissue types. Similarly, in non-woody species, leaf N and SLA were always together in one module (leaf P was part of this module, except in tropical climates), while plant height and leaf area were together in a separate module (SSD formed part of this module except in polar climates). The importance of these two modules in the strategies of plants is supported by the LES and the global spectrum of plant form and function (Díaz et al., 2016; Reich et al., 1999, 1997). Thus, much in agreement with studies focusing on defining the dimensionality of plant strategies, we find the trait network has one module whose function is carbon uptake represented by leaf physiological traits, and a second module whose function is more related to plant reproductive and architectural traits. While trait networks are integrated to maximize the efficiency of resource use and acquisition as suggested by Reich (2014) and supported by our centrality analyses and others (e.g. Kramer-Walter et al., 2016; Messier et al., 2017), trait networks are at the same time segregated into modules of more highly interacting traits that may provide adaptive advantages under varying environmental conditions.

The degree of modularity of trait networks varies across environmental gradients for woody, but not for non-woody species. This suggests plant growth forms differ in their strategies to cope with environmental conditions. The lack of variation in the modularity of non-woody species across environments suggests they experience weaker environmental filtering compared to woody species (Šimová et al., 2018). Weaker environmental filtering in non-woody species perhaps results from their shorter life span, dormancy (and other bet-hedging) strategies and lack of long-persisting, reinforced stems above ground, which in combination may allow them to avoid much environmental heterogeneity. Within woody species, modularity is greater in polar than in tropical or temperate climates, meaning that modules in the trait network in polar regions are more independent from one another compared to other climate regions. This suggests that less coordination between trait dimensions (modules) is advantageous in polar environments. Higher independence among modules in polar regions may give plants more flexibility to adjust functions and help better manage risk of component failure (Gilarranz, Rayfield,

Liñán-Cembrano, Bascompte, & Gonzalez, 2017). This may be in response to greater variation in water availability between seasons (Feddema, 2005), greater temporal asynchrony between resources in polar environments, narrower range of variation in traits under harsher environmental conditions making relationship between traits weaker and thus increasing modularity (e.g. de la Riva et al., 2018), or more variable and/or different selection forces acting more strongly on each independent trait module (Laughlin et al., 2010).

4.3 | Highly connected traits

High centrality – having a high number of connections to other units in a network – suggests that a variable tends to be influential in terms of regulating critical functions or be involved in the regulation of more functions, thereby having greater impact on higher level properties such as fitness (Koschützki & Schreiber, 2008). LLS and SSD were the traits with most connections to other traits across land plants. Indeed, LLS is a crucial trait in the resource acquisition strategy of plants, influencing the carbon construction cost and carbon gain of leaves (Reich et al., 1999, 1992), while SSD impacts plant hydraulic and mechanical properties and influences the nutrient, carbon and water economy of stems (Chave et al., 2009). Further, variation in these traits has been shown to impact growth and fecundity (Adler et al., 2014), and also to contribute to the structuring of woody communities (Kunstler et al., 2016) and to influence ecosystem level processes (Reich, 2014; Reich et al., 1997). However, trait centrality was labile within growth forms across climate regions. Changes in the centrality of traits across growth forms and climate regions may simply reflect the scale-dependent nature of the selective, genetic and biophysical forces acting on traits. We conjecture that trait centrality may indeed reflect some of the functional relevance of traits in the trait network, thus providing crucial information about what traits are critical to the definition and understanding of plant strategies.

4.4 | Trait connections across growth forms and climate regions

Connections between seed mass–leaf area, leaf N–SLA, leaf P–leaf N and plant height–leaf area stand out because they were always present in analyses of growth forms, and across different climate types (Table 3). Some of these connections are well known (e.g. leaf N–SLA, leaf P–leaf N), and previous correlation analyses have identified their importance in understanding compromises among traits and their impact on plant function (Reich et al., 1999, 1992, 1997; Wright et al., 2004). However, some others are connections that previous correlation analyses suggested have no or weak relationships across habitats, and therefore their importance has been downplayed (e.g. seed mass–leaf area, plant height–leaf area; Table 3). In the first case – where we detect a connection and previous studies show a strong correlation – evidence suggests that these connections are maintained through selective pressure of biophysical constraints and natural selection (Table 3). For example, an increase in SLA will generally be linked to an increase in concentration of leaf N and other

cytoplasmic molecules (Meziane & Shipley, 2001; Reich et al., 1997). At the same time, natural selection reinforces a strong relationship between SLA and leaf N through processes such as herbivory and competition (Reich et al., 1997), limiting the trait space where optimal combinations of these traits occur. In the second case – where conditional dependency between two traits exists, but previous correlation tests suggested a weak relationship – we propose that these trait connections are maintained in the plant phylogeny through neutral or selective processes, but contradictory selective forces across habitats weaken the correlation among these traits. Some studies have reported a triangular relationship between seed mass and leaf area in temperate woody species (i.e. big leaves have big or small seeds, but small leaves only have small seeds; Cornelissen, 1999); other studies report positive rather than triangular relationships across woody sclerophyll species (Westoby & Wright, 2003), while others show no relationship among these traits across woody tropical species (Wright et al., 2007). In both cases, our analyses suggest that these trait connections are robust after accounting for all other traits, and across climates and growth forms, but the forces maintaining these connections might differ.

Some well-known, strong trait–trait correlations across plants stand out because they do have robust connections globally but not across growth forms or climate regions in our study. For instance, the connection between SLA–LLS is direct in the global data and in woody plants but not in non-woody plants. Moreover, this connection was observed in four of five climate regions for woody plants, but only one of five for non-woody plants. This weaker connection in subsets of the global data could suggest that although a strong correlation exists between these two traits, the connection between these traits could be mechanistically mediated through other traits in some cases, or the connection only exists when the absolute range in LLS is large, as is the case for woody plants. Additionally, given that the vast majority of global data on LLS for graminoids likely reflects longevity of the entire leaf blade (much of which may be senescing or dead) rather than of active photosynthetic tissue per se (Craine, Berin, Reich, Tilman, & Knops, 1999; Craine & Reich, 2001), the majority of global data on graminoid LLS may not be as comparable to LLS in woody and non-woody dicots as the research community recognizes (or wishes). This may contribute to differences in connection of SLA–LLS in woody versus non-woody plants.

Previous plant trait studies have focused on understanding the interdependence between pairs of traits or among multiple traits using correlation analyses in combination with ordination techniques (e.g. Supporting Information Table S1.2). These studies have shown that the strength and form of the relationships between traits among co-occurring species vary across environments (Ackerly, 2004; Reich et al., 1999), and among functional groups (Reich et al., 2003), with axes of trait variation shifting, collapsing or arising across environmental gradients and different growth forms. The analyses presented here suggest that direct and indirect interactions among traits within and between tissue types also shift across growth forms and climate regions. We found that LLS and SSD are the most central traits globally, but were not always connected within all climate

TABLE 3 Trait connections that are robust (i.e. common across groups) across growth forms and climate regions and proposed mechanisms that maintain this connection

Robust connection	Woody/non-woody	Woody across climate	Non-woody across climate	Mechanisms	Details
Leaf P-leaf N	.62	.51-.70	.47-.72	Biochemical and biogeochemical constraints. Selection through competition and herbivory	Selection through biochemical constraints reinforced through compromises between allocation of resources to metabolism (investment in genetic material especially ribosomal RNA) versus growth (investment in proteins); biogeochemical constraints mediated by adaptation/acclimation to soil conditions; unviable/low fitness strategies may be selected out by herbivory and competition based processes (Kerkhoff, Fagan, Elser, & Enquist, 2006; Reich, 2014; Reich & Oleksyn, 2004)
SLA-leaf N	.52-.64	.59-.67	.41-.57	Biophysical constraint; selection of viable strategies through competition and herbivory	Changes in leaf N results from increases in LMA, increase in SLA results in an increase in cytoplasmic compounds, including N; structural investment, competition and herbivory select certain combinations of these traits (Meziane & Shipley, 2001; Reich et al., 1999, 1992, 1997)
Plant height-leaf area	.50-.54	.29-.57	.36-.70	Biophysical constraint	Decrease in leaf size with increasing light demands correlates weakly, selection through light competition in certain floras; small trunks with big leaves are physiologically unviable; large-leaf crown is more efficient because it requires less woody support investment (Givnish, 1979; Niinemets & Kull, 1994; Niklas, 1994)
Seed mass-leaf area	.41-.42	.15-.50	.36-.41	Unclear if this correlation is general across floras or what maintains it. Possible controls by biophysical constraints or vascular/meristematic demands	Correspondence between axis and appendage size (Corner's rule). Triangular relationship – plants with big leaves can have either very small or large seeds, but plants with small leaves only have small seeds in woody European species; positive but not triangular relationship in sclerophyll vegetation in Australia; no relationship among these traits in tropical forests in the Americas (Cornelissen, 1999; Westoby & Wright, 2003; Wright et al., 2007)

Note: LMA = leaf mass per area; SLA = specific leaf area. The range of R^2 values observed across growth form, and then by growth form across climate regions in this study is provided in the second to fourth columns. We provided proposed mechanisms that maintain these trait connection as well as specific hypothesis about each mechanism (Details column).

zones, especially for non-woody plants (but see above); moreover, few trait–trait connections exist robustly across all growth forms and climate gradients (Table 3). Despite the difference in statistical approach compared with previous research (e.g. Ackerly, 2004; Fortunel et al., 2012; Kramer-Walter et al., 2016), our study supports the existence of two distinguishable dimensions or functional modules across land plants and climate regions. One module is related to physiological leaf traits related to carbon uptake and economy, and another related to reproductive strategy and plant architecture. The findings presented here contribute to the fundamental understanding of dependencies between plant traits and their variation across environmental gradients. Our approach represents an important step forward on the collective path to understanding the causal links among multiple traits across multiple tissue types, and within and across different climate zones and plant growth forms.

ACKNOWLEDGMENTS

HFM, PBR, MC, EEB, KRW, AB were funded by the United States Department of Energy (DE-SL0012677). HFM was partially funded by a Discovery grant from the Institute on the Environment at University of Minnesota. AB and PBR were supported by a National Science Foundation (NSF) grant IIS-1563950. PBR was funded by two University of Minnesota Institute on the Environment Discovery Grants. OKA acknowledges funding by the Australian Research Council (CE140100008). KK was supported by the European Union Seventh Framework programme (EU-FP7) project Biodiversity And Climate Change, A Risk Analysis (BACCARA; 226299) and the national project *Resilient Forests* (KB-29-009-003). JP acknowledges the financial support from the European Research Council Synergy grant ERC-SyG-2013-610028 IMBALANCE-P. VO thanks the Russian Science Foundation (RSF) for financial support (# 19-14-00038). The study was supported by the TRY initiative on plant traits (<http://www.try-db.org>). The TRY initiative and database is hosted, developed and maintained at the Max Planck Institute for Biogeochemistry, Jena, Germany. TRY is currently supported by DIVERSITAS/Future Earth and the German 22 Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig. Discussions with Santiago Soliveres, Matthew Michalska-Smith, and comments from Andrew Kerkhoff, Enrique de la Riva and two anonymous referees contributed greatly to this paper.

DATA AVAILABILITY STATEMENT

The data used for the analyses are publicly available: climate zones (https://webmap.ornl.gov/ogc/wcsdown.jsp?dg_id=10012_1) and plants (www.try-db.org; upon request to TRY).

ORCID

Habacuc Flores-Moreno  <https://orcid.org/0000-0002-7083-0005>

Ethan E. Butler  <https://orcid.org/0000-0003-3482-1950>

Ülo Niinemets  <https://orcid.org/0000-0002-3078-2192>

Rhiannon L. Dalrymple  <https://orcid.org/0000-0002-2660-1424>

REFERENCES

- Ackerly, D. (2004). Functional strategies of chaparral shrubs in relation to seasonal water deficit and disturbance. *Ecological Monographs*, *74*, 25–44. <https://doi.org/10.1890/03-4022>
- Adler, P. B., Salguero-Gómez, R., Compagnoni, A., Hsu, J. S., Ray-Mukherjee, J., Mbeau-Ache, C., & Franco, M. (2014). Functional traits explain variation in plant life history strategies. *Proceedings of the National Academy of Sciences USA*, *111*, 740–745. <https://doi.org/10.1073/pnas.1315179111>
- Alon, U. (2003). Biological networks: The tinkerer as an engineer. *Science*, *301*, 1866–1867. <https://doi.org/10.1126/science.1089072>
- Barabasi, A.-L., & Oltvai, Z. N. (2004). Network biology: Understanding the cell's functional organization. *Nature Reviews Genetics*, *5*, 101–113. <https://doi.org/10.1038/nrg1272>
- Baraloto, C., Timothy Paine, C. E., Poorter, L., Beauchene, J., Bonal, D., Domenach, A. M., ... Chave, J. (2010). Decoupled leaf and stem economics in rain forest trees. *Ecology Letters*, *13*, 1338–1347. <https://doi.org/10.1111/j.1461-0248.2010.01517.x>
- Besag, J., Green, P., Higdon, D., & Mengersen, K. (1995). Bayesian computation and stochastic systems. *Statistical Science*, *10*, 3–41. <https://doi.org/10.1214/ss/1177010123>
- Blonder, B., Enquist, B. J., Graae, B. J., Kattge, J., Maitner, B. S., Morueta-Holme, N., ... Violle, C. (2018). Late Quaternary climate legacies in contemporary plant functional composition. *Global Change Biology*, *24*, 4827–4840. <https://doi.org/10.1111/gcb.14375>
- Cai, T. T., Li, H., Liu, W., & Xie, J. (2012). Covariate-adjusted precision matrix estimation with an application in genetical genomics. *Biometrika*, *100*, 139–156. <https://doi.org/10.1093/biomet/ass058>
- Caplan, J. S., Meiners, S. J., Flores-Moreno, H., & McCormack, M. L. (2018). Fine-root traits are linked to species dynamics in a successional plant community. *Ecology*, *100*, e02588.
- Chave, J., Coomes, D., Jansen, S., Lewis, S. L., Swenson, N. G., & Zanne, A. E. (2009). Towards a worldwide wood economics spectrum. *Ecology Letters*, *12*, 351–366. <https://doi.org/10.1111/j.1461-0248.2009.01285.x>
- Cheng, J., Chu, P., Chen, D., & Bai, Y. (2015). Functional correlations between specific leaf area and specific root length along a regional environmental gradient in Inner Mongolia grasslands. *Functional Ecology*, *30*, 985–997. <https://doi.org/10.1111/1365-2435.12569>
- Clauset, A., Newman, M. E. J., & Moore, C. (2004). Finding community structure in very large networks. *Physical Review E*, *70*, 066111. <https://doi.org/10.1103/PhysRevE.70.066111>
- Cornelissen, J. H. C. (1999). A triangular relationship between leaf size and seed size among woody species: Allometry, ontogeny, ecology and taxonomy. *Oecologia*, *118*, 248–255. <https://doi.org/10.1007/s004420050725>
- Craine, J., Berin, D., Reich, P., Tilman, D., & Knops, J. (1999). Measurement of leaf longevity of 14 species of grasses and forbs using a novel approach. *The New Phytologist*, *142*, 475–481. <https://doi.org/10.1046/j.1469-8137.1999.00411.x>
- Craine, J., Froehle, J., Tilman, D., Wedin, D., & Chapin, III, F. S. (2001). The relationships among root and leaf traits of 76 grassland species and relative abundance along fertility and disturbance gradients. *Oikos*, *93*, 274–285. <https://doi.org/10.1034/j.1600-0706.2001.930210.x>
- Craine, J., & Lee, W. (2003). Covariation in leaf and root traits for native and non-native grasses along an altitudinal gradient in New Zealand. *Oecologia*, *134*, 471–478. <https://doi.org/10.1007/s00442-002-1155-6>
- Craine, J. M., & Reich, P. B. (2001). Elevated CO₂ and nitrogen supply alter leaf longevity of grassland species. *New Phytologist*, *150*, 397–403. <https://doi.org/10.1046/j.1469-8137.2001.00116.x>
- Csardi, G., & Nepusz, T. (2006). The igraph software package for complex network research. *InterJournal, Complex Systems*, *1695*, 1–9.
- de la Riva, E. G., Tosto, A., Pérez-Ramos, I. M., Navarro-Fernández, C. M., Olmo, M., Anten, N. P. R., ... Villar, R. (2016). A plant economics

- spectrum in Mediterranean forests along environmental gradients: Is there coordination among leaf, stem and root traits? *Journal of Vegetation Science*, 27, 187–199. <https://doi.org/10.1111/jvs.12341>
- de la Riva, E. G., Violle, C., Pérez-Ramos, I. M., Marañón, T., Navarro-Fernández, C. M., Olmo, M., & Villar, R. (2018). A multidimensional functional trait approach reveals the imprint of environmental stress in Mediterranean woody communities. *Ecosystems*, 21, 248–262. <https://doi.org/10.1007/s10021-017-0147-7>
- Díaz, S., Kattge, J., Cornelissen, J. H. C., Wright, I. J., Lavorel, S., Dray, S., ... Gorné, L. D. (2016). The global spectrum of plant form and function. *Nature*, 529, 167–171. <https://doi.org/10.1038/nature16489>
- Ebert-Uphoff, I., & Deng, Y. (2012). Causal discovery for climate research using graphical models. *Journal of Climate*, 25, 5648–5665. <https://doi.org/10.1175/JCLI-D-11-00387.1>
- Fazayeli, F., Banerjee, A., Kattge, J., Schrod, F., & Reich, P. B. (2014). Uncertainty quantified matrix completion using Bayesian hierarchical matrix factorization. In X. Chen, G. Qu, P. Angelov, C. Ferri, J. Lai, & M. Arif Wani (Eds.), *13th International Conference on Machine Learning and Applications (ICMLA)* (pp. 312–317). Los Alamitos, CA: IEEE Computer Society.
- Feddema, J. J. (2005). A revised Thornthwaite-type global climate classification. *Physical Geography*, 26, 442–466. <https://doi.org/10.2747/0272-3646.26.6.442>
- Fortunel, C., Fine, P. V. A., & Baraloto, C. (2012). Leaf, stem and root tissue strategies across 758 Neotropical tree species. *Functional Ecology*, 26, 1153–1161. <https://doi.org/10.1111/j.1365-2435.2012.02020.x>
- Freschet, G. T., Cornelissen, J. H. C., Van Logtestijn, R. S. P., & Aerts, R. (2010). Evidence of the 'plant economics spectrum' in a subarctic flora. *Journal of Ecology*, 98, 362–373. <https://doi.org/10.1111/j.1365-2745.2009.01615.x>
- Friedman, J., Hastie, T., & Tibshirani, R. (2008). Sparse inverse covariance estimation with the graphical lasso. *Biostatistics*, 9, 432–441. <https://doi.org/10.1093/biostatistics/kxm045>
- Gilarranz, L. J., Rayfield, B., Liñán-Cembrano, G., Bascompte, J., & Gonzalez, A. (2017). Effects of network modularity on the spread of perturbation impact in experimental metapopulations. *Science*, 357, 199–201. <https://doi.org/10.1126/science.aal4122>
- Givnish, T. (1979). On the adaptive significance of leaf form. In O.T. Solbrig (Ed.), *Topics in plant population biology* (pp. 375–407). London, UK: The Macmillan Press.
- Horvath, S. (2011). *Weighted network analysis: Applications in genomics and systems biology*. New York, NY: Springer Science & Business Media.
- Ishida, A., Nakano, T., Yazaki, K., Matsuki, S., Koike, N., Lauenstein, D. L., ... Yamashita, N. (2008). Coordination between leaf and stem traits related to leaf carbon gain and hydraulics across 32 drought-tolerant angiosperms. *Oecologia*, 156, 193–202. <https://doi.org/10.1007/s00442-008-0965-6>
- Jager, M. M., Richardson, S. J., Bellingham, P. J., Clearwater, M. J., & Laughlin, D. C. (2015). Soil fertility induces coordinated responses of multiple independent functional traits. *Journal of Ecology*, 103, 374–385. <https://doi.org/10.1111/1365-2745.12366>
- Jankova, J., & van de Geer, S. (2015). Confidence intervals for high-dimensional inverse covariance estimation. *Electronic Journal of Statistics*, 9, 1205–1229. <https://doi.org/10.1214/15-EJS1031>
- Kattge, J., Díaz, S., Lavorel, S., Prentice, I. C., Leadley, P., Bönsch, G., ... Wirth, C. (2011). TRY—a global database of plant traits. *Global Change Biology*, 17, 2905–2935. <https://doi.org/10.1111/j.1365-2486.2011.02451.x>
- Kerckhoff, A. J., Fagan, W. F., Elser, J. J., Enquist, B. J. (2006). Phylogenetic and growth form variation in the scaling of nitrogen and phosphorus in the seed plants. *The American Naturalist*, 168(4), E103–E122.
- Kleyer, M., Trinogga, J., Cebrián-Piqueras, M. A., Trenkamp, A., Fløjgaard, C., Ejrnaes, R., ... Blasius, B. (2018). Trait correlation network analysis identifies biomass allocation traits and stem specific length as hub traits in herbaceous perennial plants. *Journal of Ecology*, 107, 829–842. <https://doi.org/10.1111/1365-2745.13066>
- Koschützki, D., & Schreiber, F. (2008). Centrality analysis methods for biological networks and their application to gene regulatory networks. *Gene Regulation and Systems Biology*, 2, 193–201. <https://doi.org/10.4137/GRSB.S702>
- Kramer-Walter, K. R., Bellingham, P. J., Millar, T. R., Smissen, R. D., Richardson, S. J., & Laughlin, D. C. (2016). Root traits are multidimensional: Specific root length is independent from root tissue density and the plant economic spectrum. *Journal of Ecology*, 104, 1299–1310. <https://doi.org/10.1111/1365-2745.12562>
- Kunstler, G., Falster, D., Coomes, D. A., Hui, F., Kooyman, R. M., Laughlin, D. C., ... Westoby, M. (2016). Plant functional traits have globally consistent effects on competition. *Nature*, 529, 204–207. <https://doi.org/10.1038/nature16476>
- Laughlin, D. C., Leppert, J. J., Moore, M. M., & Sieg, C. H. (2010). A multi-trait test of the leaf-height-seed plant strategy scheme with 133 species from a pine forest flora. *Functional Ecology*, 24, 493–501. <https://doi.org/10.1111/j.1365-2435.2009.01672.x>
- Li, F. L., & Bao, W. K. (2015). New insights into leaf and fine-root trait relationships: Implications of resource acquisition among 23 xerophytic woody species. *Ecology and Evolution*, 5, 5344–5351. <https://doi.org/10.1002/ece3.1794>
- Li, L., McCormack, M. L., Ma, C., Kong, D., Zhang, Q., Chen, X., ... Guo, D. (2015). Leaf economics and hydraulic traits are decoupled in five species-rich tropical-subtropical forests. *Ecology Letters*, 18, 899–906. <https://doi.org/10.1111/ele.12466>
- Li, X., Zhao, T., & Liu, H. (2013). *camel: Calibrated machine learning*. R package version 0.2.0. retrieved from <https://cran.r-project.org/src/contrib/Archive/camel/>
- Lipson, H., Pollack, J. B., Suh, N. P., & Wainwright, P. (2002). On the origin of modular variation. *Evolution*, 56, 1549–1556. <https://doi.org/10.1111/j.0014-3820.2002.tb01466.x>
- Liu, G., Freschet, G. T., Pan, X., Cornelissen, J. H. C., Li, Y., & Dong, M. (2010). Coordinated variation in leaf and root traits across multiple spatial scales in Chinese semi-arid and arid ecosystems. *New Phytologist*, 188, 543–553. <https://doi.org/10.1111/j.1469-8137.2010.03388.x>
- Messier, J., Lechowicz, M. J., McGill, B. J., Violle, C., & Enquist, B. J. (2017). Interspecific integration of trait dimensions at local scales: The plant phenotype as an integrated network. *Journal of Ecology*, 105, 1775–1790. <https://doi.org/10.1111/1365-2745.12755>
- Meziane, D., & Shipley, B. (2001). Direct and indirect relationships between specific leaf area, leaf nitrogen and leaf gas exchange. Effects of irradiance and nutrient supply. *Annals of Botany*, 88, 915–927. <https://doi.org/10.1006/anbo.2001.1536>
- Moles, A. T., Ackerly, D. D., Webb, C. O., Tweddle, J. C., Dickie, J. B., & Westoby, M. (2005). A brief history of seed size. *Science*, 307, 576–580. <https://doi.org/10.1126/science.1104863>
- Moles, A. T., Warton, D. I., Warman, L., Swenson, N. G., Laffan, S. W., Zanne, A. E., ... Leishman, M. R. (2009). Global patterns in plant height. *Journal of Ecology*, 97, 923–932. <https://doi.org/10.1111/j.1365-2745.2009.01526.x>
- Moles, A. T., & Westoby, M. (2006). Seed size and plant strategy across the whole life cycle. *Oikos*, 113, 91–105. <https://doi.org/10.1111/j.0030-1299.2006.14194.x>
- Moreno-Martínez, Á., Camps-Valls, G., Kattge, J., Robinson, N., Reichstein, M., van Bodegom, P., ... Running, S. W. (2018). A methodology to derive global maps of leaf traits using remote sensing and climate data. *Remote Sensing of Environment*, 218, 69–88. <https://doi.org/10.1016/j.rse.2018.09.006>
- Niklas, K. J. (1994). *Plant allometry: The scaling of form and process*. Chicago, IL: University of Chicago Press.
- Niinemets, Ü., & Kull, K. (1994). Leaf weight per area and leaf size of 85 Estonian woody species in relation to shade tolerance and light availability. *Forest Ecology and Management*, 70(1–3), 1–10.

- Osnas, J. L. D., Lichstein, J. W., Reich, P. B., & Pacala, S. W. (2013). Global leaf trait relationships: Mass, area, and the leaf economics spectrum. *Science*, *340*, 741–744. <https://doi.org/10.1126/science.1231574>
- Peel, M. C., Finlayson, B. L., & McMahon, T. A. (2007). Updated world map of the Köppen-Geiger climate classification. *Hydrology and Earth System Sciences*, *4*, 439–473. <https://doi.org/10.5194/hessd-4-439-2007>
- Pennell, M. W., FitzJohn, R. G., & Cornwell, W. K. (2016). A simple approach for maximizing the overlap of phylogenetic and comparative data. *Methods in Ecology and Evolution*, *7*, 751–758. <https://doi.org/10.1111/2041-210X.12517>
- Poorter, H., Lambers, H., & Evans, J. R. (2014). Trait correlation networks: A whole-plant perspective on the recently criticized leaf economic spectrum. *New Phytologist*, *201*, 378–382. <https://doi.org/10.1111/nph.12547>
- Ravasz, E., Somera, A. L., Mongru, D. A., Oltvai, Z. N., & Barabási, A. L. (2002). Hierarchical organization of modularity in metabolic networks. *Science*, *297*, 1551–1555. <https://doi.org/10.1126/science.1073374>
- R Core Team. (2019). *R: A language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing. Retrieved from <http://www.R-project.org/>.
- Reich, P. B. (2014). The world-wide 'fast-slow' plant economics spectrum: A traits manifesto. *Journal of Ecology*, *102*, 275–301. <https://doi.org/10.1111/1365-2745.12211>
- Reich, P. B., Buschena, C., Tjoelker, M. G., Wrage, K., Knops, J., Tilman, D., & Machado, J.-L. (2003). Variation in growth rate and ecophysiology among 34 grassland and savanna species under contrasting N supply: A test of functional group differences. *New Phytologist*, *157*, 617–631. <https://doi.org/10.1046/j.1469-8137.2003.00703.x>
- Reich, P. B., Ellsworth, D. S., Walters, M. B., Vose, J. M., Gresham, C., Volin, J. C., & Bowman, W. D. (1999). Generality of leaf trait relationships: A test across six biomes. *Ecology*, *80*, 1955–1969. [https://doi.org/10.1890/0012-9658\(1999\)080\[1955:GOLTRA\]2.0.CO;2](https://doi.org/10.1890/0012-9658(1999)080[1955:GOLTRA]2.0.CO;2)
- Reich, P. B., & Oleksyn, J. (2004). Global patterns of plant leaf N and P in relation to temperature and latitude. *Proceedings of the National Academy of Sciences USA*, *101*(30), 11001–11006.
- Reich, P. B., Walters, M. B., & Ellsworth, D. S. (1992). Leaf life-span in relation to leaf, plant, and stand characteristics among diverse ecosystems. *Ecological Monographs*, *62*, 365–392. <https://doi.org/10.2307/2937116>
- Reich, P. B., Walters, M. B., & Ellsworth, D. S. (1997). From tropics to tundra: Global convergence in plant functioning. *Proceedings of the National Academy of Sciences USA*, *94*, 13730–13734. <https://doi.org/10.1073/pnas.94.25.13730>
- Reichardt, J., & Bornholdt, S. (2006). Statistical mechanics of community detection. *Physical Review E*, *74*, Abstract 016110. <https://doi.org/10.1103/PhysRevE.74.016110>
- Santini, B. A., Hodgson, J. G., Thompson, K., Wilson, P. J., Band, S. R., Jones, G., ... Rees, M. (2017). The triangular seed mass-leaf area relationship holds for annual plants and is determined by habitat productivity. *Functional Ecology*, *31*, 1770–1779. <https://doi.org/10.1111/1365-2435.12870>
- Schrodt, F., Kattge, J., Shan, H., Fazayeli, F., Joswig, J., Banerjee, A., ... Reich, P. B. (2015). BHPMF—a hierarchical Bayesian approach to gap-filling and trait prediction for macroecology and functional biogeography. *Global Ecology and Biogeography*, *24*, 1510–1521. <https://doi.org/10.1111/geb.12335>
- Shan, H., Kattge, J., Reich, P., Banerjee, A., Schrodt, F., & Reichstein, M. (2012). Gap filling in the plant kingdom—Trait prediction using hierarchical probabilistic matrix factorization. arXiv preprint arXiv:1206.6439.
- Shibles, R., & Sundberg, D. N. (1998). Relation of leaf nitrogen content and other traits with seed yield of soybean. *Plant Production Science*, *1*, 3–7. <https://doi.org/10.1626/pp.1.3>
- Shipley, B. (2016). *Cause and correlation in biology: A user's guide to path analysis, structural equations and causal inference with R*. Cambridge, England: Cambridge University Press.
- Shipley, B., Lechowicz, M. J., Wright, I., & Reich, P. B. (2006). Fundamental trade-offs generating the worldwide leaf economics spectrum. *Ecology*, *87*, 535–541. <https://doi.org/10.1890/05-1051>
- Šimová, I., Violle, C., Svenning, J.-C., Kattge, J., Engemann, K., Sandel, B., ... Enquist, B. J. (2018). Spatial patterns and climate relationships of major plant traits in the New World differ between woody and herbaceous species. *Journal of Biogeography*, *45*, 895–916. <https://doi.org/10.1111/jbi.13171>
- The Plant List (2013) *The Plant List version 1.1*. Available at: <http://www.theplantlist.org/>
- Wang, Y. P., Lu, X. J., Wright, I. J., Dai, Y. J., Rayner, P. J., & Reich, P. B. (2012). Correlations among leaf traits provide a significant constraint on the estimate of global gross primary production. *Geophysical Research Letters*, *39*(19), L19405. <https://doi.org/10.1029/2012GL053461>
- Westoby, M., Falster, D. S., Moles, A. T., Vesk, P. A., & Wright, I. J. (2002). Plant ecological strategies: Some leading dimensions of variation between species. *Annual Review of Ecology and Systematics*, *33*, 125–159. <https://doi.org/10.1146/annurev.ecolsys.33.010802.150452>
- Westoby, M., Reich, P. B., & Wright, I. J. (2013). Understanding ecological variation across species: Area-based vs mass-based expression of leaf traits. *New Phytologist*, *199*, 322–323. <https://doi.org/10.1111/nph.12345>
- Westoby, M., & Wright, I. J. (2003). The leaf size–twig size spectrum and its relationship to other important spectra of variation among species. *Oecologia*, *135*, 621–628. <https://doi.org/10.1007/s00442-003-1231-6>
- Wright, I. J., Ackerly, D. D., Bongers, F., Harms, K. E., Ibarra-Manriquez, G., Martinez-Ramos, M., ... Wright, S. J. (2007). Relationships among ecologically important dimensions of plant trait variation in seven neotropical forests. *Annals of Botany*, *99*, 1003–1015. <https://doi.org/10.1093/aob/mcl066>
- Wright, I. J., Reich, P. B., Westoby, M., Ackerly, D. D., Baruch, Z., Bongers, F., ... Villar, R. (2004). The worldwide leaf economics spectrum. *Nature*, *428*, 821–827. <https://doi.org/10.1038/nature02403>
- Yang, Z., Algesheimer, R., & Tessone, C. J. (2016). A comparative analysis of community detection algorithms on artificial networks. *Scientific Reports*, *6*, 30750. <https://doi.org/10.1038/srep30750>

BIOSKETCH

Habacuc Flores-Moreno undertook this research as a post-doc in Peter Reichs' lab at the University of Minnesota. His work focuses on linking the role of plant functional traits to community and ecosystem processes in different environmental and ecological contexts.

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of the article.

How to cite this article: Flores-Moreno H, Fazayeli F, Banerjee A, et al. Robustness of trait connections across environmental gradients and growth forms. *Global Ecol Biogeogr*. 2019;28:1806–1826. <https://doi.org/10.1111/geb.12996>

APPENDIX

DATA SOURCES: LIST OF STUDIES INCLUDED IN
THE SUBSET OF THE TRY DATABASE USED IN THE
CURRENT STUDY

- Adler, P. B., Milchunas, D. G., Lauenroth, W. K., Sala, O. E., & Burke, I. C. (2004). Functional traits of graminoids in semi-arid steppes: A test of grazing histories. *Journal of Applied Ecology*, 41(4), 653–663.
- Adriaenssens S. (2012). *Dry deposition and canopy exchange for temperate tree species under high nitrogen deposition* (PhD thesis), Ghent University, Ghent, Belgium.
- Auger, S., & Shipley, B. (2012). Interspecific and intraspecific trait variation along short environmental gradients in an old-growth temperate forest. *Journal of Vegetation Science*, 24, 419–428.
- Bahn, M., G. Wohlfahrt, E. Haubner, I. Horak, W. Michaeler, K. Rottmar, U. Tappeiner, & Cernusca, A. (1999). Leaf photosynthesis, nitrogen contents and specific leaf area of 30 grassland species in differently managed mountain ecosystems in the Eastern Alps. In A. Cernusca, U. Tappeiner, & N. Bayfield (Eds.), *Land-use changes in European mountain ecosystems. ECOMONT-Concept and Results* (pp. 247–255). Berlin, Germany: Blackwell Wissenschaft.
- Bakker, C., Rodenburg, J., & Bodegom, P. (2005). Effects of Ca- and Fe-rich seepage on P availability and plant performance in calcareous dune soils. *Plant and Soil*, 275, 111–122.
- Bakker, C., Van Bodegom, P. M., Nelissen, H. J. M., Ernst, W. H. O., & Aerts, R. (2006). Plant responses to rising water tables and nutrient management in calcareous dune slacks. *Plant Ecology*, 185, 19–28.
- Baraloto, C., Paine, C. E. T., Poorter, L., Beauchene, J., Bonal, D., Domenach, A.-M., ... Chave, J. (2010). Decoupled leaf and stem economics in rainforest trees. *Ecology Letters*, 13, 1338–1347.
- Blonder, B., Buzzard, B., Sloat, L., Simova, I., Lipson, R., Boyle, B., ... Enquist, B. (2012). The shrinkage effect biases estimates of paleoclimate. *American Journal of Botany*, 99(11), 1756–1763.
- Blonder, B., Vasseur, F., Violle, C., Shipley, B., Enquist, B., & Vile, D. (2015). Testing models for the leaf economics spectrum with leaf and whole-plant traits in *Arabidopsis thaliana*. *Annals of Botany*, 7(plv049).
- Blonder, B., Violle, C., & Enquist, B. (2013). Assessing the causes and scales of the leaf economics spectrum using venation networks in *Populus tremuloides*. *Journal of Ecology*, 101(4), 981–989.
- Blonder, B., Violle, C., Bentley, L. P., & Enquist, B. J. (2011). Venation networks and the origin of the leaf economics spectrum. *Ecology Letters*, 14(2), 91–100.
- Bond-Lamberty, B., Wang, C., & Gower, S. T. (2002). Above- and below-ground biomass and sapwood area allometric equations for six boreal tree species of northern Manitoba. *Canadian Journal of Forest Research*, 32(8), 1441–1450.
- Bond-Lamberty, B., Wang, C., & Gower, S. T. (2002). Leaf area dynamics of a boreal black spruce fire chronosequence. *Tree Physiology*, 22(14), 993–1001.
- Bond-Lamberty, B., Wang, C., & Gower, S. T. (2004). Net primary production and net ecosystem production of a boreal black spruce fire chronosequence. *Global Change Biology*, 10(4), 473–487.
- Brown, K. A., Flynn, D. F., Abram, N. K., Ingram, J. C., Johnson, S. E., & Wright, P. (2011). Assessing natural resource use by forest-reliant communities in Madagascar using functional diversity and functional redundancy metrics. *PLoS ONE*, 6(9), e24107.
- Burrascano S., Del Vico E., Fagiani S., Giarrizzo E., Mei M., Mortelliti A., Sabatini F. M., & Blasi C. (2015). Wild boar rooting intensity determines shifts in understorey composition and functional traits. *Community Ecology*, 16(2), 244–253.
- Butterfield, B. J., & Briggs, J. M. (2011). Regeneration niche differentiates functional strategies of desert woody plant species. *Oecologia*, 165, 477–487.
- Campetella, G., Botta-Dukát, Z., Wellstein, C., Canullo, R., Gatto, S., Chelli, S., ... Bartha, S. (2011). Patterns of plant trait–environment relationships along a forest succession chronosequence. *Agriculture, Ecosystems & Environment*, 145(1), 38–48.
- Carswell, F. E., Meir, P., Wandelli, E. V., Bonates, L. C. M., Kruijt, B., Barbosa, E. M., Nobre, A. D., & Jarvis, P. G. (2000). Photosynthetic capacity in a central Amazonian rain forest. *Tree Physiology*, 20(3), 179–186.
- Cavender-Bares, J., Keen, A., & Miles, B. (2006). Phylogenetic structure of floridian plant communities depends on taxonomic and spatial scale. *Ecology*, 87(sp7), S109–S122.
- Cavender-Bares, J., Sack, L., & Savage, J. (2007). Atmospheric and soil drought reduce nocturnal conductance in live oaks. *Tree Physiology*, 27, 611–620.
- Cerabolini B. E. L., Brusa G., Ceriani R. M., De Andreis R., Luzzaro A., & Pierce S. (2010). Can CSR classification be generally applied outside Britain? *Plant Ecology*, 210, 253–261.
- Chen, Y., Han, W., Tang, L., Tang, Z., & Fang, J. (2013). Leaf nitrogen and phosphorus concentrations of woody plants differ in responses to climate, soil and plant growth form. *Ecography*, 36(2), 178–184.
- Choat, B., Jansen, S., Brodribb, T. J., Cochard, H., Delzon, S., Bhaskar, R., ... Jacobsen, A. L. (2012). Global convergence in the vulnerability of forests to drought. *Nature*, 491(7426), 752.
- Coomes, D. A., Heathcote, S., Godfrey, E. R., Shepherd, J. J., & Sack, L. (2008). Scaling of xylem vessels and veins within the leaves of oak species. *Biology Letters*, 4, 302–306.
- Cornelissen, J. H. C. (1996). An experimental comparison of leaf decomposition rates in a wide range of temperate plant species and types. *Journal of Ecology*, 84, 573–582.
- Cornelissen, J. H. C., Cerabolini, B., Castro-Diez, P., Villar-Salvador, P., Montserrat-Marti, G., Puyravaud, J. P., ... Aerts, R. (2003). Functional traits of woody plants: Correspondence of species rankings between field adults and laboratory-grown seedlings? *Journal of Vegetation Science*, 14, 311–322.
- Cornelissen, J. H. C., Queded, H. M., Gwynn-Jones, D., Van Logtestijn, R. S. P., De Beus, M. A. H., Kondratchuk, A., Callaghan, T. V., & Aerts, R. (2004). Leaf digestibility and litter decomposability are related in a wide range of subarctic plant species and types. *Functional Ecology*, 18, 779–786.
- Cornelissen, J. H. C., Perez-Harguindeguy, N., Diaz, S., Grime, J. P., Marzano, B., Cabido, M., Vendramini, F., & Cerabolini, B. (1999). Leaf structure and defence control litter decomposition rate across species and life forms in regional floras on two continents. *New Phytologist*, 143, 191–200.
- Cornelissen, J. H. C., Diez, P. C., & Hunt, R. (1996). Seedling growth, allocation and leaf attributes in a wide range of woody plant species and types. *Journal of Ecology*, 84, 755–765.
- Cornwell, W. K., Cornelissen, J. H., Amatangelo, K., Dorrepaal, E., Eviner, V. T., Godoy, O., ... Queded, H. M. (2008). Plant species traits are the predominant control on litter decomposition rates within biomes worldwide. *Ecology Letters*, 11(10), 1065–1071.
- Cornwell, W. K., Bhaskar, R., Sack, L., Cordell, S., & Lurch, C. K. (2007). Adjustment of structure and function of Hawaiian *Metrosideros polymorpha* at high vs. low precipitation. *Functional Ecology*, 21(6), 1063–1071.
- Craine, J. M., Nippert, J. B., Towne, E. G., Tucker, S., Kembel, S. W., Skibbe, A., & McLaughlan, K. K. (2011). Functional consequences of climate change-induced plant species loss in a tallgrass prairie. *Oecologia*, 165(4), 1109–1117.
- Craine, J. M., Ocheltree, T. W., Nippert, J. B., Towne, E. G., Skibbe, A. M., Kembel, S. W., & Fargione, J. E. (2013). Global diversity of drought tolerance and grassland climate-change resilience. *Nature Climate Change*, 3(1), 63.
- Craine, J. M., Towne, E. G., Ocheltree, T. W., & Nippert, J. B. (2012). Community traitscape of foliar nitrogen isotopes reveals N availability patterns in a tallgrass prairie. *Plant and Soil*, 356(1–2), 395–403.

- Craine, J. M., Elmore, A. J., Aida, M. P., Bustamante, M., Dawson, T. E., Hobbie, E. A., ... Nardoto, G. B. (2009). Global patterns of foliar nitrogen isotopes and their relationships with climate, mycorrhizal fungi, foliar nutrient concentrations, and nitrogen availability. *New Phytologist*, 183(4), 980–992.
- Diaz, S., Hodgson, J. G., Thompson, K., Cabido, M., Cornelissen, J. H., Jalili, A., ... Band, S. R. (2004). The plant traits that drive ecosystems: Evidence from three continents. *Journal of Vegetation Science*, 15(3), 295–304.
- Demey, A., Staelens, J., Baeten, L., Boeckx, P., Hermy, M., Kattge, J., & Verheyen, K. (2013). Nutrient input from hemiparasitic litter favors plant species with a fast-growth strategy. *Plant and Soil*, 371(1–2), 53–66.
- Domingues, T. F., Martinelli, L. A., & Ehleringer, J. R. (2007). Ecophysiological traits of plant functional groups in forest and pasture ecosystems from eastern Amazonia, Brazil. *Plant Ecology*, 193(1), 101–112.
- Domingues TF, Meir P, Feldpausch TR, et al. (2010). Co-limitation of photosynthetic capacity by nitrogen and phosphorus in West Africa woodlands. *Plant, Cell & Environment*, 33, 959–980.
- Dunbar-Co, S., Sporck, M. J., & Sack, L. (2009). Leaf trait diversification and design in seven rare taxa of the Hawaiian *Plantago* radiation. *International Journal of Plant Sciences*, 170(1), 61–75.
- Fonseca, C. R., Overton, J. M., Collins, B., & Westoby, M. (2000). Shifts in trait-combinations along rainfall and phosphorus gradients. *Journal of Ecology*, 88(6), 964–977.
- Frenette-Dussault, C., Shipley, B., Léger, J. F., Meziane, D., & Hingrat, Y. (2012). Functional structure of an arid steppe plant community reveals similarities with Grime's C-S-R theory. *Journal of Vegetation Science*, 23(2), 208–222.
- Freschet, G. T., Cornelissen, J. H., Van Logtestijn, R. S., & Aerts, R. (2010). Evidence of the 'plant economics spectrum' in a subarctic flora. *Journal of Ecology*, 98(2), 362–373.
- Fyllas, N. M., Patino, S., Baker, T. R., Bielefeld Nardoto, G., Martinelli, L. A., Quesada, C. A., ... Santos, A. (2009). Basin-wide variations in foliar properties of Amazonian forest: Phylogeny, soils and climate. *Biogeosciences*, 6, 2677–2708.
- Garnier, E., Lavorel, S., Ansquer, P., Castro, H., Cruz, P., Dolezal, J., ... Grigulis, K. (2006). Assessing the effects of land-use change on plant traits, communities and ecosystem functioning in grasslands: A standardized methodology and lessons from an application to 11 European sites. *Annals of Botany*, 99(5), 967–985.
- Givnish, T. J., Montgomery, R. A., & Goldstein, G. (2004). Adaptive radiation of photosynthetic physiology in the Hawaiian lobeliads: Light regimes, static light responses, and whole-plant compensation points. *American Journal of Botany*, 91(2), 228–246.
- Guerin G. R., Wen H., & Lowe A. J. (2012). Leaf morphology shift linked to climate change. *Biology Letters*, 8, 882–886.
- Gutiérrez, A. G., & Huth, A. (2012). Successional stages of primary temperate rainforests of Chiloé Island, Chile. *Perspectives in Plant Ecology, Evolution and Systematics*, 14(4), 243–256.
- Guy, A. L., Mischkolz, J. M., & Lamb, E. G. (2012). Limited effects of simulated acidic deposition on seedling survivorship and root morphology of endemic plant taxa of the Athabasca Sand Dunes in well-watered greenhouse trials. *Botany*, 91(3), 176–181.
- Han, W., Chen, Y., Zhao, F. J., Tang, L., Jiang, R., & Zhang, F. (2012). Floral, climatic and soil pH controls on leaf ash content in China's terrestrial plants. *Global Ecology and Biogeography*, 21(3), 376–382.
- Han, W., Fang, J., Guo, D., & Zhang, Y. (2005). Leaf nitrogen and phosphorus stoichiometry across 753 terrestrial plant species in China. *New Phytologist*, 168(2), 377–385.
- Hao, G. Y., Sack, L., Wang, A. Y., Cao, K. F., & Goldstein, G. (2010). Differentiation of leaf water flux and drought tolerance traits in hemiepiphytic and non-hemiepiphytic *Ficus* tree species. *Functional Ecology*, 24(4), 731–740.
- Hoof, J., Sack, L., Webb, D. T., & Nilsen, E. T. (2008). Contrasting structure and function of pubescent and glabrous varieties of Hawaiian *Metrosideros polymorpha* (Myrtaceae) at high elevation. *Biotropica*, 40(1), 113–118.
- Powers, J. S., & Tiffin, P. (2010). Plant functional type classifications in tropical dry forests in Costa Rica: Leaf habit versus taxonomic approaches. *Functional Ecology*, 24(4), 927–936.
- Kattge, J., Knorr, W., Raddatz, T., & Wirth, C. (2009). Quantifying photosynthetic capacity and its relationship to leaf nitrogen content for global-scale terrestrial biosphere models. *Global Change Biology*, 15(4), 976–991.
- Kazakou, E., Vile, D., Shipley, B., Gallet, C., & Garnier, E. (2006). Co-variations in litter decomposition, leaf traits and plant growth in species from a Mediterranean old-field succession. *Functional Ecology*, 20(1), 21–30.
- Kichenin, E., Wardle, D. A., Peltzer, D. A., Morse, C. W., & Freschet, G. T. (2013). Contrasting effects of plant inter- and intraspecific variation on community-level trait measures along an environmental gradient. *Functional Ecology*, 27(5), 1254–1261.
- Kleyer, M., Bekker, R. M., Knevel, I. C., Bakker, J. P., Thompson, K., Sonnenschein, M., ... Klotz, S. R. G. M. (2008). The LEDA Traitbase: A database of life-history traits of the Northwest European flora. *Journal of Ecology*, 96(6), 1266–1274.
- Kraft, N. J., Valencia, R., & Ackerly, D. D. (2008). Functional traits and niche-based tree community assembly in an Amazonian forest. *Science*, 322(5901), 580–582.
- Kurokawa, H., & Nakashizuka, T. (2008). Leaf herbivory and decomposability in a Malaysian tropical rain forest. *Ecology*, 89(9), 2645–2656.
- Laughlin, D. C., Leppert, J. J., Moore, M. M., & Sieg, C. H. (2010). A multi-trait test of the leaf-height-seed plant strategy scheme with 133 species from a pine forest flora. *Functional Ecology*, 24(3), 493–501.
- Laughlin, D. C., Fule, P. Z., Huffman, D. W., Crouse, J., & Laliberté, E. (2011). Climatic constraints on trait-based forest assembly. *Journal of Ecology*, 99(6), 1489–1499.
- Louault, F., Pillar, V. D., Aufrere, J., Garnier, E., & Soussana, J. F. (2005). Plant traits and functional types in response to reduced disturbance in a semi-natural grassland. *Journal of Vegetation Science*, 16(2), 151–160.
- Markesteyn, L., Poorter, L., Paz, H., Sack, L., & Bongers, F. (2011). Ecological differentiation in xylem cavitation resistance is associated with stem and leaf structural traits. *Plant, Cell & Environment*, 34(1), 137–148.
- Martin, R. E., Asner, G. P., & Sack, L. (2007). Genetic variation in leaf pigment, optical and photosynthetic function among diverse phenotypes of *Metrosideros polymorpha* grown in a common garden. *Oecologia*, 151(3), 387–400.
- Medlyn, B. E., Badeck, F. W., De Pury, D. G. G., Barton, C. V. M., Broadmeadow, M., Ceulemans, R., ... Laitat, E. (1999). Effects of elevated [CO₂] on photosynthesis in European forest species: A meta-analysis of model parameters. *Plant, Cell & Environment*, 22(12), 1475–1495.
- Meir, P., Levy, P. E., Grace, J., & Jarvis, P. G. (2007). Photosynthetic parameters from two contrasting woody vegetation types in West Africa. *Plant Ecology*, 192(2), 277–287.
- Meir, P., Kruijt, B., Broadmeadow, M., Barbosa, E., Kull, O., Carswell, F., ... Jarvis, P. G. (2002). Acclimation of photosynthetic capacity to irradiance in tree canopies in relation to leaf nitrogen concentration and leaf mass per unit area. *Plant, Cell & Environment*, 25(3), 343–357.
- Messier, J., McGill, B. J., & Lechowicz, M. J. (2010). How do traits vary across ecological scales? A case for trait-based ecology. *Ecology Letters*, 13(7), 838–848.
- Meziane, D., & Shipley, B. (1999). Interacting determinants of specific leaf area in 22 herbaceous species: Effects of irradiance and nutrient availability. *Plant, Cell & Environment*, 22(5), 447–459.

- Milla, R., & Reich, P. B. (2011). Multi-trait interactions, not phylogeny, fine-tune leaf size reduction with increasing altitude. *Annals of Botany*, 107(3), 455–465.
- Minden, V., & Kleyer, M. (2011). Testing the effect–response framework: Key response and effect traits determining above-ground biomass of salt marshes. *Journal of Vegetation Science*, 22(3), 387–401.
- Minden, V., Andratschke, S., Spalke, J., Timmermann, H., & Kleyer, M. (2012). Plant trait–environment relationships in salt marshes: Deviations from predictions by ecological concepts. *Perspectives in Plant Ecology, Evolution and Systematics*, 14(3), 183–192.
- Müller, S. C., Overbeck, G. E., Pfadenhauer, J., & Pillar, V. D. (2007). Plant functional types of woody species related to fire disturbance in forest–grassland ecotones. *Plant Ecology*, 189(1), 1–14.
- Nakahashi, C. D., Frole, K., & Sack, L. (2005). Bacterial leaf nodule symbiosis in *Ardisia* (Myrsinaceae): Does it contribute to seedling growth capacity?. *Plant Biology*, 7(05), 495–500.
- Niinemets, Ü. (2001). Global-scale climatic controls of leaf dry mass per area, density, and thickness in trees and shrubs. *Ecology*, 82(2), 453–469.
- Ogaya, R., & Peñuelas, J. (2003). Comparative field study of *Quercus ilex* and *Phillyrea latifolia*: Photosynthetic response to experimental drought conditions. *Environmental and Experimental Botany*, 50(2), 137–148.
- Onoda, Y., Westoby, M., Adler, P. B., Choong, A. M., Clissold, F. J., Cornelissen, J. H., ... Fine, P. V. (2011). Global patterns of leaf mechanical properties. *Ecology Letters*, 14(3), 301–312.
- Ordóñez, J. C., van Bodegom, P. M., Witte, J. P. M., Bartholomeus, R. P., van Hal, J. R., & Aerts, R. (2009). Plant strategies in relation to resource supply in mesic to wet environments: Does theory mirror nature?. *The American Naturalist*, 175(2), 225–239.
- Pahl, A. T., Kollmann, J., Mayer, A., & Haider, S. (2013). No evidence for local adaptation in an invasive alien plant: Field and greenhouse experiments tracing a colonization sequence. *Annals of Botany*, 112(9), 1921–1930.
- Peco, B., de Pablos, I., Traba, J., & Levassor, C. (2005). The effect of grazing abandonment on species composition and functional traits: The case of dehesa grasslands. *Basic and Applied Ecology*, 6(2), 175–183.
- Peñuelas, J., Sardans, J., Llusia, J., Owen, S. M., Carnicer, J., Giambelluca, T. W., ... Niinemets, Ü. (2010). Faster returns on 'leaf economics' and different biogeochemical niche in invasive compared with native plant species. *Global Change Biology*, 16(8), 2171–2185.
- Pierce S., Brusa G., Sartori M., & Cerabolini B. E. L. (2012). Combined use of leaf size and economics traits allows direct comparison of hydrophyte and terrestrial herbaceous adaptive strategies. *Annals of Botany*, 109(5), 1047–1053.
- Pierce, S., Brusa, G., Vagge, I., & Cerabolini, B. E. (2013). Allocating CSR plant functional types: The use of leaf economics and size traits to classify woody and herbaceous vascular plants. *Functional Ecology*, 27(4), 1002–1010.
- Pierce, S., Ceriani, R. M., De Andreis, R., Luzzaro, A., & Cerabolini, B. (2007). The leaf economics spectrum of Poaceae reflects variation in survival strategies. *Plant Biosystems*, 141(3), 337–343.
- Pierce, S., Luzzaro, A., Caccianiga, M., Ceriani, R. M., & Cerabolini, B. (2007). Disturbance is the principal α -scale filter determining niche differentiation, coexistence and biodiversity in an alpine community. *Journal of Ecology*, 95(4), 698–706.
- Pillar, V. D., & Sosinski Jr, E. E. (2003). An improved method for searching plant functional types by numerical analysis. *Journal of Vegetation Science*, 14(3), 323–332.
- Prentice, I. C., Meng, T., Wang, H., Harrison, S. P., Ni, J., & Wang, G. (2011). Evidence of a universal scaling relationship for leaf CO₂ drawdown along an aridity gradient. *New Phytologist*, 190(1), 169–180.
- Preston, K. A., Cornwell, W. K., & DeNoyer, J. L. (2006). Wood density and vessel traits as distinct correlates of ecological strategy in 51 California coast range angiosperms. *New Phytologist*, 170(4), 807–818.
- Price, C. A., & Enquist, B. J. (2007). Scaling mass and morphology in leaves: An extension of the WBE model. *Ecology*, 88(5), 1132–1141.
- Pyankov, V. I., Kondratchuk, A. V., & Shipley, B. (1999). Leaf structure and specific leaf mass: The alpine desert plants of the Eastern Pamirs, Tadjikistan. *The New Phytologist*, 143(1), 131–142.
- Quero, J. L., Villar, R., Marañón, T., Zamora, R., Vega, D., & Sack, L. (2008). Relating leaf photosynthetic rate to whole-plant growth: Drought and shade effects on seedlings of four *Quercus* species. *Functional Plant Biology*, 35(8), 725–737.
- Quested, H. M., Cornelissen, J. H. C., Press, M. C., Callaghan, T. V., Aerts, R., Trosien, F., ... Jonasson, S. E. (2003). Decomposition of sub-arctic plants with differing nitrogen economies: A functional role for hemiparasites. *Ecology*, 84(12), 3209–3221.
- Reich, P. B., Oleksyn, J., & Wright, I. J. (2009). Leaf phosphorus influences the photosynthesis–nitrogen relation: A cross-biome analysis of 314 species. *Oecologia*, 160(2), 207–212.
- Reich, P. B., Tjoelker, M. G., Pregitzer, K. S., Wright, I. J., Oleksyn, J., & Machado, J. L. (2008). Scaling of respiration to nitrogen in leaves, stems and roots of higher land plants. *Ecology Letters*, 11(8), 793–801.
- Sack, L. (2004). Responses of temperate woody seedlings to shade and drought: Do trade-offs limit potential niche differentiation?. *Oikos*, 107(1), 110–127.
- Sack, L., & Frole, K. (2006). Leaf structural diversity is related to hydraulic capacity in tropical rain forest trees. *Ecology*, 87(2), 483–491.
- Sack, L., Tyree, M. T., & Holbrook, N. M. (2005). Leaf hydraulic architecture correlates with regeneration irradiance in tropical rainforest trees. *New Phytologist*, 167(2), 403–413.
- Sack, L., Cowan, P. D., Jaikumar, N., & Holbrook, N. M. (2003). The 'hydrology' of leaves: Co-ordination of structure and function in temperate woody species. *Plant, Cell & Environment*, 26(8), 1343–1356.
- Sack, L., Melcher, P. J., Liu, W. H., Middleton, E., & Pardee, T. (2006). How strong is intracanalopy leaf plasticity in temperate deciduous trees?. *American Journal of Botany*, 93(6), 829–839.
- Sandel, B., Corbin, J. D., & Krupa, M. (2011). Using plant functional traits to guide restoration: A case study in California coastal grassland. *Ecosphere*, 2(2), 1–16.
- Scherer-Lorenzen, M., Schulze, E. D., Don, A., Schumacher, J., & Weller, E. (2007). Exploring the functional significance of forest diversity: A new long-term experiment with temperate tree species (BIOTREE). *Perspectives in Plant Ecology, Evolution and Systematics*, 9(2), 53–70.
- Schweingruber, F. H., & Landolt, W. (2005). The xylem database. *Swiss Federal Research Institute WSL Updated*. Available at: <https://www.wsl.ch/en/services-and-products/software-websites-and-apps/the-xylem-database.html>
- Scoffoni, C., Pou, A., Aasamaa, K., & Sack, L. (2008). The rapid light response of leaf hydraulic conductance: New evidence from two experimental methods. *Plant, Cell & Environment*, 31(12), 1803–1812.
- Shipley, B. (1995). Structured interspecific determinants of specific leaf area in 34 species of herbaceous angiosperms. *Functional Ecology*, 9(2), 312–319.
- Shipley, B. (2002). Trade-offs between net assimilation rate and specific leaf area in determining relative growth rate: Relationship with daily irradiance. *Functional Ecology*, 16(5), 682–689.
- Shipley, B., & Lechowicz, M. J. (2000). The functional co-ordination of leaf morphology, nitrogen concentration, and gas exchange in 40 wetland species. *Ecoscience*, 7(2), 183–194.
- Shipley, B., & Parent, M. (1991). Germination responses of 64 wetland species in relation to seed size, minimum time to reproduction and seedling relative growth rate. *Functional Ecology*, 5(1), 111–118.
- Shipley, B., & Vu, T. T. (2002). Dry matter content as a measure of dry matter concentration in plants and their parts. *New Phytologist*, 153(2), 359–364.

- Soudzilovskaia, N. A., Elumeeva, T. G., Onipchenko, V. G., Shidakov, I. I., Salpagarova, F. S., Khubiev, A. B., ... Cornelissen, J. H. (2013). Functional traits predict relationship between plant abundance dynamic and long-term climate warming. *Proceedings of the National Academy of Sciences USA*, 110(45), 18180–18184.
- Spasojevic, M. J., & Suding, K. N. (2012). Inferring community assembly mechanisms from functional diversity patterns: The importance of multiple assembly processes. *Journal of Ecology*, 100(3), 652–661.
- Swaine, E. K. (2007). *Ecological and evolutionary drivers of plant community assembly in a Bornean rain forest* (PhD thesis), University of Aberdeen, Aberdeen.
- Tucker, S. S., Craine, J. M., & Nippert, J. B. (2011). Physiological drought tolerance and the structuring of tallgrass prairie assemblages. *Ecosphere*, 2(4), 1–19.
- van Bodegom, P. M., Sorrell, B. K., Oosthoek, A., Bakker, C., & Aerts, R. (2008). Separating the effects of partial submergence and soil oxygen demand on plant physiology. *Ecology*, 89(1), 193–204.
- van Bodegom, P. M., De Kanter, M., & Aerts, C. B. R. (2005). Radial oxygen loss, a plastic property of dune slack plant species. *Plant and Soil*, 271(1–2), 351–364.
- Vergutz, L., S. Manzoni, Porporato, A., Novais, R. F., & Jackson, R. B. (2012). A global database of carbon and nutrient concentrations of green and senesced leaves. Data set. Available on-line [http://daac.ornl.gov] from Oak Ridge National Laboratory Distributed Active Archive Center, Oak Ridge, Tennessee, U.S.A.
- Vile, D. (2005). *Significations fonctionnelle et écologique des traits des espèces végétales: Exemple dans une succession post-culturale méditerranéenne et généralisations* (PhD thesis, Montpellier 2).
- Von Holle, B., & Simberloff, D. (2004). Testing Fox's assembly rule: Does plant invasion depend on recipient community structure?. *Oikos*, 105(3), 551–563.
- Waite, M., & Sack, L. (2010). How does moss photosynthesis relate to leaf and canopy structure? Trait relationships for 10 Hawaiian species of contrasting light habitats. *New Phytologist*, 185(1), 156–172.
- Williams, M., Y. E. Shimabokuro, and E. B. Rastetter. "LBA-ECO CD-09 Soil and Vegetation Characteristics, Tapajos National Forest, Brazil." Oak Ridge National Laboratory Distributed Active Archive Center, Oak Ridge, Tennessee, USA. [http://dx.doi.org/10.3334/ORNLDAAC/1104\(2012\)](http://dx.doi.org/10.3334/ORNLDAAC/1104(2012)).
- Wilson, K. B., Baldocchi, D. D., & Hanson, P. J. (2000). Spatial and seasonal variability of photosynthetic parameters and their relationship to leaf nitrogen in a deciduous forest. *Tree Physiology*, 20(9), 565–578.
- Wirth, C., & Lichstein, J. W. (2009). The imprint of species turnover on old-growth forest carbon balances—insights from a trait-based model of forest dynamics. In C. Wirth, G. Gleixner, & M. Heimann (Eds.), *Old-growth forests* (pp. 81–113). Springer, Berlin, Heidelberg.
- Wright, I. J., Ackerly, D. D., Bongers, F., Harms, K. E., Ibarra-Manriquez, G., Martinez-Ramos, M., ... Poorter, L. (2006). Relationships among ecologically important dimensions of plant trait variation in seven Neotropical forests. *Annals of Botany*, 99(5), 1003–1015.
- Wright, I. J., Reich, P. B., Westoby, M., Ackerly, D. D., Baruch, Z., Bongers, F., ... Flexas, J. (2004). The worldwide leaf economics spectrum. *Nature*, 428(6985), 821.
- Wright, S. J., Kitajima, K., Kraft, N. J., Reich, P. B., Wright, I. J., Bunker, D. E., ... Engelbrecht, B. M. (2010). Functional traits and the growth–mortality trade-off in tropical trees. *Ecology*, 91(12), 3664–3674.
- Yguel, B., Bailey, R., Tosh, N. D., Vialatte, A., Vasseur, C., Vitrac, X., ... Prinzing, A. (2011). Phytophagy on phylogenetically isolated trees: Why hosts should escape their relatives. *Ecology Letters*, 14(11), 1117–1124.