



Universiteit
Leiden
The Netherlands

How Sedna and family were captured in a close encounter with a solar sibling

Jilkova, L.; Portegies Zwart, S.F.; Pijloo, J.T.; Hammer, M.

Citation

Jilkova, L., Portegies Zwart, S. F., Pijloo, J. T., & Hammer, M. (2015). How Sedna and family were captured in a close encounter with a solar sibling. *Monthly Notices Of The Royal Astronomical Society*, 453(3), 3157-3162. doi:10.1093/mnras/stv1803

Version: Publisher's Version

License: [Licensed under Article 25fa Copyright Act/Law \(Amendment Taverne\)](#)

Downloaded from: <https://hdl.handle.net/1887/3198545>

Note: To cite this publication please use the final published version (if applicable).

How Sedna and family were captured in a close encounter with a solar sibling

Lucie Jílková,^{1★} Simon Portegies Zwart,^{1★} Tjibbaria Pijloo^{1,2} and Michael Hammer³

¹Leiden Observatory, Leiden University, PO Box 9513, NL-2300 RA Leiden, the Netherlands

²Department of Astrophysics/IMAPP, Radboud University, PO Box 9010, NL-6500 GL Nijmegen, the Netherlands

³Cornell University, 614 Space Sciences Building, Ithaca, NY 14853, USA

Accepted 2015 August 4. Received 2015 August 4; in original form 2015 March 9

ABSTRACT

The discovery of 2012 VP₁₁₃ initiated the debate on the origin of the Sedna family of planetesimals in orbit around the Sun. Sednitos roam the outer regions of the Solar system between the Egeworth–Kuiper belt and the Oort Cloud, in extraordinary wide ($a > 150$ au) orbits with a large perihelion distance of $q > 30$ au compared to the Earth's ($a \equiv 1$ au and eccentricity $e \equiv (1 - q/a) \simeq 0.0167$ or $q \simeq 1$ au). This population is composed of a dozen objects, which we consider a family because they have similar perihelion distance and inclination with respect to the ecliptic $i = 10^\circ\text{--}30^\circ$. They also have similar argument of perihelion $\omega = 340^\circ \pm 55^\circ$. There is no ready explanation for their origin. Here we show that these orbital parameters are typical for a captured population from the planetesimal disc of another star. Assuming that the orbital elements of Sednitos have not changed since they acquired their orbits, we reconstruct the encounter that led to their capture. We conclude that they might have been captured in a near miss with a $1.8 M_\odot$ star that impacted the Sun at $\simeq 340$ au at an inclination with respect to the ecliptic of $17^\circ\text{--}34^\circ$ with a relative velocity at infinity of $\sim 4.3 \text{ km s}^{-1}$. We predict that the Sednitos region is populated by 930 planetesimals and the inner Oort Cloud acquired ~ 440 planetesimals through the same encounter.

Key words: celestial mechanics – planetary systems – minor planets, asteroids: general – minor planets, asteroids: individual: Sedna, 2012 VP₁₁₃ – open clusters and associations.

1 INTRODUCTION

Upon its discovery, 90377 Sedna (Brown, Trujillo & Rabinowitz 2004) was proposed to originate from the Edgeworth–Kuiper belt; a population of rocks in an almost planar disc aligned with the ecliptic and much closer to the Sun ($q = 30\text{--}50$ au) than the Oort Cloud. According to this model, the violent and rather sudden migration of Uranus and Neptune would have excited the cold Kuiper belt (Brasser et al. 2012). This reorganization of the outer giant planets would have initiated the hot Kuiper belt (Levison et al. 2008) that, due to the long local relaxation time, cools down only very slowly (Punzo, Capuzzo-Dolcetta & Portegies Zwart 2014). Subsequent chaotic diffusion (Morbideilli et al. 2008), perturbations in the Sun's birth environment in close flybys (Davies et al. 2014), or more distant encounters could have caused further migration of the Kuiper belt objects to orbits similar to that of Sedna (Brasser et al. 2012).

Recently, Trujillo & Sheppard (2014) discovered the object 2012 VP₁₁₃, a second member of the inner Oort Cloud, which they defined as a family of planetesimals with $q \gtrsim 50$ au and $a \approx 150\text{--}$

1500 au. They furthermore identified a population of planetesimals between the Edgeworth–Kuiper belt and the Oort Cloud that share similar orbital elements (see also de la Fuente Marcos & de la Fuente Marcos 2014), namely the large perihelion and semimajor axis ($q > 30$ au and $a > 150$ au, respectively), inclination with respect to the ecliptic ($i = 10^\circ\text{--}30^\circ$), and the argument of perihelion ($\omega = 340^\circ \pm 55^\circ$). Currently 13 such objects have been observed in the outer Solar system and it was suggested that their characteristics resulted from a common origin (de la Fuente Marcos & de la Fuente Marcos 2014; Trujillo & Sheppard 2014). Here we consider this group of object a family which we call *Sednitos*.

The perihelion distances of Sedna and 2012 VP₁₁₃ are too large for any Kuiper belt object and their aphelion distances are too short for them to be Oort Cloud objects (Brasser & Schwamb 2015). It is therefore hard to explain them as members of either population. In principle, chaotic diffusion could cause sufficient internal migration, but at the distance of Sedna, the time-scale for this process exceeds the age of the Solar system (Sussman & Wisdom 1988). If Sedna stood alone, such an exotic explanation could be satisfactory. However, this cannot explain the entire population of the inner Oort Cloud which, when taking selection effects into account, amounts to 430^{+400}_{-240} members brighter than $r = 24.3$ mag (Trujillo &

* E-mail: jilkova@strw.leidenuniv.nl (LJ); spz@strw.leidenuniv.nl (SPZ)

Sheppard 2014). In the model of Brasser et al. (2012), the inner Oort objects are scattered from the Kuiper belt and decoupled to larger pericentres by perturbations in the Sun’s birth cluster. The size of the population produced by this mechanism is consistent with the one predicted from observations.

According to an alternative scenario, Sedna could have been captured from the outer disc of a passing star, as suggested by Morbidelli & Levison (2004) and Kenyon & Bromley (2004). They showed that capturing a planetesimal into a Sedna-like orbit is possible, but they did not carry out a detailed parameter space study. The model of Kenyon & Bromley (2004) could account for at most 10 per cent of the Sednitos and it was tuned at producing the outer edge of the Edgeworth–Kuiper belt at the currently observed 50 au; however, this is inconsistent with the *Nice* model that requires an edge at ~ 35 au (Gomes, Morbidelli & Levison 2004). The capture of planetesimals by the Solar system was further studied by Levison et al. (2010), who simulated the Sun’s birth cluster considering the transfer of planetesimals among stars. However, the study was aimed to explain the origin of the Oort Cloud and the orbits of most of the captured objects have large semimajor axes ($a \gtrsim 10^3$ au) and perihelia ($q \gtrsim 10^2$ au), not representative for Sedna and 2012 VP₁₁₃.

1.1 Argument of perihelion of Sednitos

Sednitos are characterized by a clustered distribution of their observed argument of perihelion ω . The precession period of ω depends on the semimajor axis, eccentricity, and inclination of the precessing orbit. The precession periods for all Sednitos excluding Sedna range from about 40 Myr up to 650 Myr, while Sedna has the longest precession period of about 1.5 Gyr (Brasser, Duncan & Levison 2006; Gomes, Matese & Lissauer 2006; Trujillo & Sheppard 2014). Therefore, the clustering of ω must have happened relatively recently (less than few Myr ago) or a dynamical mechanism must have been constraining the distribution of ω since it was established. Trujillo & Sheppard (2014) suggested that an outer Solar system perturber of $5\text{--}15 M_{\oplus}$ orbiting the Sun between 200 and 300 au is restricting the Sednitos’ evolution in ω by the Kozai–Lidov mechanism (Kozai 1962; Lidov 1962). Based on further analysis of the Sednitos’ orbital elements, de la Fuente Marcos & de la Fuente Marcos (2014) suggested that at least two planetary-mass trans-Neptunian perturbers exist at approximately 200 and 250 au.

The perturbing object (possibly more than one object) is assumed to be on a low-inclination almost circular orbit. When, in such a configuration, the ratio of semimajor axis with respect to the perturbed objects (i.e. Sednitos) is close to 1, the relative argument of perihelion of the perturbed and perturbing orbits can librate around 0° or 180° due to the Kozai–Lidov mechanism (see the Extended materials of Trujillo & Sheppard 2014, for an example where ω of 2012 VP₁₁₃ librates in the range $0^\circ \pm 60^\circ$). However, depending on the initial relative inclination and argument of perihelion of the perturbing and perturbed orbits, the argument of perihelion can also circulate (i.e. periodically change values from -180° to 180°). The libration around 0° will occur if the initial relative ω ranges from -90° to 90° (e.g. Mardling 2007). Therefore, the ω of Sednitos relative to the perturber ($\omega - \omega_{\text{perturber}}$) needs to be constrained at the beginning of the dynamical interaction with the perturber. Brasser et al. (2006) showed that for the Sedna-like orbits produced during the early evolution of the Solar system, when the Sun was still residing in its birth star cluster, preferentially $\omega = 0^\circ$ or 180° . This mechanism could therefore explain the initial clustering of Sednitos’ argument of perihelion, although it is not clear why the

orbits initially obtained ω about 0° and not 180° (see also Trujillo & Sheppard 2014).

The presence of the perturbing object(s) in the outer Solar system is currently the only mechanism suggested to explain how the clustering in ω is preserved on time-scales longer than the precession periods of Sednitos. At the same time, Iorio (2014) ruled out the presence of super-Earth planet of $2\text{--}15 M_{\oplus}$ with $a \approx 200\text{--}300$ au using the current constraints on the anomalous secular precession of the argument of perihelion of some of the known planets in the Solar system. Therefore, the existence of an outer planet is still under debate.

Irrespective of the mechanism that preserves the clustering of the argument of perihelion, we present that such clustering is a general characteristic of the population transferred during a stellar encounter. The constrained distribution in ω can then be shepherded by some other process. We argue that Sednitos are found in the Parking zone of the Solar system where their semimajor axis and eccentricity have been unaffected once the Sun left its birth cluster (Portegies Zwart & Jílková 2015). We therefore use the current semimajor axis and eccentricity to constrain the encounter that might have introduced the Sednitos into the Solar system.

2 METHODS

The encounter between the Sun and another star, here called Q, with a planetesimal disc can be simulated by integrating the equations of motion of the two stars using a symplectic N -body code. We use HUAYNO (Pelupessy, Jänes & Portegies Zwart 2012) for this. As long as the two stars are well separated (at least three times the disc size) we integrate the planetesimals using SAKURA (Gonçalves Ferrari, Boekholt & Portegies Zwart 2014), in which Kepler’s equations are solved in the potential of the Q coupled with the perturbations from the Sun. The planetesimals are represented by zero-mass particles which do not affect each other and neither the motion of the two stars, while the planetesimals themselves are affected by the two stars. Both integrators (HUAYNO and SAKURA) are coupled via BRIDGE (Fujii et al. 2007), which is an extension of the mixed variable symplectic scheme (Wisdom & Holman 1991) and is used in this context to couple two different dynamical regimes within one self-gravitating system. The coupling of codes is realized using the Astronomical Multi-purpose Software Environment (Portegies Zwart et al. 2013).¹ When the stars move close enough that the planetesimals orbits can no longer be considered Keplerian – i.e. the motion is no longer dominated by Q and both stars have a substantial influence on the orbits – the planetesimals are integrated directly using HUAYNO. We introduce this transition from hybrid to direct integration when the time since the beginning of the encounter equals half of the time for the two stars to reach their closest approach (which always results to a separation of the two stars larger than three times the disc size).

The initial conditions for the encounter are as follows. The distance between the two stars is determined by the condition that the magnitude of the gravitational force from the Sun at the outer edge of Q’s disc equals 10 per cent of Q’s force. We tested that increasing the initial separation does not change the results. In all our calculations we adopted the mass of the Sun of $1 M_{\odot}$. Planetesimals in Q’s disc have initially planar distribution and their radial distance from Q, r , follows a uniform random distribution, i.e. the surface density profile of Q’s disc $\propto (1/r)$. However, since the planetesimals are

¹ All source code is available at <http://amusecode.org>

Table 1. Reconstructed encounter parameters for the star that delivered the Sednitos into the Solar system. The first column lists the five parameters of the encounter: M_Q is the mass of the impactor star Q, q_Q the closest approach of the Sun and Q, e_Q the eccentricity of their orbit, i_Q the inclination of the orbital plane with respect to Q's disc and ω_Q the argument of periastron. We further give the impact parameter b , the relative velocity of the encounter at infinity v and the limits for the outer edges for Q's and Sun's disc, $r_{\max, Q}$ and $r_{\max, \odot}$, respectively. The orientation of the encounter with respect to the ecliptic is specified by i_{enc} and ω_{enc} . The second column gives the range considered in the Markov chain simulations, followed by the range of parameters that led to a satisfactory solution. The parameters of the preferred encounter are listed in the right most column (we give the constrained range for i_{enc} and ω_{enc} together with the individual values used in the presented example in the parenthesis).

Parameter	Parameter range	Viable range	Preferred encounter
M_Q	0.2–2.0 M_\odot	1.0–2.0 M_\odot	1.8 M_\odot
q_Q	200–393 au	210–320 au	227 au
e_Q	1.001–4.0	1.9–3.8	2.6
i_Q	0°–180°	2°–44°	35°
ω_Q	0°–180°	0°–180°	175°
b	265–2071 au	280–450 au	340 au
v	0.4–6.0 km s ^{−1}	3.1–5.4 km s ^{−1}	4.3 km s ^{−1}
$r_{\max, Q}$		130–200 au	$\gtrsim 161$ au
$r_{\max, \odot}$			$\lesssim 70$ au
i_{enc}		0°–70°	17°–34° (28°)
ω_{enc}		0°–360°	154°–197° (170°)

represented by zero-mass particles, the surface density profile can be adjusted in post-processing (see Section 3). The planetesimals are initially on circular orbits. The inner edge of Q's disc is 10 au. We set the upper limit on the outer edge of Q's disc, $r_{\max, Q}$, to 200 au and determine the actual value from the minimal requirement of producing planetesimals in the range of $q = 30$ –85 au; we do the same for Sun's disc, see below.

The encounter between the Sun and Q is characterized by the five parameters (also listed in Table 1) – the mass of the encountering star, M_Q , the closest approach of the stars, q_Q , the eccentricity of the orbit, e_Q , the inclination of the encounter plane with respect to Q's disc, i_Q , and the argument of periastron of the orbit, ω_Q . We have the computer map this parameter space automatically using the affine-invariant, parallel stretch-move algorithm for Markov Chain Monte Carlo (Hastings 1970) with specific optimizations (Goodman & Weare 2010) using EMCEE (Foreman-Mackey et al. 2013).

We run more than 10 000 realizations of possible encounters. Each calculation is performed with up to 20 000 particles (in chunks of 500) in the planetesimal disc until the number of particles captured by the Sun amounts to at least 13 objects with a perihelion distance between 30 and 85 au. To account for the observability of orbits with different eccentricities, we weigh each particle by the time it spends within 85 au from the Sun measured as a fraction of the orbital period. The weight w is calculated using the mean anomaly at the 85 au from the Sun, $M(85 \text{ au})$,

$$w = \begin{cases} M(85 \text{ au})/\pi & \text{if aphelion} > 85 \text{ au,} \\ 1 & \text{if } 30 \text{ au} < \text{aphelion} < 85 \text{ au.} \end{cases} \quad (1)$$

This weighting favours finding relatively low-eccentricity orbits with a small perihelion, as is consistent with how the Sednitos were discovered (Trujillo & Sheppard 2014).

The resulting distribution of the planetesimals in semimajor axis and eccentricity is subsequently compared with the observed dozen Sednitos listed in table 2 of Trujillo & Sheppard (2014) and table 1 in de la Fuente Marcos & de la Fuente Marcos (2014) for 2003 SS₄₂₂. The 13 observed Sednitos provide only low number statistics and we investigate what constraints can we draw based on the limited data. As a first step in the statistical comparison, we perform a consistency analysis between the simulated objects and the observed Sednitos under the hypothesis that the latter is a random sub-sample of the former using multivariate analysis. We calculate the ranking of the Henze statistic (Henze 1988; Koen & Siluyele 2007) using the nearest neighbours and based on 500 randomly pooled data (see Koen & Siluyele 2007, for more detail) and we require the final rank (or the p -value) of the actual data sets to be >0.05 to consider the samples consistent. In these cases, we measure the separation distance between these two distributions in the plane of a versus e using the Hellinger distance of binned kernel-smoothed distributions (Pak & Basu 1998). We use a grid of 20×20 bins scaled on the observed data with a symmetric Gaussian kernel with a relative width (corresponding to the standard deviation) of 0.08. The separation that emerges from this analysis is used as the posterior probability in the Markov Chain Monte Carlo.

3 RESULTS

We identify a region in the five-dimensional encounter parameter space where the a versus e distribution of the simulated transferred particles is statistically indistinguishable from the observed distribution. We constrain the viable range by the Hellinger distance <0.6 . We further divide the Sednitos a versus e region into three sections: the inner Oort Cloud ($70 \text{ au} < q < 85 \text{ au}$), the region where so far no objects have been observed ($50 \text{ au} < q < 75 \text{ au}$, or $q < 50 \text{ au}$ and $a < 140 \text{ au}$), and the remaining region. For the encounters in the viable range, we require that the weighted number of captured particles is higher than 1.0 in the inner Oort Cloud region, and lower than 6.0 for the region without any observed objects.

The limits of such viable range for individual parameters are listed in Table 1. We also select a preferred encounter as a representative example of the viable range; its parameters are also listed Table 1. The Sednitos' distribution produced by the preferred encounter compares well with the observed one: the rank of the Henze statistics is 0.5 and the distance between the distributions is ~ 0.5 (here 0 corresponds to identical binned kernel-smoothed distributions, while 1 corresponds to distributions with no overlap; see Section 2).

After the Markov chain calculation we constrain the orientation of Q's disc with respect to the ecliptic. While the inclination i_Q and the argument of periastron ω_Q of the orbital plane of the encounter with respect to Q's disc are constrained by the Markov chain calculations, the orientation of the orbital plane with respect to the ecliptic is unconstrained. We constrain the orientation using two-sample Kolmogorov–Smirnov tests comparing each of the distributions in the argument of perihelion, inclination and the longitude of the ascending node of the observed and simulated Sednitos. The clustering of inclination and argument of perihelion is a general feature of the transferred population and an orientation of the coordinate system where the simulated distributions are consistent with the observed ones is found for almost all viable encounters. Using a grid with a step size of 2° for each of the three Euler angles, we rotate the coordinate system centred on the Sun until the p -value > 0.05 for each of the three compared distributions. We derive the inclination i_{enc} and the argument of periastron ω_{enc} of the




Figure 1. Distribution of the planetesimals captured around the Sun during the preferred encounter with star Q (rightmost column of Table 1). Along the x -axis we present the semimajor axis a of the captured planetesimals. The top panel gives the inclination i along the y -axis, and the bottom panel gives the eccentricity e . The colour scale maps the initial radius in Q's disc, $r_{\text{ini},Q}$. Note that the simulated particles are not weighted here. The red diamonds give the observed positions of the Sednitos.

orbit of the encounter with respect to the ecliptic (the longitude of ascending node is a free parameter due to assumed symmetry of the Sun's disc, see below). This procedure results in the inclination i_{enc} and the argument of perihelion ω_{enc} of the encounter with respect to the ecliptic for individual encounters. These parameters are typically constrained within intervals of $\pm 10^\circ$ and $\pm 20^\circ$, for i_{enc} and ω_{enc} , respectively. We summarize the values in Table 1 (note that while always constrained within limited intervals for individual encounters, ω_{enc} have values in a wide range, unlike i_{enc}).

After constraining the initial conditions that reproduce the Sednitos, we rerun the preferred encounter with 100 000 particles in the disc around the encountering star and 100 000 particles around the Sun. The disc of the Sun extends from 1 to 200 au. Some perturbed planetesimals of such disc are members of a native population of Sednitos; we compare this population with the transferred population below. The results of this calculation are presented in Figs 1 and 2. In Fig. 1, we compare the observed Sednitos with the captured planetesimals from our best reconstruction of the encounter. The orbital distributions of the native and the captured planetesimals are presented in Fig. 2.

To estimate the number of planetesimals in the captured and the perturbed native population of Sednitos, we adopt a surface density profile $\propto r^{-3/2}$ and a mass of $10^{-3} M_\odot$ for both discs. We further assume that 10 per cent of such discs is in the form of Sedna-mass

objects (for which we assume 2×10^{21} kg). In that case, the Sun captured a total of ~ 2600 planetesimals, 884 of which accreted within the orbit of Neptune (with $q < 30$ au), but most of these are probably ejected by interacting with the planets. A total of 936 planetesimals are captured in orbits similar to the observed Sednitos ($q = 30\text{--}50$ au or $q = 75\text{--}85$ au and $a > 150$ au), and 441 in region between $q = 50$ and 75 au. The inner Oort Cloud ($q > 75$ au, $150 < a \lesssim 1500$ au) acquired 434 planetesimals, which is consistent with estimates of the current population of 430^{+400}_{-240} (Trujillo & Sheppard 2014). This would require the planetesimal disc of the encountering star to extend at least to 161 au, which is a reasonable disc size for an $\sim 1.8 M_\odot$ star (Booth et al. 2013).

If before the encounter the Sun's disc extended beyond ~ 90 au, some of its planetesimals are perturbed to a and e consistent with those of the observed Sednitos. Assuming the same surface density profile as for Q's disc, 307 Sun's planetesimals would be perturbed in orbits similar to the observed Sednitos ($q = 30\text{--}50$ au or $q = 75\text{--}85$ au and $a > 150$ au), 169 planetesimals would be scattered in the inner Oort Cloud ($q > 75$ au, $150 < a \lesssim 1500$ au), and the region between $q = 50$ and 75 au would be populated by about 319 native scattered planetesimals. We use the best encounter parameters for producing the Sednitos from Table 1 to calculate how many of the Sun's planetesimals would be transferred to the encountering star. If the solar disc extended to 90 au, it would have lost ~ 2.3 per cent of its planetesimals and ~ 92 per cent of those were captured by the other star. All the lost planetesimals originate from $a > 70$ au. In the right-hand panel of Fig. 2, we present the distributions of orbital parameters of the Q's own disc particles, and those of the planetesimals it stole from the Solar system. These captured objects are in rather curious orbits in the outer parts of the disc around the other star. Their inclination is about 14° with respect to Q's planetesimal disc and their argument of periastron is clustered around $0^\circ \pm 50^\circ$.

4 DISCUSSION

During the encounters not only Q's disc is perturbed, but at the same time Q also perturbs the Sun's disc. In particular, the preferred encounter (Table 1) excites the Sun's disc beyond ~ 30 au – see the left-hand panel of Fig. 2 – in agreement with the disc truncation radius estimate of Kobayashi & Ida (2001). Interestingly, the *Nice* model (Gomes et al. 2005; Morbidelli et al. 2005; Tsiganis et al. 2005) requires a truncation of the planetesimal disc at ~ 35 au (Gomes et al. 2004). Because both values are sufficiently close to be causal and a later subsequent encounter that would truncate the disc at 35 au would also annihilate the population of Sednitos, the capture must have happened before the resonant planetary swap.

The observed Sednitos cluster in the argument of perihelion around $\omega = 340^\circ \pm 55^\circ$ (de la Fuente Marcos & de la Fuente Marcos 2014; Trujillo & Sheppard 2014). Such clustering is a general characteristic of an exchanged population. As discussed in Section 1.1, the secular evolution due to the giant planets would cause a precession of ω on time-scales shorter than the age of the Solar system. If the clustering of ω is real, i.e. it is not a result of an observational bias (see below for more discussion on this issue), a mechanism preserving the distribution of ω is needed. The only scenario suggested so far involves a distant planetary-mass object (possibly more than one object) that causes libration of ω through the Kozai–Lidov mechanism (Gomes et al. 2006; de la Fuente Marcos & de la Fuente Marcos 2014; Trujillo & Sheppard 2014). Formation channels for such a super-Earth-mass planet were investigated by Kenyon & Bromley (2015), who analysed three mechanisms: planetary




Figure 2. Orbital distributions of planetesimals for the Sun (left) and for the encountering star Q (right) using the preferred encounter parameters (see Table 1). The top panels give inclination i as a function of semimajor axis a , the bottom panels give the orbital eccentricity e . The red (dark shade) bullets give the orbital distributions of the planetesimals native of the Sun (assuming its disc extended to 90 au), the light blue (light shade) bullets are native to Q. Both initial planetesimal discs are strongly perturbed beyond about 30 au, but within this distance they are hardly affected (see also Kobayashi & Ida 2001). Note that the simulated particles are not weighted here.

migration from the inner disc, scattering from the inner disc, and *in situ* formation. All three mechanisms require a disc extending up to the orbit of the planet – disc of gas or small planetesimals circularizes the orbit in the former two scenarios; while the later mechanism requires a reservoir of solid material with a mass $\approx 15 M_{\oplus}$ to form a planet at $a \lesssim 300$ au. However, disc and orbits of any objects beyond ~ 30 au would have been substantially perturbed by the encounter that would deposit Sednitos (see Section 3 and Fig. 2). Hence, the capture scenario as presented here appears inconsistent with the presence of the outer perturbing planet(s). The outer companion might have formed later than the Sednitos were transferred, e.g. by a capture of a free floating planet (Perets & Kouwenhoven 2012). However, in such case it is difficult to explain how the planet acquired the predicted almost circular low-inclination orbit.

Another caveat concerns our assumption that Sednitos are found in the parking zone of the Solar system (Portegies Zwart & Jílková 2015), i.e. that their eccentricities and semimajor axes have been the same since when they acquired their orbits. However if the outer perturber is present, the Kozai–Lidov oscillations it induces will also affect the eccentricity and semimajor axis of the Sednitos. The orbits of the Sednitos could also have been affected by encounters that occurred in the Sun’s birth cluster after their delivery. An encounter as close as we require to deliver the Sednitos (210–320 au, see Table 1) may have caused the Sun to escape its birth cluster, after which it becomes extremely unlikely to have further close encounters.

It should also be noted that the clustering in ω might not be a real dynamical feature of Sednitos. Trujillo & Sheppard (2014) explored possible observational biases and did not identify any that would lead to discovering objects clustered around $\omega = 0^\circ$. Nevertheless, 13 observed objects still provide only small number statistics and therefore we also discuss the possibility that the clustering in ω is not a real feature of the Sednitos family. The capture mechanism can still explain the existence of a population of objects in the inner Oort Cloud. The constraints on the population we find are determined by the encounter that is calibrated to deliver the Sednitos assuming that it is a family. Even if this assumption is wrong, and the 13 objects are not all part of the same family, Sedna and 2012 VP₁₁₃ can still be explained by a capture.

5 CONCLUSIONS

The origin of the inner Oort Cloud of the Solar system, which is defined as family of planetesimals with $q \gtrsim 50$ au and $a \approx 150$ –1500 au (Trujillo & Sheppard 2014), and which currently includes two observed objects – Sedna (Brown et al. 2004) and 2012 VP₁₁₃ (Trujillo & Sheppard 2014) – is still not well understood. Here we investigate the scenario where the inner Oort Cloud was captured from another star during a close encounter that occurred when both stars, the Sun and its sibling, were still members of their birth cluster (Kenyon & Bromley 2004; Morbidelli & Levison 2004; Levison et al. 2010). We assume that there are 13 extrasolar objects

currently observed in the outer Solar system (with $q > 30$ au, $a > 150$ au), which also share similar inclinations and argument of perihelion ($i = 10^\circ\text{--}30^\circ$, $\omega = 340^\circ \pm 55^\circ$, de la Fuente Marcos & de la Fuente Marcos 2014; Trujillo & Sheppard 2014), which we call Sednitos. Assuming that the orbits of Sednitos have not changed since they were acquired, we reconstruct the encounter that lead to their capture. The population of objects transferred from a planetesimal disc of the other star during the encounter has in general specific distributions of orbital elements around the star to which it was transferred to. We use this feature of the captured population and we carry out a Markov Chain Monte Carlo search of the parameter space typical for stellar encounters expected in the Sun's birth cluster. We provide constraints on the encounters that result in a population of the planetesimals transferred to the Solar system that is consistent with the observed objects.

Understanding the origin of Sednitos and testing the theories for an outer planetary-mass object requires additional observations. The Gaia astrometric mission is expected to discover ~ 50 objects in the outer Solar system. Being a solar sibling (Portegies Zwart 2009), the encountering star may also be discovered in the coming years in the Gaia catalogues. Having been formed in the same molecular cloud, one naively expects that the chemical composition of this star is similar to that of the Sun (Brown, Portegies Zwart & Bean 2010). Finding back our own planetesimals in the predicted orbits around this sibling (see Fig. 2) would expose the accused robber and would put strong constraints on the extend of the Sun's planetesimal disc. However, by now the other star has probably turned into a $\gtrsim 0.6 M_\odot$ carbon–oxygen white dwarf, which for an $\sim 1.8 M_\odot$ star happens within 2 Gyr. In that case, our stolen stones are probably lost to become free floating planetesimals due to the copious mass-loss in the post-asymptotic giant branch phase of the host (Veras et al. 2011).

ACKNOWLEDGEMENTS

We thank Inti Pelulessy, Tjarda Boekholt, Adrian Hamers and Jarle Brinchmann for enriching discussions. We also thank the referee, Hal Levison, for his useful comments which helped to improve the manuscript. We acknowledge the Leiden/ESA Astrophysics Program for Summer Students (LEAPS) for the support that made this work possible. This work was supported by the Interuniversity Attraction Poles Programme initiated by the Belgian Science Policy Office (IAP P7/08 CHARM) and by the Netherlands Research Council NWO (grants 643.200.503, 639.073.803 and 614.061.608) and by the Netherlands Research School for Astronomy (NOVA). The numerical computations were carried out using the Little Green Machine at Leiden University, and the distributed calculations were performed with the grid-version of *AMUSE* which was developed with support of the NLeSC.

REFERENCES

Booth M. et al., 2013, *MNRAS*, 428, 1263
 Brasser R., Schwamb M. E., 2015, *MNRAS*, 446, 3788
 Brasser R., Duncan M. J., Levison H. F., 2006, *Icarus*, 184, 59

Brasser R., Duncan M. J., Levison H. F., Schwamb M. E., Brown M. E., 2012, *Icarus*, 217, 1
 Brown M. E., Trujillo C., Rabinowitz D., 2004, *ApJ*, 617, 645
 Brown A. G. A., Portegies Zwart S. F., Bean J., 2010, *MNRAS*, 407, 458
 Davies M. B., Adams F. C., Armitage P., Chambers J., Ford E., Morbidelli A., Raymond S. N., Veras D., 2014, in Beuther H., Klessen R. S., Dullemond C. P., Henning T., eds, *Protostars and Planets VI*. Univ. Arizona press, Tucson, AZ, p. 787
 de la Fuente Marcos C., de la Fuente Marcos R., 2014, *MNRAS*, 443, L59
 Foreman-Mackey D., Hogg D. W., Lang D., Goodman J., 2013, *PASP*, 125, 306
 Fujii M., Iwasawa M., Funato Y., Makino J., 2007, *PASJ*, 59, 1095
 Gomes R. S., Morbidelli A., Levison H. F., 2004, *Icarus*, 170, 492
 Gomes R., Levison H. F., Tsiganis K., Morbidelli A., 2005, *Nature*, 435, 466
 Gomes R. S., Matese J. J., Lissauer J. J., 2006, *Icarus*, 184, 589
 Gonçalves Ferrari G., Boekholt T., Portegies Zwart S. F., 2014, *MNRAS*, 440, 719
 Goodman J., Weare J., 2010, *Commun. Appl. Math. Comput. Sci.*, 5, 65
 Hastings W. K., 1970, *Biometrika*, 57, 97
 Henze N., 1988, *Ann. Stat.*, 16, 772
 Iorio L., 2014, *MNRAS*, 444, L78
 Kenyon S. J., Bromley B. C., 2004, *Nature*, 432, 598
 Kenyon S. J., Bromley B. C., 2015, *ApJ*, 806, 42
 Kobayashi H., Ida S., 2001, *Icarus*, 153, 416
 Koen C., Siluyele I., 2007, *MNRAS*, 377, 1281
 Kozai Y., 1962, *AJ*, 67, 591
 Levison H. F., Morbidelli A., Van Laerhoven C., Gomes R., Tsiganis K., 2008, *Icarus*, 196, 258
 Levison H. F., Duncan M. J., Brasser R., Kaufmann D. E., 2010, *Science*, 329, 187
 Lidov M. L., 1962, *Planet. Space Sci.*, 9, 719
 Mardling R. A., 2007, *MNRAS*, 382, 1768
 Morbidelli A., Levison H. F., 2004, *AJ*, 128, 2564
 Morbidelli A., Levison H. F., Tsiganis K., Gomes R., 2005, *Nature*, 435, 462
 Morbidelli A., Levison H. F., Van Laerhoven C., Gomes R., Tsiganis K., 2008, in *AAS/Division of Dynamical Astronomy Meeting Vol. 39, Chaotic Capture of Planetesimals into Regular Regions of the Solar System. I: The Kuiper Belt*. Am. Astron. Soc., Washington, DC, p. 12.04
 Pak J. P., Basu A., 1998, *Ann. Stat.*, 50, 503
 Pelulessy F. I., Jänes J., Portegies Zwart S., 2012, *New Astron.*, 17, 711
 Perets H. B., Kouwenhoven M. B. N., 2012, *ApJ*, 750, 83
 Portegies Zwart S. F., 2009, *ApJ*, 696, L13
 Portegies Zwart S. F., Jílková L., 2015, *MNRAS*, 451, 144
 Portegies Zwart S. F., McMillan S. L. W., van Elteren A., Pelulessy F. I., de Vries N., 2013, *Comput. Phys. Commun.*, 184, 456
 Punzo D., Capuzzo-Dolcetta R., Portegies Zwart S., 2014, *MNRAS*, 444, 2808
 Sussman G. J., Wisdom J., 1988, *Science*, 241, 433
 Trujillo C. A., Sheppard S. S., 2014, *Nature*, 507, 471
 Tsiganis K., Gomes R., Morbidelli A., Levison H. F., 2005, *Nature*, 435, 459
 Veras D., Wyatt M. C., Mustill A. J., Bonsor A., Eldridge J. J., 2011, *MNRAS*, 417, 2104
 Wisdom J., Holman M., 1991, *AJ*, 102, 1528

This paper has been typeset from a \LaTeX file prepared by the author.