



Universiteit
Leiden
The Netherlands

Targeting HLA class II in allogeneic stem cell transplantation

Balen, P. van

Citation

Balen, P. van. (2021, October 27). *Targeting HLA class II in allogeneic stem cell transplantation*. Retrieved from <https://hdl.handle.net/1887/3221294>

Version: Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/3221294>

Note: To cite this publication please use the final published version (if applicable).



CHAPTER 3

CD4 donor lymphocyte infusion can cause conversion of chimerism without GVHD by inducing immune responses targeting minor histocompatibility antigens in HLA class II

P. van Balen, C.A.M. van Bergen, S.A.P. van Luxemburg-Heijs, W. de Klerk, H.M. van Egmond, S.A.J. Veld, C.J.M. Halkes, J.J. Zwaginga, M. Griffioen, I. Jedema, J.H.F. Falkenburg

ABSTRACT

Under non-inflammatory conditions HLA class II is predominantly expressed on hematopoietic cells. Therefore, donor CD4 T cells after allogeneic stem cell transplantation (alloSCT) may mediate graft-versus-leukemia reactivity without graft-versus-host disease (GVHD). We analyzed immune responses in four patients converting from mixed to full donor chimerism without developing GVHD upon purified CD4 donor lymphocyte infusion (DLI) from their HLA-identical sibling donor after T-cell depleted alloSCT. In vivo activated T cells were clonally isolated after CD4 DLI. Of the alloreactive T-cell clones, 96% were CD4 positive, illustrating the dominant role of CD4 T cells in the immune responses. We identified 9 minor histocompatibility antigens (MiHA) as targets for alloreactivity, of which 8 were novel HLA class II restricted MiHA. In all patients, MiHA specific CD4 T cells were found that were capable to lyse hematopoietic cells and to recognize normal and malignant cells. No GVHD was induced in these patients. Skin fibroblasts forced to express HLA class II, were recognized by only two MiHA specific CD4 T-cell clones. Of the 7 clones that failed to recognize fibroblasts, two targeted MiHA were encoded by genes not expressed in fibroblasts, presentation of one MiHA was dependent on HLA-DO, which is absent in fibroblasts, and T cells recognizing the remaining 4 MiHA had an avidity that was apparently too low to recognize fibroblasts, despite clear recognition of hematopoietic cells. In conclusion, purified CD4 DLI from HLA-identical sibling donors can induce conversion from mixed to full donor chimerism with graft-versus-malignancy reactivity, but without GVHD, by targeting HLA class II restricted MiHA.

INTRODUCTION

Allogeneic stem cell transplantation (alloSCT) provides a potentially curative therapy for patients with a variety of hematologic malignancies. However, acute graft-versus-host Disease (GVHD) and treatment of GVHD remain major causes of transplant related morbidity and mortality. The most efficient method to prevent GVHD is T-cell depletion (TCD) of the graft. AlloSCT regimens using infusion of positively selected CD34 cells or using the CD52 antibody Alemtuzumab for TCD have demonstrated efficient engraftment and reduced acute GVHD.¹⁻⁴ However, TCD substantially impairs post-transplant anti-viral and anti-tumor immunity.⁴⁻⁶ Due to the reduced antitumor effect of TCD grafts, pre-emptive post-transplant donor lymphocyte infusion (DLI) may be needed for treatment of mixed chimerism or persistent disease. Indeed, DLI after alloSCT can mediate the beneficial graft-versus-leukemia (GVL) reactivity, but frequently still at the cost of GVHD.⁷⁻¹⁰

Separation of GVL from GVHD may be achieved by infusion of alloreactive donor T cells that recognize patient hematopoietic cells, including the malignant cells, but not other tissue cells of the patient or donor hematopoietic cells.¹¹ Following HLA matched alloSCT, infusion of donor CD8 T cells recognizing minor histocompatibility antigens (MiHA) selectively expressed in hematopoietic cells may result in destruction of patient hematopoietic cells including the malignant cells, without harming normal tissues.¹²⁻¹⁵ MiHA are polymorphic peptides derived from genes containing single nucleotide polymorphisms which differ between donor and recipient and can be recognized in the context of (self) HLA. Although infusions of patient hematopoiesis directed donor CD8 T cells are being explored, the numbers of known hematopoiesis restricted MiHA that can be targeted are too limited for broad application.¹⁶⁻¹⁹

In contrast to HLA class I, constitutive expression of HLA class II molecules is predominantly restricted to normal and malignant hematopoietic cells.²⁰⁻²³ Therefore, infusion of donor CD4 T cells recognizing HLA class II restricted MiHA may result in selective recognition of recipient normal and malignant hematopoietic cells, thereby inducing GVL without GVHD even if MiHA are targeted that are encoded by genes that are broadly expressed in recipient tissues.²⁴⁻²⁸ This is supported by previous findings that, following an HLA 10/10 matched, but HLA-DPB1 mismatched TCD alloSCT, allo-HLA-DP directed CD4 T cells can cause GVL without GVHD.^{29,30} However, under inflammatory circumstances, expression of HLA class II is significantly upregulated on non-hematopoietic cells, making these tissues susceptible to recognition by CD4 T cells. Indeed, during viral infections, allo-HLA-DP directed CD4 T cells can induce GVHD³¹, suggesting that also CD4 T cells directed against broadly expressed HLA class II restricted MiHA may cause GVHD when HLA class II expression is upregulated on non-hematopoietic cells. However, we recently demonstrated that not all MiHA encoded by broadly expressed genes are adequately presented in HLA class II

on non-hematopoietic cells due to absence of HLA-DO, the natural inhibitor of HLA-DM.³² Furthermore, we demonstrated that selective GVL also depends on the magnitude and diversity of alloreactive T-cell responses and not only on tissue distribution of the MiHA that are targeted.³³ Thus, even under inflammatory conditions, GVHD target tissues may not always be damaged by CD4 T cells recognizing MiHA encoded by broadly expressed genes.

Within an ongoing clinical trial, initiated in the Leiden University Medical Center, treating patients three months after TCD alloSCT with an HLA-identical sibling donor with purified donor CD4 T cells with primary aim to improve immune reconstitution³⁴, we observed hematopoiesis restricted immune responses, as illustrated by conversion from mixed to full donor chimerism, without GVHD in four patients. We identified alloreactive CD4 T cells recognizing HLA class II restricted MiHA in all four patients without the presence of alloreactive CD8 T cells in 3 of 4 patients. Using whole genome association scanning, 9 HLA class II MiHA were identified as targets for recognition. No GVHD was induced by these MiHA specific CD4 T cells, corresponding to lack of HLA class II expression on GVHD target tissues under steady state conditions. Even after upregulation of HLA class II on non-hematopoietic cells, recognition was relatively restricted to hematopoietic cells, although the majority of the identified MiHA were encoded by genes that are broadly expressed. MiHA specific CD4 T cells from all four patients were able to recognize malignant cells. Therefore, CD4 DLI after TCD alloSCT may be an attractive strategy to separate GVL from GVHD in patients transplanted with an HLA-identical sibling donor.

MATERIAL AND METHODS

PATIENTS

At the Leiden University Medical Center (LUMC) patients to be transplanted with mobilized peripheral blood stem cells from HLA-identical sibling donors are treated with a conditioning regimen consisting of fludarabine (50 mg/m² orally from day -10 to -5), busulfan (0.8 mg/kg iv four times a day on day -7 and -6) and alemtuzumab (15 mg iv on day -4 and -3) in case of a non-myeloablative conditioning, or consisting of cyclophosphamide (60 mg/kg iv on day -6 and -5) and total body irradiation (9 Gy on day -1) in case of a myeloablative conditioning regimen. Stem cell grafts are T-cell depleted by addition of 20 mg alemtuzumab to the bag before administration and no post-transplant immune suppression is applied.^{4,35} Three months after TCD alloSCT, in the absence of overall grade II or more GVHD according to the Glucksberg-Seattle classification,³⁶ patients are eligible for treatment with infusion of purified donor CD4 T cells at a dose of 10⁶ cells per kilogram body weight in the setting of a randomized clinical trial (EudraCT Number: 2008-001447-19). This ongoing phase II open-label single-center randomized clinical trial was approved by the LUMC Institutional Review Board and national authorities. Purified CD4 DLI cell products were manufactured by positive selection using CD4 Reagents (Miltenyi Biotec, Bergisch Gladbach, Germany) and CliniMACS System (Miltenyi Biotec) according to the manufacturer's instructions. The study aims to evaluate the immunological effects of prophylactic DLI of purified CD4 T cells early after TCD alloSCT. The primary objective is to evaluate whether CD4 DLI improves immunological recovery within 6 months after alloSCT. Secondary objectives are the evaluations of influence on chimerism and disease status after CD4 DLI. Since the trial is still ongoing, we do not describe the results of the entire clinical trial in this manuscript, but only the results of an in depth analysis of immune responses occurring in a selection of trial participants.

ISOLATION, EXPANSION AND SELECTION OF T-CELL CLONES

Peripheral blood, bone marrow and skin biopsies for the generation of fibroblasts were obtained from the patients, their donors and third party healthy individuals after approval by the LUMC Institutional Review Board and informed consent according to the Declaration of Helsinki. Mononuclear cells were isolated using Ficoll separation and cryopreserved.

To isolate *in vivo* activated T cells, peripheral blood mononuclear cells (PBMC) obtained after CD4 DLI or six weeks after randomization in case patients did not receive CD4 DLI, were stained with antibodies against CD8 (Alexa Fluor, Invitrogen/Caltag, Buckingham, UK), CD4 (FITC, BD/Pharmingen, Breda, Netherlands), CD14 (APC, ITK/Biolegend, Uithoorn, Netherlands) and HLA-DR (PE, BD). HLA-DR⁺ CD8 and HLA-DR⁺ CD4 T cells were sorted single cell into 96-well U-bottomed plates (Corning, Amsterdam, Netherlands) or 384-well flat bottomed plates (Greiner Bio-One, Alphen a/d Rijn, Netherlands). T-cell clones were

expanded using Iscove's modified Dulbecco's medium (IMDM, Lonza BioWhittaker, Verviers, Belgium) with 5% pooled human serum, 5% fetal bovine serum (FBS, Gibco Invitrogen, Bleiswijk, Netherlands), 100 IU/ml Interleukin 2 (Chiron, Amsterdam, Netherlands), 2 ng/ml Interleukin 7 (Miltenyi Biotec), 2 ng/ml Interleukin 15 (Miltenyi Biotec), 0.8 μ g/ml phytohemagglutinin (Murex Biotec Limited, Dartford, UK) and $25\text{-}50 \times 10^3$ irradiated third party PBMC as feeder cells. Proliferating T-cell clones were restimulated every 10-14 days and tested for reactivity against patient and donor derived EBV-LCL. After overnight incubation of 2×10^4 patient or donor derived EBV-LCL with 2×10^3 T cells, recognition was measured by IFN γ ELISA according to the manufacturer's instructions (Sanquin Reagents, Amsterdam, Netherlands). A T-cell clone was determined to be alloreactive when at least 500 pg/ml IFN γ was produced after incubation with patient derived EBV-LCL and no IFN γ was produced after incubation with donor derived EBV-LCL.

HLA RESTRICTION AND TCRBV USAGE OF ALLOREACTIVE T CELLS

To determine whether HLA-DR, HLA-DQ or HLA-DP was the HLA restriction molecule for recognition by alloreactive CD4 T cells, patient derived EBV-LCL were pre-incubated with saturating concentrations of monoclonal antibodies (MoAb) against HLA class II (PdV5.2), HLA-DR (B8.11.2), HLA-DQ (SPVL3) or HLA-DP (B7.21) for 30 minutes at room temperature before addition of the T cells, and inhibition of IFN γ production was determined. T-cell receptor- β variable chain (TCRBV) usage of the T-cell clones was investigated by flow cytometry using specific monoclonal antibodies as provided with the TCRBV repertoire kit (Beckman Coulter).

MiHA IDENTIFICATION BY WHOLE GENOME ASSOCIATION SCANNING

The method of whole genome association scanning (WGAS) using an HLA transduced panel of third party EBV-LCL was described earlier.³⁷ In short, 48-116 third-party EBV-LCL were transduced with one of the possible HLA restriction molecules. The transduced EBV-LCL were incubated with the alloreactive CD4 T cells and IFN γ production was measured using ELISA. The presence or absence of recognition of the different EBV-LCL was compared with the EBV-LCL genotype data of over one million single nucleotide polymorphisms (SNPs) in order to find an association between the recognition and the presence of a certain SNP. If association with a missense SNP was found, patient and donor variant peptides encoded by the SNP region were synthesized. If incubation of donor derived EBV-LCL loaded with patient variant peptide, titrated in a concentration from 10^{-4} to 10^{-10} M, resulted in IFN γ production by the T-cell clone, this peptide was confirmed to be the MiHA.

CYTOTOXICITY OF MiHA SPECIFIC CD4 T CELLS

Cytotoxic capacities of alloreactive CD4 T cells was analyzed following incubation of 2.5×10^5 T cells with 2.5×10^4 EBV-LCL target cells labeled with PKH26 Red Fluorescent (Sigma-Aldrich, Zwijndrecht, Netherlands). Target cell survival after 24 hours was measured by

flowcytometric cell counting of the target cells using Flow-Count fluorospheres (Beckman Coulter, Woerden, Netherlands).³⁸ Percentage cell lysis was calculated by the formula $100 - 100 \times (\text{total number of surviving target cells after incubation with T cells} / \text{total number of target cells without incubation with T cells})$. Statistical analysis was performed using Mann-Whitney U test and significance was defined by $p < 0.05$.

RECOGNITION OF PHA BLASTS, MALIGNANT CELLS AND SKIN DERIVED FIBROBLASTS

To investigate the recognition of patient derived activated T cells, T cells isolated from the patient before alloSCT were treated with phytohemagglutinin (PHA blasts) and incubated with MiHA specific CD4 T cells in a stimulator to responder ratio of 5:1. To investigate the recognition of malignant cells, MiHA specific CD4 T cells were incubated with primary malignant cells preferably of the patient. If not available, third party cells representative for the malignancy of the patient were used, like bone marrow cells of a patient suffering from chronic myeloid leukemia and multiple myeloma cell lines UM-6 (CVCL-W395), UM-3 (CVCL-W394), RPMI8226 (ATCC CRM-CCL-155) or U266 (ATCC-TIB-196). The malignant cells expressed the MiHA and HLA restriction molecule of interest endogenously or after retroviral transduction. To investigate recognition of skin derived fibroblasts, 4×10^3 fibroblasts were incubated with 2×10^3 MiHA specific CD4 T cells. Fibroblasts were induced to express HLA class II molecules on the cell surface by culturing them with 200 IU/ml IFN γ (Boehringer Ingelheim, Rijnland-Palts, Germany) for five days in Dulbecco's Modified Eagle Medium (DMEM, Lonza BioWhittaker) supplemented with 10% FBS. Because peptide processing in HLA class II could be different in fibroblasts compared to EBV-LCL due to the absence of HLA-DO in fibroblasts, the role of HLA-DO in the recognition of fibroblasts by alloreactive CD4 T cells was studied using fibroblasts that were retrovirally transduced with HLA-DO α and HLA-DO β as described previously.³² HLA class II expression on target cells was analyzed using flowcytometry with monoclonal antibodies against HLA-DR (PE, IgG2a, L-243, BD), HLA-DQ (PE, IgG2a, 1a3, Bio-Connect, Huissen, The Netherlands) and HLA-DP (PE, IgG3, B7.21, Bio-Connect) and T-cell recognition was analyzed by measurement of IFN γ production.

GENE EXPRESSION PROFILES

To investigate whether the MiHA encoding genes are expressed in hematopoietic cells only or also in non-hematopoietic cells, quantitative RT-PCR was performed using TaqMan Assays (ThermoFisher, Breda, Netherlands). Expression of the gene of interest was corrected for expression of household reference genes GAPDH and B-actin.

RESULTS

PATIENT CHARACTERISTICS

The first 15 patients who were treated with CD4 DLI in the intervention arm of the clinical trial, did not develop GVHD, except for patient D, who developed mild skin GVHD (overall grade I) 3 months after infusion, which resolved completely after start with topical steroids. To investigate whether CD4 DLI could induce an allo-immune response following the infusion in the absence of GVHD, peripheral blood subset chimerism was analyzed. In 11 of 15 patient no conversion of chimerism occurred. However, 4 of 15 patients converted from markedly mixed to full donor chimerism within three months after infusion of donor CD4 T cells, indicating the development of an allo-immune response in a selection of patients after CD4 DLI (Table 1). These four patients were selected for further analysis. Also four patients from the control group were selected. In the control group, no conversion of chimerism was observed, although in patient G improvement of chimerism occurred (Table 1).

Table 1 Patients characteristics and chimerism data on different time points from CD4 DLI (patient A-D) or from randomization (patients E-H)

Patient	Age, gender	Disease	Conditioning regimen	Days from CD4 DLI / randomisation	Absolute leucocyte count (x 10 ⁹ /L)	Absolute CD3 count (x 10 ⁶ /L)	Chimerism (% cells from recipient origin)			
							CD8 T cells	CD4 T cells	B cells	Granulocytes
A	46, female	Plasma cell myeloma	NMA	-45	3.6	529	57	9	1	<1
				-3	5.8	274	77	28	4	4
				46	4.2	211	60	16	<1	<1
				95	5.4	420	<1	<1	<1	<1
B	57, male	Plasma cell myeloma	NMA	-56	3.5	128	74	4	<1	<1
				-7	5.6	359	2	<1	<1	<1
				35	4.6	529	1	<1	<1	<1
				91	6.2	574	<1	<1	<1	<1
C	48, male	Chronic myeloid leukemia (blastic phase)	MA	-48	8.6	88	26	5	<1	2
				0	3.8	389	90	13	<1	5
				50	3.2	254	55	5	<1	8
				92	2.7	258	<1	<1	<1	<1
D	62, male	Acute myeloid leukemia	NMA	-54	3.2	73	90	15	<1	<1
				-5	3.7	100	86	53	<1	<1
				45	4.4	196	17	8	<1	<1
				94	7.7	275	<1	1	<1	<1
E	62, male	Plasma cell myeloma	NMA	0	1.5	326	60	7	<1	<1
				28	3.6	652	48	6	<1	<1
				91	4.2	571	88	60	<1	<1
F	55, male	Plasma cell myeloma	NMA	0	4.1	91	55	20	<1	<1
				47	4.8	128	38	22	<1	<1
				90	5.0	222	37	16	<1	<1
G	55, male	Acute myeloid leukemia	NMA	0	4.6	85	4	27	1	1
				49	10.6	67	<1	2	2	<1
				98	5.3	62	2	2	2	1
H	51, female	Plasma cell myeloma	NMA	0	2.9	976	<1	1	n.a.	1
				43	3.0	695	<1	<1	<1	<1
				97	5.8	454	2	1	<1	1

NMA: non-meloablative. MA: myeloablative. N.a.: not available.

ISOLATION OF ALLOREACTIVE T CELLS

To characterize the allo-immune response observed after infusion of donor CD4 T cells in these four patients, peripheral blood samples were taken during conversion of chimerism. In vivo activated, HLA-DR expressing CD4 and CD8 T cells were clonally isolated using flowcytometric cell sorting (Figure 1) and 5-25% of the single T cells expanded, resulting in 295-1396 growing T cell clones per patient (Table 2). These T-cell clones were tested for allo-reactivity, defined as IFN γ production of at least 500 pg/ml after overnight incubation with patient derived EBV-LCL and absence of IFN γ production after incubation with donor derived EBV-LCL (Figure 2). Reactivity to both patient and donor derived EBV-LCL was interpreted as EBV specific T cells present in the in vivo activated T-cell compartment (Table 2). Total frequencies of 1-28% of the expanded CD4 T cells were alloreactive as compared to only 0-3% of the expanded CD8 T cells (Table 2). As controls, four patients randomized to the control arm of the study, who did therefor not receive CD4 DLI and in which conversion of chimerism did not occur, were analyzed six weeks after randomization. In these patients 1-5% of expanded CD4 T-cell clones and 3-28% of expanded CD8 T-cell clones were EBV specific. However, only 0-0.3% of expanded CD4 T-cell clones and 0-1% of expanded CD8 T-cell clones were alloreactive (Table 2). These data illustrate that conversion from mixed to full donor chimerism after CD4 DLI was associated with development of dominant alloreactive CD4 T-cell responses.




Figure 1. Sorted cell populations. Marked cell populations, being in vivo activated HLA-DR positive CD4 or CD8 T cells from four patients were single cell sorted using flowcytometry using HLA-DR PE, CD4 FITC and CD8 AF700 monoclonal antibodies.

Table 2. Numbers of expanded and alloreactive CD4 and CD8 T cells after isolation of in vivo activated T cells six weeks after CD4 DLI (patient A-D) or randomization (patient E-H).

Patient	A	B	C	D	E	F	G	H
				Day 45 after CD4 DLI	Day 94 after CD4 DLI			
Total number of expanded CD4 T-cell clones	332	205	341	384	452	516	360	384
Alloreactive CD4 T-cell clones	10	2	41	106	4	1	0	0
EBV-LCL reactive CD4 T-cell clones	15	25	1	3	0	20	18	4
Total number of expanded CD8 T-cell clones	99	90	66	416	144	538	250	173
Alloreactive CD8 T-cell clones	1	0	0	1	4	0	0	2
EBV-LCL reactive CD8 T-cell clones	59	33	0	0	1	24	69	5




Figure 2. IFN γ production of alloreactive CD4 T-cell clones. Recognition of patient and donor derived EBV-LCL by CD4 T cells measured by IFN γ production after overnight incubation. Each dot represents the IFN γ release by one alloreactive CD4 T-cell clone after overnight incubation in a responder to stimulator ratio of 1:10. All CD4 T-cell clones did recognize patient derived EBV-LCL and did not recognize donor derived EBV-LCL and are therefore defined as alloreactive.

IDENTIFICATION OF HLA CLASS II RESTRICTED MIHA USING WGAS

To characterize the MiHA that are recognized by the alloreactive CD4 T-cell clones, WGAS was applied. We first determined the HLA class II restriction molecules of the T-cell clones using monoclonal antibodies against HLA-DR, DQ or DP. In addition, TCRBV usage of each T-cell clone was analyzed using flowcytometry. If T-cell clones recognized a target antigen in the context of the same HLA molecule and expressed the same TCRBV, it was considered likely that these clones recognized the same MiHA, and therefore initially one clone was selected for WGAS (Table 3). A representative example of WGAS with a T-cell clone from patient C is shown in Figure 3. When association was found between T-cell recognition and one or more SNPs in the genotypes of a panel of EBV-LCL, patient and donor allelic variants of the peptides encoded by the polymorphic region were synthesized, loaded on donor EBV-LCL and tested for recognition by the T-cell clone. When the MiHA was identified by WGAS and confirmed by T-cell recognition of peptide-pulsed donor EBV-LCL, all other T-cell clones from the same patient were tested for recognition of the identified MiHA.




Figure 3. Representative example of the identification of a MiHA by WGAS. (A) IFN γ production of clone P07-066 from patient C upon incubation with a panel of 71 SNP genotyped EBV-LCL which were transduced with HLA-DRB1*15:01. **(B)** Identification of associating SNPs on chromosome 11 that are present in EBV-LCL that were recognized by clone P07-066 and absent in EBV-LCL that were not recognized. **(C)** Identification of missense SNP rs2271001 encoding the possible MiHA recognized by clone P07-066. **(D)** IFN γ production by clone P07-066 after incubation with donor EBV-LCL loaded with donor or patient derived allelic peptide variants at titrated concentrations. The patient, but not donor, peptide was recognized by the T cells, thereby validating the peptide as MiHA.

From the T-cell clones that failed to recognize the identified MiHA, one T-cell clone was selected for additional WGAS analysis attempting to further identify additional MiHA. A total number of 16 WGAS experiments were performed, resulting in the identification of 9 MiHA. Eight out of these nine antigens were novel MiHA. In addition to alloreactive CD4 T cells, 5 alloreactive CD8 T-cell clones were isolated from PBMC of patient D (Table 2). The 4 CD8 T-cell clones isolated 94 days after CD4 DLI all used the same TCRBV, which was different from the TCRBV usage of the CD8 T-cell clone isolated 45 days after CD4 DLI. WGAS resulted in identification of the novel HLA-A*03:01 restricted MiHA LB-NADK-1K, which was the target for all 5 isolated CD8 T-cell clones. Table 4 gives an overview of the identified MiHA and their HLA restriction molecules. The HLA-DRB1*15:01 restricted MiHA from gene SLC19A1 was earlier described²⁶, but we found that this MiHA could also be presented and recognized in HLA-DRB1*11:01. For 4 T-cell clones from patient A and for 3 T-cell clones from patient D, as well as for 1 T-cell clone from control patient E and 3 T-cell clones from control patient G, MiHA could not be identified due to lack of associating SNPs in WGAS. For all other T-cell clones, however, MiHA were successfully identified.

In summary, using WGAS, a total of 9 HLA class II and one HLA class I restricted MiHA were identified as targets for alloreactive T cells isolated from patients who converted from mixed to full donor chimerism after CD4 DLI.

Table 3. Classification of alloreactive T-cell clones according to HLA restriction molecule and TCR V β usage.

Patient	HLA class II restriction molecule [#]	TCR V β usage	Identified MiHA	Number of clones
Patient A	undetermined	unknown	-	1 [†]
	DQ	2	LB-LILRB1-1I	6 [†]
	DQ	2	-	2 [†]
	DQ	9	-	1 [†]
Patient B	DR	8	LB-ABCA5-1R	1 [†]
	DR	13.2	LB-RPS4Y	1 [†]
Patient C	DR	2	SCL19A1	21 [†]
	DR	5.3	SCL19A1	1 [†]
	DR	unknown	LB-ZDHHHC13-1K	10 [†]
	DR	unknown	LB-KHNYN-1K	3 [†]
	DR	7.1	LB-KHNYN-1K	1 [†]
	DR	9	LB-CTSB-1G	1 [†]
	DR	13.2	LB-CTSB-1G	1 [†]
	DR	9	LB-LGALS8-1C	2 [†]
	DR	unknown	LB-LGALS8-1C	1 [†]
Patient D	undetermined	3	-	2 [†]
	DQ	5.1	-	1
	DP	2	LB-LY75-2R	38 [†]
	DP	3	LB-LY75-2R	1 [†]
	DP	4	LB-LY75-2R	32 [†]
	DP	8	LB-LY75-2R	6 [†]
	DP	13.2	LB-LY75-2R	3 [†]
	DP	unknown	LB-LY75-2R	27 [†]

[#] HLA class II restriction molecule was determined using monoclonal antibodies against pan HLA class II (PdV5.2), HLA-DR (B8.11.2), HLA-DQ (SPVL3) and HLA-DP (B7.21).

[†] WGAS was performed using one of these T-cell clones.

[‡] MiHA specificity of these T-cell clones was determined by recognition of donor EBV-LCL loaded with the MiHA identified using WGAS analysis of another T-cell clone.

Table 4. Overview of identified MIHAs and their characteristics.

MIHA name	SNP	Peptide polymorphism*	Restriction	Number of different TCRBV usages of MIHA T-cell clones	T cells isolated from patient	Contribution to immune response [#]
LB-LILRB1-1I	rs1061680	PSPVNSGGN[V]TLQDCSQVA	DQB1*06:02	1	A	60 %
LB-RPS4Y	- [†]	EKTGEHRLVYDTKGRFAVH	DRB1*03:01	1	B	50 %
LB-ABCA5-1R	rs17686569	GKEAIRSIG[R]QJKTYRKKGEN	DRB1*11:01	1	B	50 %
LB-KHYN-1K	rs3742520	ARGDTYAVEKEGG[K]TQGGPREMDWG	DRB5*01:01	2	C	9.8 %
LB-ZDHHCT3-1K	rs2271001	INNRLDLV[K]RFYISKGAVVDQ	DRB1*15:01	1	C	24 %
LB-CTSB-1G	rs1803250	HNFYNDM[G]SYLKRLCGTF	DRB1*11:01	2	C	4.9 %
LB-LGALS8-1C	rs1041935	DQLDPGTLIV[C]RjGHVPSADRF	DRB1*11:01	2	C	7.3 %
SLC19A1	rs1051266	DPELRSWR[R]HJLVCYLCFYG	DRB1*15:01	1-2	C	54 %
SLC19A1	rs1051266	DPELRSWR[R]HJLVCYLCFYG	DRB1*11:01	1-2	C	54 %
LB-LY75-2R	rs17827158	RAGRPTIKNE[R]KFLAGLSTDG	DPB1*04:01	6	D	97 %
LB-NADK-1K	rs4751	AVHNGLGEK[N]GSSQA	A*03:01	1	D	-

* Patient type amino acid residues are underlined.

[#] Proportion of alloreactive CD4 T-cell clones recognizing this MIHA as a percentage of all alloreactive CD4 T cells isolated in that particular patient.

[†] Male specific peptide.

REACTIVITY OF MIHA SPECIFIC CD4 T CELLS AGAINST HEMATOPOIETIC CELLS

Since the induction of MiHA specific CD4 T cells correlated with conversion from mixed to full donor chimerism as a result of disappearing lymphohematopoietic cells from recipient origin, we investigated whether the isolated MiHA specific CD4 T cells were capable of exerting cytotoxic activity against recipient hematopoietic cells. MiHA specific CD4 T-cell clones were incubated with patient derived EBV-LCL at an effector to target ratio of 10:1, and after 24 hours of co-incubation the numbers of surviving target cells were quantified by flow cytometry. For all MiHA, individual MiHA specific CD4 T-cell clones were identified that exerted cytotoxic activity against EBV-LCL, resulting in 19-51% specific lysis, although not all differences of the mean lysis of all MiHA specific CD4 T-cell clones with background lysis were statistically significant due to low number of experiments (Figure 4). Since conversion of chimerism was most prominent in the T-cell compartment (Table 1), recognition of PHA-stimulated patient T cells (PHA blasts) by MiHA specific CD4 T cells was investigated. CD4 T cells specific for LB-RPS4Y, LB-ABCA5-1R and LB-LY75-2R as well as CD8 T cells specific for LB-NADK-1K all recognized PHA blasts as illustrated by high IFN γ release. CD4 T cells specific for SLC19A1 produced less IFN γ upon incubation with PHA blasts as compared to EBV-LCL, whereas the remaining T-cell clones specific for the other MiHA did not recognize PHA blasts (Figure 5A).




Figure 4. Cytotoxic capacities of MiHA specific CD4 T cells. Cell lysis calculated after measurement of target cell survival using flowcytometric cell counting after 24 hours of incubation with the MiHA specific CD4 T cells in an effector to target ratio of 10:1. Target cells were patient EBV-LCL and effector cells were MiHA specific CD4 T cells. CD4 T cells with irrelevant specificity were used as negative controls. EBV-LCL from patient A, B, C and D are represented by black, dark grey, light grey and blocked bars, respectively. For all MiHA, individual MiHA specific CD4 T-cell clones were identified that exerted cytotoxic activity against EBV-LCL, resulting in 19-51% specific lysis, although not all differences of the mean lysis by MiHA specific CD4 T-cell clones with background lysis were statistically significant (Mann-Whitney U test) due to low number of experiments.




Figure 5. Recognition of PHA blasts and malignant cells by MiHA specific T cells. Recognition of PHA blasts and malignant cells by MiHA specific CD4 T cells was measured by IFN γ production, represented as percentage of production after incubation with patient EBV-LCL. Surface expression of HLA class II was confirmed by flow cytometry (data not shown). **(A)** Recognition of patient derived PHA blasts. CD4 T cells specific for LB-RPS4Y, LB-ABCA5-1R and LB-LY75-2R as well as CD8 T cells specific for LB-NADK-1K produced high amounts of IFN γ upon incubation with patient derived PHA blasts. SLC19A1 specific CD4 T cells produced less IFN γ after incubation with PHA blasts than with EBV-LCL and the other MiHA specific CD4 T cells did not recognize PHA blasts. T-cell clones directed against the HLA restriction molecule were used as positive controls **(B)** Recognition of MiHA and HLA restriction molecule expressing malignant cells by MiHA specific T cells. LB-LILRB1-1I specific T cells were tested against myeloma cell-lines UM-6, UM-3, RPMI8226 and U266. LB-RPS4Y specific T cells against myeloma cell-lines RPMI8226 and U266 and LB-ABCA5-1R against myeloma cell-lines UM-6 and RPMI8226. T cells derived from patient C were tested against third party CML cells and T cells derived from patient D against patient derived AML cells. MiHA specific CD4 T cells recognizing malignant cells could be detected in all four patients.

Although all patients were in complete remission of their disease at the time of infusion of CD4 DLI, recognition of malignant cells by MiHA specific CD4 T cells was investigated to assess their potential to mediate GVL reactivity. Primary malignant cells were only available from patient D and for this reason, third party target cells were used as representative for the malignancies of patient A, B and C. These target cells expressed the MiHA and HLA restriction element of interest. LB-LILRB1-1I specific CD4 T cells were incubated with myeloma cell-lines UM-6, UM-3, RPMI8226 and U266 transduced with HLA-DQB1*06:02. LB-RPS4Y specific CD4 T cells were incubated with myeloma cell-lines RPMI8226 (HLA-DRB1*03:01 positive³⁹) and U266 transduced with HLA-DRB1*03:01. LB-ABCA5-1R specific CD4 T cells were incubated

with myeloma cell-lines UM-6 and RPMI8226 transduced with HLA-DRB1*11:01. MiHA specific CD4 T cells isolated from patient C were incubated with third party bone marrow cells from a patient with CML. Since these third party CML cells were negative for LB-CTSB-1G, recognition of malignant cells by CD4 T cells specific for this MiHA could not be tested. Reactivity of MiHA specific T cells from patient D was tested against autologous purified primary AML cells. As shown in Figure 5B, in all four patients, allo-reactive MiHA specific CD4 T cells were detected that were able to recognize malignant cells.

In conclusion, the results showed that all isolated MiHA specific CD4 T cells had cytolytic capacity to eliminate recipient hematopoietic cells and that PHA blasts were recognized to a lesser extent than EBV-LCL. Moreover, in all four patients, MiHA specific CD4 T cells with the potential to recognize malignant cells were found.

REACTIVITY OF MIHA SPECIFIC CD4 T CELLS AGAINST SKIN FIBROBLASTS

To investigate whether non-hematopoietic cells from GVHD target tissues could be targeted under inflammatory conditions due to upregulation of HLA class II expression, we tested recognition of skin derived fibroblasts by the MiHA specific T cells. Skin fibroblasts were available for two patients (C and D). No fibroblasts were available from patients A and B and we therefore used fibroblasts expressing the MiHA and HLA restriction molecule from third party individuals. Expression of HLA class II molecules on fibroblasts was induced by addition of IFN γ to the culture medium mimicking inflammatory conditions (Supplementary Figure 1). CD4 T cells directed against allo HLA class II alleles were used as positive controls and all fibroblasts tested were recognized by these clones. Since HLA class I is constitutively expressed on fibroblasts, as expected, CD8 T cells specific for LB-NADK-1K recognized fibroblasts already without IFN γ pretreatment (Figure 6A). CD4 T cells specific for LB-RPS4Y and LB-ZDHHC13-1K were able to recognize fibroblasts after IFN γ pretreatment (Figure 6B), whereas all other MiHA specific T-cell clones did not recognize (IFN γ pretreated) fibroblasts. CD4 T cells specific for LB-LILRB1-1I and LB-LY72-2R failed to recognize fibroblasts (Figure 6C) due to the lack of expression of the MiHA encoding genes in fibroblasts (data not shown), while CD4 T cells specific for the other 5 MiHA did not recognize fibroblasts pretreated with IFN γ (Figure 6D) despite expression of the encoding genes (data not shown). To investigate whether lack of recognition of IFN γ pretreated fibroblasts is caused by absence of HLA-DO, fibroblasts were transduced with HLA-DO α/β and tested for T-cell recognition. After transduction and pretreatment with IFN γ , fibroblasts were recognized by CD4 T cells specific for LB-LGALS8-1C, indicating that presentation of this MiHA is dependent on HLA-DO.³²

These results illustrate that not only restricted expression of HLA class II on hematopoietic cells defines hematopoiesis specific recognition by MiHA specific CD4 T-cell clones, but that even under inflammatory conditions, hematopoiesis restricted recognition is a common phenomenon.




Figure 6. Recognition of fibroblasts by MiHA specific T cells. T-cell recognition of EBV-LCL, fibroblasts and fibroblasts forced to express HLA class II molecules using interferon gamma (IFN) pretreatment. **(A)** CD8 T cells specific for LB-NADK-1K recognized fibroblasts already without IFN γ pretreatment. **(B)** CD4 T cells specific for LB-RPS4Y and LB-ZDHHHC13-1K recognized fibroblasts only after IFN γ pretreatment. **(C)** CD4 T cells specific for LB-LILRB1-1I and LB-LY75-2R failed to recognize fibroblasts due to lack of expression of the MiHA encoding genes. **(D)** CD4 T cells specific for the remaining 5 MiHA failed to recognize fibroblasts despite detectable gene expression and pretreatment with IFN γ . After transduction with HLA-DO α/β fibroblasts were recognized by LB-LGALS8-1C specific CD4 T cells, indicating that presentation of this MiHA was HLA-DO dependent. * = not tested.

DISCUSSION

Our results illustrate that purified CD4 DLI, administered with the intention to promote immune reconstitution after TCD alloSCT, can result in conversion from mixed to full donor chimerism without occurrence of GVHD, associated with the development of strong alloreactive CD4 T-cell responses against HLA class II restricted MiHA. Using WGAS we identified 9 MiHA targeted by CD4 T cells. These MiHA specific CD4 T cells were likely to be responsible for the hematopoiesis specific immune response. Three months after TCD alloSCT, in the absence of immune suppression, only purified CD4 T cells were infused, and conversion of chimerism occurred in the first weeks after CD4 DLI in all patients except for patient B in which conversion of chimerism started before, but continued after infusion. During this hematopoiesis specific immune response only alloreactive CD4 T cells recognizing HLA class II restricted MiHA could be identified in PBMC from 3 of 4 patients. These CD4 T cells were able to exert direct cytotoxicity against hematopoietic target cells in the absence of CD8 T cells. Reactivity against patient derived activated T cells (PHA blasts) could be demonstrated by MiHA specific CD4 T cells in all four patients, strongly suggesting that conversion to full donor chimerism in the T-cell compartment was caused by the alloreactive CD4 T cells. Although T cells do not constitutively express HLA class II molecules, they can become targets for CD4 T cells due to upregulation of HLA class II upon activation.⁴⁰ From all four patients, MiHA specific CD4 T cells could be isolated recognizing malignant cells in vitro, illustrating the capacity of these T cells to mediate graft-versus-malignancy reactivity.

Immune responses by MiHA specific CD4 T cells are likely to preferentially target cells of hematopoietic origin, since under non-inflammatory conditions HLA class II is predominantly expressed on hematopoietic cells. Under inflammatory conditions, however, non-hematopoietic tissues can become targets for CD4 T cells due to upregulation of HLA class II expression.^{31,40} LB-LILRB1-II and LB-LY75-2R are MiHA encoded by genes that are not expressed in fibroblasts, and therefore, even under inflammatory circumstances fibroblasts will not be targeted by CD4 T cells specific for these MiHA. Although all other identified HLA class II restricted MiHA were encoded by broadly expressed genes, only two MiHA specific CD4 T-cell clones were reactive with HLA class II expressing fibroblasts. Apparently, not only expression of the HLA restriction molecule and MiHA encoding gene in GVHD target tissue determine whether these non-hematopoietic cells are targeted by specific CD4 T cells. We previously showed that, due to the lack of HLA-DO in non-hematopoietic cells, there is a difference in physiology of peptide presentation in HLA class II in hematopoietic cells compared to GVHD target cells.^{32,41} Some peptides can only be presented in HLA class II in the presence of HLA-DO, which is expressed in hematopoietic cell types, but not in the majority of non-hematopoietic cells. LB-LGALS8-1C is an example of an HLA-DO dependent antigen since fibroblasts were only recognized by specific CD4 T cells after

enforced expression of HLA-DO. All other MiHA specific CD4 T-cell clones recognized EBV-LCL, but not HLA class II expressing fibroblasts, which may be due to differential avidity of the T-cell clones for these tissues.

Recently, we showed that GVL reactivity without GVHD by MiHA specific CD8 T cells is not solely determined by tissue distribution of the recognized MiHA, but predominantly depends on magnitude and diversity of alloreactive T-cell responses.³³ In the four patients analyzed here, conversion to full donor chimerism after CD4 DLI occurred without GVHD. In patients A and B, T-cell responses with low magnitude and diversity were induced and the majority of T cells in patient A were specific for an hematopoiesis restricted MiHA. In patient C, magnitude and diversity were higher, but absence of HLA-DO or low avidity of the T cells for fibroblasts explained lack of GVHD. In patient D there was a high magnitude of a T-cell response, which was directed against an hematopoiesis restricted MiHA. Thus, none of the patients experienced an immune response with both high magnitude and diversity against MiHA that are broadly expressed, resulting in a low risk of GVHD even after upregulation of HLA class II expression.

Patient D developed mild skin GVHD three months after CD4 DLI. At this time point, CD8 T cells specific for LB-NADK-1K were isolated that recognized fibroblasts, while at the earlier time point of conversion of chimerism, more than 99% of alloreactive T cells isolated were CD4 T cells that lacked reactivity against fibroblasts. These data suggest that LB-NADK-1K specific CD8 T cells were responsible for development of this mild GVHD, while LB-LY75-2R specific CD4 T cells may have mediated conversion of chimerism.

In conclusion, purified CD4 DLI administered three months after TCD alloSCT from an HLA-identical sibling donor can lead to conversion of mixed to full donor chimerism due to graft-versus-patient hematopoiesis reactivity without GVHD by mediating a MiHA specific HLA class II restricted immune response against patient hematopoietic cells, which can also target the malignant hematopoietic counterpart. Even under inflammatory conditions GVL may be separated from GVHD due to the higher susceptibility of hematopoietic cells compared to non-hematopoietic cells from GVHD tissues to be targeted by MiHA specific CD4 T cells and the limited diversity and magnitude of the induced immune responses.


REFERENCES

1. Pasquini MC, Devine S, Mendizabal A, et al. Comparative outcomes of donor graft CD34+ selection and immune suppressive therapy as graft-versus-host disease prophylaxis for patients with acute myeloid leukemia in complete remission undergoing HLA-matched sibling allogeneic hematopoietic cell transplantation. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*. 2012;30(26):3194-3201.
2. Bayraktar UD, de Lima M, Saliba RM, et al. Ex vivo T cell-depleted versus unmodified allografts in patients with acute myeloid leukemia in first complete remission. *Biology of blood and marrow transplantation : journal of the American Society for Blood and Marrow Transplantation*. 2013;19(6):898-903.
3. Kottaridis PD, Milligan DW, Chopra R, et al. In vivo CAMPATH-1H prevents graft-versus-host disease following nonmyeloablative stem cell transplantation. *Blood*. 2000;96(7):2419-2425.
4. Barge RM, Starrenburg CW, Falkenburg JH, Fibbe WE, Marijt EW, Willemze R. Long-term follow-up of myeloablative allogeneic stem cell transplantation using Campath "in the bag" as T-cell depletion: the Leiden experience. *Bone marrow transplantation*. 2006;37(12):1129-1134.
5. Chakrabarti S, Mackinnon S, Chopra R, et al. High incidence of cytomegalovirus infection after nonmyeloablative stem cell transplantation: potential role of Campath-1H in delaying immune reconstitution. *Blood*. 2002;99(12):4357-4363.
6. Chakrabarti S, Mautner V, Osman H, et al. Adenovirus infections following allogeneic stem cell transplantation: incidence and outcome in relation to graft manipulation, immunosuppression, and immune recovery. *Blood*. 2002;100(5):1619-1627.
7. Collins RH, Jr., Shpilberg O, Drobyski WR, et al. Donor leukocyte infusions in 140 patients with relapsed malignancy after allogeneic bone marrow transplantation. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*. 1997;15(2):433-444.
8. Peggs KS, Thomson K, Hart DP, et al. Dose-escalated donor lymphocyte infusions following reduced intensity transplantation: toxicity, chimerism, and disease responses. *Blood*. 2004;103(4):1548-1556.
9. Eefting M, de Wreede LC, Halkes CJ, et al. Multi-state analysis illustrates treatment success after stem cell transplantation for acute myeloid leukemia followed by donor lymphocyte infusion. *Haematologica*. 2016;101(4):506-514.
10. Eefting M, von dem Borne PA, de Wreede LC, et al. Intentional donor lymphocyte-induced limited acute graft-versus-host disease is essential for long-term survival of relapsed acute myeloid leukemia after allogeneic stem cell transplantation. *Haematologica*. 2014;99(4):751-758.
11. Falkenburg JH, Jedema I. Allo-reactive T cells for the treatment of hematological malignancies. *Molecular oncology*. 2015;9(10):1894-1903.
12. Griffioen M, van Bergen CA, Falkenburg JH. Autosomal Minor Histocompatibility Antigens: How Genetic Variants Create Diversity in Immune Targets. *Frontiers in immunology*. 2016;7:100.
13. Pont MJ, Hobo W, Honders MW, et al. LB-ARHGDI1B-1R as a novel minor histocompatibility antigen for therapeutic application. *Haematologica*. 2015;100(10):e419-422.
14. Norde WJ, Overes IM, Maas F, et al. Myeloid leukemic progenitor cells can be specifically targeted by minor histocompatibility antigen LRH-1-reactive cytotoxic T cells. *Blood*. 2009;113(10):2312-2323.
15. Hobo W, Broen K, van der Velden WJ, et al. Association of disparities in known minor histocompatibility antigens

- with relapse-free survival and graft-versus-host disease after allogeneic stem cell transplantation. *Biology of blood and marrow transplantation : journal of the American Society for Blood and Marrow Transplantation*. 2013;19(2):274-282.
16. Meij P, Jedema I, van der Hoorn MA, et al. Generation and administration of HA-1-specific T-cell lines for the treatment of patients with relapsed leukemia after allogeneic stem cell transplantation: a pilot study. *Haematologica*. 2012;97(8):1205-1208.
 17. Marijt WA, Heemskerk MH, Kloosterboer FM, et al. Hematopoiesis-restricted minor histocompatibility antigens HA-1- or HA-2-specific T cells can induce complete remissions of relapsed leukemia. *Proc Natl Acad Sci U S A*. 2003;100(5):2742-2747.
 18. Warren EH, Fujii N, Akatsuka Y, et al. Therapy of relapsed leukemia after allogeneic hematopoietic cell transplantation with T cells specific for minor histocompatibility antigens. *Blood*. 2010;115(19):3869-3878.
 19. van Loenen MM, de Boer R, van Liempt E, et al. A Good Manufacturing Practice procedure to engineer donor virus-specific T cells into potent anti-leukemic effector cells. *Haematologica*. 2014;99(4):759-768.
 20. Aglietta M, Piacibello W, Stacchini A, et al. Expression of HLA class II (DR, DQ) determinants by normal and chronic myeloid leukemia granulocyte/monocyte progenitors. *Cancer Res*. 1986;46(4 Pt 1):1783-1787.
 21. Falkenburg JH, Jansen J, van der Vaart-Duinkerken N, et al. Polymorphic and monomorphic HLA-DR determinants on human hematopoietic progenitor cells. *Blood*. 1984;63(5):1125-1132.
 22. Orgad S, Yunis EJ, Zaizov R, Ramot B, Altshuler R, Gazit E. Expression of HLA-DR alloantigens on acute lymphoblastic leukemia lymphoblasts. *Hum Immunol*. 1984;9(2):67-74.
 23. Kobayashi KS, van den Elsen PJ. NLR5: a key regulator of MHC class I-dependent immune responses. *Nature reviews Immunology*. 2012;12(12):813-820.
 24. Griffioen M, van der Meijden ED, Slager EH, et al. Identification of phosphatidylinositol 4-kinase type II beta as HLA class II-restricted target in graft versus leukemia reactivity. *Proc Natl Acad Sci U S A*. 2008;105(10):3837-3842.
 25. Stumpf AN, van der Meijden ED, van Bergen CA, Willemze R, Falkenburg JH, Griffioen M. Identification of 4 new HLA-DR-restricted minor histocompatibility antigens as hematopoietic targets in antitumor immunity. *Blood*. 2009;114(17):3684-3692.
 26. Spaapen RM, de Kort RA, van den Oudenalder K, et al. Rapid identification of clinical relevant minor histocompatibility antigens via genome-wide zygosity-genotype correlation analysis. *Clin Cancer Res*. 2009;15(23):7137-7143.
 27. Spaapen RM, Lokhorst HM, van den Oudenalder K, et al. Toward targeting B cell cancers with CD4+ CTLs: identification of a CD19-encoded minor histocompatibility antigen using a novel genome-wide analysis. *The Journal of experimental medicine*. 2008;205(12):2863-2872.
 28. Meyer RG, Wagner EM, Konur A, et al. Donor CD4 T cells convert mixed to full donor T-cell chimerism and replenish the CD52-positive T-cell pool after alemtuzumab-based T-cell-depleted allo-transplantation. *Bone marrow transplantation*. 2010;45(4):668-674.
 29. Rutten CE, van Luxemburg-Heijs SA, Halkes CJ, et al. Patient HLA-DP-specific CD4+ T cells from HLA-DPB1-mismatched donor lymphocyte infusion can induce graft-versus-leukemia reactivity in the presence or absence of graft-versus-host disease. *Biology of blood and marrow transplantation : journal of the American Society for Blood and Marrow Transplantation*. 2013;19(1):40-48.

30. Herr W, Eichinger Y, Beshay J, et al. HLA-DPB1 mismatch alleles represent powerful leukemia rejection antigens in CD4 T-cell immunotherapy after allogeneic stem-cell transplantation. *Leukemia*. 2017;31(2):434-445.
31. Stevanovic S, van Bergen CA, van Luxemburg-Heijs SA, et al. HLA class II upregulation during viral infection leads to HLA-DP-directed graft-versus-host disease after CD4+ donor lymphocyte infusion. *Blood*. 2013;122(11):1963-1973.
32. Kremer AN, van der Meijden ED, Honders MW, et al. Endogenous HLA class II epitopes that are immunogenic in vivo show distinct behavior toward HLA-DM and its natural inhibitor HLA-DO. *Blood*. 2012;120(16):3246-3255.
33. van Bergen CA, van Luxemburg-Heijs SA, de Wreede LC, et al. Selective graft-versus-leukemia depends on magnitude and diversity of the alloreactive T cell response. *The Journal of clinical investigation*. 2017;127(2):517-529.
34. Meyer RG, Britten CM, Wehler D, et al. Prophylactic transfer of CD8-depleted donor lymphocytes after T-cell-depleted reduced-intensity transplantation. *Blood*. 2007;109(1):374-382.
35. von dem Borne PA, Starrenburg CW, Halkes SJ, et al. Reduced-intensity conditioning allogeneic stem cell transplantation with donor T-cell depletion using alemtuzumab added to the graft ('Campath in the bag'). *Current opinion in oncology*. 2009;21 Suppl 1:S27-29.
36. Glucksberg H, Storb R, Fefer A, et al. Clinical manifestations of graft-versus-host disease in human recipients of marrow from HL-A-matched sibling donors. *Transplantation*. 1974;18(4):295-304.
37. Van Bergen CA, Rutten CE, Van Der Meijden ED, et al. High-throughput characterization of 10 new minor histocompatibility antigens by whole genome association scanning. *Cancer Res*. 2010;70(22):9073-9083.
38. Jedema I, van der Werff NM, Barge RM, Willemze R, Falkenburg JH. New CFSE-based assay to determine susceptibility to lysis by cytotoxic T cells of leukemic precursor cells within a heterogeneous target cell population. *Blood*. 2004;103(7):2677-2682.
39. Adams S, Robbins FM, Chen D, et al. HLA class I and II genotype of the NCI-60 cell lines. *Journal of translational medicine*. 2005;3(1):11.
40. Neefjes J, Jongsmá ML, Paul P, Bakke O. Towards a systems understanding of MHC class I and MHC class II antigen presentation. *Nature reviews Immunology*. 2011;11(12):823-836.
41. Kremer AN, van der Meijden ED, Honders MW, et al. Human leukocyte antigen-DO regulates surface presentation of human leukocyte antigen class II-restricted antigens on B cell malignancies. *Biology of blood and marrow transplantation : journal of the American Society for Blood and Marrow Transplantation*. 2014;20(5):742-747.

SUPPLEMENTARY FIGURE



Supplementary Figure 1. Expression of HLA class II restriction molecule for different MiHA on fibroblasts measured by flowcytometry.