**Universiteit
Leiden**
The Netherlands

Reversal Agents for the Direct Factor Xa Inhibitors: Biochemical Mechanisms of Current and Newly Emerging Therapies

Schreuder, M.; Reitsma, P.H.; Bos, M.H.A.

Citation

Schreuder, M., Reitsma, P. H., & Bos, M. H. A. (2020). Reversal Agents for the Direct Factor Xa Inhibitors: Biochemical Mechanisms of Current and Newly Emerging Therapies. *Seminars In Thrombosis And Hemostasis*. Retrieved from <https://hdl.handle.net/1887/3230119>

Version: Publisher's Version

License: [Leiden University Non-exclusive license](#)

Downloaded from: <https://hdl.handle.net/1887/3230119>

Note: To cite this publication please use the final published version (if applicable).

Reversal Agents for the Direct Factor Xa Inhibitors: Biochemical Mechanisms of Current and Newly Emerging Therapies

Mark Schreuder, MSc¹ Pieter H. Reitsma, PhD¹ Mettine H. A. Bos, PhD¹

¹ Division of Thrombosis and Hemostasis, Einthoven Laboratory for Vascular and Regenerative Medicine, Leiden University Medical Center, Leiden, The Netherlands

Semin Thromb Hemost

Address for correspondence Mettine H. A. Bos, PhD, Division of Thrombosis and Hemostasis, Einthoven Laboratory for Vascular and Regenerative Medicine, Leiden University Medical Center, Room C7-14, Albinusdreef 2, 2333 ZA, Leiden, The Netherlands (e-mail: M.H.A.Bos@lumc.nl).

Abstract

The direct oral anticoagulants targeting coagulation factor Xa or thrombin are widely used as alternatives to vitamin K antagonists in the management of venous thromboembolism and nonvalvular atrial fibrillation. In case of bleeding or emergency surgery, reversal agents are helpful to counteract the anticoagulant therapy and restore hemostasis. While idarucizumab has been established as an antidote for the direct thrombin inhibitor dabigatran, reversal strategies for the direct factor Xa inhibitors have been a focal point in clinical care over the past years. In the absence of specific reversal agents, the off-label use of (activated) prothrombin complex concentrate and recombinant factor VIIa have been suggested as effective treatment options during inhibitor-induced bleeding complications. Meanwhile, several specific reversal agents have been developed. In this review, an overview of the current state of nonspecific and specific reversal agents for the direct factor Xa inhibitors is provided, focusing on the biochemistry and mechanism of action and the preclinical assessment of newly emerging therapies.

Keywords

- ▶ factor Xa inhibitors
- ▶ hemorrhage
- ▶ anticoagulants
- ▶ antidotes

The direct oral anticoagulants (DOACs) are a class of synthetic anticoagulant drugs that consist of a direct thrombin inhibitor (dabigatran) and several direct factor (F) Xa (FXa) inhibitors (apixaban, rivaroxaban, edoxaban, and betrixaban). These drugs are characterized by a superior pharmacodynamic profile relative to the classic oral anticoagulant vitamin K antagonists (VKAs), and their use is therefore considered as the preferred anticoagulant therapy.¹ The DOACs are prescribed for the prevention of venous thrombosis following surgery or during atrial fibrillation and for the treatment of venous thromboembolism.² Unfortunately, as with all anticoagulant treatment options, bleeding complications associated with DOAC use remain the main concern in clinical practice. Therefore, rapid reversal of the anticoagulant DOAC activity may be required in case of life-threatening hemorrhage, major trauma, emergency surgery, or DOAC overdose. DOAC reversal can be achieved by the use of nonspecific or specific reversal

agents. In the absence of specific reversal agents, nonspecific plasma protein concentrates and recombinant FVIIa have been suggested to overcome the anticoagulant effects of the DOACs. Specific reversal of the thrombin-targeting DOAC dabigatran is effectively achieved by administration of monoclonal antibody fragment idarucizumab, which is approved by the U.S. Food and Drug Administration (FDA) and the European Medicines Agency (EMA).³ More recently, andexanet alfa, a specific antidote for the FXa-targeting DOACs, was approved for the treatment of uncontrollable bleeds. In addition, several specific antidotes for the FXa-targeting DOACs are currently in preclinical development, which include the synthetic small molecule ciraparantag, a modified FXa- α 2 macroglobulin complex, and several prohemostatic FX-based reversal agents. Since the direct FXa inhibitors apixaban and rivaroxaban are increasingly prescribed rather than the direct thrombin inhibitor dabigatran,⁴ the importance of an effective reversal agent

Issue Theme Recent Advances in Thrombosis and Hemostasis—Part VI; Guest Editor: Sam Schulman, MD, PhD.

Copyright © by Thieme Medical Publishers, Inc., 333 Seventh Avenue, New York, NY 10001, USA.
Tel: +1(212) 760-0888.

DOI <https://doi.org/10.1055/s-0040-1709134>.
ISSN 0094-6176.

for the direct FXa inhibitors is substantial. In this review, we will highlight the reversal of the FXa-targeting DOACs by specific and nonspecific reversal agents, with the main focus on the biochemical mechanisms of the newly emerging reversal therapies that are in preclinical development.

Mechanistic Principles of the Direct Factor Xa Inhibitors

The FXa-targeting DOACs are small synthetic molecules with overall similar chemical structures that have identical mechanisms of action. Their anticoagulant effect stems from the fact that these inhibitors interact with the active site pocket of FXa, thereby effectively blocking the natural substrate prothrombin from accessing the active site (► **Fig. 1**) and preventing thrombin formation. The characteristic L shape of these small molecule inhibitors mediates the high-affinity interaction with the FXa active site, resulting in a potent subnanomolar FXa inhibitory activity (K_i of 0.08–0.56 nM).^{5–7} Although the overall composition of the active site of serine proteases is generally conserved,⁸ the direct FXa inhibitors have a more than 10,000-fold greater selectivity for FXa compared with other human serine proteases.⁹ Binding to FXa is reversible and regulated through the S1 and S4 active site subpockets.^{5,6} The *p*-methoxyphenyl P_1 moiety of apixaban and the chlorothiophene moiety of rivaroxaban are inserted deep into the S1 specificity pocket. Although the P_1 moiety of apixaban does not appear to directly

interact with FXa, the choline substituent of rivaroxaban interacts with the aromatic ring of Tyr228 (chymotrypsinogen numbering) that is located at the bottom of the S1 subpocket and provides a high-affinity interaction.^{5,6} Apixaban interacts with the Gln192 backbone through its pyrazole N-2 and with Gly216 through a carboxyl interaction.⁶ Rivaroxaban forms two hydrogen bonds with Gly219 that serve an important role by directing its substituents into the S1 and S4 subpockets.⁵ The S4 subsite of FXa is a narrow hydrophobic pocket defined by the aromatic rings of residues Tyr99, Phe174, and Trp215.¹⁰ In this binding pocket, the P_4 ring of apixaban and the morpholinone ring of rivaroxaban are sandwiched between the residues Tyr99 and Phe174, thereby providing important binding stability and enabling high-affinity binding.^{5,6} While structural information on the interaction of the direct FXa inhibitors edoxaban or betrixaban with the FXa active site is lacking, these are considered to also engage the FXa S1 and S4 subsites.¹¹

The pharmacokinetic and pharmacodynamic profiles of the direct FXa inhibitors have several key advantages over that of the VKAs. Their rapid onset makes them ideal for clinical use as no bridging therapies are needed and they sustain their anticoagulant effect up to 24 hours, with a half-life of 6 to 14 hours.^{12,13} Unlike the VKAs, the direct FXa inhibitors are characterized by linear and predictable pharmacodynamics, no known interactions with food components, and limited drug–drug interactions.¹⁴ It is for these reasons that patients treated with a DOAC do not require routine monitoring.

Fig. 1 Binding mode of the direct factor Xa (FXa) inhibitors. (A) Surface representation of the FXa X-ray structure in complex with the direct FXa inhibitor apixaban shown in red. (B,C) Close-up view of apixaban (B) and rivaroxaban (C) binding to FXa. Apixaban and rivaroxaban are depicted in orange, with the oxygen atoms indicated in red and nitrogen atoms in blue. Essential amino acids in FXa are shown in stick configuration, in which the catalytic triad residues His57, Asp102, and Ser195 are colored in red. All FXa residues are indicated in chymotrypsinogen numbering. Hydrogen bond interactions are shown in teal dashed lines.

Reversal of the Direct Factor Xa Inhibitors

Bleeding complications remain a major concern in anticoagulant therapy. Compared with the conventional VKAs, the direct FXa inhibitors are associated with comparable or slightly lower risks of major bleeding, stroke, and fatality. Still, major bleeding is observed in 1 to 5% of the patients receiving DOACs.^{15–17} Since millions of people worldwide require anticoagulant treatment, and this is projected to increase significantly in the coming decades, bleeding complications will remain a major health care burden.^{18,19} To treat these DOAC-associated bleeding complications, several nonspecific reversal agents used in hemophilia-associated and VKA-induced bleedings have been suggested as potential candidates for reversal therapy, although supporting evidence is lacking or incomplete. In addition to these nonspecific treatment options, the specific antidote for the direct FXa inhibitors andexanet alfa has recently been approved by the FDA and EMA and could significantly improve the management of major bleeding events.^{20,21} Furthermore, various specific reversal agents with different modes of action are currently in preclinical development. In the following sections, we will discuss the efficacy of the nonspecific and specific reversal strategies and focus on their molecular details and mechanisms of action.

Nonspecific Reversal Strategies

Prothrombin Complex Concentrate

Prothrombin complex concentrates (PCCs) are derived from human plasma and contain vitamin K-dependent coagulation factors that are partly purified through ion-exchange chromatography.^{22,23} Strong ion exchangers produce four-factor concentrates (4F-PCC), consisting of FVII, FIX, FX, and prothrombin, whereas weak ion exchangers produce three-factor concentrates (3F-PCC) that do not contain FVII. Most PCC products have been supplemented with heparin to prevent factor activation²⁴ and may contain various concentrations of the vitamin K-dependent anticoagulants protein C and protein S. PCCs are used for the reversal of VKAs by replenishing the lowered and defective circulating vitamin K-dependent factors if patients present with an urgent need to restore hemostasis.²⁵ In contrast, the direct FXa inhibitors do not affect the level or posttranslational modification of coagulation factors, thereby obviating the necessity for factor supplementation and replacement. PCC-mediated reversal of FXa inhibition has been studied in several animal models and small-sized human studies. In rabbit models of apixaban- or rivaroxaban-induced bleeding, 4F-PCC partially improved laboratory parameters but was not able to reduce blood loss.^{26,27} In contrast, 4F-PCC significantly shortened the bleeding time of rivaroxaban-induced bleeding in a rat model and could reduce the prothrombin time (PT) in rivaroxaban-treated baboon plasma.²⁸ 4F-PCC was also able to dose-dependently reverse hematoma expansion in a murine intracerebral hemorrhage model following pretreatment with rivaroxaban.²⁹ In humans, the efficacy of PCC has mostly been assessed in healthy nonbleeding volunteers using surrogate laboratory tests. Similar to the animal models, these studies have led to variable reports on the efficacy of direct FXa reversal

by PCC. While both 3F-PCC and 4F-PCC administrations were demonstrated to restore some parameters of coagulation (PT and endogenous thrombin potential [ETP] in thrombin generation [TG]) in rivaroxaban-treated healthy individuals, other parameters such as the activated partial thromboplastin time (aPTT), anti-FXa activity, and the TG lag time were unaffected.^{30–33} Similar results were obtained following 3F-PCC and 4F-PCC administration to apixaban- or edoxaban-dosed healthy individuals.^{34–37} A more clinically relevant study, in which bleeding following a punch biopsy in edoxaban-treated healthy individuals was assessed, revealed that 4F-PCC reversed edoxaban's effect on the duration of bleeding and on the ETP, whereas the PT and total blood loss could only be partially restored.³⁸ It is important to consider that restoration of laboratory parameters has to be interpreted with caution as these do not accurately represent the clinical setting. To overcome this, the efficacy of PCC has been evaluated in the management of major bleeding events in patients using the direct FXa inhibitors. In clinical studies, 4F-PCC proved to be effective in 65 to 75% of the major bleeding events in patients treated with apixaban or rivaroxaban.^{39–42} Collectively, the available clinical data indicate that both 3F-PCC and 4F-PCC may be effective reversal agents for the majority of patients presenting with major bleeding events associated with direct FXa inhibitor therapy. However, larger studies would be required to evaluate the true potential of this nonspecific reversal strategy in a real-world patient population. When specific reversal agents are unavailable or unattainable, PCC is considered as the most effective reversal agent and has therefore been included as the first treatment of choice in life-threatening bleeding events.⁴³

Activated Prothrombin Complex Concentrate

Activated PCC (aPCC), or factor eight inhibitor bypassing activity (FEIBA), is a PCC product that contains small amounts of FVIII and the activated FVIIa, FIXa, and FXa, and thrombin. This product was originally used as a bypassing agent for the treatment of bleeding complications in hemophilia patients with inhibitors.^{44–46} Given that aPCC elevates the circulating levels of several coagulation factors, thereby stimulating coagulation, it has the potential to be a reversal agent in DOAC-induced bleeding. Perzborn et al reported that aPCC significantly reduced the bleeding time and partially restored the PT in rivaroxaban-induced bleeding in rat and baboon models.²⁸ Similarly, aPCC was found to dose-dependently reduce the bleeding time in edoxaban-treated rats.⁴⁷ Furthermore, in vitro and ex vivo studies have shown significant but variable improvements of surrogate end points, including PT, aPTT, anti-FXa activity, and several TG parameters.^{30,48–51} The administration of aPCC as a reversal agent for direct FXa inhibitors has also been evaluated in a few patient studies. In two small-scale studies, aPCC-mediated reversal of apixaban and rivaroxaban was associated with a reduction in intracranial hemorrhage (ICH) expansion.^{52,53} In two larger studies, administration of aPCC led to an overall stable hemorrhage progression and a 14 to 29% mortality rate.^{54,55} Collectively, aPCC is associated with the reversal of direct FXa inhibitors in major bleeding events. Therefore, the off-label use of aPCC has

been included in the guidelines for the management of uncontrolled bleeding in patients on DOACs.⁴³ However, as more (pre)clinical data have become available, pointing to a beneficial effect of PCCs in this setting, these are currently preferred over aPCC as nonspecific reversal agents.

Recombinant Activated Factor VII

Recombinant human activated FVII (rFVIIa), or NovoSeven (eptacog alfa), was initially developed for the treatment of hemophilia in patients presenting with inhibitors.^{56,57} However, its off-label use has increased significantly for uncontrolled bleeding following trauma and surgery,⁵⁸ and rFVIIa has been suggested as a treatment option in DOAC-related bleeding. Evaluation of the efficacy of rFVIIa in animal models and healthy volunteers has shown variable results. In edoxaban-treated rats, rFVIIa dose-dependently reduced the bleeding time and restored the PT.⁴⁷ Comparable results were obtained for rivaroxaban reversal employing rat and baboon models.²⁸ Conversely, rFVIIa did not affect the PT or blood loss in an apixaban- and rivaroxaban-induced bleeding model in rabbits, whereas the aPTT and several rotational thrombelastography (ROTEM) parameters were at least par-

tially restored.^{26,27} These inconsistent and thus far unexplained effects reflect in vitro reversal studies, displaying mostly marginal effects of rFVIIa on the restoration of laboratory parameters.^{30,48,50,51,59} Thus far, no reports on the efficacy of rFVIIa in major bleeding events that are associated with DOAC therapy are available. Therefore, the benefit of rFVIIa therapy in a nonhemophilia setting remains uncertain, and as such, clinical guidelines state that rFVIIa might be considered for the reversal of the direct FXa inhibitors on an individual basis.⁶⁰

Mechanistic Principles of the Nonspecific Reversal Agents

Although the exact mechanism by which the nonspecific reversal agents PCC, aPCC, and rFVIIa bypass the anticoagulant effect of the DOACs is currently not fully understood, it is generally acknowledged to result from an increase in factor concentrations that promote TG (►Fig. 2). The latter is also at the basis of their FVIII bypassing activity that supports their use in hemophilia treatment. The plasma concentrates contain a variety of procoagulant clotting factors, with coagulation FX and prothrombin being considered the main procoagulant components.⁶¹ Turecek et al revealed that aPCC was unable

Fig. 2 Schematic representation of the procoagulant actions exerted by nonspecific reversal agents. The nonspecific reversal agents promote various procoagulant reactions of the coagulation cascade to drive thrombin generation in the presence of the direct factor Xa inhibitors. The overlapping actions of the PCC and aPCC are indicated in green arrows. The distinct effects of aPCC are shown in red and that of rFVIIa are shown in blue. PCC, nonactivated prothrombin complex concentrate; aPCC, activated prothrombin complex concentrate; rFVIIa, recombinant activated factor VIIa; TF, tissue factor.

to generate thrombin in FV-deficient plasma, indicating that prothrombinase activity is essential.⁶² Furthermore, a highly purified complex of FXa and prothrombin was able to shorten the prolonged aPTT in hemophilia A plasma in a similar manner as aPCC. The fact that both factors are essential was illustrated by TG assessment, in which FXa addition induced a slight improvement in thrombin formation, whereas subsequent prothrombin supplementation dramatically increased the amount of thrombin formed.^{62,63} Consequently, administration of FXa and prothrombin in equivalent amounts similar to aPCC administration, normalized blood loss in a rabbit hemophilia model.^{62,63} These results show that the administration of FXa and its substrate prothrombin is sufficient to initiate blood coagulation. As FXa is capable of directly activating FV,⁶⁴ a positive feedback loop is generated through the assembly of the prothrombinase complex and the conversion of prothrombin to thrombin.⁶⁵ Additionally, the thrombin formed will initiate the intrinsic pathway by the activation of factors VIII and XI, thereby further propagating FXa and TG. While the latter mechanism describes the mode of action of aPCC, PCC is considered to mainly function by elevating the circulating levels of several coagulation factor substrates. In general, increasing the substrate concentration for the appropriate enzymes will enhance the catalytic turnover if the substrate concentration is limited. As such, PCC likely achieves its procoagulant effect by supplementation of the FIX (substrate of factors VIIa and XIa) and FX (substrate of FVIIa and FIXa) following PCC administration.

While the aforementioned mode of action predicts a procoagulant response under conditions where bypassing of FVIII is required, the question remains how this mechanism explains effectiveness in the presence of DOACs given that the latter will target the newly formed FXa or thrombin. A possible explanation can be found in the reversible binding nature of these inhibitors. The direct inhibitors interact with the active site pocket of their respective targets and, as a result, compete with the natural and irreversible inhibitor antithrombin that engages the FXa and thrombin active site through the S1 subsite.^{66,67} In the absence of the DOACs, antithrombin rapidly inhibits free FXa or thrombin. However, Thalji et al described that with the direct FXa inhibitor rivaroxaban present, the majority of FXa reversibly interacts with rivaroxaban rather than forming the slower but irreversible FXa-antithrombin complex.⁶⁸ As such, an equilibrium is formed between the FXa-antithrombin and the FXa-rivaroxaban complexes, in addition to the generation of a pool of free and uninhibited FXa. As the free FXa reaches steady state, a persistent amount of free FXa is established under these conditions. While in most scenarios, this pool of free FXa is likely functionally irrelevant, the enhanced factor levels and the subsequent increase in FXa formed ensuing supplementation with the nonspecific reversal agents could significantly elevate the steady-state levels of free FXa. Thalji et al suggest that these elevated levels of free FXa restore hemostasis in the presence of rivaroxaban.⁶⁸

These mechanistic principles might also provide an explanation for the reduced effectiveness of rFVIIa in DOAC reversal. In hemophilia, rFVIIa stimulates blood clot formation by FXa generation through tissue factor localized at the site of injury as

well as on the surface of platelets independent of tissue factor.⁶⁹ While PCC and aPCC promote the generation of multiple active serine proteases, rFVIIa focuses largely on the formation of FXa. The rate of FXa and subsequent TG could therefore be lower and prove insufficient in the presence of the DOACs.

Specific Reversal Strategies

Andexanet Alfa

The specific antidote andexanet alfa (andexanet) is a universal reversal agent for the direct FXa inhibitors and is currently the only FDA- and EMA-approved treatment option for major bleeding events induced by these anticoagulants (►Fig. 3). Andexanet is a recombinant modified human FXa variant that is expressed in its mature FXa form due to the replacement of the activation peptide by a double Arg-Lys-Arg linker (RKRRKR), which is intracellularly and proteolytically removed by proprotein convertases.²¹ Furthermore, andexanet lacks the membrane-binding γ -carboxyglutamic acid (GLA) domain of native FXa, which prevents its assembly into the prothrombinase complex. Additionally, andexanet is enzymatically inactive due to the replacement of the catalytic triad residue serine 195 by alanine.²¹ Despite this active site substitution, andexanet displays a similar apparent binding affinity for the direct FXa inhibitors compared with wild-type (wt) FXa.²¹ Consequently, andexanet will compete with endogenous FXa for the direct inhibitor binding and as such, when administered in molar excess, will effectively bind and clear the circulation of inhibitors, thereby restoring TG through liberated endogenous FXa (►Fig. 3). This mechanism has been proven effective against all currently approved direct FXa inhibitors, as andexanet dose-dependently reversed the anti-FXa activity in both purified and plasma systems.^{21,70} Andexanet was also able to restore hemostasis in an inhibitor-infused rat model and was shown to reduce blood loss following liver laceration in a rabbit bleeding model.^{21,70,71} Reversal of the direct FXa inhibitors was not associated with major adverse effects in monkeys, as no histopathological evidence of andexanet-related thrombosis was observed.⁷¹ Based on these data, a randomized placebo-controlled study was performed in healthy volunteers treated with apixaban (ANNEXA-A) or rivaroxaban (ANNEXA-R).⁷² Within minutes after a bolus injection or a bolus injection followed by a 1- to 2-hour infusion of andexanet, the anti-FXa activity was reversed by more than 90% and the ETP was restored in both apixaban- or rivaroxaban-treated volunteers. These parameters returned gradually to placebo levels within 1 to 2 hours as andexanet was eliminated from the circulation. Moreover, in a safety and dose escalation study in healthy individuals, andexanet was well tolerated and thrombotic events were not observed despite the detection of increased levels of D-dimer and prothrombin fragment 1.2.⁷³

The ANNEXA-4 study was designed to examine the use of andexanet in patients with potential life-threatening bleeding.²⁰ Recently, the study was completed and included 352 patients with major bleeding complications. While in 92% of the patients, treatment with andexanet resulted in excellent or good efficacy as determined by anti-FXa activity, this primary outcome did not correlate with the overall

Fig. 3 Schematic representation of human factor Xa (FXa) and the reversal agent andexanet alfa. **(A)** The light chain of FXa consisting of the GLA (Gla), the endothelial growth factor 1 and 2 domains, and the heavy chain comprising the SP and AS are shown linked by a disulfide bond (S-S). The heavy-chain NT, formed upon proteolytic activation of factor X, is shown as an insertion into the active site pocket, representing the salt bridge between Ile16 and Asp194. **(B)** Schematic representation of andexanet alfa. Despite being enzymatically inactive due to a Ser195Ala substitution, andexanet alfa encompasses a high binding affinity for the direct FXa inhibitors, which are shown in purple. Removal of the GLA domain prevents incorporation of andexanet alfa into the prothrombinase complex. AS, active site pocket; EGF, endothelial growth factor domain; GLA, γ -carboxyglutamic acid domain; NT, N-terminus; SP, serine protease domain.

hemostatic efficacy (good efficacy in 82% of the evaluated patients). Additionally, thrombotic events occurred in 10% of the patients within 30 days following andexanet treatment, which was relatively higher than idarucizumab-mediated reversal of dabigatran (4.8%)³ and other anticoagulant reversal studies (3.9–7%).^{25,74} A possible explanation for the relatively high percentage of thrombosis might be the interaction of andexanet with the natural anticoagulant tissue factor pathway inhibitor (TFPI). Ersayin et al showed that andexanet demonstrates high-affinity binding to TFPI, similar to GLA-domainless FXa.⁷⁵ The TFPI-neutralizing capacity of andexanet was further underscored by the observation that the addition of supratherapeutic levels of andexanet to hemophilic or normal pooled plasma (in the absence of an FXa inhibitor) resulted in a dose-dependent increase in the thrombin peak height in a TFPI-sensitive TG assay.^{75–77} Importantly, a major limitation of the ANNEXA-4 study is the absence of a randomized comparison with PCC, the standard treatment option in the absence of a specific reversal agent. Interestingly, such a randomized comparison with PCC is also lacking for idarucizumab. A randomized controlled trial evaluating andexanet with a PCC-control group is underway (NCT03661528).

Specific Reversal Strategies in Clinical Development

Ciraparantag

Ciraparantag (aripazine, PER 977) is a synthetic small molecule that consists of two L-arginine units connected with a piperazine containing linker chain, which is considered a universal anticoagulant antidote capable of reversing the

effects of low-molecular-weight heparins and the direct thrombin and FXa inhibitors^{78–80} (**Fig. 4A**). Data on this antidote are scarce and mainly derived from abstracts. As such, the mechanistic principles remain largely unknown. Yet, it has been stated that ciraparantag binds directly to the DOACs and heparins through noncovalent hydrogen bonding based on *in silico* modeling and dynamic light scattering data, whereas binding to any human coagulation factor or albumin was not observed.⁸¹ Therefore, the mode of action of ciraparantag appears relatively similar to that of andexanet: interaction with and removal of the inhibitors from their intended target to generate free endogenous FXa. In line with this, ciraparantag was observed to completely reverse the anti-FXa activity in apixaban- and rivaroxaban-spiked human plasma.⁸² Additionally, *in vivo* assessments employing rat tail transection and rat and rabbit liver laceration models revealed that ciraparantag significantly decreased apixaban-, rivaroxaban-, and edoxaban-induced bleeding.^{81–84} Moreover, ciraparantag was well tolerated in rats and dogs in doses up to 20 mg/kg per day, and no toxicity was observed.⁸⁵ Data from a first-in-human study in 80 edoxaban-treated healthy volunteers who received an intravenous injection of 100 or 300 mg of ciraparantag demonstrated a shortening of the whole blood clotting time to within 10% of baseline levels and restoration of the fibrin clot diameter.^{78,86} In addition, ciraparantag did not induce serious adverse effects or an increase in the procoagulant markers D-dimer or prothrombin fragment 1.2, nor were alterations in TFPI levels observed.

One of the main concerns when addressing the therapeutic potential of ciraparantag is the fact that no data concerning the

Fig. 4 Schematic representation of the universal reversal agent ciraparantag and the GLA-domainless factor Xa (FXa) in complex with $\alpha 2$ -macroglobulin. (A) Ciraparantag (the green squares represent two L-arginine units at each end that are connected through a linker consisting of three units, of which a piperazine comprises the middle unit) reportedly binds directly to the direct FXa inhibitors (shown in purple) through hydrogen bonding, thereby sequestering the inhibitors from the circulation. (B) $\alpha 2$ -Macroglobulin consists of four identical subunits that allow for the interaction with two FXa proteases. Despite being trapped, GLA-domainless FXa retains its ability to bind the direct FXa inhibitors (shown in purple), thereby clearing the inhibitors from the circulation. Complex formation with $\alpha 2$ -macroglobulin further prevents macromolecular interactions with FXa.

direct binding of ciraparantag to the DOACs have been published thus far. This complicates our understanding of the mechanisms that are at the basis of the apparent selectivity of ciraparantag for heparins and DOACs and prevent its interaction with other proteins and molecules. These concerns are also highlighted by studies in which ciraparantag was shown to be incapable of restoring the anti-FXa activity of either apixaban, rivaroxaban, or edoxaban in plasma or a purified system.^{87,88} Furthermore, ciraparantag was found to be unable to restore the augmented clotting time induced by rivaroxaban and edoxaban but could normalize the clot structure and fibrin fiber diameter.⁸⁷ More importantly, using isothermal titration calorimetry, direct binding of ciraparantag to either edoxaban or rivaroxaban could not be established.⁸⁷ In contrast, a direct interaction with FIXa, but not FIX, was observed,⁸⁷ corroborating earlier reports on ciraparantag-dependent stimulation of the FIXa-dependent activation of FX.⁸⁸ These results may suggest a role in FX activation for ciraparantag, as opposed to direct binding and interfering with anticoagulant molecules. It is essential that its exact mechanistic principles are elucidated and potential off-target effects are fully characterized in order for ciraparantag to be used in clinical practice.

Specific Reversal Strategies in Preclinical Development

GLA-Domainless Factor Xa in Complex with $\alpha 2$ -Macroglobulin

Similar to andexanet, Jourdi et al have generated a specific antidote for the direct FXa inhibitors based on GLA-domainless

FXa.⁸⁹ As andexanet might interact with several pro- and anticoagulant macromolecules such as FVa, prothrombin, antithrombin, and most importantly TFPI,^{21,75–77} Jourdi et al hypothesized that a complex between GLA-domainless FXa and $\alpha 2$ -macroglobulin (GDFXa- $\alpha 2$ M) would prevent these macromolecular interactions while retaining its ability to sequester the direct FXa inhibitors (**Fig. 4B**). $\alpha 2$ M is a molecular trap inhibitor that targets a broad spectrum of proteases in the circulation without directly blocking the protease active site.^{90,91} The inhibitor is composed of four identical subunits, each including a bait region targeted by the protease, although only two proteases are able to interact with the tetramer. Upon interaction and cleavage by the protease, $\alpha 2$ M undergoes an irreversible conformational change, which traps the protein within a cage-like structure.^{90,91} By making use of GLA-domainless FXa that retains its native active site, as opposed to catalytically inactive andexanet due to the Ser195Ala substitution, incubation with $\alpha 2$ M results in cleavage of the $\alpha 2$ M bait region, generating the irreversible GDFXa- $\alpha 2$ M complex. Evaluation of the constants of apixaban and rivaroxaban inhibition revealed somewhat lower binding affinities for interaction with the GDFXa- $\alpha 2$ M complex,^{89,92} although still in the low nanomolar range (1–2.5 nM), similar to wt-FXa. Furthermore, GDFXa- $\alpha 2$ M could fully reverse the effects of supraphysiological levels of rivaroxaban and apixaban in human plasma, and it restored blood loss and bleeding time in a rivaroxaban-treated mouse model following lateral tail vein transection.⁸⁹ Importantly, the GDFXa- $\alpha 2$ M complex did not have any detectable effect on a clot waveform assay and was found to be resistant to heparin-catalyzed antithrombin

and TFPI, indicating that the trapped FXa does not interact with any of the naturally occurring anticoagulation proteins. This was corroborated by the observation that the D-dimer and thrombin-antithrombin (TAT) complex levels in mice following GDFXa- α 2M administration were comparable to vehicle-treated controls. Surprisingly, GDFXa- α 2M also partially neutralized the anticoagulant effect of dabigatran.⁸⁹ Although the mechanism that lies at the basis of this observation has not been defined thus far, it could involve a nonspecific dabigatran binding site in α 2M. Together, the GDFXa- α 2M complex provides an alternative for andexanet and could potentially be a safer option as it does not interact with relevant macromolecular molecules, that is, prothrombin, antithrombin, and TFPI. However, since α 2M is a large 720-kDa protein, large-scale recombinant production may prove difficult, requiring purification from human plasma.

Zymogen-like Factor Xa (FXa-Ile16Leu)

Besides specific bait antidotes that bind and sequester the direct FXa inhibitors, several prohemostatic bypassing agents have been developed that are able to enhance TG in the presence of the FXa-targeting inhibitors. One such reversal agent is an FXa variant comprising an Ile16Leu substitution (chymotrypsin numbering)^{68,93} (► Fig. 5A). Upon activation through proteolytic cleavage at the highly conserved Arg15-Ile16 site, serine proteases are converted into their protease state by the insertion of the newly formed N-terminus in the active site, thereby forging an internal salt bridge of Ile16 with Asp194. This transition induces maturation of the serine protease domain, specifically following conformational rearrangement of the S1 specificity pocket and oxyanion hole.⁹⁴ Disrupting the zymo-

gen-to-protease transition by the substitution of Ile16Leu induces a zymogen-like conformation, in which the active site of FXa retains a partially immature state.⁹³ As a result, FXa-Ile16Leu displays an impaired chromogenic substrate conversion and is less sensitive to small molecule inhibitors. Although FXa-Ile16Leu also demonstrated weaker FVa binding and prothrombin activation, saturating concentrations of the cofactor and anionic membranes partly restored thrombin conversion, suggesting that FVa acts as a stabilizing factor in the zymogen-to-protease transition of this variant. Further assessments revealed a 20- to 40-fold reduced sensitivity for the natural inhibitors antithrombin and TFPI, whereas prothrombinase-assembled FXa-Ile16Leu was similarly inhibited by antithrombin compared with wt-FXa.⁹⁵ Due to the reduced sensitivity, the FXa-Ile16Leu half-life was extended to approximately 40 minutes in plasma and approximately 5 minutes in mice relative to wt-FXa (half-life of ~ 1 minute and < 20 seconds, respectively).⁹⁵⁻⁹⁷ These data indicate that FXa-Ile16Leu is protected from being targeted by natural anticoagulants in the circulation but still retains the ability to catalyze thrombin formation upon binding to FVa on an anionic membrane layer. As such, FXa-Ile16Leu was examined for its bypassing potential in various clinical bleeding settings and was found to be effective in hemophilia (in the presence and absence of inhibitors) and capable of reversing the effects of warfarin.⁹⁵⁻⁹⁸ Moreover, it was hypothesized that the properties of FXa-Ile16Leu might enable this variant to bypass the DOACs. In vitro evaluation by TG and ROTEM assays demonstrated that FXa-Ile16Leu dose-dependently reversed the anti-FXa effects of rivaroxaban-spiked plasma and whole blood.⁶⁸ These results could be translated to murine in vivo models as mouse FXa-Ile16Leu completely restored hemostasis in several thrombosis and bleeding models in the presence of rivaroxaban.⁶⁸ Infusion

Fig. 5 Schematic representation of the prohemostatic reversal agents in preclinical development. (A) Substitution of Ile16 for leucine (shown in red) disrupts the insertion of the newly formed N-terminus upon factor X activation, thereby destabilizing the formation of the characteristic internal salt bridge with Asp194. As a consequence, the active site of FXa-Ile16Leu is partly matured (indicated as a smaller active site), thereby inducing a zymogen-like state. This zymogen-like state prevents the inhibition of FXa-Ile16Leu by the direct factor Xa (FXa) inhibitors (shown in purple). Binding to factor Va rescues the maturation of the serine protease domain. (B) FXa-C is resistant to the direct FXa inhibitors (shown in purple) due to an insertion within the active site 99-loop. This structural element (highlighted in yellow) sterically hinders the binding of the direct FXa inhibitors to the active site. (C) FXa-F174 comprises an alanine or serine substitution of phenylalanine 174 (shown in red). Although the direct FXa inhibitors (shown in purple) are able to engage the active site, substitution of phenylalanine 174 destabilizes inhibitor binding, resulting in a loss of affinity.

of this FXa variant in mice did increase the circulating TAT levels, but only in the absence of rivaroxaban, whereas D-dimer levels remained unaltered.⁶⁸ The possible mechanism by which FXa-Ile16Leu reverses the effect of the direct FXa inhibitors might be similar to that of previously mentioned bypassing agents. In the presence of rivaroxaban, antithrombin and rivaroxaban compete for FXa binding. As a consequence, an equilibrium is formed that is characterized by a steady-state pool of free FXa.⁶⁸ Computational simulations revealed that the free FXa-Ile16Leu steady-state levels were substantially higher compared with those of wt-FXa. However, the net effect on TG was identical to that of wt-FXa due to the reduced catalytic activity of FXa-Ile16Leu. This suggests that besides FXa-Ile16Leu, wt-FXa could also be able to reverse the direct FXa inhibitor-induced anticoagulant state. Yet, FXa-Ile16Leu is dependent on the binding of FVa to rescue its catalytic activity, which could be beneficial in terms of safety and prothrombotic risks. As such, administration of FXa-Ile16Leu would only result in local clot formation at the sites of injury. Collectively, these studies highlight the potential efficacy of FXa-Ile16Leu as a rescue agent in a variety of bleeding disorders including direct FXa inhibitor-induced bleeding complications. This prohemostatic agent therefore provides a rapid-onset alternative for the bait antidotes, thereby requiring significantly lower amounts of the reversal agent to control bleeding.

The clinical development of FXa-Ile16Leu is currently ongoing. Assessment of the pharmacokinetics and pharmacodynamics in rats and monkeys displayed a half-life of 2 to 3 minutes.⁹⁹ Furthermore, FXa-Ile16Leu administration resulted in a dose-dependent shortening of the aPTT in nonhuman primates but simultaneously led to increased TAT levels. A first-in-human dose escalation study in 49 healthy volunteers revealed that FXa-Ile16Leu was well tolerated, with no observed toxicity or serious adverse events.¹⁰⁰ The pharmacodynamics displayed shortening of the aPTT and TG lag time, as well as an increase in TG peak height, D-dimer, TAT, and prothrombin fragments 1.2, with all parameters returning to baseline within 24 hours. Finally, a clinical trial evaluating the safety and tolerability of FXa-Ile16Leu in subjects with spontaneous ICH is underway (NCT02687191). Whether this reversal agent will be further tested in the setting of the direct FXa inhibitors is unknown.

The 99-Loop Factor X Variant

The 99-loop FX variant, also known as FX-C, is a chimeric FX protein comprising a structural modification derived from the so-called isoform FX variant that is expressed in the venom and liver of the eastern common brown snake (*Pseudonaja textilis*)¹⁰¹ (►Fig. 5B). Characterization of the isoform and venom FXa proteases revealed a high resistance for the direct FXa inhibitors rivaroxaban and apixaban. Sequence analysis showed that these FXa variants comprise an extended 99-loop (His91-Asp102) located N-terminal to the S4 subsite. Since the S4 pocket was demonstrated to be a primary binding site for the FXa inhibitors and stabilized their positioning in the active site,^{5,6,101} this nonconserved structural element was hypothesized to be responsible for the observed resistance toward the direct FXa inhibitors. To

further investigate the reduced sensitivity, the 99-loop extension of isoform FX was recombinantly inserted into human FX, thereby generating FX-C. Insertion of this structural element significantly reduced the FXa sensitivity for apixaban and edoxaban (up to 700-fold), although this came at the cost of a threefold lower catalytic efficiency toward the natural substrate prothrombin.¹⁰¹ To examine whether the reduced inhibitor sensitivity of FXa-C was sufficient to restore the thrombin formation, TG assays were employed in apixaban- and edoxaban-spiked plasma. These experiments revealed that the addition of zymogen FX-C resulted in a dose-dependent increase in TG. Moreover, complete TG normalization was observed with the addition of 20 to 40 µg/mL of the FX-C variant. Interestingly, molecular dynamics simulations revealed that the extended 99-loop in FXa-C adopts a helical conformation, which displays substantial mobility. This structural mobility results in the rapid displacement of the apixaban molecule, pointing to steric hindrance leading to impaired inhibitor binding. As such, significant levels of free FXa-C are formed that, despite its reduced prothrombin conversion rate, are able to restore TG parameters to near-normal conditions, indicative of a restored hemostasis. The inhibitor-resistant variant FX-C is currently in preclinical and clinical development.

Phe174-Substituted Factor X Variants

Arising from the FX-C study, a Phe174-substituted FXa variant displayed potential as a bypassing agent for the direct FXa inhibitors (►Fig. 5C). Molecular dynamics simulations of the human FXa-apixaban complex demonstrated a tight inhibitor interaction with the FXa S4 subsite pocket.¹⁰¹ In this pocket, the P₄ moiety of apixaban is sandwiched between the aromatic side-chains of S4 subsite residues Tyr99 and Phe174. Replacing the S4 subsite residues Tyr99 and/or Phe174 with alanine-disrupted apixaban binding results in a markedly decreased FXa sensitivity for the inhibitor. While the Tyr99Ala substitution was accompanied by an almost complete diminished FXa activity, substitution of Phe174 retained the intrinsic clotting potential. As such, human FX variants were generated comprising an alanine (FX-F174A) or serine (FX-F174S) substitution at position 174¹⁰² (►Fig. 5C). These variants displayed a 10-fold reduced sensitivity toward apixaban and a 4- to 7-fold reduced sensitivity for inhibition by rivaroxaban and edoxaban. Similar effects were demonstrated in human plasma spiked with the direct FXa inhibitors, revealing a 6- to 17-fold loss of inhibitor sensitivity. Furthermore, the addition of 30 to 45 µg/mL of these variants completely rescued the thrombin peak height in the presence of 1 µM apixaban and rivaroxaban, whereas an 80% improvement was found in identical conditions with edoxaban. Although further research is needed to fully elucidate the effects of the Phe174-substituted FX variants, these agents might prove to be a potential treatment option.

Conclusion

Bleeding complications associated with DOAC use remain a major concern, creating a critical need for rapid-onset reversal

agents. Although a specific antidote is available for the direct thrombin inhibitor in idarucizumab, counteracting the effect of the direct FXa inhibitors has been a point of focus over the last years. In the absence of specific antidotes for the direct FXa inhibitors, off-label use of prohemostatic agents has been recommended as standard of care. In the meantime, considerable efforts have been made by researchers to fill the unmet need for a specific reversal agent, leading to the recent FDA and EMA approval of andexanet alfa. Furthermore, several specific reversal agents are currently in (pre)clinical development. Since the potential interaction of andexanet with TFPI might raise some concerns, other specific reversal agents that are currently in various stages of (pre)clinical development may prove to be alternative options for reversal. For instance, the GDFXa- α 2M complex did not display interactions with endogenous inhibitors. Another potential option is ciraparantag, a small synthetic molecule that reportedly binds and sequesters the FXa-targeting inhibitors, although its mode of action seems unresolved at this point. Interestingly, several prohemostatic bypassing agents have been developed, including FXa-Ile16Leu, FX-C, and the Phe174-substituted variants FX-F174A and FX-F174S. In contrast to the stoichiometric bait antidotes, their key advantage resides in the limited amount of protein that is needed to correct inhibitor-induced bleeding. Concomitantly, these prohemostatic bypassing agents may increase the risk for thrombotic complications, and, as such, more caution might be required in clinical practice. Additionally, the protein modifications in the FXa variants may induce an antibody response, limiting the efficacy of repeated use. Further characterization of these specific agents is therefore imperative, but a future with specific, safe, and effective reversal agents for the direct FXa inhibitors seems imminent.

Authors' Contribution

M. S. drafted the manuscript, P. H. R. reviewed the manuscript, and M. H. A. B. revised the manuscript.

Funding

Our work in this area has been financially supported by the Bayer Hemophilia Awards Program (Special Project Award) and Landsteiner Foundation for Blood Transfusion (LSBR grant. no. 1451). The funding agencies had no role in the preparation, review, or approval of the manuscript.

Conflicts of Interest

P. H. R. owns equity in VarmX. The remaining authors declare no competing financial interests.

Acknowledgments

We thank Dr. F.A. Klok for the critical evaluation of this manuscript and M. Zuurmond for generating the images.

References

- Cuker A, Siegal D. Monitoring and reversal of direct oral anticoagulants. *Hematology (Am Soc Hematol Educ Program)* 2015; 2015:117–124
- Barnes GD, Lucas E, Alexander GC, Goldberger ZD. National trends in ambulatory oral anticoagulant use. *Am J Med* 2015; 128(12):1300–5.e2
- Pollack CV Jr, Reilly PA, van Ryn J, et al. Idarucizumab for dabigatran reversal - full cohort analysis. *N Engl J Med* 2017;377(05):431–441
- Loo SY, Dell'Aniello S, Huiart L, Renoux C. Trends in the prescription of novel oral anticoagulants in UK primary care. *Br J Clin Pharmacol* 2017;83(09):2096–2106
- Roehrig S, Straub A, Pohlmann J, et al. Discovery of the novel antithrombotic agent 5-chloro-N-((5S)-2-oxo-3- [4-(3-oxomorpholin-4-yl)phenyl]-1,3-oxazolidin-5-ylmethyl)thiophene-2-carboxamide (BAY 59-7939): an oral, direct factor Xa inhibitor. *J Med Chem* 2005;48(19):5900–5908
- Pinto DJ, Orwat MJ, Koch S, et al. Discovery of 1-(4-methoxyphenyl)-7-oxo-6-(4-(2-oxopiperidin-1-yl)phenyl)-4,5,6,7-tetrahydro-1H-pyrazolo[3,4-c]pyridine-3-carboxamide (apixaban, BMS-562247), a highly potent, selective, efficacious, and orally bioavailable inhibitor of blood coagulation factor Xa. *J Med Chem* 2007;50(22):5339–5356
- Furugohri T, Isobe K, Honda Y, et al. DU-176b, a potent and orally active factor Xa inhibitor: in vitro and in vivo pharmacological profiles. *J Thromb Haemost* 2008;6(09):1542–1549
- Hedstrom L. Serine protease mechanism and specificity. *Chem Rev* 2002;102(12):4501–4524
- Wong PC, Pinto DJ, Zhang D. Preclinical discovery of apixaban, a direct and orally bioavailable factor Xa inhibitor. *J Thromb Thrombolysis* 2011;31(04):478–492
- Padmanabhan K, Padmanabhan KP, Tulinsky A, et al. Structure of human des(1-45) factor Xa at 2.2 Å resolution. *J Mol Biol* 1993; 232(03):947–966
- Zhang P, Huang W, Wang L, et al. Discovery of betrixaban (PRT054021), N-(5-chloropyridin-2-yl)-2-(4-(N,N-dimethylcarbamimidoyl)benzamido)-5-methoxybenzamide, a highly potent, selective, and orally efficacious factor Xa inhibitor. *Bioorg Med Chem Lett* 2009;19(08):2179–2185
- Samama MM. The mechanism of action of rivaroxaban—an oral, direct Factor Xa inhibitor—compared with other anticoagulants. *Thromb Res* 2011;127(06):497–504
- Masotti L, Campanini M. Pharmacology of new oral anticoagulants: mechanism of action, pharmacokinetics, pharmacodynamics. *Ital J Med* 2013;7(08):1–7
- Gómez-Outes A, Suárez-Gea ML, Lecumberri R, Terleira-Fernández AI, Vargas-Castrillón E. Direct-acting oral anticoagulants: pharmacology, indications, management, and future perspectives. *Eur J Haematol* 2015;95(05):389–404
- Connolly SJ, Ezekowitz MD, Yusuf S, et al; RE-LY Steering Committee and Investigators. Dabigatran versus warfarin in patients with atrial fibrillation. *N Engl J Med* 2009;361(12):1139–1151
- Patel MR, Mahaffey KW, Garg J, et al; ROCKET AF Investigators. Rivaroxaban versus warfarin in nonvalvular atrial fibrillation. *N Engl J Med* 2011;365(10):883–891
- Agnelli G, Buller HR, Cohen A, et al; AMPLIFY Investigators. Oral apixaban for the treatment of acute venous thromboembolism. *N Engl J Med* 2013;369(09):799–808
- Deitelzweig SB, Johnson BH, Lin J, Schulman KL. Prevalence of clinical venous thromboembolism in the USA: current trends and future projections. *Am J Hematol* 2011;86(02):217–220
- Chugh SS, Havmoeller R, Narayanan K, et al. Worldwide epidemiology of atrial fibrillation: a Global Burden of Disease 2010 Study. *Circulation* 2014;129(08):837–847
- Connolly SJ, Crowther M, Eikelboom JW, et al; ANNEXA-4 Investigators. Full study report of andexanet alfa for bleeding associated with factor Xa inhibitors. *N Engl J Med* 2019;380(14): 1326–1335
- Lu G, DeGuzman FR, Hollenbach SJ, et al. A specific antidote for reversal of anticoagulation by direct and indirect inhibitors of coagulation factor Xa. *Nat Med* 2013;19(04):446–451

- 22 Hellstern P. Production and composition of prothrombin complex concentrates: correlation between composition and therapeutic efficiency. *Thromb Res* 1999;95(4, Suppl 1):S7–S12
- 23 Franchini M, Lippi G. Prothrombin complex concentrates: an update. *Blood Transfus* 2010;8(03):149–154
- 24 Cada DJ, Leven TL, Baker DE. Prothrombin complex concentrate. *Hosp Pharm* 2013;48(11):951–957
- 25 Sarode R, Milling TJ Jr, Refaai MA, et al. Efficacy and safety of a 4-factor prothrombin complex concentrate in patients on vitamin K antagonists presenting with major bleeding: a randomized, plasma-controlled, phase IIIb study. *Circulation* 2013;128(11):1234–1243
- 26 Godier A, Miclot A, Le Bonniec B, et al. Evaluation of prothrombin complex concentrate and recombinant activated factor VII to reverse rivaroxaban in a rabbit model. *Anesthesiology* 2012;116(01):94–102
- 27 Martin AC, Le Bonniec B, Fischer AM, et al. Evaluation of recombinant activated factor VII, prothrombin complex concentrate, and fibrinogen concentrate to reverse apixaban in a rabbit model of bleeding and thrombosis. *Int J Cardiol* 2013;168(04):4228–4233
- 28 Perzborn E, Gruber A, Tinel H, et al. Reversal of rivaroxaban anticoagulation by haemostatic agents in rats and primates. *Thromb Haemost* 2013;110(01):162–172
- 29 Zhou W, Zorn M, Nawroth P, et al. Hemostatic therapy in experimental intracerebral hemorrhage associated with rivaroxaban. *Stroke* 2013;44(03):771–778
- 30 Herrmann R, Thom J, Wood A, Phillips M, Muhammad S, Baker R. Thrombin generation using the calibrated automated thromboscope to assess reversibility of dabigatran and rivaroxaban. *Thromb Haemost* 2014;111(05):989–995
- 31 Levi M, Moore KT, Castillejos CF, et al. Comparison of three-factor and four-factor prothrombin complex concentrates regarding reversal of the anticoagulant effects of rivaroxaban in healthy volunteers. *J Thromb Haemost* 2014;12(09):1428–1436
- 32 Eerenberg ES, Kamphuisen PW, Sijpkens MK, Meijers JC, Buller HR, Levi M. Reversal of rivaroxaban and dabigatran by prothrombin complex concentrate: a randomized, placebo-controlled, crossover study in healthy subjects. *Circulation* 2011;124(14):1573–1579
- 33 Barco S, Whitney Cheung Y, Coppens M, Hutten BA, Meijers JC, Middeldorp S. In vivo reversal of the anticoagulant effect of rivaroxaban with four-factor prothrombin complex concentrate. *Br J Haematol* 2016;172(02):255–261
- 34 Song Y, Wang Z, Perlstein I, et al. Reversal of apixaban anticoagulation by four-factor prothrombin complex concentrates in healthy subjects: a randomized three-period crossover study. *J Thromb Haemost* 2017;15(11):2125–2137
- 35 Brown KS, Wickremasingha P, Parasrampur DA, et al. The impact of a three-factor prothrombin complex concentrate on the anticoagulatory effects of the factor Xa inhibitor edoxaban. *Thromb Res* 2015;136(04):825–831
- 36 Nagalla S, Thomson L, Oppong Y, Bachman B, Chervoneva I, Kraft WK. Reversibility of apixaban anticoagulation with a four-factor prothrombin complex concentrate in healthy volunteers. *Clin Transl Sci* 2016;9(03):176–180
- 37 Cheung YW, Barco S, Hutten BA, Meijers JC, Middeldorp S, Coppens M. In vivo increase in thrombin generation by four-factor prothrombin complex concentrate in apixaban-treated healthy volunteers. *J Thromb Haemost* 2015;13(10):1799–1805
- 38 Zahir H, Brown KS, Vandell AG, et al. Edoxaban effects on bleeding following punch biopsy and reversal by a 4-factor prothrombin complex concentrate. *Circulation* 2015;131(01):82–90
- 39 Schulman S, Gross PL, Ritchie B, et al; Study Investigators. Prothrombin complex concentrate for major bleeding on factor Xa inhibitors: a prospective cohort study. *Thromb Haemost* 2018;118(05):842–851
- 40 Majeed A, Ågren A, Holmström M, et al. Management of rivaroxaban- or apixaban-associated major bleeding with prothrombin complex concentrates: a cohort study. *Blood* 2017;130(15):1706–1712
- 41 Arachchillage DRJ, Alavian S, Griffin J, et al. Efficacy and safety of prothrombin complex concentrate in patients treated with rivaroxaban or apixaban compared to warfarin presenting with major bleeding. *Br J Haematol* 2019;184(05):808–816
- 42 Gerner ST, Kuramatsu JB, Sembill JA, et al; RETRACE II (German-Wide Multicenter Analysis of Oral Anticoagulation-Associated Intracerebral Hemorrhage II) Investigators. Association of prothrombin complex concentrate administration and hematoma enlargement in non-vitamin K antagonist oral anticoagulant-related intracerebral hemorrhage. *Ann Neurol* 2018;83(01):186–196
- 43 Tomaselli GF, Mahaffey KW, Cuker A, et al. 2017 ACC expert consensus decision pathway on management of bleeding in patients on oral anticoagulants: a report of the American College of Cardiology Task Force on Expert Consensus Decision Pathways. *J Am Coll Cardiol* 2017;70(24):3042–3067
- 44 Sjamsoedin LJ, Heijnen L, Mauser-Bunschoten EP, et al. The effect of activated prothrombin-complex concentrate (FEIBA) on joint and muscle bleeding in patients with hemophilia A and antibodies to factor VIII. A double-blind clinical trial. *N Engl J Med* 1981;305(13):717–721
- 45 Hilgartner MW, Knatterud GL. The use of factor eight inhibitor by-passing activity (FEIBA immuno) product for treatment of bleeding episodes in hemophiliacs with inhibitors. *Blood* 1983;61(01):36–40
- 46 Hilgartner M, Aledort L, Andes A, Gill J; FEIBA Study Group. Efficacy and safety of vapor-heated anti-inhibitor coagulant complex in hemophilia patients. *Transfusion* 1990;30(07):626–630
- 47 Fukuda T, Honda Y, Kamisato C, Morishima Y, Shibano T. Reversal of anticoagulant effects of edoxaban, an oral, direct factor Xa inhibitor, with haemostatic agents. *Thromb Haemost* 2012;107(02):253–259
- 48 Halim AB, Samama MM, Mendell J. Ex vivo reversal of the anticoagulant effects of edoxaban. *Thromb Res* 2014;134(04):909–913
- 49 Schultz NH, Tran HTT, Bjørnsen S, Henriksson CE, Sandset PM, Holme PA. The reversal effect of prothrombin complex concentrate (PCC), activated PCC and recombinant activated factor VII against anticoagulation of Xa inhibitor. *Thromb J* 2017;15:6
- 50 Perzborn E, Heitmeier S, Laux V, Buchmüller A. Reversal of rivaroxaban-induced anticoagulation with prothrombin complex concentrate, activated prothrombin complex concentrate and recombinant activated factor VII in vitro. *Thromb Res* 2014;133(04):671–681
- 51 Marlu R, Hodaj E, Paris A, Albaladejo P, Cracowski JL, Pernod G. Effect of non-specific reversal agents on anticoagulant activity of dabigatran and rivaroxaban: a randomised crossover ex vivo study in healthy volunteers. *Thromb Haemost* 2012;108(02):217–224
- 52 Mao G, King L, Young S, Kaplan R. Factor eight inhibitor bypassing agent (FEIBA) for reversal of target-specific oral anticoagulants in life-threatening intracranial bleeding. *J Emerg Med* 2017;52(05):731–737
- 53 Dibu JR, Weimer JM, Ahrens C, Manno E, Frontera JA. The role of FEIBA in reversing novel oral anticoagulants in intracerebral hemorrhage. *Neurocrit Care* 2016;24(03):413–419
- 54 Dager WE, Roberts AJ, Nishijima DK. Effect of low and moderate dose FEIBA to reverse major bleeding in patients on direct oral anticoagulants. *Thromb Res* 2019;173:71–76
- 55 Engelbart JM, Zepeski A, Galet C, Policeni B, Skeete DA, Faine BA. Safety and effectiveness of Factor Eight Inhibitor Bypassing Activity for direct oral anticoagulant-related hemorrhage reversal. *Am J Emerg Med* 2019;37(02):214–219

- 56 Lentz SR, Ehrenforth S, Karim FA, et al; adept™2 investigators. Recombinant factor VIIa analog in the management of hemophilia with inhibitors: results from a multicenter, randomized, controlled trial of vatreptacog alfa. *J Thromb Haemost* 2014;12(08):1244–1253
- 57 Lusher JM, Roberts HR, Davignon G, et al. A randomized, double-blind comparison of two dosage levels of recombinant factor VIIa in the treatment of joint, muscle and mucocutaneous haemorrhages in persons with haemophilia A and B, with and without inhibitors. *rFVIIa Study Group. Haemophilia* 1998;4(06):790–798
- 58 Goodnough LT, Levy JH. The judicious use of recombinant factor VIIa. *Semin Thromb Hemost* 2016;42(02):125–132
- 59 Schmidt K, Krüger K, Langer E, et al. Reversal of apixaban induced alterations in haemostasis by different coagulation factor concentrates in patients after hip or knee replacement surgery. *Blood Transfus* 2019;17(02):157–162
- 60 Hemphill JC III, Greenberg SM, Anderson CS, et al; American Heart Association Stroke Council; Council on Cardiovascular and Stroke Nursing; Council on Clinical Cardiology. Guidelines for the management of spontaneous intracerebral hemorrhage: a guideline for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke* 2015;46(07):2032–2060
- 61 Turecek PL, Váradi K, Gritsch H, Schwarz HP. FEIBA: mode of action. *Haemophilia* 2004;10(Suppl 2):3–9
- 62 Turecek PL, Varadi K, Gritsch H, et al. Factor Xa and prothrombin: mechanism of action of FEIBA. *Vox Sang* 1999;77(Suppl 1):72–79
- 63 Himmelsbach M, Richter G, Muhr E, et al. A fully recombinant partial prothrombin complex effectively bypasses FVIII in vitro and in vivo. *Thromb Haemost* 2002;88(06):1003–1011
- 64 Schuijt TJ, Bakhtiari K, Daffre S, et al. Factor Xa activation of factor V is of paramount importance in initiating the coagulation system: lessons from a tick salivary protein. *Circulation* 2013;128(03):254–266
- 65 Schreuder M, Reitsma PH, Bos MHA. Blood coagulation factor Va's key interactive residues and regions for prothrombinase assembly and prothrombin binding. *J Thromb Haemost* 2019;17(08):1229–1239
- 66 Johnson DJ, Li W, Adams TE, Huntington JA. Antithrombin-S195A factor Xa-heparin structure reveals the allosteric mechanism of antithrombin activation. *EMBO J* 2006;25(09):2029–2037
- 67 Li W, Johnson DJ, Esmon CT, Huntington JA. Structure of the antithrombin-thrombin-heparin ternary complex reveals the antithrombotic mechanism of heparin. *Nat Struct Mol Biol* 2004;11(09):857–862
- 68 Thalji NK, Ivanciu L, Davidson R, Gimotty PA, Krishnaswamy S, Camire RM. A rapid pro-hemostatic approach to overcome direct oral anticoagulants. *Nat Med* 2016;22(08):924–932
- 69 Monroe DM, Hoffman M, Oliver JA, Roberts HR. Platelet activity of high-dose factor VIIa is independent of tissue factor. *Br J Haematol* 1997;99(03):542–547
- 70 Lu G, Pine P, Leeds JM, et al. Andexanet alfa effectively reverses edoxaban anticoagulation effects and associated bleeding in a rabbit acute hemorrhage model. *PLoS One* 2018;13(03):e0195122
- 71 Lu G, Hollenbach SJ, Baker DC, et al. Preclinical safety and efficacy of andexanet alfa in animal models. *J Thromb Haemost* 2017;15(09):1747–1756
- 72 Siegal DM, Curnutte JT, Connolly SJ, et al. Andexanet alfa for the reversal of factor Xa inhibitor activity. *N Engl J Med* 2015;373(25):2413–2424
- 73 Siegal D, Lu G, Leeds JM, et al. Safety, pharmacokinetics, and reversal of apixaban anticoagulation with andexanet alfa. *Blood Adv* 2017;1(21):1827–1838
- 74 Goldstein JN, Refaai MA, Milling TJ Jr, et al. Four-factor prothrombin complex concentrate versus plasma for rapid vitamin K antagonist reversal in patients needing urgent surgical or invasive interventions: a phase 3b, open-label, non-inferiority, randomised trial. *Lancet* 2015;385(9982):2077–2087
- 75 Ersayin A, Thomas A, Seyve L, et al. Catalytically inactive Gla-domainless factor Xa binds to TFPI and restores *ex vivo* coagulation in hemophilia plasma. *Haematologica* 2017;102(12):e483–e485
- 76 Marlu R, Polack B. Gla-domainless factor Xa: molecular bait to bypass a blocked tenase complex. *Haematologica* 2012;97(08):1165–1172
- 77 Lu G, Lin JP, Curnutte JT, Conley PB. Effect of andexanet-TFPI interaction on in vitro thrombin formation and coagulation markers in the TF-pathway [Abstract]. *Blood* 2017;130(629):629
- 78 Ansell JE, Bakhru SH, Lailicht BE, et al. Single-dose ciraparantag safely and completely reverses anticoagulant effects of edoxaban. *Thromb Haemost* 2017;117(02):238–245
- 79 Weitz JI, Eikelboom JW. Ciraparantag for enoxaparin reversal: adding to the evidence. *Thromb Res* 2016;146:106–107
- 80 Ansell JE, Lailicht BE, Bakhru SH, Hoffman M, Steiner SS, Costin JC. Ciraparantag safely and completely reverses the anticoagulant effects of low molecular weight heparin. *Thromb Res* 2016;146:113–118
- 81 Lailicht BE, Bakhru SH, Jiang X, et al. Antidote for new oral anticoagulants: mechanism of action and binding specificity of PER977 [Abstract]. *J Thromb Haemost* 2013;11:75
- 82 Lailicht BE, Bakhru SH, Lee C, et al. Small molecule antidote for anticoagulants [Abstract]. *Circulation* 2018;126(Suppl 21):A11395
- 83 Bakhru SH, Lailicht BE, Jiang X, et al. Reversal of anticoagulant-induced bleeding in external and internal bleeding models by PER977, a small molecule anticoagulant antidote [Abstract]. *Circulation* 2018;130(Suppl 2):A19361
- 84 Hollenbach SJ, Lu G, DeGuzman F, et al. Andexanet-alfa and PER977 (arapazine) correct blood loss in a rabbit liver laceration model – only andexanet reverses markers of fXa-mediated anticoagulation [Abstract]. *Circulation* 2018;130(Suppl 2):A14657
- 85 Sullivan DW Jr, Gad SC, Lailicht B, Bakhru S, Steiner S. Nonclinical safety assessment of PER977: a small molecule reversal agent for new oral anticoagulants and heparins. *Int J Toxicol* 2015;34(04):308–317
- 86 Ansell JE, Bakhru SH, Lailicht BE, et al. Use of PER977 to reverse the anticoagulant effect of edoxaban. *N Engl J Med* 2014;371(22):2141–2142
- 87 Kalathottukaren MT, Creagh AL, Abbina S, et al. Comparison of reversal activity and mechanism of action of UHRA, andexanet, and PER977 on heparin and oral FXa inhibitors. *Blood Adv* 2018;2(16):2104–2114
- 88 Lu G, Kotha J, Cardenas JM, et al. Abstract 18218: In vitro characterization of andexanet alfa (PRT064445), a specific fXa inhibitor antidote versus aripazine (PER977), a non-specific reversal agent. *Circulation* 2018;130(02):
- 89 Jourdi G, Gouin-Thibault I, Siguret V, Gandrille S, Gaussem P, Le Bonniec B. FXa-α2-macroglobulin complex neutralizes direct oral anticoagulants targeting FXa in vitro and in vivo. *Thromb Haemost* 2018;118(09):1535–1544
- 90 Qazi U, Kolodziej SJ, Gettins PG, Stoops JK. The structure of the C949S mutant human α(2)-macroglobulin demonstrates the critical role of the internal thiol esters in its proteinase-entrapping structural transformation. *J Struct Biol* 2000;131(01):19–26
- 91 Rehman AA, Ahsan H, Khan FH. α-2-Macroglobulin: a physiological guardian. *J Cell Physiol* 2013;228(08):1665–1675
- 92 Jourdi G, Siguret V, Martin AC, et al. Association rate constants rationalise the pharmacodynamics of apixaban and rivaroxaban. *Thromb Haemost* 2015;114(01):78–86
- 93 Toso R, Zhu H, Camire RM. The conformational switch from the factor X zymogen to protease state mediates exosite expression and prothrombinase assembly. *J Biol Chem* 2008;283(27):18627–18635

- 94 Huber R, Bode W. Structural basis of the activation and action of trypsin. *Acc Chem Res* 1978;11(11):114–122
- 95 Bunce MW, Toso R, Camire RM. Zymogen-like factor Xa variants restore thrombin generation and effectively bypass the intrinsic pathway in vitro. *Blood* 2011;117(01):290–298
- 96 Ivanciu L, Camire RM. Hemostatic agents of broad applicability produced by selective tuning of factor Xa zymogenicity. *Blood* 2015;126(01):94–102
- 97 Ivanciu L, Toso R, Margaritis P, et al. A zymogen-like factor Xa variant corrects the coagulation defect in hemophilia. *Nat Biotechnol* 2011;29(11):1028–1033
- 98 Greene LA, Thalji NK, Raffini LJ, Camire RM. Abstract: Zymogen-like FXa variant as a short-acting warfarin reversal agent: pre-clinical evaluation and mechanism of action. *Blood* 2014;124(4262):
- 99 Parng C, Markiewicz V, Chen J, et al. Preclinical pharmacokinetics, pharmacodynamics, tissue distribution, and interspecies scaling of recombinant human coagulation factor Xa^{I16L}. *J Pharm Sci* 2017;106(08):2136–2143
- 100 Parsons-Rich D, Hua F, Li G, Kantaridis C, Pittman DD, Arkin S. Phase 1 dose-escalating study to evaluate the safety, pharmacokinetics, and pharmacodynamics of a recombinant factor Xa variant (FXa^{I16L}). *J Thromb Haemost* 2017;15(05):931–937
- 101 Verhoef D, Visscher KM, Vosmeer CR, et al. Engineered factor Xa variants retain procoagulant activity independent of direct factor Xa inhibitors. *Nat Commun* 2017;8(01):528
- 102 Schreuder M, Verhoef D, Cheung KL, et al. Abstract: Phe174-mutated human factor X as bypassing agent to the direct FXa inhibitors. *Blood* 2017;130(363):