



Universiteit
Leiden
The Netherlands

Rational empiric antibiotic therapy in clinical practice and policy making: uncertainties, probabilities, and ethics

Lambregts, M.M.C.

Citation

Lambregts, M. M. C. (2021, September 28). *Rational empiric antibiotic therapy in clinical practice and policy making: uncertainties, probabilities, and ethics*. Retrieved from <https://hdl.handle.net/1887/3239271>

Version: Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/3239271>

Note: To cite this publication please use the final published version (if applicable).






Part I

Uncertainties and probabilities in
empiric antimicrobial therapy





2

Time to positivity of blood cultures supports early re-evaluation of empiric broad-spectrum antimicrobial therapy

Merel M.C. Lambregts, Alexandra T. Bernards, Martha T. van der Beek, Leo G. Visser, Mark G. J. de boer

ABSTRACT

Background: Blood cultures are considered the gold standard to distinguish bacteraemia from non-bacteraemic systemic inflammation. In current clinical practice, bacteraemia is considered unlikely if blood cultures have been negative for 48-72 hours. Modern BC systems have reduced this time-to-positivity (TTP), questioning whether the time frame of 48-72 hrs is still valid. This study investigates the distribution of TTP, the probability of blood culture positivity after 24 hours, and identifies clinical predictors of prolonged TTP.

Methods: Adult patients with monomicrobial bacteraemia in an academic hospital were included retrospectively over a three-year period. Clinical data were retrieved from the medical records. Predictors of TTP >24 hours were determined by uni- and multivariable analyses. The residual probability of bacteraemia was estimated for the scenario of negative BCs at 24 hours after bedside collection.

Results: The cohort consisted of 801 patients, accounting for 897 episodes of bacteraemia. Mean age was 65 years (IQR 54-73), 534 (59.5%) patients were male. Median TTP was 15.7 (IQR 13.5-19.3) hours. TTP was ≤24 hours in 85.3% of episodes. Antibiotic pre-treatment (adjusted OR 1.77; 95%CI 1.14-2.74, $p < 0.01$) was independently associated with prolonged TTP. The probability of bacteraemia, if BC had remained negative for 24 hours, was 1.8% (95% CI 1.46-2.14).

Conclusion: With adequate hospital logistics, the probability of positive blood cultures after 24 hours of negative cultures was low. Combined with clinical reassessment, knowledge of this low probability may contribute to prioritization of the differential diagnosis and decisions on antimicrobial therapy. As a potential antibiotic stewardship tool, this strategy warrants further prospective investigation.

INTRODUCTION

Empirical in-hospital antibiotic prescription forms a significant proportion of broad-spectrum antibiotic consumption. The in-hospital use of antibiotics is expected to increase even further due to advancing life-expectancy and an increase in the application of immunosuppressive therapies, resulting in an increasing incidence of bacterial infections.¹⁻³ Appropriate empirical treatment for severe bacterial infections improves survival.^{1,4} However, upon presentation, the clinical diagnosis is often uncertain, and the presence of bacterial infection is not always evident. There is a broad differential diagnosis for fever, including viral infections and inflammatory states of non-infectious origin such as pancreatitis. Furthermore, thromboembolic events and severe drug reactions can mimic the symptoms of bacterial infection.⁵

Identifying patients without bacterial infection at an early time point is an important component of antimicrobial stewardship. Prolonged administration of broad-spectrum antibiotics may cause adverse events in the individual patient. Duration of antibiotic therapy is associated with toxicity, *Clostridioides difficile* infection and increased mortality rates.⁶⁻⁹ Furthermore, antibiotic consumption, especially the use of broad-spectrum agents, is one of the major drivers of the increasing antimicrobial resistance worldwide.^{10,11} Because of these individual and societal risks, guidelines recommend to de-escalate broad-spectrum antimicrobial treatment based on the source of infection and culture results. De-escalation of empirical therapy is defined as a reduction in number and/or narrowing of spectrum of antimicrobial agents.¹² When infection is found not to be present, the recommendation is to discontinue antimicrobial therapy.¹

Diagnostics directed at the possible source of infection, i.e. radiographic exams and urine analysis, can be completed within hours. In contrast, differentiating bacteraemia from non-bacteraemic infection is still time consuming as reliable alternatives for conventional blood culture incubation are not yet available in clinical practice. Biomarkers for exclusion of bacteraemia lack sensitivity or have practical limitations.¹³⁻¹⁶ Historically, the consensus is to await blood culture results for at least 48 to 72 hours, before bacteraemia is deemed unlikely.^{17,18} Because of the modernisation of blood culture methods, and especially the development of continuous monitoring systems, the time to positivity (TTP) of blood cultures has been reduced substantially.¹⁹⁻²¹

Knowledge of the distribution of blood culture TTP is of clinical benefit in the re-evaluation of patients with a clinical syndrome consistent with infection. A low probability of bacteraemia when blood cultures have remained negative after 24 hours, may have impact on the differential diagnosis and subsequent diagnostic and therapeutic actions.

Our aim was to determine the distribution of the TTP of blood cultures in adult patients and assess the probability of bacteraemia when blood cultures have remained negative for 24 hours. In addition, we aimed to identify clinical characteristics that predict late (i.e. >24 hours) positivity.

METHODS

Setting and study participants

The retrospective cohort study was performed at the Leiden University Medical Center, a tertiary care and teaching hospital in the Netherlands.

All patients aged 18 years and older, with mono-microbial bacteraemia in 2013 and 2014 were identified. An additional 100 patients that presented in the year 2015 were randomly included, by case identification code. Patients with polymicrobial bacteraemia were excluded as the time to positivity of the individual pathogens was unknown. Furthermore the relevance of the individual pathogens to the TTP of the polymicrobial culture can not be determined.

The blood culture database of the Department of Medical Microbiology was used to identify eligible patients. Patients admitted to clinical wards, including medium and intensive care unit and cases presenting at the emergency department were eligible for inclusion. Multiple episodes of bacteraemia per patient were allowed if the antimicrobial therapy for the previous episode had been completed and clinical and microbiological cure had been achieved. All blood cultures with coagulase-negative staphylococci (CoNs) were excluded, because the likelihood that these cultures represent contamination is high. For other possible contaminants (including anaerobes) the differentiation between true bacteraemia and contamination was based on the number of positive vials and the documented assessment of the microbiologist and responsible physician.

Standard empiric therapy for sepsis of unknown origin in the study centre was a second generation cephalosporin combined with an aminoglycoside.

The study was approved by the Institutional Ethics Review Board of the Leiden University Medical Center. The requirement to obtain informed consent was waived because of the retrospective nature of the study.

Data collection

Demographic data, data about pre-existing medical conditions, clinical parameters at presentation, the most likely source of bacteraemia and the outcome data were retrieved from the electronic medical records. The classification of the source of infection was based on review of the available clinical, radiological and microbiological information. Outcome measurements included admission to the intensive care, length of hospitalisation and 30-day mortality. Microbiological data, including pathogen identification and TTP, were retrieved from the database of the Department of Medical Microbiology.

Antibiotic pre-treatment was defined as treatment with one or more antibiotic agents, administered intravenously, intramuscularly or orally, within the 24 hours preceding collection of the first blood culture. Oral antibiotics without systemic absorption, such as vancomycin, were excluded from this definition.

Blood culture handling procedures and laboratory techniques

TTP was defined as the time between collection of the blood cultures and the positive signal in the BACTEC FX continuous monitoring system (Becton Dickinson B.V., Breda). The institutional protocol is to collect both an aerobic and anaerobic vial, and to collect 8–10 ml of blood per vial.^{22,23} A quality assessment in 100 individual vials showed a median blood volume of 9 ml (IQR 7–11) per vial (Supplementary files, Table S3).

The time of bedside blood culture collection was recorded in the electronic medical records, as part of the ordering procedure. Cultures were transported to the in-hospital microbiology department by dedicated hospital transportation employees. During day-hours, transportation is performed every 3 hours. Quality assessment at the beginning of the study period showed a median time from collection to placement in the incubator of 94 minutes (IQR 63–137). Outside working hours the maximum time to transportation is 5 hours.

Upon arrival at the Department of Medical Microbiology the blood cultures were directly placed in the BACTEC FX continuous monitoring system (Becton Dickinson B.V., Breda), for a minimum of seven days. The time of the positive signal was automatically recorded. During evening and night hours, blood cultures were directly placed in the BACTEC, but registration in the system was performed the following morning between 8 and 9 a.m. If the threshold for positivity was reached between placement and this registration, the culture was recorded positive at the time of registration, instead of upon positive signalling. This technical limitation leads to an overestimation of the TTP in ‘unregistered’ bottles. Therefore, median TTP was additionally calculated excluding ‘unregistered’ episodes (Supplementary files).

If multiple separate blood cultures from one patient were collected within a time frame of two hours, the shortest TTP was used for the statistical analyses.

Blood culture positivity rate

To calculate the probability of positive blood cultures when they have remained negative for 24 hours, information on the institutional blood culture positivity rate is required (see statistical analysis). To estimate the overall blood culture positivity rate, the proportion of bacteraemia was determined during two separate months, June and December 2014. During this period, all patients in whom blood cultures were obtained because of fever or (suspected) sepsis were included. True bacteraemia was defined as growth of a pathogenic bacterial species in ≥ 1 blood culture bottle. Definition of contamination was identical to the definition applied in the main cohort. Patients were only included for the first episode of suspected infection, subsequent episodes were excluded.

Statistical analyses

Median TTP and interquartile ranges (IQR) were determined for the complete cohort and for the most frequently isolated pathogens in patients with bacteraemia. Median TTP was additionally calculated excluding 'unregistered' episodes, because of the potential overestimation of TTP (S1 Table). Normally and non-normally distributed continuous variables were compared by Student's t test and Mann-Whitney U test, respectively. Univariate risk factor analysis was performed for short (<16 hours) and prolonged TTP (>24 hours), using Chi-square statistical tests. Results were reported as risk ratios (RR) with 95% confidence intervals (95%CI). A multivariable analysis for prolonged TTP was performed and results were reported as adjusted odds ratios (OR with 95%CI). Determinants for the multivariable analysis were selected based on <0.25 in the univariate analysis.

We applied a generalized estimating equation model to assess the potential effect of repeated measurements by inclusion of multiple episodes of bacteraemia for a proportion of patients.

The residual risk of detection of bacteraemia after 24 hours was calculated applying a previously published mathematical equation.²⁰ (Supplementary files, Box 1). This equation is based on the proportion of positive blood cultures in suspected sepsis and the proportion of blood cultures with prolonged TTP.

All statistical analyses were performed with the IBM SPSS Statistics, version 23.

RESULTS

Study population characteristics

After exclusion of polymicrobial and contaminated blood cultures, a total of 801 individual adult patients was included, representing 897 episodes of bacteraemia. Mean age was 65 years (IQR 54–73), 534 (59.5%) patients were male.

The majority of bacteraemia episodes (511 episodes, 57.0%) was caused by a Gram-negative pathogen, predominantly *Escherichia coli* (263/511, 51.5%). *Streptococcus spp* were the most common Gram-positive isolates (163/386, 42%). The demographic and clinical characteristics of the 897 episodes of bacteraemia are summarized in Table 1. In 450/897 (50.2%) episodes ≥ 2 blood culture sets were obtained within a time frame of two hours.

Table 1. Demographic and clinical characteristics among 897 episodes with bacteraemia.

Characteristic	n = 897 (100%)
<i>Patient demographics</i>	
Male gender	534 (59.5)
Age (years) , (median, IQR)	65 (54-73)
<i>Medical history</i>	
Diabetes mellitus	188 (21.0)
Corticosteroid therapy (prior 6 months)	276 (30.8)
Neutropenia	113 (12.6)
Solid organ transplantation	116 (12.9)
Solid malignancy	170 (19.0)
haematological malignancy	96 (10.7)
Dialysis (haemodialysis/peritoneal dialysis)	20 (2.2)
<i>Clinical presentation</i>	
Fever (temperature>38.5 °C)	538 (60.0)
Systolic blood pressure (mmHg) (median, IQR)	125 (107-142)
Pulse rate (bpm) (median, IQR)	101 (88-115)
EMV <15	173 (19.3)
PITT Bacteraemia score (median, IQR)	1 (0-2)
Quick SOFA-score (median, IQR)	1 (1-2)
<i>Antibiotic pre-treatment</i>	264 (29.4)
<i>Location of presentation</i>	
Emergency department	507 (56.5)
General ward	340 (37.9)
ICU/MCU	50 (5.6)
Hospitalization before BC (hours) (median IQR)	3.0 (0.4-136.8)

Table 1. Demographic and clinical characteristics among 897 episodes with bacteraemia. (continued)

Characteristic	n = 897 (100%)
<i>Microbiological parameters</i>	
Gram-positive bacteraemia:	386 (43.0)
Gram-negative bacteraemia	511 (57.0)
Anaerobic bacteraemia	37 (4.1)
<i>Source of infection</i>	
Gastro-intestinal	245 (27.3)
Respiratory	89 (9.9)
Endovascular (e.g. thrombus)	111 (12.4)
Urinary tract	232 (25.9)
Skin and soft tissue	71 (7.9)
Other	56 (6.2)
Not identified	84 (9.4)
<i>Outcome</i>	
ICU/MCU admission during hospitalization	180 (20.1)
Hospitalization after BC (days) (median IQR)	8.9 (3.9-19.0)
30-day mortality	134 (14.9)

Legend: BC=blood culture, ICU/MCU = intensive care unit / medium care unit, IQR= interquartile range


Time to positivity

The median TTP was 15.7 hours (IQR 13.5-19.3). The TTP was below 24 hours in 765 episodes (85.3%). In 34 (3.8%) episodes and 18 (2.0%) episodes TTP was longer than 48 hours and 72 hours, respectively (Fig 1).

Anaerobic bacteraemia was frequent in the prolonged TTP group, 28/132 (21.2%) episodes. After exclusion of anaerobic bacteraemia, there was no statistically significant difference in TTP between Gram-negative and Gram-positive bacteraemia (TTP 18.6 h vs 19.4 h, $p = 0.48$). The TTP of the most common pathogens is illustrated in Fig 2. All episodes of *Streptococcus pneumoniae* bacteraemia were diagnosed within 24 hours (median 13.4 h, IQR 11.3-15.5 h). TTP was long in bacteraemia caused by *Proteus mirabilis* (median 18.6 hr, IQR 14.8-34.9 h). All cases ($n=3$) of *Propionibacterium acnes* bacteraemia were diagnosed after 72 hours.


In 87 of the 132 (65.9%) episodes with prolonged TTP, the isolated pathogen was susceptible to the institutions empirical sepsis therapy (2nd generation cephalosporin and an aminoglycoside).

Fig 1. Distribution of time to blood culture positivity (TTP) in 897 episodes of bacteraemia.



Legend: Fig. 1A illustrates the distribution of TTP in patients with and without antibiotic pre-treatment at the time blood cultures were collected. Fig 1B. illustrates the distribution of TTP, short (≤ 24) versus prolonged (>24) TTP, according to isolated pathogen. The group ‘Other’ comprises *Citrobacter spp*, *Haemophilus spp*, *Listeria spp*, *Achromobacter spp*, *Acinetobacter spp*, *Moraxella catarrhalis*, *Morganella morganii*, *Propionibacterium acnes*, *Rothia mucilaginosa*, *Salmonella spp*, *Stenotrophomonas maltophilia*, *Lactobacillus spp*, *Prevotella spp*, *Fusobacterium spp*.

Fig 2. Pathogens and time to positivity (TTP) distributions.



Legend: The boxplot figure illustrates the distribution of TTP (median, interquartile range) for the most frequently isolated pathogens. The ends of the whiskers represent one and a half times the interquartile range. The group 'Other' comprises *Citrobacter spp.*, *Haemophilus spp.*, *Listeria spp.*, *Achromobacter spp.*, *Acinetobacter spp.*, *Moraxella catarrhalis*, *Morganella morganii*, *Propionibacterium acnes*, *Rothia mucilaginosa*, *Salmonella spp.*, *Stenotrophomonas maltophilia*, *Lactobacillus spp.*, *Vibrio spp.*

In 108 (12.0%) episodes blood cultures were placed in the incubator 'unregistered' and reached the threshold for positivity before registration. TTP analysis excluding these episodes did not have an important effect on the results (Supplementary files, S1/2).

Predictors of short versus prolonged time to positivity

Neutropenia (RR 0.22, 95%CI 0.08-0.58, $p < 0.01$) and corticosteroid therapy (RR 0.66, 95%CI 0.45-0.97, $p = 0.03$) were associated with short TTP (≤ 24 hours) in univariate analysis (Table 2). The source of infection was not a predictor of short versus prolonged TTP. In multivariable analysis, antibiotic pre-treatment (adjusted OR 1.71 95%CI 1.11-2.65, $p < 0.01$) was associated with prolonged TTP (> 24 hours). Neutropenia (adjusted OR 0.15 95%CI 0.05-0.43, $p < 0.01$), was associated with short TTP. Application of a generalized estimating equations model did not detect a relevant effect of including > 1 episode in a proportion of patients.

Table 2. Univariate and multivariable analysis for long time-to-positivity (>24 hours) in 897 episodes of bacteraemia.

Characteristic	Univariate analysis			Multivariable analysis		
	RR	95% CI	p-value	OR	95% CI	p-value
<i>Patient demographics</i>						
Male gender	0.98	0.71 - 1.35	0.91			
Age > 70 years	1.22	0.88 - 1.69	0.23	1.15	0.76-1.72	0.52
<i>Medical history</i>						
Immunocompetent	1.00	-	-			
Neutropenia	0.21	0.08 - 0.56	<0.01	0.15	0.05-0.43	<0.01
Corticosteroid therapy	0.64	0.44 - 0.94	0.02	0.71	0.45-1.14	0.16
<i>Clinical presentation</i>						
Temperature>38.5 °C	0.80	0.58 - 1.11	0.18	0.79	0.54-1.17	0.24
Systolic blood pressure <100 mmHg	1.29	0.86 - 1.94	0.23	1.09	0.65-1.80	0.73
PITT bacteraemia score≥2	0.85	0.58 - 1.25	0.41			
Quick SOFA score >1	1.10	0.69 - 1.75	0.78			
Antibiotic pre-treatment	1.23	0.92 - 1.77	0.15	1.71	1.11-2.65	0.01
Emergency department	0.77	0.56 - 1.05	0.10	0.71	0.47-1.07	0.10
<i>Source of infection</i>						
Gastro-intestinal	1.29	0.92 - 1.79	0.14	1.45	0.95-2.22	0.08
Respiratory tract	0.66	0.35 - 1.26	0.17	0.79	0.38-1.66	0.53
Endovascular	0.78	0.45 - 1.32	0.34			
Urinary tract	0.99	0.69 - 1.43	0.97			
Skin and soft tissue	0.95	0.52 -1.73	0.88			

Legend: CI= confidence interval. RR= relative risk. OR = adjusted odds ratio. CI = confidence interval. The PITT bacteraemia score is calculated from temperature of 35.1–36.0°C or 39.0–39.9°C (1 point), temperature of ≤35°C or ≥40°C (2 points), mental status (alert, 0 points; disoriented, 1 point; stuporous, 2 points; comatose, 4 points), hypotension (2 points), receipt of mechanical ventilation (2 points) and cardiac arrest (4 points). The Quick SOFA score is calculated from glasgow coma scale <15 (1 point) , Respiratory rate ≥22 (1 point), systolic blood pressure ≤100 (1 point).

Probability of bacteraemia at T=24 hours

The two determinants in the equation of the probability of bacteraemia at T = 24 hours are the blood culture positivity rate and the proportion of blood cultures that is positive within 24 hours (S1 Box). The rate of blood culture positivity was determined during two separate months, June and December 2014. In this period 2,099 blood cultures in 778 patients were obtained because of suspected bacterial infection. In 83/778 episodes one or multiple blood cultures were positive, resulting in a positivity rate of 10.7%.

The probability of bacteraemia after 24 hours was calculated using the above estimated overall a priori risk of bacteraemia in patients with suspected infection (10.7%), and the fraction of blood cultures that were positive within 24 hours (85.3%).²⁰ The probability of

bacteraemia when blood cultures had remained negative after 24 hours was 1.8% (95% CI 1.46–2.14%).

DISCUSSION

We found that under the condition of adequate hospital logistics and by using modern, continuously monitoring blood culture systems, 85.3% of blood cultures is positive within 24 hours. Neutropenia was a predictor of short TTP in our study and antibiotic pre-treatment was a predictor of prolonged TTP. These predictors are in line with results from a study by Martinez *et al.*²¹ Most previous studies have defined TTP as the time between incubation and positivity. To permit clinical applicability of the results, we here defined TTP as time between collection of the blood samples and blood culture positivity, taking into account the transportation and laboratory logistics during and outside office hours. As a result, median TTP in our study is longer than in most previous studies^{21,24}, but applicable to real-life clinical settings.

For daily practice, the proportion of blood cultures that becomes positive after different periods of elapsed time is more relevant than median TTP. Two previous studies, that included smaller numbers of patients, found similar results on TTP distribution, despite the above mentioned differences in definition.^{18,25} The authors of these studies conclude that their findings support antibiotic de-escalation after 48 hours. However, to decide on the optimal timing of re-evaluation of the differential diagnosis, the more relevant question is how probable bacteraemia still is when blood cultures have remained negative at different time points. For that purpose, knowledge about the overall blood culture positivity rate, i.e. the pre-test probability, is essential. The blood culture positivity rate in our centre is 10.7%. This is in line with literature on the prevalence of bacteraemia in localised bacterial infection and systemic inflammatory response syndrome (SIRS).^{26–28} By using the previously published mathematical equation (S1 Box), the probability of blood culture positivity after 24 hours is below 2 percent in our institution.²⁰

This probability is centre specific, as both variables in the mathematical equation may vary between institutions. The first variable, the overall blood culture positivity rate, is dependent on the patient population and the criteria that are applied by doctors to order blood cultures. For example, a ‘culture of culturing’ will result in low blood culture positivity rates.

The second variable, the proportion of blood cultures that is positive within 24 hours, is dependent on hospital logistics. If there is an important delay in transportation of the

cultures to the laboratory or placement in the incubator, TTP according to our definition, will be longer. The mathematical equation, allows for the calculation of an institution specific probability of blood culture positivity at $T = 24$ hours.

With adequate hospital logistics, the overall probability of positive blood cultures at $T = 24$ hours is low. This knowledge is valuable for the differential diagnosis and management of patients with suspected bacterial infection. For example, in the scenario of a confirmed source of infection (e.g. pneumonia), and clinical recovery, preliminary negative blood culture results may support an early intravenous-oral switch.²⁹ Alternatively, when there are no signs of localised infection and blood cultures are still negative after 24 hours, bacteraemia becomes unlikely. This knowledge should prompt timely diagnostic steps into non-bacterial causes of fever that require interventions, as e.g. Influenza, thrombo-embolic events or drug-reactions. Despite low probabilities a blood culture may incidentally become positive after more than 24 hours. Furthermore, negative blood cultures do not exclude bacteraemia. Nor does the absence of positive blood cultures exclude non-bacteraemic infections. Despite this level of uncertainty, re-evaluation of empirical therapy is in place when the probability of bacterial bloodstream infection changes. Re-evaluation of clinical stability, response to empirical therapy and an update of the differential diagnosis, is essential when balancing the potential costs and benefits of de-escalating empiric therapy.

For the application of the findings to clinical practice, it is also important to emphasize that the pre-test probability of bacteraemia is variable, not only between institutions, but between patients as well.²⁸ For example, in the severely ill patient with septic shock, the blood culture positivity rate is higher, and TTP may be shorter, both affecting the bacteraemia probability after 24 hours.³⁰ In the severely ill patient without an alternative diagnosis, even a low probability of bacteraemia or non-bacteraemic bacterial infection may warrant continuation or even escalation of broad-spectrum antibiotic therapy.

To the best of our knowledge, we present the largest cohort of patients investigating the distribution of TTP. More importantly, this is the first study to approach TTP of blood cultures from a clinical perspective, providing insight into the probability of bacteraemia at the 24 hour time point. A limitation of the present study is that patients with polymicrobial bacteraemia were excluded. Based on previous research^{31, 32} and on theoretical grounds, the TTP of polymicrobial episodes is comparable to monomicrobial bacteraemia, possibly even shorter. Therefore, inclusion of these episodes would at most reduce the probability of blood culture positivity at $T > 24$ hours.

Secondly, the volume of blood collected in the vials was not recorded in the individual cases, and in a proportion of patients only 1 vial-set was collected. In daily practice TTP and the yield of blood cultures could be improved by further optimising specimen collection; specifically vial filling and number of vials.^{22,23,33}

Thirdly, we were not informed about the individual vial transportation times to the microbiology laboratory. However, our institutional transport logistics and transport times are in line with current guidelines and comparable to other institutions.^{34,35} Previous research has shown that transport and incubation of blood cultures outside laboratory reduces turnaround time and accelerates therapeutic interventions.³⁶ As blood culture collection and transportation procedures impact TTP, audit of blood culture logistics is probably a prerequisite for translation of our results to other institutions.

In conclusion, if modern blood culture systems are used in combination with adequate logistics, the probability of positivity when blood cultures are negative after 24 hours is very low. Postponing re-evaluation of the differential diagnosis, solely for the reason of pending blood culture results, is not rational at this time point. The search for alternative causes of fever can be initiated more rapidly if the probability of bacteraemia is incorporated in clinical reasoning. This may lead to better timed de-escalation, iv to oral switch and earlier hospital discharge. The safety as well as the benefits of this antibiotic stewardship opportunity should be subject of future clinical trials.

REFERENCES

1. Rhodes A, Evans LE, Alhazzani W, Levy MM, Antonelli M, Ferrer R, et al. Surviving Sepsis Campaign: International Guidelines for Management of Sepsis and Septic Shock: 2016. *Crit Care Med*. 2017;45(3):486-552.
2. Wiersinga WJ, van Zanten ARH, Kaasjager KA, van der Poll T. Sepsis: new insights, new definition. *Ned Tijdschr Geneeskd*. 2017;161(0):D1551.
3. Esper AM, Martin GS. The impact of comorbid conditions on critical illness. *Crit Care Med*. 2011;39(12):2728-35.
4. Paul M, Shani V, Muchtar E, Kariv G, Robenshtok E, Leibovici L. Systematic review and meta-analysis of the efficacy of appropriate empiric antibiotic therapy for sepsis. *Antimicrob Agents Chemother*. 2010;54(11):4851-63.
5. Rangel-Frausto MS, Pittet D, Costigan M, Hwang T, Davis CS, Wenzel RP. The natural history of the systemic inflammatory response syndrome (SIRS). A prospective study. *JAMA*. 1995;273(2):117-23.
6. Srigley JA, Brooks A, Sung M, Yamamura D, Haider S, Mertz D. Inappropriate use of antibiotics and *Clostridium difficile* infection. *Am J Infect Control*. 2013;41(11):1116-8.
7. Kibe S, Adams K, Barlow G. Diagnostic and prognostic biomarkers of sepsis in critical care. *J Antimicrob Chemother*. 2011;66 Suppl 2:ii33-40.
8. Zilahi G, McMahon MA, Povia P, Martin-Loeches I. Duration of antibiotic therapy in the intensive care unit. *J Thorac Dis*. 2016;8(12):3774-80.
9. Barlam TF, Soria-Saucedo R, Cabral HJ, Kazis LE. Unnecessary Antibiotics for Acute Respiratory Tract Infections: Association With Care Setting and Patient Demographics. *Open Forum Infect Dis*. 2016;3(1):ofw045.
10. Paterson DL, Bonomo RA. Extended-spectrum beta-lactamases: a clinical update. *Clin Microbiol Rev*. 2005;18(4):657-86.
11. Safdar N, Maki DG. The commonality of risk factors for nosocomial colonization and infection with antimicrobial-resistant *Staphylococcus aureus*, *enterococcus*, gram-negative bacilli, *Clostridium difficile*, and *Candida*. *Ann Intern Med*. 2002;136(11):834-44.
12. Garnacho-Montero J, Gutierrez-Pizarra A, Escolera-Ortega A, Corcia-Palomo Y, Fernandez-Delgado E, Herrera-Melero I, et al. De-escalation of empirical therapy is associated with lower mortality in patients with severe sepsis and septic shock. *Intensive Care Med*. 2014;40(1):32-40.
13. Jones AE, Fiechtel JF, Brown MD, Ballew JJ, Kline JA. Procalcitonin test in the diagnosis of bacteraemia: a meta-analysis. *Ann Emerg Med*. 2007;50(1):34-41.
14. Hoenigl M, Raggam RB, Wagner J, Valentin T, Leitner E, Seeber K, et al. Diagnostic accuracy of soluble urokinase plasminogen activator receptor (suPAR) for prediction of bacteraemia in patients with systemic inflammatory response j. *Clin Biochem*. 2013;46(3):225-9.
15. Wildi K, Tschudin-Sutter S, Dell-Kuster S, Frei R, Bucher HC, Nuesch R. Factors associated with positive blood cultures in outpatients with suspected bacteraemia. *Eur J Clin Microbiol Infect Dis*. 2011;30(12):1615-9.
16. Morrison J, Zweitzig D, Riccardello N, Rubino J, Axelband J, Sodowich B, et al. Retrospective analysis of clinical data associated with patients enrolled in a molecular diagnostic feasibility study highlights the potential utility for rapid detection of bloodstream infection. *Am J Emerg Med*. 2014;32(6):511-6.

17. Centers for Disease Control and Prevention Core elements of hospital antibiotic stewardship programs. <https://www.cdc.gov/antibiotic-use/healthcare/implementation/core-elements.html>. accessed November 2018
18. Pardo J, Klinker KP, Borgert SJ, Trikha G, Rand KH, Ramphal R. Time to positivity of blood cultures supports antibiotic de-escalation at 48 hours. *Ann Pharmacother*. 2014;48(1):33-40.
19. Reimer LG, Wilson ML, Weinstein MP. Update on detection of bacteraemia and fungemia. *Clin Microbiol Rev*. 1997;10(3):444-65.
20. Lambregts MMC, Warreman EB, Bernards AT, Veelken H, von dem Borne PA, Dekkers OM, et al. Distribution and clinical determinants of time-to-positivity of blood cultures in patients with neutropenia. *Eur J Haematol*. 2017.
21. Martinez JA, Pozo L, Almela M, Marco F, Soriano A, Lopez F, et al. Microbial and clinical determinants of time-to-positivity in patients with bacteraemia. *Clin Microbiol Infect*. 2007;13(7):709-16.
22. Lin HH, Liu YF, Tien N, Ho CM, Hsu LN, Lu JJ. Evaluation of the blood volume effect on the diagnosis of bacteraemia in automated blood culture systems. *J Microbiol Immunol Infect*. 2013;46(1):48-52.
23. Kim SC, Kim S, Lee DH, Choi SR, Kim JS. Effect of blood volume in standard anaerobic blood culture bottles of the BacT/ALERT 3D system used for the detection of pathogens and time to detection. *PLoS One*. 2015;10(2):e0116728.
24. Zhang Q, Li D, Bai C, Zhang W, Zheng S, Zhang P, et al. Clinical prognostic factors for time to positivity in cancer patients with bloodstream infections. *Infection*. 2016;44(5):583-8.
25. Ning Y, Hu R, Yao G, Bo S. Time to positivity of blood culture and its prognostic value in bloodstream infection. *Eur J Clin Microbiol Infect Dis*. 2016;35(4):619-24.
26. Chirouze C, Schuhmacher H, Rabaud C, Gil H, Khayat N, Estavoyer JM, et al. Low serum procalcitonin level accurately predicts the absence of bacteraemia in adult patients with acute fever. *Clin Infect Dis*. 2002;35(2):156-61.
27. Metersky ML, Ma A, Bratzler DW, Houck PM. Predicting bacteraemia in patients with community-acquired pneumonia. *Am J Respir Crit Care Med*. 2004;169(3):342-7.
28. Coburn B, Morris AM, Tomlinson G, Detsky AS. Does this adult patient with suspected bacteraemia require blood cultures? *JAMA*. 2012;308(5):502-11.
29. Halm EA, Switzer GE, Mittman BS, Walsh MB, Chang CC, Fine MJ. What factors influence physicians' decisions to switch from intravenous to oral antibiotics for community-acquired pneumonia? *J Gen Intern Med*. 2001;16(9):599-605.
30. Bates DW, Sands K, Miller E, Lanken PN, Hibberd PL, Graman PS, et al. Predicting bacteraemia in patients with sepsis syndrome. Academic Medical Center Consortium Sepsis Project Working Group. *J Infect Dis*. 1997;176(6):1538-51.
31. Shah SS, Downes KJ, Elliott MR, Bell LM, McGowan KL, Metlay JP. How long does it take to "rule out" bacteraemia in children with central venous catheters? *Pediatrics*. 2008;121(1):135-41.
32. McGowan KL, Foster JA, Coffin SE. Outpatient pediatric blood cultures: time to positivity. *Pediatrics*. 2000;106(2 Pt 1):251-5.
33. Tarai B, Jain D, Das P, Budhiraja S. Paired blood cultures increase the sensitivity for detecting pathogens in both inpatients and outpatients. *Eur J Clin Microbiol Infect Dis*. 2018;37(3):435-41.
34. Baron EJ, Miller JM, Weinstein MP, Richter SS, Gilligan PH, Thomson RB, Jr., et al. A guide to utilization of the microbiology laboratory for diagnosis of infectious diseases: 2013 recommendations by the Infectious Diseases Society of America (IDSA) and the American Society for Microbiology (ASM)(a). *Clin Infect Dis*. 2013;57(4):e22-e121.

35. Bignardi GE, Berrington A, Koerner R, Settle C. Immediate incubation of blood cultures is critical to bring about better bacteraemia outcomes. *J Infect.* 2015;70(5):546.
36. Kerremans JJ, van der Bij AK, Goessens W, Verbrugh HA, Vos MC. Immediate incubation of blood cultures outside routine laboratory hours of operation accelerates antibiotic switching. *J Clin Microbiol.* 2009;47(11):3520-3.

SUPPLEMENTARY DATA

Pathogen distribution and median time to positivity excluding and including ‘unregistered’ episodes

In 108 (12.0%) episodes blood cultures were placed in the incubator ‘unregistered’ and reached the threshold for positivity before registration. TTP analysis excluding these episodes did not have an important effect on the results compared to the analysis that included all episodes.

Table S1. Pathogen distribution and median time to positivity excluding unregistered episodes.

Pathogen	Number of episodes (%)	Median TTP (hours) (IQR)
Overall	790 (100.0)	16.1 (13.3-20.0)
Gram-positive pathogen ^a :	346 (43.8)	16.3 (13.6-19.8)
<i>Streptococcus spp</i>	152 (19.2)	14.5 (12.4-18.5)
<i>Strept. pneumoniae</i>	52 (6.6)	13.4 (11.4-15.2)
<i>Enterococcus spp.</i>	88 (11.1)	16.5 (14.5-18.5)
<i>Staphylococcus aureus</i>	91 (11.5)	17.7 (14.4-21.7)
<i>Other</i>	15 (1.9)	26.4 (21.3-94.8)
Gram-negative pathogen ^a :	417 (52.8)	15.5 (13.1-19.3)
<i>Escherichia coli</i>	218 (27.6)	14.1 (12.5-17.5)
<i>Pseudomonas aeruginosa</i>	38 (4.8)	19.2 (17.6-22.4)
<i>Enterobacter spp.</i>	26 (3.3)	15.0 (12.8-20.5)
<i>Klebsiella spp.</i>	66 (8.4)	15.5 (13.2-20.3)
<i>Serratia marcesens</i>	25 (3.2)	15.7 (13.8-16.9)
<i>Proteus mirabilis</i>	15 (1.9)	15.8 (14.7-35.9)
<i>Other</i>	29 (3.7)	18.2 (15.9-24.2)
Anaerobic pathogen	27 (3.4)	32.6 (24.1-46.6)

Legend: TTP= Time to positivity, IQR= inter quartile range. ^a: aerobes only.

Table S2. Pathogen distribution and median time to positivity including unregistered episodes.

Pathogen	Number of episodes (%)	Median TTP (hours) (IQR)	Maximum TTP (hours)
Overall	897 (100)	15.3 (13.3-19.5)	303.6
Gram-positive pathogen ^a :	374 (41.7)	15.7 (13.5-19.3)	303.6
<i>Streptococcus spp</i>	163 (18.2)	14.5 (12.5-18.3)	55.3
<i>Strept. pneumoniae</i>	63 (7.0)	13.4 (11.3-15.5)	23.2
<i>Enterococcus spp.</i>	99 (11.0)	16.0 (14.3-18.2)	41.3
<i>Staphylococcus aureus</i>	96 (10.1)	17.4 (14.4-21.4)	74.8
Other	16 (1.8)	26.2 (21.1-92.2)	303.6
Gram-negative pathogen ^a :	486 (54.2)	15.0 (13.0-18.5)	162.3
<i>Escherichia coli</i>	263 (29.3)	14.1 (12.6-16.7)	162.3
<i>Pseudomonas aeruginosa</i>	40 (4.6)	18.6 (17.5-22.3)	50.5
<i>Enterobacter spp.</i>	28 (32.2)	14.7 (13.1-20.2)	34.7
<i>Klebsiella spp.</i>	77 (8.6)	15.0 (13.1-19.6)	116.1
<i>Serratia marcesens</i>	30 (3.3)	15.3 (13.8-16.5)	41.3
<i>Proteus mirabilis</i>	17 (1.9)	18.6 (14.8-34.9)	43.6
Other	31 (3.5)	18.1 (15.5-24.1)	48.6
Anaerobic pathogen	37 (4.1)	32.6 (24.1-46.6)	129.4

Legend: TTP= Time to positivity, IQR= inter quartile range. ^a: Aerobes only.

Results of the quality assessment of blood volume in blood culture vials

Vial volume was measured in 50 anaerobic vials and 50 aerobic vials. Vial volume was below 8 ml in 32/100 (32.0%) of vials and below 7 ml in 15/100 vials (15.0%).

Table S3. Results of the quality assessment of blood volume in blood culture vials.

Blood volume/vial in milliliters	Frequency (%)	Cumulative Percentage
4	3 (3.0)	3.0
5	7 (7.0)	10.0
6	5 (5.0)	15.0
7	17 (17.0)	32.0
8	15 (15.0)	47.0
9	13 (13.0)	60.0
10	5 (5.0)	65.0
11	13 (13.0)	78.0
12	5 (5.0)	83.0
13	4 (4.0)	87.0
14	4 (4.0)	91.0
15	4 (4.0)	95.0
16	3 (3.0)	98.0
17	1 (1.0)	99.0
18	1 (1.0)	100.0

Legend: Volume of blood in 100 BACTEC blood culture vials (Becton Dickinson B.V., Breda).

Residual risk of detection of bacteraemia after 24 hours

Box S1. Formula for the Estimation of probability of bacteraemia after 24 hours.

$$P = \frac{(1 - TTP_e) \cdot X}{1 - (TTP_e \cdot X)} \cdot 100\%$$

P = Probability of a positive blood culture when the sets have remained negative 24 hours after bedside collection.

X = The proportion of patients with positive blood cultures among all patients in whom blood cultures are obtained for suspected bacterial infection (centre-specific).

TTP_e = proportion of positive blood cultures that are positive in ≤24 hours.

Example

In the Leiden University Medical Center an estimated 10.6% of patients in whom blood cultures were obtained have true bacteremia and 85.3% (95%CI 83.0-87.6) of blood cultures is positive within 24 hours.

Then (use formula, X = 0.11 TTP_e = 0.85) If BCs are still negative after 24 hours of incubation, the probability that the culture will become positive is approximately 1.8% (95%CI 1.5-2.1).

This formula is reproduced from a publication in the European Journal of Haematology.

