

Plasmonic enhancement of single-molecule fluorescence under one- and two-photon excitation

Lu, X.

Citation

Lu, X. (2021, December 8). *Plasmonic enhancement of single-molecule fluorescence under one- and two-photon excitation. Casimir PhD Series.* Retrieved from https://hdl.handle.net/1887/3245677

Version:	Publisher's Version
License:	<u>Licence agreement concerning inclusion of doctoral</u> <u>thesis in the Institutional Repository of the University</u> <u>of Leiden</u>
Downloaded from:	https://hdl.handle.net/1887/3245677

Note: To cite this publication please use the final published version (if applicable).

Stellingen

behorende bij het proefschrift

Plasmonic Enhancement of Single-Molecule Fluorescence under Oneand Two-Photon Excitation

- 1. Compared to strong emitters, weak emitters with low quantum yields experience a stronger fluorescence enhancement by plasmonic nanoantennas, yet it doesn't mean they can be brighter. *Chapter 2, 3 of this thesis.*
- A trade-off between the excitation rate and the fluorescence enhancement is normally needed to optimize the enhanced brightness for the ultra-weak emitters with large Stokes shifts.

Chapter 3 of this thesis.

3. A simple strategy of sandwiching gold nanorod assemblies with two slides can "freeze" the assembly process at the dimer stage, enriching the fraction of dimers on the glass surfaces.

Chapter 4, 5 of this thesis.

- End-to-end gold nanorod dimers with open gaps of ~ 5 nm make it possible to detect the two-photon-excited single-molecule fluorescence, through an enhancement of up to seven orders of magnitude. *Chapter 2, 5 of this thesis.*
- Nanoparticle-on-mirror systems offer sub-nm cavities with enhancement factors up to 10⁸ for two-photon-excitation, yet the hotspots are inaccssible to fresh molecules. O. S. Ojambati et al., Nano Lett. 20, 4653 (2020)
- Plasmonic nanoantennas can not only change the intensity but also the shape of the emission of fluorescent molecules. *M. Ringler et al.*, *PRL 100*, 203002 (2008).
- 7. A hierarchical assembly approach consisting of template-particle and particle-particle interactions ensures the fabrication of heterodimer with precise addressability. *J. Li et al., Nano Lett. 19, 4314 (2019).*
- Quantum effects should be considered seriously when the plasmonic particles are placed apart at the atomic length scale.
 K. J. Savage et al., Nature 491, 574-577 (2012).
- 9. "Perfect" is sometimes the enemy of "good enough" in optical experiments.

Xuxing Lu Leiden, December 08, 2021