



Universiteit
Leiden
The Netherlands

Ionization by galaxy cluster photons alters the ionization state of the nearby warm-hot intergalactic medium

Štofánová, L.; Simionescu, A.; Wijers, N.A.; Schaye, J.; Kaastra, J.S.

Citation

Štofánová, L., Simionescu, A., Wijers, N. A., Schaye, J., & Kaastra, J. S. (2021). *Ionization by galaxy cluster photons alters the ionization state of the nearby warm-hot intergalactic medium*. *Galaxy Cluster Formation II* (p. 73). doi:10.5281/zenodo.5497702

Version: Publisher's Version

License: [Creative Commons CC BY 4.0 license](https://creativecommons.org/licenses/by/4.0/)

Downloaded from: <https://hdl.handle.net/1887/3250318>

Note: To cite this publication please use the final published version (if applicable).

Ionization by galaxy cluster photons alters the ionization state of the nearby warm-hot intergalactic medium

SRON

Netherlands Institute for Space Research






Universiteit
Leiden

ABSTRACT

The physical properties of the faint and extremely tenuous plasma in the far outskirts of **galaxy clusters**, the **circumgalactic media** of normal galaxies, and **filaments of the cosmic web**, remain one of the biggest unknowns in our story of large-scale structure evolution. Modeling the spectral features due to emission and absorption from this very diffuse plasma poses unique challenges, as both collisional and photo-ionisation processes must be accounted for. In this paper we study the **ionisation by galaxy cluster photons** in addition to the photo-ionisation by the cosmic UV/X-ray background and its impact on the **ionisation balance**. We model realistic spectra by taking into account the cosmic UV/X-ray background together with the emission from three different cool-core galaxy clusters: A262, A1795 and A2029, and illuminate the photo-ionised gas in the galaxy cluster vicinity. We assume the gas has temperatures between 10^{-3} –1 keV ($\approx 10^4$ – 10^7 K), densities between 10^{-6} – 10^{-1} cm^{-3} and can be located between r_{500} and ~ 13 Mpc from the cluster center. We find that depending on the distance from the galaxy cluster and the plasma properties, **the total photo-ionisation rate can be higher than 10% or even 100% of the total ionization rate**. We show how this affects the **ionisation fractions of O VI, O VII, O VIII, Ne IX, Fe XVII, N VII, C V and C VI ions** and compare it with the ionization fractions with photo-ionization from cosmic background only as well as with plasma that is in collisional ionisation equilibrium. We assume a simplified model of a cosmic web filament and predict **the column densities** for different lines of sight.

Lýdia Štofánová*, Aurora Simionescu, Nastasha A. Wijers, Joop Schaye and Jelle S. Kaastra

*stofanova@strw.leidenuniv.nl



Stay tuned!
On arXiv this summer.