



Universiteit  
Leiden  
The Netherlands

## Novel imaging strategies in venous thromboembolism

Dam, L.F. van

### Citation

Dam, L. F. van. (2022, January 27). *Novel imaging strategies in venous thromboembolism*. Retrieved from <https://hdl.handle.net/1887/3254464>

Version: Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/3254464>

**Note:** To cite this publication please use the final published version (if applicable).



6



# CHAPTER

## **Magnetic resonance thrombus imaging to differentiate acute from chronic portal vein thrombosis**

Lisette F. van Dam, Frederikus A. Klok, Maarten E.  
Tushuizen, Walter Ageno, Sarwa Darwish Murad, Guido R.  
van Haren, Menno V. Huisman, Mandy N. Lauw, Antonio  
Iglesias del Sol, Martin N.J.M. Wasser, Ysbrand Willink,  
Lucia J.M. Kroft

*TH Open. 2020 Sep 23;4(3):224-e230*

## ABSTRACT

*Background and aim:* Timely diagnosis and treatment of portal vein thrombosis (PVT) is crucial to prevent morbidity and mortality. However, current imaging tests cannot always accurately differentiate acute from chronic (non-occlusive) PVT. Magnetic resonance non-contrast thrombus imaging (MR-NCTI) has been shown to accurately differentiate acute from chronic venous thrombosis at other locations and may also be of value in the diagnostic management of PVT. This study describes the first phase of the Rhea study (NTR 7061). Our aim was to select and optimize MR-NCTI sequences that would be accurate for differentiation of acute from chronic PVT.

*Methods:* The literature was searched for different MRI sequences for portal vein and acute thrombosis imaging. The most promising sequences were tested in a healthy volunteer, followed by one patient with acute PVT and two patients with chronic PVT, all diagnosed on (repetitive) contrast enhanced computed tomography venography, to optimize the MR-NCTI sequences. All images were evaluated by an expert panel.

*Results:* Several MR-NCTI sequences were identified and tested. Differentiation of acute from chronic PVT was achieved with 3-dimensional T1 Turbo Field Echo (3D T1-TFE) and 3D T1 Fast Field Echo (3D T1 Dixon FFE) sequences with best image-quality. The expert panel was able to confirm the diagnosis of acute PVT on the combined two MR-NCTI sequences and to exclude acute PVT in the two patients with chronic PVT.

*Conclusion:* Using 3D T1-TFE and 3D T1 Dixon FFE sequences we were able to distinguish acute from chronic PVT. This clinically relevant finding will be elucidated in clinical studies to establish their test performance.



## INTRODUCTION

Acute portal vein thrombosis (PVT) has a poor short- and long-term prognosis<sup>1</sup> and a higher mortality rate than the 'usual site' venous thromboembolism, i.e. acute pulmonary embolism and deep vein thrombosis (DVT). Therefore, timely diagnosis and anticoagulant treatment are crucial for the patients' prognosis.<sup>2-4</sup> Distinguishing between a fresh thrombus, in which anticoagulant treatment is indicated, and a chronic (organized, non-resolvable) thrombus is of paramount importance for treatment decision.<sup>5</sup> However, currently available imaging tests are not always able to accurately differentiate acute from chronic PVT, especially when it concerns an organized non-occlusive chronic thrombus without signs of cicatrization of the affected vessel.

Magnetic resonance non-contrast thrombus imaging (MR-NCTI), also referred to as Magnetic Resonance Direct Thrombus Imaging (MRDTI) in previous publications, is a non-contrast-enhanced magnetic resonance imaging (MRI) technique which allows for direct visualization of acute thrombi.<sup>6</sup> This technique is based on the formation of methemoglobin in a fresh thrombus which appears as high signal intensity on a T1 weighted MRI sequence by shortening the T1 relaxation time.<sup>6</sup> This increased signal intensity fades when the thrombus ages over a period of 3-6 months.<sup>7</sup> MR-NCTI has been shown to accurately differentiate acute from chronic DVT and to safely exclude acute recurrent ipsilateral DVT of the legs.<sup>7-9</sup> The diagnostic accuracy of MR-NCTI has not yet been evaluated for the diagnosis of PVT. Because imaging of abdominal veins differs in many ways from imaging of veins in the extremities, e.g. artefacts from ascites and intestinal movement and gas, the MR-NCTI sequences used in the extremities need to be adjusted for optimal visualization of PVT. We set out to select and optimize MR-NCTI sequences for differentiation between acute and chronic portal vein thrombosis.

## MATERIAL & METHODS

This study is the first phase of the Rhea study (NTR 7061), a prospective diagnostic study to evaluate the diagnostic accuracy of MR-NCTI for distinguishing acute from chronic PVT. In the Rhea study our aim is to include 70 PVT patients, including 35 patients with confirmed acute PVT (partial or occlusive), i.e. acute symptoms (<2 weeks) characteristic for PVT, diagnosed with doppler ultrasound (Doppler US),

computed tomography (CT) venography or MRI and 35 patients with confirmed non-symptomatic chronic PVT (partial or occlusive), i.e. unchanged chronic thrombi on 2 serial imaging tests with an at least 3-month interval. The study protocol and its amendments were approved by the institutional review board of the Leiden University Medical Center, Leiden, the Netherlands, and the University of Insubria, Varese, Italy. All patients will be asked for written informed consent.

For this first phase of the Rhea study, we aimed to select and develop MR-NCTI sequences that would be accurate for differentiation of acute from chronic PVT. First, a literature search for different MRI sequences for portal vein and acute thrombosis imaging was performed. The literature search was conducted in PubMed for papers published in English and in humans on December 16<sup>th</sup> 2019 (search strategy detailed in **Appendix A**). Because imaging of the abdomen is hampered by intestinal movements and gas, search terms included techniques with correction for respiratory motion artefacts, good spatial and contrast resolution and fat suppression.

The most promising sequences were adjusted for abdominal imaging and tested in a healthy volunteer using a MRI 3.0 Tesla unit (Philips Ingenia, Philips Medical Systems, Best, The Netherlands). A 55 cm receive-only body multi coil (combination of posterior and anterior coil) was used. Image assessment involved acquiring images in the coronal and axial plane with standard image reconstruction techniques. To gain knowledge of the performance of the sequence to detect fresh blood clots in addition to the visualisation of the venous anatomy, two tests coagulation tubes were attached to the abdomen of the volunteer; one was filled with water (control) and one with clotted blood of a healthy volunteer, prepared and stored at room temperature at least 48 hours before each scanning session. The test sequence scan parameters were adjusted until adequate image quality of the veins with high signal intensity for thrombus in the coagulation tube was achieved.

MR-NCTI sequences showing the best image quality and contrast resolution were tested in three patients with confirmed PVT, including one patient with acute PVT and two patients with chronic PVT. Scan optimization was performed until a clear distinct signal intensity was achieved between acute and chronic PVT.

Finally, the MRI images of the three PVT patients were evaluated by an expert panel, consisting of two radiologists with over 20 years of experience with vascular MRI (LK and MW), one radiology technician with over 20 years of experience with vascular MRI acquisition (GH), one internist with 6 years (FK) and one researcher with 2 years (LD) of experience with vascular MRI interpretation. The evaluation



of the scans was performed and compared with the clinical presentation and the results of other imaging (Doppler US, CT venography or MRI). The scan results were assessed for image quality, venous location, and either presence or absence of acute thrombosis.

## RESULTS

### *Literature search*

We identified the following MRI sequences: 3-dimensional T1 Turbo Field Echo (3D T1 TFE), 3D Turbo Spin Echo with Spectral Attenuated Inversion Recovery (3D TSE SPAIR) and T1 high-resolution isotropic volume excitation (THRIVE) and techniques: black-blood, Dixon and Principle of Selective Excitation Technique (ProSET) which could be suitable for portal vein imaging.

A 3D T1 TFE sequence was shown to be highly accurate for the diagnosis of first deep vein thrombosis and the differentiation of acute deep vein thrombosis from chronic residual vascular abnormalities in the leg.<sup>8-10</sup> Hence, this sequence seemed promising for the differentiation of acute from chronic PVT. Furthermore, in a pilot study 3D T1 TFE and 3D TSE SPAIR sequences successfully confirmed the diagnosis of arm vein thrombosis when compared to ultrasonography or contrast venography.<sup>11</sup> The advantages of 3D TSE-SPAIR over 3D T1 TFE were a higher spatial resolution of the vessel wall and less high signal artefacts in arteries caused by inflow effects. Therefore, both 3D T1 TFE and 3D TSE SPAIR may be suitable for portal vein imaging.

THRIVE sequences may also be a promising MR-NCTI technique in PVT with potential good contrast resolution between high signal intensity of an acute thrombus and low signal intensity intravascular.<sup>12</sup>

The black-blood technique was successfully used to identify deep vein thrombosis of the leg and in cerebral vein thrombosis.<sup>13,14</sup> With this technique the signal from flowing blood is nullified and highlights static anatomy including thrombi.<sup>15</sup> Because of known good contrast resolution between thrombus and flowing blood, we hypothesized that combining the black-blood technique with 3D TSE SPAIR may be suitable for portal vein imaging.

Since it can be difficult to distinguish an acute thrombus from other tissues with a short T1 relaxation time, e.g. fat-tissue which is present in the liver, adding a fat-suppression technique that suppresses the fat signal may help depicting thrombosis. With this technique, selective pulses cause signal from fat to be nulled (saturated) while the water signal is relatively unaffected. The Dixon technique is an often-used fat suppression technique that can be used with several sequences such as T1- and T2-weighted imaging, and gradient echo MRI<sup>16,17</sup>. An alternative to fat suppression methods is water excitation by means of a spectral spatial pulse. With this technique, only water is excited by using section-selective composite pulses, while lipid spins are left in equilibrium, thereby producing no signal.<sup>18</sup> ProSET, a water excitation technique, was successfully used in a pregnant patient with proximal iliac vein thrombosis<sup>19</sup> and may therefore be applicable in portal vein imaging as well.

### *Sequence testing and optimization*

All three selected imaging sequences (3D T1 TFE, 3D TSE SPAIR and THRIVE) and black-blood, Dixon and ProSET techniques were tested in a healthy volunteer.

**Table 1.** 3D T1 TFE and T1 Dixon FFE scan parameters

	<b>3D T1 TFE</b>	<b>3D T1 Dixon FFE</b>
Technique	T1 TFE	T1 FFE
Orientation	Coronal	Coronal/Transversal
Respiratory motion suppression	Respiratory gating	Breath hold
Slices	80	160
Slice thickness (mm)	4.0	3.0
Slice distance (mm)	2.0	1.5
FOV	400	400
Voxel size (mm)	1.56 x 2.24 x 4	1.7 x 1.7 x 3.5
Scan time (ms)	02:36	00:19
Echo time (ms)	3.73	-
Repetition time (ms)	7.41	3.75
Flip angle	10	15


*3D T1 Dixon FFE, three-dimensional T1 fast field echo sequence; 3D T1 TFE, three-dimensional T1 turbo field echo; FOV, field of view.*



3D T1 TFE sequence combined with ProSET technique and the Dixon technique added to a 3D T1 Fast Field Echo sequence (3D T1 Dixon FFE) showed good contrast resolution between the liver parenchyma and intrahepatic veins. Moreover, a high signal intensity was acquired from the coagulation tube (**Figure 1**). These two sequences were thus the most promising for portal vein imaging and were evaluated in three PVT patients (**Table 1**).

3D TSE SPAIR, with and without the black-blood technique, and THRIVE sequence were shown to be suboptimal for abdominal direct thrombus imaging, because of a high signal intensity from blood flow in splanchnic veins and intestines despite saturation slab and low image quality due to motion artifacts. These sequences were excluded for further analysis.


**Figure 1A-D.** MR images of the abdomen in coronal view of a healthy volunteer with a coagulation tube attached to the abdomen; A and C. 3D T1 TFE images at two levels showing a low signal intensity in intrahepatic veins and good contrast resolution between the liver parenchyma and veins. High signal intensity is present in the coagulation tube with clotted blood (arrow). B. and D. 3D T1 Dixon FFE (water-only) images at two levels showing a low signal intensity in the intrahepatic veins and intermediate signal intensity in the coagulation tube (arrow).



### *MR-NCTI optimization in PVT patients*

3D T1 TFE and 3D T1 Dixon FFE sequences were used to evaluate the patient with acute PVT and two patients with chronic PVT (**Table 2**). 3D T1 TFE and 3D T1 Dixon FFE showed a high signal intensity in all abdominal vein segments with acute thrombosis diagnosed on CT venography (**Figure 2**). In the two patients with chronic PVT both sequences showed no increased signal intensity in the portal or mesenteric veins (**Figure 3**). The expert panel was able to confirm the diagnosis of acute PVT on the combined two MR-NCTI sequences and to exclude acute PVT in the two patients with chronic thrombosis. The combination of 3D T1 TFE and 3D T1 Dixon FF was thus judged optimal for locating and differentiating acute from chronic PVT with good image quality and short scanning time (10-15 minutes).

**Figure 2A-F.** Coronal computed tomography (CT) and magnetic resonance imaging (MRI) of the abdomen of a patient diagnosed with acute thrombosis in portal and mesenteric veins; A. CT image after intravenous contrast administration in portal-venous contrast phase shows a large luminal filling defect in the portal vein (arrow). B. MRI, 3D T1 TFE image shows a high signal intensity in the portal vein compatible with acute thrombus (arrow). C. MRI, 3D T1 Dixon FFE (water-only) image shows a high signal intensity in the portal vein (arrow). D. CT image after intravenous contrast administration shows extensive filling defects in mesenteric veins (arrow) with increased attenuation of the surrounding mesenteric fat (encircled). E. MRI, 3D T1 TFE image shows a high signal intensity in the mesenteric veins compatible with acute thrombus (encircled). F. MRI, 3D T1 Dixon FFE (water-only) image shows a high signal intensity in the mesenteric veins (encircled).






**Table 2.** Patient characteristics of the three patients diagnosed with portal vein thrombosis

	<i>Diagnosis at inclusion</i>	<i>Sex</i>	<i>Age (years)</i>	<i>Relevant patient history</i>	<i>Previous venous thromboembolism (VTE)</i>	<i>Anticoagulants at inclusion</i>	<i>Risk factors for VTE</i>
Patient 1	Acute PVT	Female	35	None	No	No	Hormonal contraception use
Patient 2	Chronic PVT	Male	29	Myeloproliferative neoplasia complicated by thrombosis of mesenteric veins, portal and splenic vein. Portal hypertension with esophageal varices.	Yes	No	Myeloproliferative neoplasia (JAK2 mutation)
Patient 3	Chronic PVT	Female	67	Non-alcoholic steatohepatitis (NASH) liver cirrhosis for which liver transplantation. Thrombosis of superior mesenteric vein with ischemic small bowel for which ileum resection.	Yes	Yes	NASH liver cirrhosis

NASH, nonalcoholic steatohepatitis; PVT, portal vein thrombosis; VTE, venous thromboembolism; JAK2, janus kinase 2.

**Figure 3.** A. Computed tomography, axial image of the abdomen in portal-venous contrast phase after intravenous contrast administration, showing a central luminal contrast defect caused by chronic thrombus in the mesenteric vein (arrow). B. Magnetic resonance imaging (MRI), axial 3D T1 Dixon FFE image of the same patient showing isointense signal intensity in the mesenteric vein (arrow). On MRI, the signal intensity of the filling defect in the mesenteric vein is not increased, indicating chronic thrombus.



## DISCUSSION

In this first phase of the Rhea study the combination of 3D T1 TFE and 3D T1 Dixon FFE sequences was found to be the most optimal for diagnosing and differentiating acute from chronic PVT. Differentiation between acute and chronic PVT is essential as current guidelines recommend different anticoagulant strategies in patients with acute or chronic PVT, though based on (very) low level of evidence.<sup>5</sup> However, with currently available imaging tests, Doppler US, CT venography, and MRI, it is not always possible to accurately differentiate acute from chronic thrombosis. The diagnosis of chronic PVT can be made in cases of morphologic changes such as an atretic portal vein, inflow branches and cavernous transformation of the portal vein, also called portal cavernoma.<sup>20,21</sup> However, it may be impossible to differentiate acute from chronic PVT in the presence of an organized non-occlusive chronic PVT, without these morphologic changes. Furthermore, up to 30% of the PVT cases are detected incidentally in imaging studies performed for other indications in which it is very challenging to determine whether the incidentally observed thrombosis is acute, chronic or even an imaging artefact.<sup>22</sup>

MR-NCTI has been shown to be a valuable diagnostic test for the diagnosis of upper extremity DVT, pelvic vein and cerebral vein thrombosis<sup>11,19,23</sup> and was shown to be an accurate, simple, feasible and reproducible diagnostic test in suspected recurrent ipsilateral DVT of the leg.<sup>9</sup> There are some important differences between

venous thrombosis of the portal vein and at other locations. For instance, portal vein imaging is different from imaging of veins in the extremities and brain, due to the presence of ascites, gas and bowel movements which can hamper abdominal vein imaging. Furthermore, PVT occurs in different clinical circumstances (i.e. portal hypertension, cirrhosis) that are less relevant to typical VTE.<sup>24</sup>

There are only a few case-reports available on non-contrast-enhanced MRI for the diagnosis of PVT, using different techniques and without results on the diagnostic accuracy of MRI for PVT.<sup>25,26</sup> In a study from Zirinsky et al, T1- and T2 weighted MR images of 14 patients with acute and chronic PVT on CT or ultrasonography and of 8 patients with portal hypertension but without evidence of PVT were evaluated. With MRI (sub-)acute (<5 weeks old) thrombi appeared hyperintense relative to liver and muscle on both T1- and T2 weighted sequences and older thrombi (2-18 months old) appeared hypointense in some patients, but only on T2-weighted images. Therefore, it was suggested that chronic thrombosis may lose their relative hyperintensity on T1-weighted images and thus may be used for detecting and classification of PVT.<sup>25</sup> In a case-series from Haddad et al, thrombosis on non-contrast-enhanced T1-weighted images appeared different depending on the age of the thrombus, with a high signal intensity in subacute splanchnic vein thrombosis (<6 weeks) and low signal intensity in more chronic (> 2 months old) thrombosis.<sup>26</sup>

There are some limitations of the current study. First, this study includes only three patients and therefore, the validity of the results must be evaluated in a large cohort before we can proceed to an outcome study. Such an accuracy study is currently ongoing. Furthermore, the evaluation of MR-NTCI images by the expert panel was not blinded for other imaging studies including ultrasound and CT images. Additionally, due to the small number of patients the interobserver agreement could not be established.

In conclusion, we were able to identify two MR-NTCI sequences, 3D T1 TFE and 3D T1 Dixon FFE, that were able to diagnose and differentiate acute from chronic PVT. With our previous experience based on imaging of DVT in the lower and upper extremities<sup>9,11,19</sup>, we believe that the image quality is sufficient to be of clinical value and initiate the clinical part of the Rhea study (NTR 7061) to establish the diagnostic accuracy of MR-NCTI for the differentiation of acute from chronic PVT. Furthermore, the interobserver agreement of these sequences for PVT imaging will be assessed.

## REFERENCES

1. Sogaard KK, Darvalics B, Horvath-Puho E, Sorensen HT. Survival after splanchnic vein thrombosis: A 20-year nationwide cohort study. *Thrombosis research*. 2016;141:1-7.
2. Ageno W, Riva N, Schulman S, et al. Long-term Clinical Outcomes of Splanchnic Vein Thrombosis: Results of an International Registry. *JAMA internal medicine*. 2015;175(9):1474-1480.
3. Plessier A, Darwish-Murad S, Hernandez-Guerra M, et al. Acute portal vein thrombosis unrelated to cirrhosis: a prospective multicenter follow-up study. *Hepatology*. 2010;51(1):210-218.
4. Pettinari I, Vukotic R, Stefanescu H, et al. Clinical Impact and Safety of Anticoagulants for Portal Vein Thrombosis in Cirrhosis. *The American journal of gastroenterology*. 2019;114(2):258-266.
5. Simonetto DA, Singal AK, Garcia-Tsao G, Caldwell SH, Ahn J, Kamath PS. ACG Clinical Guideline: Disorders of the Hepatic and Mesenteric Circulation. *The American journal of gastroenterology*. 2020;115(1):18-40.
6. Moody AR. Magnetic resonance direct thrombus imaging. *Journal of thrombosis and haemostasis : JTH*. 2003;1(7):1403-1409.
7. Westerbeek RE, Van Rooden CJ, Tan M, et al. Magnetic resonance direct thrombus imaging of the evolution of acute deep vein thrombosis of the leg. *Journal of thrombosis and haemostasis : JTH*. 2008;6(7):1087-1092.
8. Tan M, Mol GC, van Rooden CJ, et al. Magnetic resonance direct thrombus imaging differentiates acute recurrent ipsilateral deep vein thrombosis from residual thrombosis. *Blood*. 2014;124(4):623-627.
9. van Dam LF, Dronkers CEA, Gautam G, et al. Magnetic resonance imaging for diagnosis of recurrent ipsilateral deep vein thrombosis. *Blood*. 2020;135(16):1377-1385.
10. Fraser DG, Moody AR, Morgan PS, Martel AL, Davidson I. Diagnosis of lower-limb deep venous thrombosis: a prospective blinded study of magnetic resonance direct thrombus imaging. *Ann Intern Med*. 2002;136(2):89-98.
11. Dronkers CEA, Klok FA, van Haren GR, et al. Diagnosing upper extremity deep vein thrombosis with non-contrast-enhanced Magnetic Resonance Direct Thrombus Imaging: A pilot study. *Thrombosis research*. 2018;163:47-50.
12. Riffel P, Attenberger UI, Kannengiesser S, et al. Highly accelerated T1-weighted abdominal imaging using 2-dimensional controlled aliasing in parallel imaging results in higher acceleration: a comparison with generalized autocalibrating partially parallel acquisitions parallel imaging. *Investigative radiology*. 2013;48(7):554-561.
13. Treitl KM, Treitl M, Kooijman-Kurfuerst H, et al. Three-dimensional black-blood T1-weighted turbo spin-echo techniques for the diagnosis of deep vein thrombosis in comparison with contrast-enhanced magnetic resonance imaging: a pilot study. *Investigative radiology*. 2015;50(6):401-408.



14. Yang Q, Duan J, Fan Z, et al. Early Detection and Quantification of Cerebral Venous Thrombosis by Magnetic Resonance Black-Blood Thrombus Imaging. *Stroke*. 2016;47(2):404-409.
15. Xie G, Chen H, He X, et al. Black-blood thrombus imaging (BTI): a contrast-free cardiovascular magnetic resonance approach for the diagnosis of non-acute deep vein thrombosis. *Journal of cardiovascular magnetic resonance : official journal of the Society for Cardiovascular Magnetic Resonance*. 2017;19(1):4.
16. Del Grande F, Santini F, Herzka DA, et al. Fat-suppression techniques for 3-T MR imaging of the musculoskeletal system. *Radiographics : a review publication of the Radiological Society of North America, Inc*. 2014;34(1):217-233.
17. Guerini H, Omoumi P, Guichoux F, et al. Fat Suppression with Dixon Techniques in Musculoskeletal Magnetic Resonance Imaging: A Pictorial Review. *Seminars in musculoskeletal radiology*. 2015;19(4):335-347.
18. Hauger O, Dumont E, Chateil JF, Moinard M, Diard F. Water excitation as an alternative to fat saturation in MR imaging: preliminary results in musculoskeletal imaging. *Radiology*. 2002;224(3):657-663.
19. Dronkers CE, Sramek A, Huisman MV, Klok FA. Accurate diagnosis of iliac vein thrombosis in pregnancy with magnetic resonance direct thrombus imaging (MRDTI). *BMJ case reports*. 2016;2016.
20. Intagliata NM, Caldwell SH, Tripodi A. Diagnosis, Development, and Treatment of Portal Vein Thrombosis in Patients With and Without Cirrhosis. *Gastroenterology*. 2019;156(6):1582-1599.e1581.
21. De Gaetano AM, Lafortune M, Patriquin H, De Franco A, Aubin B, Paradis K. Cavernous transformation of the portal vein: patterns of intrahepatic and splanchnic collateral circulation detected with Doppler sonography. *AJR American journal of roentgenology*. 1995;165(5):1151-1155.
22. Amitrano L, Guardascione MA, Scaglione M, et al. Prognostic factors in noncirrhotic patients with splanchnic vein thromboses. *The American journal of gastroenterology*. 2007;102(11):2464-2470.
23. Yang Q, Duan J, Fan Z, et al. Early Detection and Quantification of Cerebral Venous Thrombosis by Magnetic Resonance Black Blood Thrombus Imaging (MRBTI). *Stroke*. 2016;47(2):404-409.
24. Denninger MH, Chait Y, Casadevall N, et al. Cause of portal or hepatic venous thrombosis in adults: the role of multiple concurrent factors. *Hepatology*. 2000;31(3):587-591.
25. Zirinsky K, Markisz JA, Rubenstein WA, et al. MR imaging of portal venous thrombosis: correlation with CT and sonography. *AJR American journal of roentgenology*. 1988;150(2):283-288.
26. Haddad MC, Clark DC, Sharif HS, al Shahed M, Aideyan O, Sammak BM. MR, CT, and ultrasonography of splanchnic venous thrombosis. *Gastrointestinal radiology*. 1992;17(1):34-40.

## Appendix A: Pubmed search string

((("Venous Thrombosis"[mesh]) OR "splanchnic vein thrombosis"[tw] OR "splanchnic vein thromboses"[tw] OR "splanchnic venous thrombosis"[tw] OR "mesenteric vein thrombosis"[tw] OR "mesenteric vein thromboses"[tw] OR "mesenteric venous thrombosis"[tw] OR "mesenteric venous thromboses"[tw] OR "portosplenomesenteric venous thrombosis"[tw] OR "splanchnic"[ti] OR "mesenteric"[ti] OR "portosplenomesenteric"[ti] OR "vein"[ti] OR "venous"[ti] OR ("abdominal"[ti] AND "imaging"[ti]) OR ("thrombosis"[ti] OR "thromboses"[ti] OR "embolism"[ti] OR "thrombus"[ti] OR "occlusion"[ti]))) AND ("MRDTI"[tw] OR MRDTI[tw] OR "magnetic resonance direct thrombus imaging"[tw] OR "magnetic resonance"[tw] OR "Magnetic Resonance Imaging"[mesh] OR "Magnetic Resonance Imaging"[tw] OR "MRI"[tw] OR mr imag\*[tw] OR "thrombus imaging"[tw]) OR ("Fat"[All Fields] AND "Suppression"[All Fields]) AND ("methods"[All Fields] OR "techniques"[All Fields] OR "methods"[mesh] OR "techniques"[All Fields])).

For online supplementary material, please scan this QR code:



