



Universiteit
Leiden
The Netherlands

Chemo-dynamical modeling of Cha-mms1 to predict new solid-phase species for detection with JWST

Jin, M.; Lam, K.; Terwisscha van Scheltinga, J.; McClure, M.; Li, Z.; Boogert, A.; ... ; Garrod, R.

Citation

Jin, M., Lam, K., Terwisscha van Scheltinga, J., McClure, M., Li, Z., Boogert, A., ... Garrod, R. (2021). Chemo-dynamical modeling of Cha-mms1 to predict new solid-phase species for detection with JWST. *Bulletin Of The Aas*, (6), 326.03. Retrieved from <https://hdl.handle.net/1887/3263679>

Version: Publisher's Version
License: [Creative Commons CC BY 4.0 license](https://creativecommons.org/licenses/by/4.0/)
Downloaded from: <https://hdl.handle.net/1887/3263679>

Note: To cite this publication please use the final published version (if applicable).

Bulletin of the AAS • Vol. 53, Issue 6

Chemo-dynamical Modeling Of Cha-mms1 To Predict New Solid-phase Species For Detection With JWST

**M. Jin¹, K. Lam¹, J. Terwisscha van Scheltinga², M. McClure³, Z. Li¹,
A. Boogert⁴, E. Herbst¹, R. Garrod¹**

¹University of Virginia, Charlottesville, VA, ²Leiden University, Leiden, Netherlands,

³Universiteit van Amsterdam, Amsterdam, Netherlands, ⁴University of Hawaii, Honolulu, HI

Published on: Jun 18, 2021

License: [Creative Commons Attribution 4.0 International License \(CC-BY 4.0\)](https://creativecommons.org/licenses/by/4.0/)

Chemical models and experiments indicate that interstellar dust grains and their ice mantles play an important role in the production of complex organic molecules (COMs). The meeting of chemical species adsorbed from the gas phase onto the grain/ice surface allows reactions to occur efficiently, and leads to the emergence of the ice mantles as a major reservoir of volatile species. To date, the most complex solid-phase molecule detected with certainty in the ISM is methanol, but it is hoped that the upcoming James Webb Space Telescope (JWST), aided by new laboratory data, will be able to identify yet larger organic species. In this study, we use a coupled chemo-dynamical model to predict new candidate species for JWST detection toward the young star-forming core Cha-MMS1, combining the gas-grain chemical kinetic model MAGICKAL with a 1-D radiative hydrodynamics (RHD) simulation. The abundances of the main ice constituents with respect to water in the model match well with observational values. The model also predicts high solid-phase abundances of intermediate radicals associated with the hydrogenation of CO to methanol ($> 0.1\%$ with respect to water ice). Four oxygen-bearing COMs (ethanol, dimethyl ether, methyl formate, and acetaldehyde) as well as formic acid, show abundances as high as $> 0.01\%$ with respect to water ice. N-bearing COMs are generally less abundant than O-bearing ones, suggesting methylamine as the only N-bearing solid-phase COM candidate for JWST detection.