



Universiteit
Leiden
The Netherlands

Cosmic evolution of the H₂ mass density and the epoch of molecular gas

Garratt, T.K.; Coppin, K.E.K.; Geach, J.E.; Almaini, O.; Hartley, W.G.; Maltby, D.T.; ... ; Werf, P.P. van der

Citation

Garratt, T. K., Coppin, K. E. K., Geach, J. E., Almaini, O., Hartley, W. G., Maltby, D. T., ... Werf, P. P. van der. (2021). Cosmic evolution of the H₂ mass density and the epoch of molecular gas. *The Astrophysical Journal*, 912(1). doi:10.3847/1538-4357/abec81

Version: Not Applicable (or Unknown)

License: [Leiden University Non-exclusive license](#)

Downloaded from: <https://hdl.handle.net/1887/3263683>

Note: To cite this publication please use the final published version (if applicable).

Cosmic evolution of the H₂ mass density and the epoch of molecular gas

T. K. GARRATT,¹ K. E. K. COPPIN,¹ J. E. GEACH,¹ O. ALMAINI,² W. G. HARTLEY,³ D. T. MALTBY,² C. J. SIMPSON,⁴
A. WILKINSON,⁵ C. J. CONSELICE,² M. FRANCO,¹ R. J. IVISON,⁶ M. P. KOPROWSKI,⁷ C. C. LOVELL,¹ A. POPE,⁸
D. SCOTT,⁹ AND P. VAN DER WERF¹⁰

¹*Centre for Astrophysics Research, University of Hertfordshire, Hatfield, AL10 9AB, UK.*

²*School of Physics and Astronomy, University of Nottingham, University Park, Nottingham, NG7 2RD, UK*

³*Department of Astronomy, University of Geneva, ch. d'Écogia 16, CH-1290 Versoix, Switzerland*

⁴*Gemini Observatory, Hilo, HI 96720, USA*

⁵*Sterrenkundig Observatorium, Universiteit Gent, Krijgslaan 281 S9, 9000 Gent, Belgium*

⁶*European Southern Observatory, Karl Schwarzschild Strasse 2, Garching, Germany*

⁷*Institute of Astronomy, Faculty of Physics, Astronomy and Informatics, Nicolaus Copernicus University, Grudziadzka 5, 87-100 Torun, Poland*

⁸*Department of Astronomy, University of Massachusetts, 710 North Pleasant Street Amherst, MA 01003, USA*

⁹*Dept. of Physics & Astronomy, University of British Columbia, Vancouver, Canada*

¹⁰*Leiden Observatory, Leiden University, P.O. Box 9513, NL-2300 RA Leiden, The Netherlands*

ABSTRACT

We present new empirical constraints on the evolution of ρ_{H_2} , the cosmological mass density of molecular hydrogen, back to $z \approx 2.5$. We employ a statistical approach measuring the average observed $850 \mu\text{m}$ flux density of near-infrared selected galaxies as a function of redshift. The redshift range considered corresponds to a span where the $850 \mu\text{m}$ band probes the Rayleigh-Jeans tail of thermal dust emission in the rest-frame, and can therefore be used as an estimate of the mass of the interstellar medium (ISM). Our sample comprises of $\approx 150,000$ galaxies in the UKIDSS-UDS field with near-infrared magnitudes $K_{\text{AB}} \leq 25$ mag and photometric redshifts with corresponding probability distribution functions derived from deep 12-band photometry. With a sample approximately 2 orders of magnitude larger than in previous works we significantly reduce statistical uncertainties on ρ_{H_2} to $z \approx 2.5$. Our measurements are in broad agreement with recent direct estimates from blank field molecular gas surveys, finding that the epoch of molecular gas coincides with the peak epoch of star formation with $\rho_{\text{H}_2} \approx 2 \times 10^7 \text{M}_{\odot} \text{Mpc}^{-3}$ at $z \approx 2$. We demonstrate that ρ_{H_2} can be broadly modelled by inverting the star-formation rate density with a fixed or weakly evolving star-formation efficiency. This “constant efficiency” model shows a similar evolution to our statistically derived ρ_{H_2} , indicating that the dominant factor driving the peak star formation history at $z \approx 2$ is a larger supply of molecular gas in galaxies rather than a significant evolution of the star-formation rate efficiency within individual galaxies.

Keywords: galaxies: evolution — galaxies: high-redshift — galaxies: ISM — galaxies: star formation

1. INTRODUCTION

Three intimately linked observational tracers broadly characterise the cosmic evolution of galaxies: the volume averaged star-formation rate density $\rho_{\dot{M}_{\star}}(z)$, the stellar mass density $\rho_{M_{\star}}(z)$, and the molecular gas density $\rho_{M_{\text{H}_2}}(z)$. Our current understanding of galaxy evolution is largely driven by comprehensive measurements

of the former two (see Madau & Dickinson 2014, for a review), with a clear empirical picture emerging of an evolution of star formation, which rises rapidly to a peak around $z \approx 2$ and then decays to the present day. Completing the triptych is important since the evolution of the molecular gas content of galaxies encodes several important pieces of astrophysics: gas consumption in star formation; gas recycling via feedback; and fresh gas accretion. Ultimately, it is the evolution of molecular gas that drives galaxy evolution as it is the fuel from which stars are assembled. Measurements of molecular

gas in galaxies are therefore needed to complete the picture, and to resolve a key outstanding question: *Was the peak of star formation history driven by a larger supply of molecular gas or because galaxies formed stars more efficiently (e.g., driven by galaxy mergers/instabilities etc.), or both?*

The bulk of the cold gas reservoir in the Universe is comprised of hydrogen gas in the form of atomic hydrogen (H_I) and molecular hydrogen (H_2). In the current model of galaxy formation gas is delivered into galaxies via hot- or cold-mode accretion (e.g., [Birnboim & Dekel 2003](#)). The cooling gas must form H_2 for star formation to occur. The two main routes to H_2 formation in galaxies are via the gas phase reaction $\text{H} + \text{e}^- \rightarrow \text{H}^- + \gamma$, $\text{H}^- + \text{H} \rightarrow \text{H}_2 + \text{e}^-$, and via a dust phase, where H_2 forms on the surface of dust grains via efficient three-body reactions ([Gould & Salpeter 1963](#)).

H_2 radiates poorly in typical ISM conditions due to the lack of a permanent dipole moment and a minimum rotational excitation temperature that is significantly higher (≈ 500 K) than typical temperatures of the cold star-forming ISM ([Wakelam et al. 2017](#)). However, H_2 can be indirectly traced through its interactions with CO, which traces the same cold, dense ISM and has a low dipole moment enabling its excitation in regions of low density ($n_{\text{crit}} \sim 10^2 \text{ cm}^{-3}$). Consequently, ^{12}CO , the second most abundant molecule in the ISM, is commonly used as a tracer of the available reservoir of molecular gas in galaxies (e.g., [Solomon & Vanden Bout 2005](#); [Carilli & Walter 2013](#)). The ground state transition $\text{CO}(J = 1 \rightarrow 0)$ is a reliable tracer of total molecular gas, with the conversion factor from CO luminosity to H_2 gas mass ($\alpha_{\text{CO}} = M_{\text{H}_2}/L'_{\text{CO}}$) calibrated locally (see [Bolatto et al. 2013](#), for a review). Observing the ground-state transition line avoids additional uncertainties inherent in observations of higher- J CO transitions, which require a correction for gas excitation to derive the equivalent $\text{CO}(J = 1 \rightarrow 0)$ luminosity.

Until recently measurements of the cosmological molecular gas mass density were hampered by a paucity of observational data. Over the past few years direct measurements of the cold molecular gas reservoirs of individual galaxies have increased rapidly, with surveys primarily targeting star-forming and lensed galaxies (e.g., [Frayer et al. 1998, 1999](#); [Coppin et al. 2007](#); [Tacconi et al. 2010](#); [Ivison et al. 2011](#); [Thomson et al. 2012](#); [Bothwell et al. 2013](#); [Riechers et al. 2013](#); [Stach et al. 2017](#); [Oteo et al. 2018](#); [Gómez-Guijarro et al. 2019](#); [Lenkić et al. 2019](#)). However, as these surveys rely on observationally-expensive detections of faint spectral lines, measurements of molecular gas mass are still dwarfed in number in comparison to the samples for

which star-formation rate (SFR) and stellar mass estimates are derived. Moreover, to properly assess the cosmological evolution of the cold gas content of galaxies a blank field survey approach is required to measure the gas mass function, rather than targeted (and therefore biased) observations of high- z galaxies as has generally been the case for cold gas observations outside the local volume.

Recently surveys using a blank field molecular line scan strategy have emerged as an alternative to targeted observations. These surveys evade many of the biases towards massive star-forming galaxies inherent in targeted approaches. The inaugural blank field CO survey employed the Plateau de Bure Interferometer in observations of the *Hubble* Deep Field North ([Walter et al. 2014](#); [Decarli et al. 2014](#)). This was followed more recently by the ALMA Spectroscopic Survey in the *Hubble* Ultra Deep Field (ASPECS; [Walter et al. 2016](#); [Decarli et al. 2016a,b](#); [Aravena et al. 2016a,b](#); [Bouwens et al. 2016](#); [Carilli et al. 2016](#)), the ASPECS Large Program (ASPECS LP; [Decarli et al. 2019](#); [González-López et al. 2019](#); [Boogaard et al. 2019](#); [Popping et al. 2019](#); [Decarli et al. 2020](#)) and the CO Luminosity Density at High Redshift survey (COLDZ; [Pavesi et al. 2018](#); [Riechers et al. 2019](#)). These surveys have presented results setting out valuable new constraints on the evolution of ρ_{H_2} over a redshift range $0 \lesssim z \lesssim 7$, obtained through blank field observations of CO line emission. However, due to low number statistics and the small survey areas (which are prone to strong clustering-enhanced sample variance) used to derive the CO luminosity functions, these measurements are hampered by large statistical uncertainties.

To combat the shortfall in direct measurements of molecular gas [Scoville \(2013\)](#); [Scoville et al. \(2014, 2016, 2017\)](#) employed a complementary approach that utilises submillimeter observations of the long wavelength dust continuum as a measure of the molecular gas mass in galaxies. The Rayleigh-Jeans (RJ) tail is nearly always optically thin, and consequently measurements of dust emission can be used as a direct probe of molecular gas mass (e.g., [Eales et al. 2012](#); [Magdis et al. 2012](#)). Whilst ordinarily a conversion from dust to gas mass would require dust emissivity and dust-to-gas abundance to be constrained, [Scoville et al. \(2014, 2016, 2017\)](#) circumvent this by deriving an empirically calibrated *RJ luminosity-to-gas mass* ratio using $\text{CO}(J = 1 \rightarrow 0)$ and submillimeter continuum observations of a sample of normal star-forming and starburst galaxies at low- z and submillimeter galaxies (SMGs) at high- z . This approach requires assumptions about dust temperature and the evolution of the gas-to-dust mass ratio, but provides molecular

gas mass (M_{mol}) estimates within factor of ≈ 2 accuracy (e.g., Scoville et al. 2016; Kaasinen et al. 2019). Since dust continuum measurements can be made in minutes (in contrast to CO line observations which can take multiple hours, e.g., Bothwell et al. 2013; Tacconi et al. 2013) this method can be used to derive molecular gas measurements for much larger samples of galaxies.

The Scoville et al. (2016) RJ luminosity-to-gas mass calibration has been used to estimate the molecular gas mass for ~ 700 ALMA-detected galaxies from the COSMOS field, with Scoville et al. (2017) deriving molecular gas masses for individual galaxies at redshifts $0.3 < z < 4.5$ and Liu et al. (2019) extending this approach to redshifts of $z \approx 6$. This method has also been used in combination with stacking methodologies to estimate average molecular gas masses for large samples of galaxies. Millard et al. (2020) use the Scoville et al. (2016) calibration and apply this to a sample of 63,658 galaxies to derive the gas mass fraction out to $z \approx 5$. Magnelli et al. (2020) use a method similar to Scoville et al. (2016, and which is equivalent at solar metallicity) and apply this to a sample of 555 galaxies to derive the molecular gas mass density to $z \approx 3$.

In this paper we contribute to the picture of cosmic galaxy evolution by building on the approach of Scoville (2013); Scoville et al. (2014, 2016, 2017) estimating the evolution of the cosmological mass density of molecular hydrogen to $z \approx 2.5$ via the average submillimeter continuum emission of a sample of 150,000 galaxies selected from a deep near-infrared survey in the well studied UKIDSS-UDS field. The Ultra-Deep Survey (UDS) is the deepest component of the UK InfraRed Telescope (UKIRT) Infrared Deep Sky Survey (UKIDSS; Lawrence et al. 2007). We limit our estimate of the molecular gas mass density to $z \approx 2.5$ as the Scoville et al. (2016) calibration has only been shown to be robust out to this redshift. Adopting a statistical approach allows us to take advantage of a near-infrared selected sample which is an order of magnitude larger than in surveys that measure dust emission (e.g., Scoville et al. 2017; Liu et al. 2019; Magnelli et al. 2020) or CO spectral line emission (e.g., Walter et al. 2016; Riechers et al. 2019; Decarli et al. 2019; Kaasinen et al. 2019) for individual sources. Our method differs from previous stacking approaches (e.g., Millard et al. 2020; Magnelli et al. 2020) as we do not use a combination of spectroscopic and photometric redshifts for our binning. Instead, in the absence of a sample complete with spectroscopic redshifts, we utilise the full photometric redshift probability distribution functions for all our sources. Our method is complementary to previous works in this field (e.g., Decarli et al. 2020; Liu et al. 2019; Magnelli et al. 2020; Riechers

et al. 2019) and allows us to reduce the statistical uncertainties on the cosmological molecular gas mass density out to $z \approx 2.5$.

This paper is organized as follows: in Section 2, we define the maps and catalogs used; in Section 3, we present a 3-dimensional stacking method which we employ to measure the average (stacked) observed $850 \mu\text{m}$ flux densities for near-infrared selected galaxies as a function of redshift; in Section 4, we show that the approach of Scoville et al. (2016) can be applied to our stacked $850 \mu\text{m}$ flux densities to derive the cosmological molecular gas density to $z \approx 2.5$. We also demonstrate that the cosmic molecular gas density can be broadly modelled by 2 complementary approaches (i) from the halo mass function assuming a constant halo mass range, and employing stellar-halo mass and ISM-stellar mass ratios, and (ii) inverting the star-formation rate density assuming a “constant efficiency” model, and in Section 5 we interpret the overall evolution of the cosmic molecular gas mass density in the context of our results and in comparison to previous works. We present our conclusions in Section 6. We assume a Planck 2015 cosmology, where $\Omega_{\text{m}} = 0.31$, $\Omega_{\Lambda} = 0.69$, $H_0 = 68 \text{ km s}^{-1} \text{ Mpc}^{-1}$ (Planck Collaboration et al. 2016), and a Chabrier (2003) Initial Mass Function. The AB magnitude system is used throughout.

2. DATA

2.1. SCUBA-2 Cosmology Legacy Survey

The UKIDSS-UDS field was mapped at $850 \mu\text{m}$ as part of the Submillimetre Common-User Bolometer Array 2 (SCUBA-2) Cosmology Legacy Survey (Geach et al. 2017). The full details of the data collection, reduction and map properties are given in Geach et al. (2017). Briefly, the beam-convolved map spans approximately 1 deg^2 covering the bulk of the multi-wavelength coverage of this field, with a uniform (instrumental) noise of $\sigma_{850} = 0.9 \text{ mJy beam}^{-1}$. Geach et al. (2017) estimate the SCUBA-2 confusion limit to be $\sigma_{\text{conf}} = 0.8 \text{ mJy beam}^{-1}$. The beam full width half maximum (FWHM) is approximately $15''$, with a full analytic description of the point spread function (PSF) given by Geach et al. (2017).

2.2. UKIDSS-UDS ultraviolet–optical–mid-infrared imaging and catalog

The UDS Data Release 11 (DR11) 12-band matched catalog is K -band selected with the 95% completeness limit estimated to be $K_{\text{AB}} = 25 \text{ mag}$. The full details of this catalog will be comprehensively provided in Almaini et al., (in prep.) and Hartley et al., (in prep.), and only a summary is given here. The catalog provides

photometry in 12 bands ($U, B, V, R, i', z', Y, J, H, K$, $3.6\ \mu\text{m}$ and $4.5\ \mu\text{m}$), where available.

The J, H , and K photometry is taken from the DR11 release of UKIDSS-UDS. The UKIDSS project, described in Lawrence et al. (2007) utilises the UKIRT Wide Field CAMera (WFCAM; Casali et al. 2007). The photometric system and calibration are outlined in Hewett et al. (2006) and Hodgkin et al. (2009), respectively, and the pipeline processing and science archive are described in Irwin et al., (in prep.) and Hambly et al. (2008). UKIDSS-UDS covers an area of $0.8\ \text{deg}^2$, reaching median depths of $J=25.6$, $H=25.1$, and $K=25.3$ (5σ , AB, estimated from $2''$ apertures in source free areas; Almaini et al. in prep.).

The B, V, R, i' , and z' optical imaging is from the Subaru/*XMM-Newton* Deep Survey, which utilises Suprime-Cam on the Subaru Telescope (Furusawa et al. 2008). U -band data are from the Canada-France-Hawaii Telescope Megacam instrument (Almaini et al. in prep.) and Y -band imaging is obtained from the VISTA Deep Extragalactic Observations survey (VISTA-VIDEO: Jarvis et al. 2013). The InfraRed Array Camera (IRAC) imaging at $3.6\ \mu\text{m}$ and $4.5\ \mu\text{m}$ is from the Spitzer UKIDSS Ultra Deep Survey (SpUDS: PI Dunlop), combined with deeper data from the Spitzer Extended Deep Survey (SEDS: Ashby et al. 2013). To expand the coverage to outer regions of the field, shallower data are also used from the SIRTf Wide-area InfraRed Extragalactic survey (SWIRE: Lonsdale et al. 2003).

UDS DR11 provides image masks, with masked regions corresponding to image boundaries, artefacts, and bright stars. We employ the UDS binary mask for “good” regions which has an unmasked area of $0.64\ \text{deg}^2$. This binary mask combines the masked regions of the photometry images detailed above (not including the deeper SEDS or SpUDS IRAC images).

We also utilise the subsets feature of the UKIDSS-UDS catalog and for our galaxy sample chose the catalog-defined “good galaxy” subset, which comprises 217,429 sources. These sources have full 12-band photometry and lie within the corresponding “good” mask regions, are not cross-talk sources (for which JHK photometry is likely compromised), and are not classified as stars.

UDS DR11 also includes photometric redshifts derived using the code EAZY (Easy and Accurate Z_{phot} from Yale; Brammer et al. 2008). To estimate the photometric redshift for each source the 12-band broadband photometry was fit with a spectral energy distribution template producing a redshift probability distribution (Hartley et al. in prep.). We utilise both the maximum-

likelihood photometric redshifts and redshift probability distributions provided with UDS DR11. EAZY performs well compared to other commonly used photometric redshift codes (e.g., ZPHOT, HYPERZ, RAINBOW; Dahlen et al. 2013), with the resulting normalized mean absolute deviation between EAZY derived photometric redshifts and spectroscopic redshifts found to be only $\sigma_{\text{nmad}} \approx 0.02$ (Hartley et al. in prep.).

3. METHODS

We employ a 3-dimensional stacking approach based on the simultaneous stacking algorithm SIMSTACK (presented in detail in Viero et al. 2013). This method allows for the simultaneous fitting of the average observed flux density for multiple populations that contribute to the flux density in the observed map (such as a population of galaxies split into bins of redshift). Importantly, this method takes into account the (usually) large beam in single-dish submillimeter maps with simulations demonstrating that this method returns an unbiased estimate of the average observed flux density for beam sizes ranging from $\text{FWHM} = 15\text{--}35''$ (Viero et al. 2013). This approach also mitigates against boosting of stacking signals from clustered galaxies (e.g., Chary & Pope 2010; Alberts et al. 2014).

Our goal is to find the average observed submillimeter flux densities at given redshift intervals for a population of near-infrared selected galaxies that best fit the observed flux density in the SCUBA-2 map, taking into account the convolution of point sources with the large beam. In this work, rather than binning galaxies by discrete photometric redshift values, we split our sample across redshift intervals according to the redshift probability distribution of each source.

First, we define our sample, performing a selection in observed K -band total magnitude, $K_{\text{AB}} \leq 25\ \text{mag}$, with the faint-end corresponding to the 95% completeness limit of the UKIDSS-UDS catalog, giving us a sample of 153,399 galaxies. At this limiting magnitude the 95% stellar mass completeness is $\approx 10^{9.5}\ M_{\odot}$ at $z = 2.5$ (Wilkinson et al. in prep.). The redshift probability distribution, $\mathcal{P}(z)$, for each source is discretized in bins of Δz (Hartley et al. in prep.). We make a completeness correction to the redshift probability distribution of each source, such that $\mathcal{P}(z)$ of a source of magnitude K integrates to $C(K)^{-1}$, where $C(K)$ is the catalog completeness at K (Hartley et al. in prep.). We assume there is no systematic redshift bias in $C(K)$ for this correction.

With the sample defined we consider a sky model in which each galaxy contributes a flux density that can be

described as

$$S_\nu = \int_0^\infty S_\nu(z) \mathcal{P}(z) dz. \quad (1)$$

where $\mathcal{P}(z)$ is the normalized redshift probability distribution function and $S_\nu(z)$ is the flux density “weighting” at redshift z . In practice we have discrete redshift probability distributions defined over R bins such that, for a population of N galaxies, the flux density in the ij th pixel of a map can be written

$$M_{ij} = \sum_p^{N_{ij}} \sum_q^R S_\nu(z_q)_{p,ij} \mathcal{P}(z_q)_{p,ij} \Delta z \quad (2)$$

Because of the PSF, the flux contribution of each galaxy is distributed over many pixels according to the convolution

$$\mathcal{M} = M \otimes \text{PSF}. \quad (3)$$

In effect equation 1 uses $\mathcal{P}(z)$ to split each of the N galaxies in our K -selected sample into R redshift bins and assumes that the galaxies in each redshift bin can be represented by an average observed flux density, $\langle S_\nu(z) \rangle$. This is effectively the “stacked” flux density.

With the model sky defined we consider an optimization problem where the set of average observed flux densities, $\langle S_\nu(z) \rangle$, per redshift interval in equation 1 are unknown coefficients. A key decision in defining our sky model is in the binning of $\mathcal{P}(z)$. The UKIDSS-UDS $\mathcal{P}(z)$ are binned in non-linear steps of $1 + z_{(n+1)} = 1.001(1 + z_n)$. This would result in hundreds of free parameters across the redshift range of interest, which is computationally impractical as well as unnecessary given the photometric redshift uncertainties. Instead we bin each $\mathcal{P}(z)$ to $\Delta z = 0.5$, giving 20 equally-sized bins across the redshift range $0 < z \lesssim 10$.

We aim to find the optimal set of average flux densities that minimizes the square of the residual flux between the model in equation 3 and the observed beam-convolved map, weighted by the noise. We use the Markov chain Monte Carlo (MCMC) sampler `emcee` (Foreman-Mackey et al. 2013) to estimate the best fit flux densities and their uncertainties. We minimize a negative log likelihood $\ln(\mathcal{L}) = -0.5\chi^2$, with

$$\chi^2 = \sum_{ij} \left(\frac{\mathcal{O}_{ij} - \mathcal{M}_{ij}}{\sigma_{\text{rms},ij}} \right)^2 \quad (4)$$

where \mathcal{O} is the observed map and σ_{rms} is the instrumental noise map. We initialise 1000 “walkers” with an uninformative prior, such that each walker is set with a vector of flux densities (representing $S_\nu(z)$) with each

flux density drawn from a Gaussian distribution of mean 0.5 mJy and width 0.05 mJy. The sampler runs for 1000 iterations with the first 500 iterations discarded as burn-in. The best fitting flux densities and the 1σ bounds are estimated from the 16th, 50th and 84th percentiles of accepted samples for 500 iterations. In Figures 1a, 1b and 1c we show the `emcee` corner plot of the posterior distributions for all our free parameters.

To estimate the additional uncertainty on the stacked flux densities due to sampling variance we employ the “delete one” jackknife technique (Tukey 1958), splitting the map into $A = 21$ approximately equal area sectors and running the MCMC fit for each jackknife. We find that the sampler chains converge quickly (within 200 steps), and tests indicate that the best fit parameters are insensitive to the initialisation parameters. The covariance matrix is given by

$$C_{ij} = \frac{A-1}{A} \sum_{i=1}^A (S_i^k - \bar{S}_i) (S_j^k - \bar{S}_j) \quad (5)$$

where S_i^k is the average flux density in the i th redshift bin, eliminating the k th sample and \bar{S}_i is the average over all samples. The 1σ uncertainties on the stacked fluxes are estimated by the square root of the diagonal elements of C .

At high- z the increase in the cosmic microwave background (CMB) temperature affects the measurement of submillimeter dust continuum in two ways (see da Cunha et al. 2013, for a detailed discussion). Firstly, the CMB provides an additional source of dust heating increasing the intrinsic dust temperature as shown in equation 6 (da Cunha et al. 2013):

$$T_{\text{dust}}(z) = \left((T_{\text{dust}}^{z=0})^{\beta+4} + (T_{\text{CMB}}^{z=0})^{\beta+4} [(1+z)^{\beta+4} - 1] \right)^{\frac{1}{\beta+4}} \quad (6)$$

Secondly, submillimeter observations of dust emission are always measured against the background of the CMB. At low- z $T_{\text{dust}}(z) \gg T_{\text{cmb}}(z)$, so essentially all the intrinsic flux is detected against the CMB. However, at high- z , as $T_{\text{cmb}}(z)$ approaches $T_{\text{dust}}(z)$, the fraction of submillimeter flux detected against the CMB decreases. Equation 7 (da Cunha et al. 2013) shows the fraction of the intrinsic dust emission from a galaxy measured at a given frequency, $v_{\text{obs}} = v_{\text{rest}}/(1+z)$ against the CMB background:

$$\frac{F_{v_{\text{obs}}}^{\text{obs against CMB}}}{F_{v_{\text{obs}}}^{\text{intrinsic}}} = 1 - \frac{B_v[T_{\text{CMB}}(z)]}{B_v[T_{\text{dust}}(z)]} \quad (7)$$

Assuming $T_{\text{CMB}}^{z=0} = 2.73$ K, $T_{\text{dust}}^{z=0} = 25$ K (in line with the T_{dust} adopted in Scoville et al. 2016) and $\beta = 2$ we derive




Figure 1a. Standard emcee corner plot showing the one- and two-dimensional posterior distributions for our parameters (the average observed flux density in μJy of galaxies in each redshift bin) for the redshift intervals Δz (0.25)– Δz (4.75). The density of the points and the contours correlate with the posterior probability distributions from a 1000-step run (with 500 steps for burn-in discarded) based on our sky model and SCUBA-2 maps, and employing the “delete one” jackknife technique with the map segment area corresponding to $A = 1$ deleted to take into account sample variance (Tukey 1958). The vertical red lines show the average $850\ \mu\text{m}$ flux density of galaxies for each redshift interval, with the dashed red lines showing the associated 1σ uncertainties (these values are not corrected for the influence of the CMB, see Table 1 for CMB corrected-estimates). As evidenced by this plot there are only very weak correlations between our parameters for the redshifts intervals Δz (0.25)– Δz (4.75).




Figure 1b. Standard *emcee* corner plot showing the one- and two-dimensional posterior distributions for our parameters (the average observed flux density in μJy of galaxies in each redshift bin) in the redshift intervals $\Delta z(5.25)$ – $\Delta z(9.75)$. Detailed description as in Figure 1a. As evidenced by this plot there are only very weak correlations between our parameters for the redshifts intervals $\Delta z(5.25)$ – $\Delta z(9.75)$.




Figure 1c. Standard emcee corner plot showing the two-dimensional posterior distributions for our parameters (the average observed flux density in μJy of galaxies in each redshift bin) for redshift intervals $\Delta z(0.25)$ – $\Delta z(4.75)$ and $\Delta z(5.25)$ – $\Delta z(9.75)$. Detailed description as in Figure 1a. As evidenced by this plot there are only very weak correlations between our parameters for the redshifts intervals $\Delta z(0.25)$ – $\Delta z(4.75)$ and $\Delta z(5.25)$ – $\Delta z(9.75)$.

the fraction of submillimeter flux observed against the CMB for the redshift range $0 < z < 10$ at $v_{\text{obs}}=353$ GHz ($\lambda_{\text{obs}}=850 \mu\text{m}$), including the extra heating contributed by the CMB. We apply this correction to our average observed $850 \mu\text{m}$ flux densities in all redshift bins to account for the impact of the CMB on our estimates.

4. RESULTS

4.1. Estimating molecular gas mass: RJ luminosity-to-gas mass relation

In Table 1 we present the average observed $850 \mu\text{m}$ flux densities for our galaxy sample as a function of redshift. We quote the uncertainties to 1σ and include the additional uncertainty due to sample variance. We note that at $z \gtrsim 6$ the UDS redshifts are untested. However, as our sample is binned according to the $\mathcal{P}(z)$ for each source, every galaxy effectively contributes to the flux in each redshift interval. Hence, we show the average $850 \mu\text{m}$ flux density estimates for our galaxy sample to $z = 10$.

We sum $\mathcal{P}(z)$ (which is completeness corrected) in each redshift bin, giving us the galaxy “weighting” for each redshift interval. The integral of the summed $\mathcal{P}(z)$ across all redshift intervals should be approximately equal to the total number of galaxies in our sample. We calculate this to be 154,839, which is consistent with our galaxy sample of 153,399 sources (taking into account the completeness corrections). With the summed $\mathcal{P}(z)$ and taking the area of our sample as the unmasked region of the SCUBA-2 $850 \mu\text{m}$ map (which corresponds to the UDS binary mask for “good galaxy” regions) we calculate the number density of galaxies as a function of redshift. By combining this with our average flux density (see Table 1 column 4 for CMB corrected values) we calculate the summed flux density for our galaxy sample in each redshift interval. In Figure 2 we present the number density and summed flux density for galaxies in our sample as a function of redshift. The distribution of the summed flux densities with redshift is broadly comparable to the redshift distribution found for SMGs, which peaks at $z \approx 2$ (e.g., Blain et al. 2002; Chapman et al. 2005; Simpson et al. 2014; Miettinen et al. 2017; Zavala et al. 2018), whilst the number density distribution generally declines with increasing redshift as expected. The difference in the evolution of these distributions demonstrates that our derived flux densities are not biased by the number density of galaxies in each redshift bin.

We adopt the approach of Scoville et al. (2016, hereafter S16) and, utilising our flux density measurements, estimate the average molecular gas mass for our galaxy sample in each redshift interval. The full details of this




Figure 2. Number density of galaxies in each redshift bin (blue bars) calculated from the galaxy “weighting” (the sum of the completeness corrected $\mathcal{P}(z)$ in each redshift bin) and the area of the unmasked region of the SCUBA-2 image. The black points show the summed flux density of galaxies as a function of redshift derived by combining the number density and the average flux density of galaxies (see Table 1 column 4 for average flux density values) in each redshift bin. This figure shows the contrasting distributions, with the summed flux density mirroring the redshift distribution found for SMGs with a peak at $z \approx 2$, whereas the number density generally shows a steady decline. At $z \approx 7$ there is an unexpected uptick in the number density and the summed flux density of galaxies, this is likely due to a combination of uncertainties in the photometric redshift fitting at high- z (UDS photometric redshifts are untested at $z \gtrsim 6$) and the small number statistics in these bins with the galaxy weighting being $\lesssim 100$ per redshift interval.

approach are given in S16; however, we provide a brief description here.

The long wavelength RJ tail of dust emission is nearly always optically thin ($\tau \ll 1$) and consequently this provides a direct probe of the total dust mass and hence the molecular gas mass. S16 utilise this to obtain an empirically calibrated RJ luminosity-to-gas mass ratio

$$\left(\frac{M_{\text{mol}}}{M_{\odot}} \right) = \frac{1}{\alpha_{850}} \left(\frac{L_{850, \text{rest}}}{\text{erg s}^{-1} \text{ Hz}^{-1}} \right) \text{ for } \lambda_{\text{rest}} \gtrsim 250 \mu\text{m} \quad (8)$$

with $\alpha_{850} = 6.7 \pm 1.7 \times 10^{19} \text{ erg s}^{-1} \text{ Hz}^{-1} M_{\odot}^{-1}$. The restriction $\lambda_{\text{rest}} \gtrsim 250 \mu\text{m}$ is required to ensure that at an observed wavelength of $850 \mu\text{m}$ the rest-frame emission stays on the RJ tail. S16 demonstrate that this luminosity-to-mass ratio is relatively constant for high-stellar mass ($M_{\text{stellar}} = (2-40) \times 10^{10} M_{\odot}$) normal star-forming and star-bursting galaxies, both locally and at high- z .

Table 1. Redshift intervals, galaxy “weighting” in each redshift interval, average 850 μm flux density, and average 850 μm CMB corrected flux density.

Δz	Galaxy weighting	$\langle S_{850} \rangle$ [$\mu\text{Jy dz}^{-1}$]	CMB corr. $\langle S_{850} \rangle$ [$\mu\text{Jy dz}^{-1}$]
0.25	90003	2.1 ± 2.0	2.1 ± 2.0
0.75	79917	32.9 ± 8.2	33.0 ± 8.2
1.25	64475	67.5 ± 10.3	68.0 ± 10.3
1.75	47922	146.9 ± 12.8	148.5 ± 12.9
2.25	31448	135.0 ± 17.9	137.2 ± 18.2
2.75	18368	224.9 ± 21.5	230.3 ± 22.0
3.25	13071	104.6 ± 23.3	108.3 ± 24.1
3.75	5271	236.5 ± 45.1	248.4 ± 47.4
4.25	2413	195.0 ± 52.9	209.1 ± 56.7
4.75	848	130.8 ± 109.6	144.3 ± 120.9
5.25	336	251.2 ± 109.8	288.4 ± 126.1
5.75	79	422.9 ± 362.7	511.9 ± 439.0
6.25	86	74.2 ± 95.7	96.2 ± 124.1
6.75	37	1606.2 ± 868.8	2276.3 ± 1231.3
7.25	49	60.2 ± 77.3	95.2 ± 122.1
7.75	38	106.0 ± 127.0	190.8 ± 228.6
8.25	91	63.2 ± 73.7	132.2 ± 154.1
8.75	121	29.0 ± 36.5	71.8 ± 90.3
9.25	73	155.2 ± 152.5	460.6 ± 452.5
9.75	70	289.5 ± 161.3	1039.0 ± 578.9

NOTE—Col. 1. Mid-point of each redshift interval (Δz); col. 2. the galaxy “weighting” which is the sum of the completeness corrected $\mathcal{P}(z)$ in each redshift bin. The summed $\mathcal{P}(z)$ across all redshift intervals integrates to 154,839, which is consistent with the number of galaxies in our sample (153,399) taking into account the completeness corrections; col. 3. average (stacked) 850 μm flux density as a function of redshift with the uncertainty quoted to 1σ ; col. 4. average 850 μm flux density corrected for the impact of the CMB as a function of redshift with the uncertainty quoted to 1σ .

We estimate the average rest-frame 850 μm luminosity density of galaxies as a function of redshift using the average flux densities detailed in Table 1, assuming a mass-weighted dust temperature of 25 K¹ and employ-

¹ The S16 calibration uses a mass-weighted temperature of 25 K, rather than a luminosity-weighted dust temperature (see Appendix A.2 of S16)

ing the relation from S16:

$$L_{\nu 850} = S_{\nu}[\text{Jy}] \times 1.19 \times 10^{27} \times \left(\frac{\nu(850 \mu\text{m})}{\nu_{\text{obs}}(1+z)} \right)^{3.8} \times \frac{(d_L[\text{Mpc}])^2}{1+z} \times \frac{\Gamma_{\text{RJ}}(25, \nu_{850 \mu\text{m}}, 0)}{\Gamma_{\text{RJ}}(25, \nu_{\text{obs}}, z)} [\text{erg s}^{-1} \text{Hz}^{-1}] \quad (9)$$

The Γ_{RJ} term in equation 10 corrects for departures from the RJ ν^2 dependence as the observed emission approaches the spectral energy distribution (SED) peak in the rest frame and where T_{dust} is the mass-weighted temperature characterizing the RJ dust emission

$$\Gamma_{\text{RJ}}(T_{\text{dust}}, \nu_{\text{obs}}, z) = \frac{h\nu_{\text{obs}}(1+z)/kT_{\text{dust}}}{e^{h\nu_{\text{obs}}(1+z)/kT_{\text{dust}}}-1} \quad (10)$$

We restrict our estimates of the average rest-frame 850 μm luminosity to $z \lesssim 2.5$ to ensure that rest-frame emission stays on the RJ tail. With the average rest-frame 850 μm luminosity density derived (detailed in Table 2) we use the *RJ luminosity-to-gas mass* ratio from equation 8 to estimate the average molecular gas mass as a function of redshift to $z \approx 2.5$. This calibration includes a factor of 1.36 to account for the associated mass of heavy elements (mostly Helium at 8% by number), so we correct our results by a factor $1/1.36$ (M_{mol}) to obtain M_{H_2} .

Since the summed photometric redshift probability distributions inform us about the galaxy “weighting” in each redshift interval and the UDS binary mask for “good” regions gives us the unmasked area of the SCUBA-2 850 μm map, we combine this information with the average molecular gas mass and differential co-moving volume element to estimate the co-moving volume density of molecular gas

$$\rho_{\text{H}_2} = \Omega \int_{z-\Delta z/2}^{z+\Delta z/2} N(z) \langle M_{\text{H}_2}(z) \rangle \frac{dV}{dz d\Omega} dz. \quad (11)$$

We present our values for $\rho_{\text{H}_2}(z)$ in Table 2 as a function of redshift, also giving this in terms of the critical mass density $\Omega_{\text{H}_2} = \rho_{\text{H}_2}(z)/\rho_{\text{crit}}(z)$. We use a Monte Carlo analysis to calculate the uncertainties for our values of $L_{850, \text{rest}}$, M_{H_2} and $\rho_{\text{H}_2}(z)$, first drawing random values for S_{ν} from a Gaussian distribution where the mean is the average flux density and width the uncertainty on the average flux density, and then drawing values for a mass-weighted T_{dust} from a Gaussian distribution with a mean of 25 K (corresponding to the constant T_{dust} assumed by S16), and width 3 K. Observations (e.g., Planck Collaboration 2011; S16) and simulations (e.g., Liang et al. 2018, 2019) find that a mass-weighted T_{dust} shows little variation with galaxy L_{850} or redshift

Table 2. Redshift intervals, average rest-frame $850\mu\text{m}$ luminosity, average molecular gas mass, co-moving molecular gas mass density as a function of z .

Δz	$\langle L_{850, \text{rest}} \rangle$ [$10^{27} \text{ erg s}^{-1} \text{ Hz}^{-1} \text{ dz}^{-1}$]	$\langle M_{\text{H}_2} \rangle$ [$10^7 M_{\odot} \text{ dz}^{-1}$]	ρ_{H_2} [$10^6 M_{\odot} \text{ Mpc}^{-3}$]	Ω_{H_2} [10^{-7}]
0.25	1.64 ± 0.02	1.80 ± 0.03	1.92 ± 1.80	115.66 ± 108.42
0.75	81.97 ± 3.89	89.95 ± 4.26	17.12 ± 4.39	569.51 ± 146.13
1.25	217.79 ± 18.32	239.01 ± 20.10	23.22 ± 4.09	430.70 ± 75.93
1.75	514.61 ± 64.72	564.77 ± 71.02	34.67 ± 5.32	380.54 ± 58.41
2.25	492.12 ± 85.19	540.08 ± 93.49	20.74 ± 4.60	143.48 ± 31.84

NOTE— Col. 1. Mid-point of each redshift interval (Δz); cols. 2 and 3. the average rest-frame $850\mu\text{m}$ luminosity and average molecular gas mass for galaxies in each redshift interval; col. 4. molecular gas mass density as a function of redshift and col. 5. molecular gas mass density in terms of the critical mass density. We restrict our results to $z \lesssim 2.5$ to ensure that the observed $850\mu\text{m}$ dust emission is tracing the RJ tail.

(cf. Behrens et al. 2018), and by utilising a temperature distribution with $\sigma = \pm 3\text{K}$ we recognise this minimal variance in our uncertainty calculations. We use these values to estimate $L_{850, \text{rest}}$, M_{H_2} and $\rho_{\text{H}_2}(z)$ from equations 9, 8, and 11 respectively for 1000 runs, with the uncertainty being taken as the standard deviation across these trials.

We note that the galaxy sample we use to derive the results in Tables 1 and 2 includes all galaxies in the “good galaxy” subset of the UDS DR11 catalog, regardless of the reliability of photometric redshifts for individual sources. If we apply a χ^2 cut to exclude galaxies with the least reliable redshifts (omitting galaxies with a reduced χ^2 value for the photometric redshift of > 10) and repeat the process outlined in Sections 3 and 4 above we find a less than 2% variation in our results with estimates consistent with those in Table 2 within the uncertainties.

We plot $\rho_{\text{H}_2}(z)$ in Figure 3, compared to direct CO line estimates from ASPECS (Decarli et al. 2016b; Decarli et al. 2020), COLDZ (Riechers et al. 2019) and VLASPECS (Riechers et al. 2020), as well as values derived using far-infrared and UV photometry (Berta et al. 2013). We fit a function of the same form as the star-formation rate density function presented in Madau & Dickinson (2014) to the log of our results, and derive the best fit parameters for our data using `emcee`. This yields:

$$\log_{10} \left(\frac{\rho_{\text{H}_2}}{M_{\odot} \text{ Mpc}^{-3}} \right) = (6.59 \pm 0.30) \times \frac{(1+z)^{0.38 \pm 0.16}}{1 + [(1+z)/5.57 \pm 1.69]^{1.78 \pm 0.61}} \quad (12)$$

which we plot in Figure 3. Our results show a peak $\rho_{\text{H}_2}(z)$ at $z \approx 2$ mirroring existing constraints.

4.2. Deriving molecular gas mass density from the halo mass function.

Using an alternative approach we derive $\rho_{\text{H}_2}(z)$ from first principles using the halo mass function from Murray et al. (2013) and assuming a constant halo mass range of $10^{11.5} - 10^{15} M_{\odot}$. We estimate the molecular gas mass density as a function of halo mass (for redshifts $0 \leq z \leq 7$) using the stellar-halo mass ratio from Moster et al. (2013) and the ISM-stellar mass relation from Scoville et al. (2017). The ISM-stellar mass relation is calibrated using a sample of high mass galaxies ($M_{\text{stellar}} \gtrsim 10^{10} M_{\odot}$), therefore we adopt a halo mass range for which the corresponding stellar masses are comparable with the Scoville et al. (2017) calibration sample. Integrating these estimates with respect to halo mass gives the total molecular gas density as a function of redshift, which we present in Figure 3.

4.3. Estimating molecular gas mass density using a “constant efficiency” model

We also estimate $\rho_{\text{H}_2}(z)$ from the star-formation rate density, $\rho_{M_{\star}}(z)$, assuming a corresponding volume averaged star-formation “efficiency”, $\eta(z) = \rho_{\text{H}_2}(z)/\psi_{\star}(z)$. We use the functional fit of Wilkins et al. (2019), a recalibration of the well-known Madau & Dickinson (2014) cosmic star formation history. We make the assumption that $\eta(z)$ is constant and that the *total* molecular gas mass per galaxy can be related to on-going star formation as $\xi M_{\text{H}_2} = \text{SFR}/\epsilon$ (Geach & Papadopoulos 2012). Here ξ is the ratio of dense, actively star-forming molecular gas to the total molecular reservoir with $\xi \approx 0.04$

for quiescent disks and $\xi > 0.5$ for starbursts (e.g., Papadopoulos & Geach 2012), while the factor ϵ describes the rate at which the dense molecular gas forms stars. Figure 3 shows the predicted $\rho_{\text{H}_2}(z)$ inferred from the Wilkins et al. (2019) fit, assuming a constant ‘‘average’’ $\eta(z) = 0.3 \text{ Gyr}$ corresponding to $\xi = 0.1$ and $\epsilon = 37 \text{ Gyr}^{-1}$ (e.g., Geach & Papadopoulos 2012).

This value for $\eta(z)$ is similar to the typical values of $t_{\text{dep}} (M_{\text{H}_2}/\text{SFR}) \approx 1 \text{ Gyr}$ (e.g., Tacconi et al. 2018) for main-sequence galaxies. Tacconi et al. (2018) find a relatively weak dependence of t_{dep} with redshift, $t_{\text{dep}} \propto (1+z)^{-0.57}$ to $z < 2.5$, consistent with our picture of a common mode of star formation in normal galaxies, at least out to the peak epoch. We use this relation from Tacconi et al. (2018) and the Wilkins et al. (2019) fit to derive an estimate of $\rho_{\text{H}_2}(z)$ which incorporates a weakly evolving star-formation efficiency. We present our predicted $\rho_{\text{H}_2}(z)$ in Figure 3.

5. DISCUSSION

Our results appear to be in reasonable agreement with existing empirical constraints, indicating that the epoch of molecular gas coincided with the peak epoch of star formation at $z \approx 2$. So what does this mean in terms of the evolving molecular gas budget? We might ask what is the *complete* picture of $\rho_{\text{H}_2}(z)$, or rather, what galaxies host the majority of the cosmic molecular gas budget across cosmic time? In the following discussion we interpret the overall evolution of the cosmic molecular gas density, in the context of our results, within the established framework of star formation in galaxies from the cosmic dawn to the present day.

5.1. Evolution of cosmic molecular gas mass density at $0 \leq z \leq 2.5$

The S16 *RJ luminosity-to-gas mass* ratio has been shown to provide molecular gas mass estimates accurate to within a factor of around 2 when compared with measurements made via direct CO ($J = 1 \rightarrow 0$) line observations (e.g., Scoville et al. 2017; Kaasinen et al. 2019), with variations in the dust emissivity index, temperature, and gas-to-dust ratios being accountable for the deviations. This factor of 2 accuracy is based on samples of galaxies with high stellar masses $M_{\text{stellar}} = (2 - 40) \times 10^{10} M_{\odot}$ as these galaxies are likely to have near-solar metallicity (Tremonti et al. 2004). This avoids probing low metallicity sources for which the dust-to-gas abundance ratio is likely to drop or the CO gas fraction is low (Bolatto et al. 2013).

In Figure 4 we show the marginalised stellar mass estimates for UDS galaxies (Almaini et al. in prep.) and the corresponding 95% stellar mass completeness (de-

rived using the method of Pozzetti et al. 2010, Wilkinson et al. in prep.) for the UDS catalog. As can be seen in Figure 4 our galaxy sample includes a proportion of galaxies with stellar masses lower than those used to derive the S16 *RJ luminosity-to-gas mass* ratio, with these sources being more abundant in lower redshift bins.

The dust-to-gas relation has been found to be relatively consistent for nearby galaxies with $M_{\text{stellar}} > 10^9 M_{\odot}$ (e.g., Groves et al. 2015), but drops for galaxies with lower stellar masses (hence lower metallicities). Cosmological galaxy formation simulations have shown that deviations from this relation become significant ($\gtrsim 0.5$ dex) at $L_{850} \lesssim 10^{28} \text{ erg s}^{-1} \text{ Hz}^{-1}$ in the redshift range $0 < z < 9.5$ (e.g., Privon et al. 2018). As shown in Figure 4 at $z \gtrsim 1$ the majority of our sample are likely to have $M_{\text{stellar}} > 10^9 M_{\odot}$ and Table 2 shows that the mean rest-frame 850 μm luminosity for our sample in all but the lowest redshift interval ($\Delta z = 0.25$) is $L_{850\mu\text{m}} \gtrsim 10^{28} \text{ erg s}^{-1} \text{ Hz}^{-1}$. Therefore, whilst the *RJ luminosity-to-gas mass* ratio has been calibrated on high stellar mass galaxies ($M_{\text{stellar}} = (2 - 40) \times 10^{10} M_{\odot}$), we make the assumption that applying this calibration to our sample at redshifts $z \gtrsim 0.5$ is likely to result in comparable uncertainties (i.e. a factor of 2). However, in the redshift bin $\Delta z = 0.25$ the mean rest-frame luminosity is $\langle L_{850\mu\text{m}} \rangle = 1.6 \times 10^{27} \text{ erg s}^{-1} \text{ Hz}^{-1}$, and as such our results are likely to be under-predicted by $\gtrsim 0.5$ dex (e.g., Privon et al. 2018) due to the abundance of lower mass (low metallicity) galaxies in this redshift interval.

5.2. Comparison of the evolution of molecular gas mass density to other studies in the literature

In Figure 3 we compare our results, which are revised to account for the influence of the CMB, to those from direct CO line surveys (e.g., Decarli et al. 2016b; Decarli et al. 2020; Riechers et al. 2019; Riechers et al. 2020). We limit our discussion to the results from these surveys at $z \lesssim 2.5$ as we are restricted to this redshift range due to the wavelength of our observations ($\lambda_{\text{obs}} = 850 \mu\text{m}$). Whilst the results from ASPECS/ASPECS LP (Decarli et al. 2016b; Decarli et al. 2020) and COLDz (Riechers et al. 2019; Riechers et al. 2020) were not corrected for the influence of the CMB, we note that this is not necessary at $z \lesssim 4.5$ as the effect of the CMB on measurements of molecular gas mass density from direct CO line observations is minimal ($\lesssim 15\%$, e.g., Decarli et al. 2019) and as such this does not impact on our analysis here.

We find that our results are broadly consistent with the estimates from direct CO line surveys within uncertainties and show notably good agreement with results obtained through observations of the ground state




Figure 3. Values for $\rho_{\text{H}_2}(z)$ (CMB corrected) derived using a 3D stacking method and the *RJ* luminosity-to-gas mass ratio of S16. The upper x-axis shows the rest-frame wavelength of observed $850\ \mu\text{m}$ emission for the redshift range shown, illustrating the range at which rest-frame emission traces the long-wavelength RJ tail (i.e. $\lambda_{\text{rest}} \gtrsim 250\ \mu\text{m}$) and the S16 calibration can be reliably applied. Our estimates are represented by the red points (uncertainties shown to 2σ) with the solid red line showing the best-fit function derived using `emcee` and a function of the same form as the star-formation rate density function from Madau & Dickinson (2014). We note that at $z \gtrsim 2.5$ our results should be considered speculative as the observed $850\ \mu\text{m}$ emission no longer traces the rest-frame RJ tail. Hence we only show an average of our estimates after this point. We do not include any estimates for $z \gtrsim 6$ as the UDS photometric redshifts beyond this are untested and as such highly uncertain. Alongside our values we show results from ASPECS (off white rectangles; Decarli et al. 2016b), ASPECS LP (blue rectangles; Decarli et al. 2020), COLDz (green rectangles; Riechers et al. 2019) and VLASPECS (green diamond; Riechers et al. 2020) which are derived from direct measurements of CO line emission. The grey points show values from Berta et al. (2013), estimated using deep far-infrared and UV data, and assuming either typical gas depletion times (Tacconi et al. 2013) or from IR luminosity and obscuration properties (Nordon et al. 2013). We plot our “constant efficiency” models derived using the re-calibrated star formation history from Wilkins et al. (2019) and assuming either a corresponding constant (e.g., Geach & Papadopoulos 2012) or weakly evolving (e.g., Tacconi et al. 2018) volume averaged star-formation “efficiency” to infer $\rho_{\text{H}_2}(z)$. The former is shown as the dark blue dot dash line and the latter illustrated by the dark blue circles. We also plot $\rho_{\text{H}_2}(z)$ to $z \approx 7$ derived from the halo mass function (Murray et al. 2013), assuming the stellar-halo mass ratio from Moster et al. (2013) and ISM-stellar-mass relation from Scoville et al. (2017). The dotted dark blue line corresponds to a halo mass range $10^{11.5} - 10^{15} M_{\odot}$, with the minimum stellar masses derived for this range ($\approx 10^{9.5} M_{\odot}$) being consistent with the lowest stellar masses probed in ASPECS LP (Boogaard et al. 2019).




Figure 4. Marginalised stellar mass estimates (Almaini et al. in prep.) for UDS galaxies in our sample as a density plot, with darker colours corresponding to higher number densities of galaxies. The red line corresponds to the UDS catalog 95% stellar mass completeness (derived using the method of Pozzetti et al. 2010, see Wilkinson et al. in prep.). This figure demonstrates that at high redshifts we are only sensitive to the most massive galaxies.

CO line (ASPECS LP at $\Delta z \simeq 0.25$ and COLDZ at $\Delta z \simeq 2.25$). Albeit, we caution that our results at $\Delta z \simeq 0.25$ are likely underestimated due to the abundance of low stellar mass galaxies in this redshift bin.

When compared with the ASPECS LP survey (Decarli et al. 2020) our results generally trace the lower boundaries of their estimates between $0.75 \lesssim z \lesssim 1.75$. The S16 *RJ luminosity-to-gas mass* ratio is calibrated using the ground state CO ($J = 1 \rightarrow 0$) line, whereas at $z > 0.75$ the ASPECS LP results are derived from observations of higher state excitation CO lines. Therefore, this offset could be explained by the uncertainties associated with translating higher excitation CO lines observations to ground state CO ($J = 1 \rightarrow 0$) luminosities. However, in the redshift interval $\Delta z = 1.25$, even if the extreme case of thermalized gas is assumed, our upper estimate (taking into account 1σ uncertainties) falls a factor of 1.42 below the lower boundary of the ASPECS LP survey (e.g., $38.90 \times 10^6 M_{\odot} \text{Mpc}^{-3}$, Table A3. Decarli et al. 2019). As such the uncertainties in CO line ratios do not fully account for the offset we see.

Building on previous studies Liu et al. (2019) derive the molecular gas mass density using a dataset comprised of ≈ 700 ALMA continuum detected galaxies and ≈ 1000 galaxies with CO observations (taken from the literature). To derive molecular gas masses for the continuum detected galaxies Liu et al. (2019) employ the Hughes et al. (2017) *luminosity-to-gas mass* calibration.

Liu et al. (2019) estimate a SMF (stellar mass function) integrated molecular gas mass density based on the SMF integrated to $M_{\text{stellar}} = 10^9 M_{\odot}$ and using a gas fraction function derived from their composite sample of ≈ 1700 galaxies. Their results trace the upper boundaries of the molecular gas mass density derived from the most recent blank field CO line surveys (e.g., Decarli et al. 2019; Riechers et al. 2019) and are ≈ 1 dex higher than our estimates. This offset with our estimates of molecular gas mass density could be in part due to assumptions made in Liu et al. (2019) to derive an SMF integrated molecular gas mass density (i.e. that all star-forming galaxies are on the main-sequence) or potentially differences in sample selection. The majority (≈ 800) of the CO detected sources in the Liu et al. (2019) composite sample are in the Local Universe (i.e. $z < 0.3$), so at $z > 0.3$ their dataset is dominated by ALMA continuum detected galaxies which are preferentially massive and dust-rich (and hence, using a dust-to-gas mass conversion, gas-rich). In contrast our galaxy sample is near-infrared selected and as such our selection is less likely to sample these luminous dust-rich SMGs, with previous studies finding that $\approx 20\%$ of SMGs are missed in optical/new-infrared surveys (e.g. Dudzevičiūtė et al. 2019).

Magnelli et al. (2020) use a stacking approach to measure the comoving gas mass density of a sample of 555 near-infrared selected galaxies, with galaxies split into bins of z and M_{stellar} . Their stacking method accounts for the metallicity of galaxies in these bins (inferred using the stellar mass-metallicity from Tacconi et al. 2018) and is equivalent to the S16 calibration at solar metallicity. Our results trace the lower boundaries of their estimates, but are inconsistent (within 1σ uncertainties) at $z > 1$. This discrepancy may be in part due to our method not accounting for the metallicity of low mass galaxies, resulting in an under-estimation of the molecular gas mass for galaxies with $M_{\text{stellar}} < 10^9 M_{\odot}$. However, if this was the sole reason for the difference in our results we would expect this to have more of an impact at $z < 1$ where this effect will be more prominent.

We caution that our results also rely solely on photometric redshifts, which despite the high quality 12 band photometry of the UDS catalog, cannot compete with the accuracy of redshifts derived via spectroscopic surveys. By utilising the redshift probability distributions in our 3D stacking approach we aim to provide mitigation against these uncertainties. However, whilst our estimates rely exclusively on the use of photometric redshifts, the results obtained in both ASPECS LP (Decarli et al. 2020; Magnelli et al. 2020) and Liu et al. (2019) benefit from the inclusion of sources with more reliable

spectroscopic redshifts. This may also play a part in deviations seen when we compare our estimates with these previous surveys.

5.3. *Contribution of the brightest submillimeter sources to the cosmic evolution of the molecular gas mass density*

In order to present the most complete view of the evolution of the molecular gas mass density we stack all sources in our near-infrared selected sample, including counterparts to the bright submillimeter sources in the SCUBA-2 UDS map. To test the contribution of these galaxies to our results we repeat our stacking analysis with the SCUBA2 UDS source subtracted map. As expected excluding the ~ 1000 UDS submillimeter sources reduces the average observed $850\ \mu\text{m}$ flux in our redshift intervals, which propagates to our estimate of the co-moving molecular gas mass density. At $z < 1.5$ the exclusion of the UDS submillimeter sources has a minimal impact on our estimates of $\rho_{\text{H}_2}(z)$ and these remain consistent within the 1σ uncertainties. However, at $z > 1.5$ our estimates of $\rho_{\text{H}_2}(z)$ drop by a factor of 2.05 and 2.33 in the redshift intervals $\Delta z = 1.75$ and $\Delta z = 2.25$ respectively. This coincides with the peak number density of SMGs at $z \approx 2$. This indicates that approximately 50% of the molecular gas mass density at the peak of the star formation rate density is locked in dust-rich SMGs. We note that our inferred contribution of SMGs is also likely under-estimated as we expect that approximately 20% of SMGs are undetected in our near-infrared selected sample (e.g., [Dudzevičiūtė et al. 2019](#)). Our finding is in keeping with [Zavala et al. \(2021\)](#) who find that bright SMGS ($L_{\text{IR}} > 10^{12}L_{\odot}$) dominate the obscured star formation rate density at $z \approx 2$ and also [Magnelli et al. \(2020\)](#) who find that the bulk of dust and gas in galaxies is locked in massive star-forming galaxies.

5.4. *Additional constraints on the evolution of molecular gas mass density*

We have added further valuable constraints to this picture of cosmic molecular gas evolution using two alternative approaches.

We estimate $\rho_{\text{H}_2}(z)$ from the halo mass function ([Murray et al. 2013](#)) assuming a constant halo mass range of $10^{11.5}-10^{15} M_{\odot}$, and using the stellar-halo mass ratio from [Moster et al. \(2013\)](#) and the ISM-stellar mass relation from [Scoville et al. \(2017\)](#). For the latter relation we make the assumption that all galaxies are on the star-forming main sequence (e.g., $\text{sSFR}/\text{sSFR}_{\text{MS}} = 1$). The evolution of our halo mass derived $\rho_{\text{H}_2}(z)$ (shown in [Figure 3](#)) follows a similar shape to the star-formation rate density, rising to a peak at $1 \lesssim z \lesssim 3$ and decreasing to the present day. The minimum halo mass

we assume corresponds to stellar masses of $\approx 10^{9.5} M_{\odot}$ (e.g., [Moster et al. 2013](#)), equivalent to the lowest stellar masses probed in the ASPECS LP survey ([Decarli et al. 2020](#)). Our $\rho_{\text{H}_2}(z)$ estimates show good agreement with the ASPECS/ASPECS LP surveys ([Decarli et al. 2016a; Decarli et al. 2020](#)) at $z \gtrsim 0.6$. However, as shown in [Figure 3](#) at $z \lesssim 0.6$ our estimate of $\rho_{\text{H}_2}(z)$ lies above the lowest redshift bins from the ASPECS LP survey ([Decarli et al. 2020](#)) and is ≈ 1 dex higher than our estimate of $\rho_{\text{H}_2}(z)$ derived from measurements of observed $850\ \mu\text{m}$ flux. To obtain an estimate of $\rho_{\text{H}_2}(z)$ from the halo mass function we make the assumption that all galaxies are star-forming. As such our estimate of $\rho_{\text{H}_2}(z)$ derived from the halo mass function can be seen as an upper limit $\rho_{\text{H}_2}(z)$ for the stellar mass range sampled. It follows that we see a more significant deviation between our estimate and observationally derived results at lower redshifts as the fraction of passive galaxies is higher at later epochs.

We also estimate $\rho_{\text{H}_2}(z)$ from the star-formation rate density ([Wilkins et al. 2019](#)), assuming a constant ([Geach & Papadopoulos 2012](#)) and weakly evolving ([Tacconi et al. 2018](#)) star-formation efficiency. These “constant efficiency” models predict a co-moving molecular gas mass density in good agreement with both measurements of molecular gas mass via observations of direct CO line emission ([Decarli et al. 2016b; Decarli et al. 2020; Riechers et al. 2019; Riechers et al. 2020](#)) and our results derived from measurements of the long-wavelength dust emission, out to a peak at $z \approx 2$. A simple conclusion is that the peak epoch of star formation at $z \approx 2$ is not driven by significantly more efficient (or starburst-like) star formation in galaxies, but by a higher abundance of molecular fuel in galaxies. We note that the estimate derived from weakly evolving star-formation is ≈ 1 dex higher than our results at $\Delta z \simeq 0.25$. This is likely a consequence of the latter being under estimated due to the abundance of low stellar mass galaxies in this redshift bin.

We recognise that our assumption of a “constant efficiency” model is at odds with [Scoville et al. \(2017\)](#), who argue that whilst cold molecular gas reservoirs increase with z (as $(1+z)^{1.84}$), the star-formation rate increases more rapidly (as $(1+z)^{2.9}$), indicating that the peak of star formation is a consequence of both increased molecular gas content in galaxies and higher star-formation efficiency. We also note that at $z \gtrsim 1$ early-type galaxies have been shown to be more compact for a given stellar mass than their local counterparts (e.g., [Daddi et al. 2005; Cappellari et al. 2009](#)), which taken in combination with the “Kennicutt-Schmidt” relation (a power-law relation between star-formation rate and gas sur-

face densities, Kennicutt 1998; Schmidt 1959) implies that star formation may be more efficient at $z \gtrsim 1$.

The 3D stacking approach we use derives the *average* properties for galaxies in our sample as a function of redshift, and thus we do not measure the molecular gas mass and star-formation rates for individual sources. Whilst the UDS DR11 catalog does include M_{stellar} estimates (which are evaluated at the peak maximum likelihood redshift) for individual galaxies, our 3D stacking method bins galaxies according to the discretized redshift probability distribution ($\mathcal{P}(z)$), and as such each galaxy in our sample effectively contributes to the flux in all redshift intervals. Hence, using this 3D stacking method precludes a M_{stellar} selection relative to our redshift bins. Therefore, we are not able to repeat the analysis from Scoville et al. (2017) to test their assertion of an evolving star-formation efficiency.

5.5. The epoch of molecular gas

Although we cannot quantify the contribution of higher star-formation efficiencies to the peak of star-formation rate density at $z \approx 2$, the symmetry between our “constant efficiency” models with our statistically derived $\rho_{\text{H}_2}(z)$ indicates a star formation history which is predominantly driven by an increased supply of molecular gas in galaxies, rather than a significant evolution in star-formation efficiency (consistent with the findings of Decarli et al. 2020; Magnelli et al. 2020). With this in mind we now turn to the formation of H_2 itself.

Cazaux & Spaans (2004) combine a microscopic model for the relative rates of gas-phase and dust H_2 production with a cosmological model to show the more efficient dust-phase production becomes the dominant route to H_2 formation at $z \approx 3\text{--}6$ for reasonable assumptions about the conditions of the interstellar medium of early galaxies. Therefore, there is a perfect storm for massive galaxy growth at $z \approx 2$: not only is the cosmic accretion rate at its peak, massive halos have had time to grow, galaxies have increased gas densities, and previous generations of stars in the progenitors of these systems have provided the metal enrichment that accelerates the formation of H_2 , which, as the fuel for star formation, drives galaxy growth; this could be described as the epoch of molecular gas.

5.6. Estimating the evolution of cosmic molecular gas mass density at $z \gtrsim 2.5$

S16 intentionally restrict their calibration sample to galaxies at $z \leq 3$ to ensure observed $850\ \mu\text{m}$ emission is from the rest-frame long wavelength RJ tail, where dust is optically thin and emission is dominated by the contribution of cold dust (which is well represented by a

mass-weighted $T_{\text{dust}} = 25\ \text{K}$). In Figure 3 we have shown an average of our $\rho_{\text{H}_2}(z)$ estimates at $2.5 \lesssim z \lesssim 6$ (the UDS redshifts are untested at earlier epochs as there are no UDS galaxies with spectroscopic redshifts at $z \gtrsim 6.5$), but note that at these redshifts estimates are less reliable due to large uncertainties in the RJ correction (see equation 10) as rest-frame emission approaches the peak of the SED.

In the optically thick regime (as rest-frame dust emission moves off the long-wavelength RJ tail) the rest-frame emission no longer correlates with the total dust mass of a galaxy and probes only the surface dust, which using the approach of S16 would result in under-estimation of L_{850} and hence the molecular gas mass. However, as the rest-frame emission approaches the peak of the SED we are increasingly sensitive to the dense, warm dust component, which significantly boosts the luminosity (with only a small mass fraction) and dominates the emission close to the SED peak. Consequently rest-frame dust emission at high- z is not well represented by a mass-weighted $T_{\text{dust}} = 25\ \text{K}$, which would result in an over-estimate of the dust and gas mass.

In addition to these competing effects, we are also likely to be missing a significant population of lower mass galaxies at $z > 2.5$. As shown in Figure 4 the 95% stellar mass completeness at $z \approx 2.5$ is predicted to be $\simeq 10^{9.5} M_{\odot}$, so we are simply not sensitive to the majority of low mass galaxies at the highest redshifts. In addition, although relatively rare (with number counts $N(> 3.5\ \text{mJy}) \simeq 3000\ \text{deg}^{-2}$; Geach et al. 2017) SMGs are dust-rich ($M_{\text{dust}} \sim 10^9 M_{\odot}$; e.g., da Cunha et al. 2015; Magnelli et al. 2019) and about 20% are undetected in optical/near-infrared surveys (e.g., Dudzevičiūtė et al. 2019). This non-detection of SMGs is unlikely to have a significant impact on our estimates at low z . However, at $z \gtrsim 2.5$ since we are significantly under-sampling the galaxy population and as the number of galaxies in our redshift bins fall the non-detection of dust-rich SMGs becomes more statistically significant, further contributing to an under-estimation of the molecular gas mass density at $z \gtrsim 2.5$.

The overall impact of the above is difficult to quantify. However, as shown by Figure 3 our results at $z \gtrsim 2.5$ are systematically lower than the estimates obtained via direct CO line emission, which suggests that the use of this method past $z \approx 2.5$ (when $\lambda_{\text{obs}} = 850\ \mu\text{m}$ no longer probes the rest-frame RJ tail) results in an under-estimation of the molecular gas mass density. In consequence, whilst our results are highly uncertain at $z > 2.5$ we suggest that to $z \lesssim 6$ these can be seen as providing a lower-limit to the molecular gas mass density.

6. CONCLUSIONS

We employ a 3-dimensional stacking method (Viero et al. 2013) and an empirically calibrated *RJ luminosity-to-gas mass* ratio (S16) to derive the average molecular gas mass as a function of redshift utilising a sample of $\approx 150,000$ galaxies in the UKIDSS-UDS field. By combining these techniques we are able to reduce the statistical uncertainties on the evolution of the molecular gas mass density, $\rho_{\text{H}_2}(z)$, within the redshift range $0.5 \lesssim z \lesssim 2.5$. We find that:

- $\rho_{\text{H}_2}(z)$ shows a clear evolution over cosmic time which traces that of the star-formation rate density with a peak $\approx 2 \times 10^7 \text{ M}_\odot \text{ Mpc}^{-3}$ at $z \approx 2$.

- Our results are consistent with those of blank field CO line surveys, albeit our estimates are systematically lower than those derived using observations of higher excitation CO lines. This may in part be a consequence of the line ratios used to translate higher excitation CO line luminosity to ground state CO line luminosity.

- Our results are an order of magnitude lower than those derived by Liu et al. (2019) who use the Hughes et al. (2017) *luminosity-to-gas mass* calibration to estimate molecular gas masses for the ALMA continuum detected galaxies in their sample. This difference in results may be in part due to selection effects, as their ALMA-selected sample preferentially selects dust-rich (and consequently gas-rich), sources, whereas by using a NIR selection we are likely to miss $\approx 20\%$ of these dust-rich SMGs.

- $\rho_{\text{H}_2}(z)$ can be broadly modelled by inverting the star-formation rate density (Wilkins et al. 2019) with a constant (Geach & Papadopoulos 2012) or weakly evolving (Tacconi et al. 2018) volume averaged star-formation efficiency. Our “constant efficiency” models closely align to our statistically derived $\rho_{\text{H}_2}(z)$.

- $\rho_{\text{H}_2}(z)$ can be derived from first principles from the halo mass function (Murray et al. 2013) in conjunction with stellar-halo mass (Moster et al. 2013) and ISM-stellar mass ratios (Scoville et al. 2017). To obtain this estimate we make the assumption that all galaxies are star-forming and hence this can be seen as an upper limit for $\rho_{\text{H}_2}(z)$ with respect to the stellar mass range sampled.

We have demonstrated that by applying a statistical method and the approach of Scoville et al. (2016) we can provide robust, statistically significant constraints

to the cosmological gas mass density to $z \lesssim 2.5$. Our results show an evolution that mirrors that of the star-formation rate density indicating that the peak of the star formation history is primarily driven by an increased supply of molecular gas rather than a significantly increased star-formation efficiency. We have shown that at $z \gtrsim 2.5$ we detect dust emission from high mass galaxies, even with our near-infrared selected sample. Hence, in the future there is potential for this approach to be extended to provide improved constraints at higher- z through 1 mm/3 mm wide-field surveys with facilities such as the Large Millimeter Telescope.

ACKNOWLEDGMENTS

We thank the anonymous referee for their helpful comments. TKG acknowledges support from a UK Science and Technology Facilities Council studentship. KEKC is supported by a Royal Society Leverhulme Senior Research Fellowship (SRF/R1/191013). KEKC and MF acknowledge the support from STFC (grant number ST/R000905/1). JEG is supported by a Royal Society University Research Fellowship. CCL acknowledges support from the Royal Society under grant RGF/EA/181016. MPK acknowledges support from the First TEAM grant of the Foundation for Polish Science No. POIR.04.04.00-00-5D21/18-00. MPK also acknowledges support from the Polish National Agency for Academic Exchange under Grant No. PPI/APM/2018/1/00036/U/001. The James Clerk Maxwell Telescope has historically been operated by the Joint Astronomy Centre on behalf of the Science and Technology Facilities Council of the United Kingdom, the National Research Council of Canada and the Netherlands Organisation for Scientific Research. The S2CLS map data were taken as part of Program ID MJLSC02. Additional funds for the construction of SCUBA-2 were provided by the Canada Foundation for Innovation. The UKIDSS project is defined in Lawrence et al. (2007). Further details on the UDS can be found in Almaini et al. (in prep.). UKIDSS uses the UKIRT Wide Field Camera (WFCAM; Casali et al. 2007). The photometric system is described in Hewett et al. (2006), and the calibration is described in Hodgkin et al. (2009). The pipeline processing and science archive are described in Irwin et al (in prep.) and Hambly et al. (2008).

REFERENCES

- Alberts, S., Pope, A., Brodwin, M., et al. 2014, MNRAS, 437, 437, doi: [10.1093/mnras/stt1897](https://doi.org/10.1093/mnras/stt1897)
- Aravena, M., Decarli, R., Walter, F., et al. 2016a, ApJ, 833, 71, doi: [10.3847/1538-4357/833/1/71](https://doi.org/10.3847/1538-4357/833/1/71)

- . 2016b, *ApJ*, 833, 68, doi: [10.3847/1538-4357/833/1/68](https://doi.org/10.3847/1538-4357/833/1/68)
- Ashby, M. L. N., Willner, S. P., Fazio, G. G., et al. 2013, *ApJ*, 769, 80, doi: [10.1088/0004-637X/769/1/80](https://doi.org/10.1088/0004-637X/769/1/80)
- Behrens, C., Pallottini, A., Ferrara, A., Gallerani, S., & Vallini, L. 2018, *MNRAS*, 477, 552, doi: [10.1093/mnras/sty552](https://doi.org/10.1093/mnras/sty552)
- Berta, S., Lutz, D., Nordon, R., et al. 2013, *A&A*, 555, L8, doi: [10.1051/0004-6361/201321776](https://doi.org/10.1051/0004-6361/201321776)
- Birnboim, Y., & Dekel, A. 2003, *MNRAS*, 345, 349, doi: [10.1046/j.1365-8711.2003.06955.x](https://doi.org/10.1046/j.1365-8711.2003.06955.x)
- Blain, A. W., Smail, I., Ivison, R. J., Kneib, J.-P., & Frayer, D. T. 2002, *PhR*, 369, 111, doi: [10.1016/S0370-1573\(02\)00134-5](https://doi.org/10.1016/S0370-1573(02)00134-5)
- Bolatto, A. D., Wolfire, M., & Leroy, A. K. 2013, *ARA&A*, 51, 207, doi: [10.1146/annurev-astro-082812-140944](https://doi.org/10.1146/annurev-astro-082812-140944)
- Boogaard, L. A., Decarli, R., González-López, J., et al. 2019, *ApJ*, 882, 140, doi: [10.3847/1538-4357/ab3102](https://doi.org/10.3847/1538-4357/ab3102)
- Bothwell, M. S., Smail, I., Chapman, S. C., et al. 2013, *MNRAS*, 429, 3047, doi: [10.1093/mnras/sts562](https://doi.org/10.1093/mnras/sts562)
- Bouwens, R. J., Aravena, M., Decarli, R., et al. 2016, *ApJ*, 833, 72, doi: [10.3847/1538-4357/833/1/72](https://doi.org/10.3847/1538-4357/833/1/72)
- Brammer, G. B., van Dokkum, P. G., & Coppi, P. 2008, *The Astrophysical Journal*, 686, 1503, doi: [10.1086/591786](https://doi.org/10.1086/591786)
- Cappellari, M., di Serego Alighieri, S., Cimatti, A., et al. 2009, *ApJL*, 704, L34, doi: [10.1088/0004-637X/704/1/L34](https://doi.org/10.1088/0004-637X/704/1/L34)
- Carilli, C. L., & Walter, F. 2013, *ARA&A*, 51, 105, doi: [10.1146/annurev-astro-082812-140953](https://doi.org/10.1146/annurev-astro-082812-140953)
- Carilli, C. L., Chluba, J., Decarli, R., et al. 2016, *ApJ*, 833, 73, doi: [10.3847/1538-4357/833/1/73](https://doi.org/10.3847/1538-4357/833/1/73)
- Casali, M., Adamson, A., Alves de Oliveira, C., et al. 2007, *A&A*, 467, 777, doi: [10.1051/0004-6361:20066514](https://doi.org/10.1051/0004-6361:20066514)
- Cazaux, S., & Spaans, M. 2004, *ApJ*, 611, 40, doi: [10.1086/422087](https://doi.org/10.1086/422087)
- Chabrier, G. 2003, *Publications of the Astronomical Society of the Pacific*, 115, 763, doi: [10.1086/376392](https://doi.org/10.1086/376392)
- Chapman, S. C., Blain, A. W., Smail, I., & Ivison, R. J. 2005, *ApJ*, 622, 772, doi: [10.1086/428082](https://doi.org/10.1086/428082)
- Chary, R.-R., & Pope, A. 2010, arXiv e-prints, arXiv:1003.1731. <https://arxiv.org/abs/1003.1731>
- Coppin, K. E. K., Swinbank, A. M., Neri, R., et al. 2007, *ApJ*, 665, 936, doi: [10.1086/519789](https://doi.org/10.1086/519789)
- da Cunha, E., Groves, B., Walter, F., et al. 2013, *ApJ*, 766, 13, doi: [10.1088/0004-637X/766/1/13](https://doi.org/10.1088/0004-637X/766/1/13)
- da Cunha, E., Walter, F., Smail, I. R., et al. 2015, *ApJ*, 806, 110, doi: [10.1088/0004-637X/806/1/110](https://doi.org/10.1088/0004-637X/806/1/110)
- Daddi, E., Renzini, A., Pirzkal, N., et al. 2005, *ApJ*, 626, 680, doi: [10.1086/430104](https://doi.org/10.1086/430104)
- Dahlen, T., Mobasher, B., Faber, S. M., et al. 2013, *ApJ*, 775, 93, doi: [10.1088/0004-637X/775/2/93](https://doi.org/10.1088/0004-637X/775/2/93)
- Decarli, R., Walter, F., Carilli, C., et al. 2014, *ApJ*, 782, 78, doi: [10.1088/0004-637X/782/2/78](https://doi.org/10.1088/0004-637X/782/2/78)
- Decarli, R., Walter, F., Aravena, M., et al. 2016a, *ApJ*, 833, 70, doi: [10.3847/1538-4357/833/1/70](https://doi.org/10.3847/1538-4357/833/1/70)
- . 2016b, *ApJ*, 833, 69, doi: [10.3847/1538-4357/833/1/69](https://doi.org/10.3847/1538-4357/833/1/69)
- Decarli, R., Walter, F., González-López, J., et al. 2019, arXiv e-prints, arXiv:1903.09164. <https://arxiv.org/abs/1903.09164>
- Decarli, R., Aravena, M., Boogaard, L., et al. 2020, *The Astrophysical Journal*, 902, 110, doi: [10.3847/1538-4357/abaa3b](https://doi.org/10.3847/1538-4357/abaa3b)
- Dudzevičiūtė, U., Smail, I., Swinbank, A. M., et al. 2019, arXiv e-prints, arXiv:1910.07524. <https://arxiv.org/abs/1910.07524>
- Eales, S., Smith, M. W. L., Auld, R., et al. 2012, *ApJ*, 761, 168, doi: [10.1088/0004-637X/761/2/168](https://doi.org/10.1088/0004-637X/761/2/168)
- Foreman-Mackey, D., Hogg, D. W., Lang, D., & Goodman, J. 2013, *PASP*, 125, 306, doi: [10.1086/670067](https://doi.org/10.1086/670067)
- Frayer, D. T., Ivison, R. J., Scoville, N. Z., et al. 1998, *ApJL*, 506, L7, doi: [10.1086/311639](https://doi.org/10.1086/311639)
- . 1999, *ApJL*, 514, L13, doi: [10.1086/311940](https://doi.org/10.1086/311940)
- Furusawa, H., Kosugi, G., Akiyama, M., et al. 2008, *ApJS*, 176, 1, doi: [10.1086/527321](https://doi.org/10.1086/527321)
- Geach, J. E., & Papadopoulos, P. P. 2012, *ApJ*, 757, 156, doi: [10.1088/0004-637X/757/2/156](https://doi.org/10.1088/0004-637X/757/2/156)
- Geach, J. E., Dunlop, J. S., Halpern, M., et al. 2017, *MNRAS*, 465, 1789, doi: [10.1093/mnras/stw2721](https://doi.org/10.1093/mnras/stw2721)
- Gómez-Guijarro, C., Riechers, D. A., Pavesi, R., et al. 2019, *ApJ*, 872, 117, doi: [10.3847/1538-4357/ab002a](https://doi.org/10.3847/1538-4357/ab002a)
- González-López, J., Decarli, R., Pavesi, R., et al. 2019, arXiv e-prints, arXiv:1903.09161. <https://arxiv.org/abs/1903.09161>
- Gould, R. J., & Salpeter, E. E. 1963, *ApJ*, 138, 393, doi: [10.1086/147654](https://doi.org/10.1086/147654)
- Groves, B. A., Schinnerer, E., Leroy, A., et al. 2015, *ApJ*, 799, 96, doi: [10.1088/0004-637X/799/1/96](https://doi.org/10.1088/0004-637X/799/1/96)
- Hambly, N. C., Collins, R. S., Cross, N. J. G., et al. 2008, *MNRAS*, 384, 637, doi: [10.1111/j.1365-2966.2007.12700.x](https://doi.org/10.1111/j.1365-2966.2007.12700.x)
- Hewett, P. C., Warren, S. J., Leggett, S. K., & Hodgkin, S. T. 2006, *Monthly Notices of the Royal Astronomical Society*, 367, 454, doi: [10.1111/j.1365-2966.2005.09969.x](https://doi.org/10.1111/j.1365-2966.2005.09969.x)
- Hodgkin, S. T., Irwin, M. J., Hewett, P. C., & Warren, S. J. 2009, *Monthly Notices of the Royal Astronomical Society*, 394, 675, doi: [10.1111/j.1365-2966.2008.14387.x](https://doi.org/10.1111/j.1365-2966.2008.14387.x)
- Hughes, T. M., Ibar, E., Villanueva, V., et al. 2017, *MNRAS*, 468, L103, doi: [10.1093/mnrasl/slx033](https://doi.org/10.1093/mnrasl/slx033)

- Ivison, R. J., Papadopoulos, P. P., Smail, I., et al. 2011, *MNRAS*, 412, 1913, doi: [10.1111/j.1365-2966.2010.18028.x](https://doi.org/10.1111/j.1365-2966.2010.18028.x)
- Jarvis, M. J., Bonfield, D. G., Bruce, V. A., et al. 2013, *Monthly Notices of the Royal Astronomical Society*, 428, 1281, doi: [10.1093/mnras/sts118](https://doi.org/10.1093/mnras/sts118)
- Kaasinen, M., Scoville, N., Walter, F., et al. 2019, *ApJ*, 880, 15, doi: [10.3847/1538-4357/ab253b](https://doi.org/10.3847/1538-4357/ab253b)
- Kennicutt, Robert C., J. 1998, *ApJ*, 498, 541, doi: [10.1086/305588](https://doi.org/10.1086/305588)
- Lawrence, A., Warren, S. J., Almaini, O., et al. 2007, *MNRAS*, 379, 1599, doi: [10.1111/j.1365-2966.2007.12040.x](https://doi.org/10.1111/j.1365-2966.2007.12040.x)
- Lenkić, L., Bolatto, A. D., Förster Schreiber, N. M., et al. 2019, arXiv e-prints, arXiv:1908.01791. <https://arxiv.org/abs/1908.01791>
- Liang, L., Feldmann, R., Faucher-Giguère, C.-A., et al. 2018, *MNRAS*, 478, L83, doi: [10.1093/mnrasl/sly071](https://doi.org/10.1093/mnrasl/sly071)
- Liang, L., Feldmann, R., Kereš, D., et al. 2019, *MNRAS*, 489, 1397, doi: [10.1093/mnras/stz2134](https://doi.org/10.1093/mnras/stz2134)
- Liu, D., Schinnerer, E., Groves, B., et al. 2019, arXiv e-prints, arXiv:1910.12883. <https://arxiv.org/abs/1910.12883>
- Lonsdale, C. J., Smith, H. E., Rowan-Robinson, M., et al. 2003, *PASP*, 115, 897, doi: [10.1086/376850](https://doi.org/10.1086/376850)
- Madau, P., & Dickinson, M. 2014, *ARA&A*, 52, 415, doi: [10.1146/annurev-astro-081811-125615](https://doi.org/10.1146/annurev-astro-081811-125615)
- Magdis, G. E., Daddi, E., Béthermin, M., et al. 2012, *ApJ*, 760, 6, doi: [10.1088/0004-637X/760/1/6](https://doi.org/10.1088/0004-637X/760/1/6)
- Magnelli, B., Karim, A., Staguhn, J., et al. 2019, *ApJ*, 877, 45, doi: [10.3847/1538-4357/ab1912](https://doi.org/10.3847/1538-4357/ab1912)
- Magnelli, B., Boogaard, L., Decarli, R., et al. 2020, *ApJ*, 892, 66, doi: [10.3847/1538-4357/ab7897](https://doi.org/10.3847/1538-4357/ab7897)
- Miettinen, O., Delvecchio, I., Smolčić, V., et al. 2017, *A&A*, 606, A17, doi: [10.1051/0004-6361/201730762](https://doi.org/10.1051/0004-6361/201730762)
- Millard, J. S., Eales, S. A., Smith, M. W. L., et al. 2020, *Monthly Notices of the Royal Astronomical Society*, 494, 293–315, doi: [10.1093/mnras/staa609](https://doi.org/10.1093/mnras/staa609)
- Moster, B. P., Naab, T., & White, S. D. M. 2013, *MNRAS*, 428, 3121, doi: [10.1093/mnras/sts261](https://doi.org/10.1093/mnras/sts261)
- Murray, S. G., Power, C., & Robotham, A. S. G. 2013, *Astronomy and Computing*, 3, 23, doi: [10.1016/j.ascom.2013.11.001](https://doi.org/10.1016/j.ascom.2013.11.001)
- Nordon, R., Lutz, D., Saintonge, A., et al. 2013, *ApJ*, 762, 125, doi: [10.1088/0004-637X/762/2/125](https://doi.org/10.1088/0004-637X/762/2/125)
- Oteo, I., Ivison, R. J., Dunne, L., et al. 2018, *ApJ*, 856, 72, doi: [10.3847/1538-4357/aaa1f1](https://doi.org/10.3847/1538-4357/aaa1f1)
- Papadopoulos, P. P., & Geach, J. E. 2012, *ApJ*, 757, 157, doi: [10.1088/0004-637X/757/2/157](https://doi.org/10.1088/0004-637X/757/2/157)
- Pavesi, R., Sharon, C. E., Riechers, D. A., et al. 2018, *ApJ*, 864, 49, doi: [10.3847/1538-4357/aacb79](https://doi.org/10.3847/1538-4357/aacb79)
- Planck Collaboration, Abergel, A., Ade, P. A. R., et al. 2011, *A&A*, 536, A25, doi: [10.1051/0004-6361/201116483](https://doi.org/10.1051/0004-6361/201116483)
- Planck Collaboration, Ade, P. A. R., Aghanim, N., et al. 2016, *A&A*, 594, A13, doi: [10.1051/0004-6361/201525830](https://doi.org/10.1051/0004-6361/201525830)
- Popping, G., Pillepich, A., Somerville, R. S., et al. 2019, *ApJ*, 882, 137, doi: [10.3847/1538-4357/ab30f2](https://doi.org/10.3847/1538-4357/ab30f2)
- Pozzetti, L., Bolzonella, M., Zucca, E., et al. 2010, *A&A*, 523, A13, doi: [10.1051/0004-6361/200913020](https://doi.org/10.1051/0004-6361/200913020)
- Privon, G. C., Narayanan, D., & Davé, R. 2018, *ApJ*, 867, 102, doi: [10.3847/1538-4357/aae485](https://doi.org/10.3847/1538-4357/aae485)
- Riechers, D. A., Bradford, C. M., Clements, D. L., et al. 2013, *Nature*, 496, 329, doi: [10.1038/nature12050](https://doi.org/10.1038/nature12050)
- Riechers, D. A., Pavesi, R., Sharon, C. E., et al. 2019, *ApJ*, 872, 7, doi: [10.3847/1538-4357/aafc27](https://doi.org/10.3847/1538-4357/aafc27)
- Riechers, D. A., Boogaard, L. A., Decarli, R., et al. 2020, *The Astrophysical Journal*, 896, L21, doi: [10.3847/2041-8213/ab9595](https://doi.org/10.3847/2041-8213/ab9595)
- Schmidt, M. 1959, *ApJ*, 129, 243, doi: [10.1086/146614](https://doi.org/10.1086/146614)
- Scoville, N., Aussel, H., Sheth, K., et al. 2014, *ApJ*, 783, 84, doi: [10.1088/0004-637X/783/2/84](https://doi.org/10.1088/0004-637X/783/2/84)
- Scoville, N., Sheth, K., Aussel, H., et al. 2016, *The Astrophysical Journal*, 820, 83, doi: [10.3847/0004-637x/820/2/83](https://doi.org/10.3847/0004-637x/820/2/83)
- Scoville, N., Lee, N., Bout, P. V., et al. 2017, *The Astrophysical Journal*, 837, 150, doi: [10.3847/1538-4357/aa61a0](https://doi.org/10.3847/1538-4357/aa61a0)
- Scoville, N. Z. 2013, *Evolution of star formation and gas*, ed. J. Falcón-Barroso & J. H. Knapen, 491
- Simpson, J. M., Swinbank, A. M., Smail, I., et al. 2014, *ApJ*, 788, 125, doi: [10.1088/0004-637X/788/2/125](https://doi.org/10.1088/0004-637X/788/2/125)
- Solomon, P. M., & Vanden Bout, P. A. 2005, *ARA&A*, 43, 677, doi: [10.1146/annurev.astro.43.051804.102221](https://doi.org/10.1146/annurev.astro.43.051804.102221)
- Stach, S. M., Swinbank, A. M., Smail, I., et al. 2017, *ApJ*, 849, 154, doi: [10.3847/1538-4357/aa93f6](https://doi.org/10.3847/1538-4357/aa93f6)
- Tacconi, L. J., Genzel, R., Neri, R., et al. 2010, *Nature*, 463, 781, doi: [10.1038/nature08773](https://doi.org/10.1038/nature08773)
- Tacconi, L. J., Neri, R., Genzel, R., et al. 2013, *ApJ*, 768, 74, doi: [10.1088/0004-637X/768/1/74](https://doi.org/10.1088/0004-637X/768/1/74)
- Tacconi, L. J., Genzel, R., Saintonge, A., et al. 2018, *ApJ*, 853, 179, doi: [10.3847/1538-4357/aaa4b4](https://doi.org/10.3847/1538-4357/aaa4b4)
- Thomson, A. P., Ivison, R. J., Smail, I., et al. 2012, *MNRAS*, 425, 2203, doi: [10.1111/j.1365-2966.2012.21584.x](https://doi.org/10.1111/j.1365-2966.2012.21584.x)
- Tremonti, C. A., Heckman, T. M., Kauffmann, G., et al. 2004, *ApJ*, 613, 898, doi: [10.1086/423264](https://doi.org/10.1086/423264)
- Tukey, J. W. 1958, *Ann. Math. Statist.*, 29, 614
- Viero, M. P., Monceli, L., Quadri, R. F., et al. 2013, *ApJ*, 779, 32, doi: [10.1088/0004-637X/779/1/32](https://doi.org/10.1088/0004-637X/779/1/32)

- Wakelam, V., Bron, E., Cazaux, S., et al. 2017, *Molecular Astrophysics*, 9, 1, doi: [10.1016/j.molap.2017.11.001](https://doi.org/10.1016/j.molap.2017.11.001)
- Walter, F., Decarli, R., Sargent, M., et al. 2014, *ApJ*, 782, 79, doi: [10.1088/0004-637X/782/2/79](https://doi.org/10.1088/0004-637X/782/2/79)
- Walter, F., Decarli, R., Aravena, M., et al. 2016, *ApJ*, 833, 67, doi: [10.3847/1538-4357/833/1/67](https://doi.org/10.3847/1538-4357/833/1/67)
- Wilkins, S. M., Lovell, C. C., & Stanway, E. R. 2019, *MNRAS*, 490, 5359, doi: [10.1093/mnras/stz2894](https://doi.org/10.1093/mnras/stz2894)
- Zavala, J. A., Aretxaga, I., Dunlop, J. S., et al. 2018, *MNRAS*, 475, 5585, doi: [10.1093/mnras/sty217](https://doi.org/10.1093/mnras/sty217)
- Zavala, J. A., Casey, C. M., Manning, S. M., et al. 2021, The Evolution of the IR Luminosity Function and Dust-obscured Star Formation in the Last 13 Billion Years. <https://arxiv.org/abs/2101.04734>