



Universiteit  
Leiden  
The Netherlands

## **Past and present: raw material identification approaches at Umhlatuzana rockshelter, South Africa**

Sifogeorgaki, I.; Os, B. van; Fratta, V.; Huisman, H.; Dusseldorp, G.L.

### **Citation**

Sifogeorgaki, I., Os, B. van, Fratta, V., Huisman, H., & Dusseldorp, G. L. (2021). *Past and present: raw material identification approaches at Umhlatuzana rockshelter, South Africa*. Retrieved from <https://hdl.handle.net/1887/3263866>

Version: Publisher's Version


License: [Leiden University Non-exclusive license](#)

Downloaded from: <https://hdl.handle.net/1887/3263866>

**Note:** To cite this publication please use the final published version (if applicable).

**Fig. 1** Raw material distribution according to the 1985 (Kaplan 1995) and 2018-2019 (current study, Sifogeorgaki et al. 2020) excavation reports (legend idem Fig. 4).

# Past and Present



**Fig. 2.** Map highlighting the location of Umhlatuzana rockshelter at KwaZulu-Natal, South Africa.



## Raw material identification approaches at Umhlatuzana rockshelter, South Africa

Irini Sifogeorgaki<sup>1</sup>, Bertil van Os<sup>2</sup>, Vi Fratta<sup>1</sup>, Hans Huisman<sup>2,3</sup>, Gerrit Dusseldorp<sup>1,4</sup>

<sup>1</sup>Faculty of Archaeology, Leiden University, NL; <sup>2</sup>Cultural Heritage Agency of Netherlands, NL; <sup>3</sup>University of Groningen, NL; <sup>4</sup>Palaeo-Research Institute, University of Johannesburg, SA

### RESULTS

Thin section analysis reveals **sandstone** (Fig. 5 A-B), **quartz** (Fig. 5. C), **hornfels** (Fig. 5. D-G), **chert** (Fig. 5. H-I), and **dolerite** (Fig. 5. J-K) raw materials.

There are pronounced **differences** on the raw material distribution between Kaplan and the current study (Fig. 1). **Quartzite** fragments were not detected.


The pXRF results indicate that we can **differentiate** between different types of raw materials (Fig. 4). Categorization of lithics that cannot be visually identified will be supplemented with pXRF.

### INTRODUCTION


Umhlatuzana is an important site documenting technological development over the past 70.000 years. This period is associated with the appearance of sophisticated lithic industries and 'modern behaviour' (Fig. 3). The site was first excavated by Jonathan Kaplan (1990) who suggested that the lithic assemblage of the Pleistocene deposits consisted mainly of quartz, hornfels, and quartzite (Kaplan 1990, Fig. 1). Renewed excavations were conducted by a team from Leiden during 2018 and 2019 (Reidsma et al. 2021, Sifogeorgaki et al. 2020). The raw material types and proportion of the renewed excavations are not in agreement with Kaplan's estimations. Here we present our first results on qualifying and quantifying raw material used at Umhlatuzana rockshelter.

### METHODS

We conducted petrographic analysis on 18 micromorphological thin sections and distinguished the raw material types present. We carried out pXRF analysis of the thin section raw material and of c. 100 artefacts. This revealed the elemental composition of the different types and allowed linkage of micromorphological and archaeological materials. Sample preparation methodologies are described in Reidsma et al. (2021) and Huisman et al. (2017).



**Fig. 3.** Middle Stone Age artefacts from Umhlatuzana. A. Backed segment; B, C. Unifacial points; D. Fragment of bifacial point.



**Fig. 4** pXRF correlation plots: A. Rb-Ba; B. Fe<sub>2</sub>O<sub>3</sub>-MgO; C. Sr-Rb.

### CONCLUSION

The reliability of visual classification of lithic raw materials is increased by the application of geoarchaeological techniques. This represents a necessary first step before behavioural questions on raw material selection and sourcing can be reliably addressed.

### ACKNOWLEDGEMENTS


We thank V. van den Brink, V. Madinani, F. Reidsma, J. Dekker, G. Halewijn, C. Thornhill, A. Sokela, and L. van Schalkwyk for assistance in the field. We are grateful to the KwaZulu-Natal Museum, Department of Human Sciences, G. Blundell, G. Laue, P. Madonda, M. Munzhedzi, D. Thloalele, and G. Whitelaw who supported our excavations. We gratefully acknowledge AMAFA aKwaZulu-Natali, SAHRA, B. Pawandiwa, and S. Uys for issuing the requisite permits. The research is funded by an NWO Vidi grant awarded to G. Dusseldorp.

Kaplan, J., 1990. The Umhlatuzana Rock Shelter sequence: 100 000 years of Stone Age history. *Natal Museum Journal of Humanities*, 2, 1-94.

Huisman, D. J., van der Laan, J., Davies, G. R., van Os, B., Roymans, N., Fermin, B., Karwowski, M., 2017. Purple haze: Combined geochemical and Pb-Sr isotope constraints on colourants in Celtic glass. *Journal of Archaeological Science*, 81, 59-78.

Sifogeorgaki, I., Klinkenberg, V., Esteban, I., Murungi, M., Carr, A. S., Van den Brink, V. B., Dusseldorp, G. L., 2020. New Excavations at Umhlatuzana Rockshelter, KwaZulu-Natal, South Africa: A Stratigraphic and Taphonomic Evaluation. *The African Archaeological Review*, 37(4), 551.

Reidsma, F., Sifogeorgaki, I., Dinckal, A., Huisman, H., Sier, M., van Os, B., Dusseldorp, G., 2021. Making the invisible stratigraphy visible: a grid-based, multi-proxy geoarchaeological study of Umhlatuzana rockshelter, South Africa. *Frontiers in Earth Science*.



**Fig. 5:** A-K: Micrographs of the main raw material types: A, B. Sandstone (PPL, XPL); C. Quartz (XPL); D, E, F, G. Hornfels (PPL, XPL, PPL, XPL); H, I. Chert (PPL, XPL); J, K. Dolerite (PPL, XPL)



**L-P: Textural view of the main materials by stereo microscope:** L. Sandstone; M, N. Hornfels; O, P. Quartz



Universiteit Leiden



rijksuniversiteit groningen



Cultural Heritage Agency  
Ministry of Education,  
Culture and Science

