



Universiteit
Leiden
The Netherlands

Exoplanets with ELT-METIS I: estimating the direct imaging exoplanet yield around nearby stars within 6.5 parsecs

Bowens, R.; Meyer, M.; Delacroix, C.; Absil, O.; Boekel, R. van; Quanz, S.; ... ; Pathak, P.

Citation

Bowens, R., Meyer, M., Delacroix, C., Absil, O., Boekel, R. van, Quanz, S., ... Pathak, P. (2021). Exoplanets with ELT-METIS I: estimating the direct imaging exoplanet yield around nearby stars within 6.5 parsecs. *European Planetary Science Congress*, EPSC2021-79. Retrieved from <https://hdl.handle.net/1887/3263978>

Version: Publisher's Version

License: [Creative Commons CC BY 4.0 license](https://creativecommons.org/licenses/by/4.0/)

Downloaded from: <https://hdl.handle.net/1887/3263978>

Note: To cite this publication please use the final published version (if applicable).



Exoplanets with ELT-METIS I: Estimating the Direct Imaging Exoplanet Yield Around Nearby Stars within 6.5 Parsecs

Rory Bowens¹, Michael Meyer², Christian Delacroix³, Olivier Absil³, Roy van Boekel⁴, Sascha Quanz⁵, Muskan Shinde⁶, Matthew Kenworthy⁷, Brunella Carlomagno³, Gilles Orban de Xivry³, Faustine Cantalloube⁴, and Prashant Pathak⁸

¹The University of Michigan, LSA, United States of America (rpbowens@umich.edu)

²The University of Michigan, LSA, United States of America

³STAR Institute, The University of Liege, Liege, Belgium

⁴MPIA, Heidelberg, Germany

⁵Institute for Particle Physics and Astrophysics, ETH Zurich, Zurich, Switzerland

⁶IISER Pune, Indore, India

⁷Leiden Observatory, Leiden University, Leiden, The Netherlands

⁸ESO, Garching, Germany

Direct imaging is a powerful exoplanet discovery technique that is complimentary to other techniques with great promise in the era of 30-meter class telescopes. Space-based transit surveys have revolutionized our understanding of the frequency of planets at small orbital radii around sun-like stars. The next generation of extremely large ground-based telescopes will have the angular resolution and sensitivity to directly image planets with $R < 4$ Earth radii around the very nearest stars. Here we predict yields from a direct imaging survey of a volume-limited sample of sun-like stars with the Mid-Infrared ELT Imager and Spectrograph (METIS) instrument, planned for the 39-m European Southern Observatory (ESO) Extremely Large Telescope (ELT) that is expected to be operational towards the end of the decade. Using Kepler occurrence rates, a sample of stars with spectral types A-K within 6.5 pc, and simulated contrast curves based on an advanced model of what is achievable from coronagraphic imaging with adaptive optics, we estimate the expected yield from METIS using Monte Carlo simulations. We find the METIS expected yield of small planets in the N2 band (10.10 - 12.40 μm) is 1.15 planets which is greater than similar observations in the L (3.70 - 3.95 μm) and M (4.70 - 4.90 μm) bands. We also determine a 42% chance of detecting at least one Jovian planet in the background limited regime assuming a 1-hour integration. We calculate the yield per star and estimate optimal observing revisit times to increase the yield. We also analyze this survey if performed in the northern hemisphere and find there are additional targets worth considering. Finally, we present an observing strategy in order to maximize the possible yield for limited telescope time, resulting in 1.52 expected planets in the N2 band.