Universiteit
Leiden
The Netherlands

Spurious heating of stellar motions in simulated galactic discs by dark matter halo particles

Ludlow, A.D.; Fall, S.M.; Schaye, J.; Obreschkow, D.

Citation

Ludlow, A. D., Fall, S. M., Schaye, J., & Obreschkow, D. (2021). Spurious heating of stellar motions in simulated galactic discs by dark matter halo particles. *Monthly Notices Of The Royal Astronomical Society*, 508(4), 5114-5137. doi:10.1093/mnras/stab2770

Version: Submitted Manuscript (under Review)

License: [Leiden University Non-exclusive license](#)

Downloaded from: <https://hdl.handle.net/1887/3273938>

Note: To cite this publication please use the final published version (if applicable).

Spurious heating of stellar motions in simulated galactic disks by dark matter halo particles

Aaron D. Ludlow^{1,*}, S. Michael Fall², Joop Schaye³ and Danail Obreschkow¹

¹*International Centre for Radio Astronomy Research, University of Western Australia, 35 Stirling Highway, Crawley, Western Australia, 6009, Australia*

²*Space Telescope Science Institute, 3700 San Martin Drive, Baltimore, MD 21218, USA.*

³*Leiden Observatory, Leiden University, PO Box 9513, 2300 RA Leiden, the Netherlands*

Accepted XXX. Received YYY; in original form ZZZ

ABSTRACT

We use idealized N-body simulations of equilibrium stellar disks embedded within course-grained dark matter haloes to study the effects of spurious collisional heating on disk structure and kinematics. Our results confirm prior suspicions that simulated disk galaxies are susceptible to collisional relaxation at a rate (and to a radial extent) that greatly exceeds that of their surrounding dark matter haloes. Collisional heating drives a systematic increase in both the vertical and radial velocity dispersions of disk stars, and leads to an artificial increase in the thickness and size of disks; the effects are felt at all galacto-centric radii, and are not limited to the central regions of galaxies. We demonstrate that relaxation is driven primarily by the coarse-grained nature of simulated dark matter haloes, with bulges, stellar haloes and disk stars contributing little to disk heating. The integrated effects of collisional heating are determined primarily by the mass of dark matter particles (or equivalently by the number of particles at fixed halo mass), their local density and characteristic velocity, but are largely insensitive to the masses of stellar particles. This suggests that the effects of numerical relaxation on simulated galaxies can be reduced by increasing the mass resolution of the dark matter in cosmological simulations, with limited benefits from increasing the baryonic (or stellar) mass resolution. We provide a simple empirical model that accurately captures the effects of collisional heating on the vertical and radial velocity dispersions of disk stars, as well as on their scale heights. We use the model to assess the extent to which spurious collisional relaxation may have affected the structure of simulated galaxy disks. For example, we find that dark matter haloes resolved with fewer than $\approx 10^6$ particles will collisionally heat stars near the stellar half-mass radius such that their vertical velocity dispersion increases by more than 10 per cent of the halo’s virial velocity in approximately one Hubble time.

Key words: Galaxy: kinematics and dynamics – Galaxy: evolution – Galaxy: structure – Galaxy: disc – methods: numerical

1 INTRODUCTION

Large-volume cosmological hydrodynamical simulations are now commonplace, and published runs span a considerable range in both force and mass resolution (e.g. Schaye et al. 2010; Dubois, et al 2014; Vogelsberger et al. 2014; Schaye et al. 2015; Dolag et al. 2016; McCarthy et al. 2017; Pillepich et al. 2019). By self-consistently co-evolving the dark matter (DM), gas and stellar particles, these simulations can in principle be used to inspect the morphological and kinematic properties of galaxies, the relation between them and their cosmological origins (e.g. Dubois et al. 2016; Correa et al. 2017; Clauwens et al. 2018; Trayford et al. 2019; Thob et al 2019; Du et al. 2020). Recent simulations have gained traction in both the theoretical and observational communities, in large part due to their ability to reproduce a variety of observed galaxy scaling relations, often interpreted as a testament to their credibility. The results of these studies are reassuring: in many respects, simulated galaxies re-

semble observed ones (e.g. Furlong et al. 2017; Ludlow et al. 2017; Trayford et al. 2017; Nelson et al. 2018; van de Sande, et al 2019).

The initial conditions for smoothed particle hydrodynamical simulations typically sample the linear density field with an equal number of DM and gas particles such that the particle mass ratio is $\mu \equiv m_{\text{DM}}/m_{\text{gas}} = \Omega_{\text{DM}}/\Omega_{\text{bar}} \approx 5.4$ (Ω_i is the cosmological density parameter of species i). Cosmological adaptive mesh refinement (AMR) or moving-mesh simulations also adopt DM particle masses that exceed those of baryonic fluid elements. Under the right conditions stellar particles form from these fluid elements, inheriting their masses. While not strictly necessary, this set-up is practical for a number of reasons: it simplifies the creation of initial conditions, it reduces compute times which in turn permits larger volumes to be simulated, and may appear to offer better resolution in the stellar components of galaxies¹ by shifting the computational effort away from their larger, rounder DM haloes.

* E-mail: aaron.ludlow@icrar.org

¹ Several simulations artificially increase the stellar mass resolution by creating multiple stellar particles per gas fluid element such that the resulting

Recently, Ludlow et al. (2019a, 2020) showed that long-range gravitational interactions between stellar and DM particles (a scattering processes often referred to as “collisions”, a vernacular we adopt throughout the paper) heat the stellar particles within galaxies resulting in spurious growth of their sizes. The effect is numerical and arises due to momentum exchange between the two particle species as galaxies and their DM haloes progress toward energy equipartition. This undesirable outcome affects both cosmological and idealized simulations; it can be mitigated (but not eliminated) by adopting baryonic and dark matter particles of approximately equal mass, i.e. by increasing the resolution of DM particles relative to baryonic ones such that $\mu \approx 1$. These results likely have broader implications for the kinematics and internal structure of galaxies in hydrodynamical simulations. For example, collisional heating will modify the structure of simulated disks, thickening them and making them appear more spheroidal than real disks, for which this heating effect is negligible.

A simple, order-of-magnitude calculation relates the collisional heating timescale t_{heat} of an initially-cold stellar system (i.e. the timescale over which the stellar velocity dispersion increases by order $\Delta\sigma$) to the mass (m), local density (ρ) and velocity (v) of perturbers (e.g. Chandrasekhar 1960; Lacey & Ostriker 1985):

$$t_{\text{heat}} \approx \frac{\Delta\sigma^2 v}{8\pi G^2 \rho m \ln\Lambda}, \quad (1)$$

where G and $\ln\Lambda$ are Newton’s constant and the Coulomb logarithm, respectively. Assuming a heating time of $t_{\text{heat}} \approx 10^{10}$ yr and typical properties of a Milky Way-like galaxy – a virial mass² $M_{200} \approx 10^{12} M_{\odot}$ and characteristic velocity $v \approx 220 \text{ km s}^{-1}$, a local density at $r = 8 \text{ kpc}$ of $\rho \approx 10^7 M_{\odot}/\text{kpc}^3$, and a Coulomb logarithm $\ln\Lambda \approx 10$ – eq. 1 yields a perturber mass of $m \approx 10^6 M_{\odot}$ for $\Delta\sigma \approx 0.25 \times v = 55 \text{ km s}^{-1}$, which is similar to the observed vertical velocity dispersion of disk stars. This mass is comparable to (or smaller than) the DM particle masses adopted for the majority of recent large-volume cosmological hydrodynamical simulations, suggesting that DM haloes and the galaxies that they contain may be affected by numerical heating.

A more rigorous theoretical framework for modelling the collisional heating of disk stars by DM particles in simulated galaxies was outlined by Lacey & Ostriker (1985, hereafter, LO85), although in a different context. These authors derived the (vertical, radial and azimuthal) heating rates brought about by a halo of “dark matter” composed of massive, point-mass black holes (at the time viable DM candidates). Based on observations of the vertical velocity dispersion of stars in the Milky Way’s disk, they conclude that black holes with a critical mass $\gtrsim 10^6 M_{\odot}$ cannot make a significant contribution to the DM; the integrated effects of black hole-stellar scattering would otherwise heat the disk to levels above observational constraints.

The extent to which numerical heating affects various estimates of the morphology and kinematics of simulated galaxies has not been established, yet the results above suggest that it should be given careful consideration. The purpose of this paper is to address a particular problem: the spurious collisional heating of initially thin stellar disks embedded in spherically-symmetric, coarse-grained haloes of

dark matter. We target the issue using N-body simulations of idealized equilibrium systems, and focus on how DM-star and star-star collisions affect the vertical and horizontal kinematics of disk stars, and their vertical scale-heights. Implications of numerical heating for the morphological transformation of simulated disk galaxies will be addressed in follow-up work.

Our paper proceeds as follows. In Section 2, we lay out the analytic groundwork that helps guide the interpretation of our simulations, which are described in Section 3. Section 4 details our main results: We provide an initial assessment of the possible importance of collisional disk heating in Section 4.1; we present an empirical model for disk heating in Section 4.2 and apply it to vertical and radial heating rates of stars measured in our simulations in Section 4.3; we use those results to interpret the evolution of disk scale heights in Section 4.4. We discuss the implementation of our model and its implications for current and future cosmological hydrodynamical simulations in Section 5 before providing some concluding remarks in Section 6. The main body of the paper contains the most important results; we remit a full discussion of a number of technical but important points to the Appendix.

2 PRELIMINARIES

This section provides a brief overview of the theoretical background required to interpret and model our simulation results. Throughout the paper, we adopt a cylindrical coordinate system with the z -axis normal to the disk plane (i.e. aligned with its angular momentum vector), and $R = \sqrt{r^2 - z^2}$ is the distance from the z -axis, where r is the radial coordinate in three dimensions.

2.1 Parameterizing the impact of collisional heating on the velocity dispersion profiles of disk stars

As mentioned above, an analytic description of the collisional heating of cold stellar disks was provided by LO85. Their detailed calculations implicitly assume a *cold* disk embedded in a *hot* isotropic halo, such that $\sigma_i \ll \sigma_{\text{DM}}$ at all times (σ_i is the stellar velocity dispersion in the i direction, and σ_{DM} the 1D velocity dispersion of halo particles; we adopt this nomenclature throughout the paper). Their model also assumes that the mass of DM particles is much greater than that of disk stars, i.e. $m_{\text{DM}} \gg m_{\star}$.

For perturbers of mass m_{DM} with local density ρ_{DM} , the heating rate is given by

$$\frac{\Delta\sigma_i^2}{\Delta t} = \sqrt{2} \pi \ln\Lambda f_i(\dots) \frac{G^2 \rho_{\text{DM}} m_{\text{DM}}}{\sigma_{\text{DM}}}, \quad (2)$$

where $\Delta\sigma_i^2(t) = \sigma_i^2(t) - \sigma_{i,0}^2$ ($\sigma_{i,0}$ is the *initial* velocity dispersion of stars in the i direction). The functions $f_i(\dots)$ (see LO85 for details), which are dimensionless constants of order unity for the cases discussed in this paper, are derived from epicyclic theory (Chandrasekhar 1960) assuming a Gaussian velocity distribution for halo particles; the values differ for different velocity components.

Equation 2 has clear implications. First, it suggests that the stellar velocity variance, $\sigma_i^2(t)$, grows linearly with time as a result of scattering. And second, that collisional heating depends linearly on the density and mass of perturbers, ρ_{DM} and m_{DM} , respectively, and inversely on their characteristic velocity, σ_{DM} . One of the goals of this paper is to compare the heating rates predicted by eq. 2 to the results of numerical simulations and, if necessary, to empirically calibrate its dependence on the density and characteristic velocity of DM particles.

stellar-to-DM particle mass ratio is $\mu_{\star} \equiv m_{\text{DM}}/m_{\star} > 5$ (e.g. Dubois, et al 2014; Revaz & Jablonka 2018; Dubois et al. 2020).

² Throughout the paper we quote halo masses as M_{200} , the mass contained within a sphere of radius r_{200} that encloses a mean density of $200 \times \rho_{\text{crit}}$ ($\rho_{\text{crit}} = 3H^2/8\pi G$ is the critical density for a closed universe and H is the Hubble constant); the circular velocity at r_{200} is $V_{200} = \sqrt{GM_{200}/r_{200}}$.

In the LO85 model, ρ_{DM} and σ_{DM} are *local* quantities; it is indeed sensible to assume that the heating rate at a given location in the disk should depend on the density and velocity of perturbers there. Their model, however, neglects the strong density gradients of DM particles across the radial extent of galactic disks, as well as the weak gradients in their characteristic velocities. Furthermore, gravitational scattering is *not* a local phenomenon: it is an integrated effect resulting from a small number of short-range interactions, and a large number of distant ones (Binney & Tremaine 2008; Ludlow et al. 2019b). It is therefore plausible that suitable halo-averaged values of ρ_{DM} and σ_{DM} may be more appropriate than local ones, or that simulation results will be best described by a suitable balance between the two extremes.

To accommodate a possible non-linear density and velocity dependence on the rate of disk heating, we generalize eq. 2 as

$$\frac{\Delta\sigma_i^2}{\Delta t} = \sqrt{2}\pi \ln \Lambda \frac{G^2 \rho_{\text{DM}} m_{\text{DM}}}{\sigma_{\text{DM}}} \times \left[k_i \left(\frac{\rho_{\text{DM}}}{\rho_{200}} \right)^\alpha \left(\frac{V_{200}}{\sigma_{\text{DM}}} \right)^\beta \right], \quad (3)$$

where ρ_{200} and V_{200} are characteristic, halo-averaged values of the density and velocity dispersion of dark matter. Note that we have absorbed the functions $f_i(\dots)$ into the dimensionless constant k_i ; its value will depend on the velocity component, and on the exponents α and β ; the latter two parameters determine the degree to which heating rates depend on local values of the density and velocity dispersion of dark matter.

Because the primary goal of this paper is to devise criteria for the importance of spurious DM heating in galactic disks in cosmological simulations, we have normalized the DM density and velocity dispersion in eq. 3 to the mean enclosed density and circular velocity at the halo’s virial radius, r_{200} , respectively. However, our DM heating criteria are equally valid for galactic disks in other contexts, in which different choices of reference density and velocity dispersion may be more appropriate. In these cases, the best-fit values of the free parameters will differ from those obtained in Section 4.2.

For some purposes it is convenient to express eq. 3 in dimensionless form. To do so, we define the normalized components of the stellar velocity dispersion,

$$\bar{\sigma}_i \equiv \frac{\sigma_i(r)}{\sigma_{\text{DM}}} \quad (4)$$

and a characteristic timescale

$$t_c \equiv \left(\frac{G^2 \rho_{\text{DM}} m_{\text{DM}}}{\sigma_{\text{DM}}^3} \right)^{-1}. \quad (5)$$

This allows us to rewrite eq. 3 as

$$\frac{\Delta\bar{\sigma}_i^2}{\Delta\tau} = \sqrt{2}\pi \ln \Lambda \times \left[k_i \delta_{\text{DM}}^\alpha v_{\text{DM}}^{-\beta} \right], \quad (6)$$

where $\tau \equiv t/t_c$, and we have defined $\delta_{\text{DM}} \equiv \rho_{\text{DM}}/\rho_{200}$ and $v_{\text{DM}} \equiv \sigma_{\text{DM}}/V_{200}$.

We consider α , β and $k_i \ln \Lambda$ to be free parameters, and calibrate their values using results from idealized simulations of secularly-evolving, equilibrium disks embedded within “live” DM haloes. Although $\ln \Lambda$ can, in principle, be calculated explicitly from the density and velocity structure of the halo and disk, fitting the combined term $k_i \ln \Lambda$ allows us to rectify the limitations of the binary scattering model (see discussion in Chandrasekhar 1960, p. 55-57) and to neglect density gradients across the disk when calculating the Coulomb logarithm. Note that the values of the free parameters are expected to differ for σ_z and σ_R .

2.2 The asymptotic velocity dispersion of disk stars

Stellar and DM particles in any simulated galaxy will progress toward energy equipartition. As discussed above, disk galaxies will be heated by this process, reaching an asymptotic end state (which may not occur in a Hubble time) for which $\sigma_i \approx \sqrt{\mu} \times \sigma_{\text{DM}}$, at which point the net exchange of energy between the two particle species ceases. It is useful to compare this asymptotic value to the halo’s escape speed which, according to the virial theorem, is $v_{\text{esc}} \approx \sqrt{2} \sigma_{\text{DM}}$. This suggests that energy equipartition cannot be reached for $\mu > 2$, and that the asymptotic velocity dispersion of disk stars will be

$$\sigma_{i,\text{max}} = \begin{cases} \sqrt{\mu} \times \sigma_{\text{DM}}, & \text{if } \mu \leq 2 \\ \sqrt{2} \times \sigma_{\text{DM}}, & \text{if } \mu > 2 \end{cases}. \quad (7)$$

In practice (see Section 4.3), we find that $\sigma_{i,\text{max}}(R) \approx \sigma_{\text{DM}}(R)$ provides an adequate description of our numerical simulations on timescales $\lesssim 10$ Gyr. We therefore assume $\sigma_{i,\text{max}} \approx \sigma_{\text{DM}}$ throughout the paper. In Appendix B we provide a detailed assessment of the *long-term* evolution of the vertical and radial velocity dispersion of disk stars for a sub-set of galaxy models, which validates eq. 7.

In the limiting case of an initially cold disk (i.e. $\sigma_{0,i} = 0$) the dimensionless timescale at which $\sigma_i = \sigma_{\text{DM}}$ is given by

$$\tau_{\text{vir}} \equiv \frac{t_{\text{vir}}}{t_c} = \left(\sqrt{2}\pi \ln \Lambda k_i \delta_{\text{DM}}^\alpha v_{\text{DM}}^{-\beta} \right)^{-1}. \quad (8)$$

We will see in Section 4.3 that t_{vir} can be considerably shorter than a Hubble time in poorly-resolved systems.

2.3 Collisional heating and the evolution of disk scale heights

The vertical density structure of galaxy disks, $\rho_*(z)$, is determined by the balance between the kinetic energy stored in vertical stellar motions and the combined vertical compressive forces of the outlying DM halo and disk (see Benítez-Llambay et al. 2018, for a recent discussion in the context of gaseous equilibrium disks). For a vertically-isothermal disk (i.e. σ_z independent of z), the structure is described by the hydrostatic equilibrium equation:

$$\frac{\sigma_z^2}{\rho_*} \frac{\partial \rho_*}{\partial z} = - \frac{\partial}{\partial z} (\Phi_{\text{DM}} + \Phi_*), \quad (9)$$

where Φ_{DM} and Φ_* are the gravitational potential due to DM and stars, respectively.

For an equilibrium disk with stellar surface density $\Sigma_*(R)$ embedded within a DM halo with circular velocity profile $V_c(r) = \sqrt{GM/r}$, eq. 9 admits analytic solutions in limiting cases where the gravitational potential is dominated by either the DM halo (the “non-self-gravitating” case, NSG) or by the disk (the “self-gravitating”, SG). In the thin disk limit ($z/R \ll 1$), the SG disk has a characteristic scale-height given by

$$h_{\text{SG}} = \frac{\sigma_z^2}{\pi G \Sigma_*(R)}; \quad (10)$$

for the NSG case,

$$h_{\text{NSG}} = \frac{\sqrt{2} \sigma_z}{V_c(R)} R, \quad (11)$$

where $V_c(R)$ is the circular velocity due to DM within a sphere of radius R . To meaningfully compare the vertical scale heights in the SG and NSG cases – and to facilitate a comparison with our simulation results – we will recast these characteristic heights into their corresponding half-mass scale-heights, denoted z_{SG} and z_{NSG} for the SG and NSG cases, respectively (the conversion factors are given

by $z_{\text{SG}} \approx 0.55 h_{\text{SG}}$ and $z_{\text{NSG}} \approx 0.47 h_{\text{NSG}}$, respectively; see Benítez-Llambay et al. 2018, for details).

As mentioned above, the characteristic timescale τ_{max} on which the components of the disk velocity dispersion grow to of order σ_{DM} can be significantly shorter than the Hubble time in poorly-resolved systems. In such instances, the half-mass scale-heights of disks can grow to of order their scale-radii R_d , and we must abandon the thin-disk approximation used to arrive at eqs. 10 and 11 (see Section 4.4). The structure of these “thick” disks can be calculated in the NSG limit by solving the hydrostatic equilibrium equation

$$\frac{\sigma_z^2}{\rho_*} \frac{\partial \rho_*}{\partial z} = - \frac{V_c^2(r)}{r} \left(\frac{z}{r} \right), \quad (12)$$

where $r^2 = R^2 - z^2$ has an explicit z -dependence (neglected in the thin disk limit, for which $r \rightarrow R$), and $V_c(r)$ is the circular velocity due to dark matter. We discuss the vertical structure of thick NSG disks in more detail in Section 4.4.

3 SIMULATIONS AND ANALYSIS

3.1 Initial conditions

Initial conditions (ICs) for all of our simulations were created using GALICS (Yurin & Springel 2014) and are solutions of the collisionless Boltzmann equation. In the absence of diffusion, scattering, or genuine disk instabilities, these collisionless structures should remain stable for many dynamical times. Axially-symmetric models that satisfy these criteria can be used to assess the heating rates of disk stars due to collisions with massive DM particles.

We model the galaxy/halo pair as a Hernquist (1990) sphere in equilibrium with a thin, rotationally-supported stellar disk. Specifically, for the DM halo we adopt a radial density distribution given by

$$\rho_{\text{DM}}(r) = \frac{M_{\text{DM}}}{2\pi} \frac{a}{r(r+a)^3}, \quad (13)$$

where a is the halo’s scale radius and M_{DM} its total mass. For an isotropic DM velocity distribution, the (one dimensional) velocity dispersion profile of the halo is given by

$$\sigma_{\text{DM}}(r) = \frac{GM_{\text{DM}}}{12a} \left\{ \frac{12r(r+a)^3}{a^4} \ln \left(1 + \frac{a}{r} \right) - \frac{r}{r+a} \left[25 + 52 \frac{r}{a} + 42 \left(\frac{r}{a} \right)^2 + 12 \left(\frac{r}{a} \right)^3 \right] \right\}. \quad (14)$$

Equation 13 scales as $\rho \propto r^{-1}$ in the inner regions ($r \ll a$) and as $\rho \propto r^{-4}$ at large radii ($r \gg a$). Over the radial extent of the stellar disk it resembles the Navarro-Frenk-White profile (Navarro et al. 1996, hereafter, NFW) that is often used to parameterize the density profiles of CDM haloes, but is benefited by its finite mass (for NFW, $\rho \propto r^{-1}$ at small radii but as $\rho \propto r^{-3}$ at large r , resulting in a divergent mass profile). For that reason, we will characterise the structural properties of haloes in terms of the circular virial velocity, V_{200} , and concentration, c , of an NFW halo whose density distribution matches that of eq. 13 in the inner regions (see Springel et al. 2005, for details). In this case, $M_{\text{DM}} = (1 - f_*) M_{200}$, where f_* is the stellar mass fraction, and $a = (r_{200}/c) \sqrt{2f(c)}$, where $f(c) = \ln(1+c) - c/(1+c)$.

The radial and vertical structure of the stellar disk are initially described by

$$\rho_*(R, z) = \frac{M_*}{4\pi z_d R_d^2} \text{sech}^2 \left(\frac{z}{z_d} \right) \exp \left(- \frac{R}{R_d} \right), \quad (15)$$

where R_d and z_d are the scale radius and height of the disk, respectively. Most models considered in this paper adopt a radially-constant initial disk scale height (i.e. constant z_d) with $z_d/R_d \approx 0.05$. The scale radius, R_d , is calculated assuming that the specific angular momentum of the disk, j_d , is a fraction $f_j \equiv j_d/j_{\text{DM}}$ of the halo’s specific angular momentum (where $j_k \equiv J_k/M_k$ is the specific angular momentum of component k), which we prescribe by means of the traditional spin parameter, λ_{DM} . We will often describe the disk scale-radius in terms of its half-mass radius, $R_{1/2}$, or analogous cylindrical radii, R_f , enclosing a fraction f of all disk stars. Similarly, $z_{1/2}$ is the vertical half-mass scale height of the disk.

As discussed above, analytic estimates of the collisional heating rates of stellar disks depend only on properties of the DM halo: the density of dark matter, the mass of the DM particles, and their characteristic velocities. To test these expectations, we have carried out a fiducial suite of simulations that vary each parameter individually. To do so, we initially hold a number of the structural properties of the halo and disk constant: we adopt $c = 10$ and $\lambda_{\text{DM}} = 0.03$ for the concentration and spin parameter of the halo, respectively; $f_* = 0.01$ is the disk mass fraction³ (relative to the total mass of the system; see Appendix A for a detailed discussion of this choice), and $f_j = 1$, i.e. the disk and halo have the same specific angular momentum.

We generate a series of models with $V_{200} = 50, 100, 200$, and 400 km s^{-1} , and for each systematically vary the DM particle mass by fixed factors of $\Delta \log m_{\text{DM}} = 0.5$, ranging from $m_{\text{DM}} = 10^4 M_\odot$ to $10^9 M_\odot$. For any particular V_{200} we simulate a subset of m_{DM} values such that the *least*-resolved haloes have $N_{\text{DM}} \approx 10^4$ particles, and the *best*-resolved have $N_{\text{DM}} \approx \text{a few} \times 10^6$. Note that for these values of c , λ_{DM} , f_* and f_j , the ratios $R_d/a = 0.12$ and $R_d/r_{200} = 0.02$ (which agree well with observational data; see, e.g. Kravtsov 2013; Huang et al. 2017) are independent of V_{200} and the local DM density is constant at all R_f at fixed f . We exploit this self-similarity when testing the dependence of the collisional heating rate on the characteristic velocity and local density of halo particles.

Although the dependence of heating rates on ρ_{DM} can be assessed by comparing those measured at different disk radii R_f , we also carried out an additional subset of runs that varied the halo’s concentration parameter, carefully adjusting the angular momentum of disk stars so that the *stellar* mass distribution remained fixed (each of these used $V_{200} = 200 \text{ km s}^{-1}$).

For each V_{200} and m_{DM} , disks are sampled with N_* stellar particles of equal mass m_* such that $\mu \equiv m_{\text{DM}}/m_* = 5$, which we take as our default value. For a Planck cosmology, this is close the DM-to-stellar particle mass ratio used for cosmological simulations that adopt equal numbers of baryonic and DM particles (i.e. $\mu_{\text{sim}} \equiv m_{\text{DM}}/m_* = \Omega_{\text{DM}}/\Omega_{\text{bar}} \approx 5.4$).

The accumulation of DM-star scattering events – if the stellar and DM particle masses are unequal, as is the case for our fiducial runs – may eventually lead to a spatial segregation of the two components, with the heavier species becoming increasingly concentrated and the lighter species more extended. This “mass segregation” is one way the system tends toward energy equipartition, and has a measurable impact on the structural evolution of galaxies in both cosmological

³ We acknowledge that this value is likely too low for Milky Way-like galaxies, perhaps considerably (Posti & Fall 2021). This choice, however, simplifies our analysis for two important reasons: 1) it ensures that the disks are not massive enough to modify the distribution of dark matter during the simulation; and 2) results in disks are Toomre stable and do not develop unwanted instabilities that may contribute to disk heating (see Appendix A for details).

Table 1. Main properties of disks and haloes in our suite of simulations. The first five rows list properties of our fiducial set of runs. The first column lists V_{200} , the circular velocity at the virial radius, r_{200} , of a Navarro-Frenk-White halo whose density profile matches our adopted Hernquist density distribution in the central regions; c is the concentration of the NFW halo; a/r_{200} is the Hernquist scale radius in units of r_{200} . The disk-to-halo specific angular momentum ratio is f_j ; the characteristic radius and scale-height of the disk are respectively R_d and z_d (see eq. 15), which have been normalized by a ; $\sigma_{z,0}(R_d)$ and $\sigma_{R,0}(R_d)$ are the initial velocity dispersion of disk stars at R_d in the vertical and radial directions, respectively (expressed in units of V_{200}). N_{DM} is the number of DM particles and μ the ratio of the DM-to-stellar particle mass. The gravitational softening lengths are expressed in units of the disk scale height, i.e. ε/z_d .

V_{200} [km s ⁻¹]	c	f_j	a/r_{200}	R_d/a	z_d/a [10 ⁻³]	$\sigma_{z,0}(R_d)$ [V_{200}]	$\sigma_{R,0}(R_d)$ [V_{200}]	N_{DM} [10 ⁵]	μ [m_{DM}/m_*]	ε/z_d
50	10	1.0	0.173	0.120	6.0	0.0686	0.0687	0.091, 0.29, 0.91, 2.9, 9.1	1, 5, 25	1
100	10	1.0	0.173	0.120	6.0	0.0716	0.0716	0.23, 0.73, 2.30, 7.28, 23.0	1, 5, 25	1
200	10	1.0	0.173	0.120	6.0	0.0724	0.0724	0.18, 0.58, 1.84, 5.82, 18.4	1, 5, 25	1
300	10	1.0	0.173	0.120	6.0	0.0728	0.0732	0.20, 0.62, 1.96, 6.21, 19.6	1, 5, 25	1
400	10	1.0	0.173	0.120	6.0	0.0732	0.0731	0.15, 0.47, 1.47, 4.66, 14.7	1, 5, 25	1
200	7	0.81	0.222	0.093	4.7	0.0708	0.0692	0.18, 1.84	5	1
200	15	1.03	0.128	0.162	8.1	0.0777	0.0784	0.18, 1.84	5	1
200	10	1.0	0.173	0.120	12.0	0.111	0.110	0.18, 1.84	5	1
200	10	1.0	0.173	0.120	24.0	0.179	0.173	0.18, 1.84	5	1
200	10	1.0	0.173	0.120	6.0	0.0724	0.0724	0.18, 1.84	5	0.25, 0.5, 2, 4

simulations and in idealised numerical experiments (see Binney & Knebe 2002; Ludlow et al. 2019a, for a discussion). For that reason, we have repeated our fiducial runs for $\mu = 1$ (which suppresses mass segregation but not energy equipartition) and $\mu = 25$.

Finally, we stress that all of our disks are bar-mode and Toomre-stable, which precludes heating due to dynamical instabilities. A summary of pertinent aspects of our runs is provided in Table 1.

3.2 The simulation code

All runs were carried out for a total of $t = 9.6$ Gyr. Particle orbits were integrated using the N-body/SPH code GADGET2 (Springel 2005) using a default value of the integration accuracy parameter `ErrToIntAcc` = 0.01 (several runs were repeated using `ErrToIntAcc` = 0.0025, which yields similar results to those presented below). For all models, we adopt fixed (Plummer equivalent) softening lengths for both DM and stellar particles that initially resolve the disk scale heights, i.e. $\varepsilon/z_d = 1$ (note that this also preserves the self-similarity of our runs, since it ensures that ε scales self-consistently with the characteristic size of the halo and disk). In Appendix D we show that our numerical results are largely insensitive to gravitational softening provided $\varepsilon \lesssim z_d$; larger values, for which gravitational forces are not properly modelled across the disk, suppress collisional heating, but do not eliminate it (see also Ludlow et al. 2019b, 2020).

For most runs, we output snapshots every $\Delta t = 96$ Myr (corresponding to a total of 100 snapshots), but for those in which DM haloes are resolved with fewer than 10^5 particles we adopt a higher cadence, $\Delta t = 24$ Myr (400 snapshots).

3.3 Analysis

We focus our analysis on the evolution of the vertical and radial velocity dispersions of stellar particles, and on their half-mass scale height. In practice, we measure these quantities in cylindrical annuli whose midpoints enclose specific fractions of the total *initial* stellar mass of the galaxy. We typically adopt $R_{1/4}$, $R_{1/2}$ and $R_{3/4}$ enclosing 1/4, 1/2 and 3/4 of the disk stars, respectively; we use a logarithmic radial bin of width $\Delta \log R = 0.2$. The local DM density at each of these radii is independent of V_{200} for our fiducial models and, as explained in Appendix A2, remains unaffected by collisional

relaxation for the duration of the simulations. Heating rates are defined after a time interval Δt by $\Delta \sigma_i^2 / \Delta t$, where $\Delta \sigma_i^2 = \sigma_i^2(t) - \sigma_i^2(t_{\text{init}})$ and $\Delta t = t - t_{\text{init}}$; we adopt $t_{\text{init}} = 0$ in all that follows (i.e. we measure the *average* heating rates since the initial time $t = t_{\text{init}} = 0$). The vertical half-mass height, $z_{1/2}$, is defined as the median value of $|z|$ within cylindrical shells at R_f .

For aesthetic purposes, results presented in several plots (Figures 4, 6 and 7) have been smoothed using a Savitzky-Golay filter.

4 RESULTS

4.1 A glimpse at the importance of collisional disk heating

The left panels of Figure 1 compare the vertical (top row) and radial (middle row) dimensionless velocity dispersion profiles of stellar particles after $\Delta t = 5$ Gyr for haloes of virial velocity $V_{200} = 200$ km s⁻¹ (and $c = 10$). Different colours show results for the range of DM particle masses spanning $10^6 M_\odot$ to $10^8 M_\odot$ (N_{DM} indicates the corresponding number of DM particles), and for three values of $\mu \equiv m_{DM}/m_*$ (different line styles). Vertical dotted lines correspond to the *initial* half-stellar mass radius $R_{1/2}$; for comparison, outsized circles mark the *measured* $R_{1/2}$ for the subset of simulations with $\mu = 5$ (similar results, not shown for clarity, are obtained for the other values of μ). The grey dotted lines show the initial profiles for the highest-resolution run (i.e. $m_{DM} = 10^6 M_\odot$ and $\mu = 25$). Profiles are plotted down to the radius that encloses at least 10 stellar particles.

As anticipated from eq. 2, the velocity dispersion profiles have increased at a rate proportional to the DM particle mass. For the highest mass resolution run (black lines; $m_{DM} = 10^6 M_\odot$, $N_{DM} = 1.8 \times 10^6$) heating rates are relatively low, but still substantial. Take the $\mu = 5$ runs for example. At the galaxy's half-stellar mass radius (vertical dashed line), the vertical (radial) velocity dispersion of stars has increased by a factor of ≈ 1.5 (2.0) after $\Delta t = 5$ Gyr. For increasing particle masses the situation becomes progressively worse: for $m_{DM} = 10^{7.5} M_\odot$ (salmon lines; equivalent to $N_{DM} = 5.8 \times 10^4$) the corresponding factor is ≈ 5.3 (6.8), and for $m_{DM} = 10^8 M_\odot$ (blue lines; $N_{DM} = 1.8 \times 10^4$) it is ≈ 7.0 (7.7).

Another important point can be inferred from Figure 1: the velocity dispersion of stellar particles does not significantly exceed the local (1-dimensional) velocity dispersion of the DM halo (plotted

Figure 1. *Left-hand panels:* Radial profiles of the vertical (top) and radial (middle) dimensionless velocity dispersions (i.e. normalized to V_{200}) of stellar particles for our fiducial stellar disk galaxy/DM halo models ($V_{200} = 200 \text{ km s}^{-1}$, $c = 10$) after $\Delta t = 5$ Gyr; the bottom left panel shows the radial dependence of the vertical half-mass scale height of the stellar disk. The initial profiles for the highest-resolution run ($m_{\text{DM}} = 10^6 M_{\odot}$ and $\mu = 25$) are shown using a grey dotted line in each panel on the left. In all cases, profiles are plotted to the innermost radius that encloses at least 10 stellar particles. Vertical dotted lines in panels on the left mark the initial half-mass radius $R_{1/2}$; for comparison, the *measured* values are shown using outsized circles (for $\mu = 5$). *Right-hand panels:* Evolution of the vertical (top) and radial (middle) velocity dispersion and the half-mass scale height, $z_{1/2}$ (bottom), measured at the initial $R_{1/2}$. In all panels, different colours correspond to different DM particle masses and different line-styles to different values of $\mu \equiv m_{\text{DM}}/m_*$. Note that the collisional heating of disk stars by DM particles is evident at *all* radii, even those that well-exceed the convergence radius of either DM or stars (not shown; see Appendix A2). Note also that the velocity dispersion of stars rarely, if ever, exceeds the local 1-dimensional velocity dispersion of the surrounding DM halo (thick, solid black line in the upper-left and middle-left panels, eq. 14).

using a thick solid black line, i.e. eq. 14) at any resolved radius. This is not unexpected: as discussed in Section 2.2, an equilibrium distribution of DM particles is cannot “heat” a cold stellar component to the extent that locally $\sigma_i > \sqrt{2} \sigma_{\text{DM}}$. Indeed, one may expect the collisional heating rate to drop significantly when σ_i approaches this value: any further injection of kinetic energy will unbind stellar particles from the halo’s potential (see Appendix B for a detailed discussion). Note that eq. 2 is valid provided $\sigma_i \ll \sigma_{\text{DM}}$.

The collisional heating of disk particles has additional consequences for the evolution of the vertical scale-height of thin disks. This is shown in the lower-left panel of Figure 1, where we plot the radial dependence of the disk half-mass scale-height, $z_{1/2}(R)$. All disks start with a uniform vertical scale height of $z_h = 0.05 R_d$ ($\approx 113 \text{ pc}$ for the runs shown here; thin, dotted grey line), but become progressively thicker with time. After $\Delta t = 5$ Gyr, $z_{1/2}$ measured at $R_{1/2}$ increases to $\approx 216 \text{ pc}$ for $m_{\text{DM}} = 10^6 M_{\odot}$ ($N_{\text{DM}} = 1.8 \times 10^6$)

but, as with σ_i , the relative increase is a strong function of N_{DM} . For example, $z_{1/2}(R_{1/2}) \approx 1.6 \text{ kpc}$ for $m_{\text{DM}} = 10^{7.5} M_{\odot}$ ($N_{\text{DM}} = 5.8 \times 10^4$), and $\approx 3.0 \text{ kpc}$ for $m_{\text{DM}} = 10^8 M_{\odot}$ ($N_{\text{DM}} = 1.8 \times 10^4$) after the same time interval (values are again quoted for $\mu = 5$).

Note also that the increase in the velocity dispersions and scale height of stellar particles depends only weakly on μ , at least over the timescale plotted in Figure 1 (but see Appendix B). This suggests that the rate of relaxation of the disk is driven primarily by incoherent potential fluctuations brought about by shot noise in the DM particle distribution, which lead to orbital deflections that also depend only weakly on μ . This is because a 2-body interaction between a DM and stellar particle will deflect latter’s orbit by the same angle provided $\mu \gg 1$; the deflection angle is at most a factor of 2 larger for $\mu = 1$ (Henon 1973), the maximum value we consider. Indeed, the same variations in potential cause the DM halo to relax, albeit at a slower rate than that of the disk (see Appendix A2). The

Figure 2. The evolution of the vertical (upper panels) and radial (lower panels) velocity dispersion of disk stars in our fiducial models (see section 3.1 for details). In all panels we plot the dimensionless velocity dispersion σ_i^2/V_{200}^2 versus $\tau = t/t_c$ (see eq. 5) measured at the radii $R_{1/4}$ (left), $R_{1/2}$ (middle) and $R_{3/4}$ (right) that enclose one quarter, one half and three quarters of the *initial* stellar mass, respectively. Different colour points correspond to different V_{200} . At each radius, the evolution of σ_i^2 can be approximated by a single power-law (solid black lines) provided $\sigma_i \ll \sigma_{DM}$ (σ_{DM} at each R_f is indicated using a horizontal grey line in each panel). At $\tau \gtrsim 0.01$, departures from power-law growth occur when σ_i begins to approach the local 1-dimensional DM velocity dispersion, a behaviour that can be approximated by eq. 16 (black dashed lines).

motions of stars may also be scattered by any coherent density or potential fluctuation (resulting from, e.g., molecular clouds or globular clusters, or those excited by disk instabilities), which also result in deflected stellar particle motions that are independent of their mass, although we have deliberately suppressed these in our simulations.

In addition to these potential fluctuations – which affect *all* particles – mass segregation leads to a slow divergence in the average energies of the heavy (DM) and light (stellar) particle species. Since energy equipartition demands $m_{DM} \sigma_{DM}^2 \approx m_* \sigma_*^2$, we can eliminate the mass segregation effect by setting $\mu = m_{DM}/m_* = 1$, but collisions between particles in a cold stellar component and a comparatively hot DM halo will nevertheless result in $\sigma_* \rightarrow \sigma_{DM}$. The latter is the dominant effect in our simulations.

It is worth stressing that collisional heating affects the scale-height and velocity dispersion of disk stars at *all* radii. Unlike the mass profiles of DM haloes of differing numerical resolution, which have well-defined radii beyond which convergence is achieved (e.g. Power et al. 2003; Ludlow et al. 2019b), *the kinematics and vertical structure of disks do not converge at any radius*.

The right-hand panels of Figure 1 show the time evolution of the vertical (top) and radial (middle) velocity dispersion, and the half-mass scale-height (bottom) measured at the initial value of $R_{1/2}$. Different lines-styles and colours have the same meaning as those used in the corresponding panels on the left.

4.2 An empirical model for collisional disk heating

Figure 2 plots the normalized (i.e. to V_{200}) velocity dispersions, $\Delta\sigma_z^2(\tau)$ and $\Delta\sigma_R^2(\tau)$ (top and bottom panels, respectively), as a func-

tion of the dimensionless time variable, $\tau = t/t_c$ (see eq. 5), for our fiducial models. Results are shown for three galacto-centric radii: $R_{1/4}$, $R_{1/2}$ and $R_{3/4}$ (left to right, respectively). We use different symbols for different V_{200} (for clarity, results are limited to those obtained for $\mu = 5$); the colour coding indicates m_{DM} , and is the same as that adopted in Figure 1. When expressed in scaled units, all runs exhibit a remarkable self-similarity, regardless of V_{200} or m_{DM} . Note also that, provided $\sigma_i \ll \sigma_{DM}$ (σ_{DM} at R_f is indicated using a horizontal grey line in each panel), $\Delta\sigma_i^2(\tau)$ exhibits an approximate power-law dependence on time, i.e. $\Delta\sigma_i^2 \propto \tau$ (the solid black lines show the best-fit power-laws obtained as described below).

However, a comparison of the results measured at different radii suggests that the local heating rate depends on the local density or velocity dispersion of DM, rather than on a single halo-averaged value (i.e. the *normalization* of $\Delta\sigma_i^2$ depends on the radius at which it is measured). This is indeed expected from the LO85 model, in which heating rates depend linearly on the local density of perturbers, and inversely on their characteristic velocities, both of which vary smoothly with radius. In practice, we determine the dependence on ρ_{DM} and σ_{DM} empirically, by fitting eq. 6 to the results of our fiducial runs. Specifically, we determine the values of α , β and $k_i \ln \Lambda$ that minimize the variance in $\Delta\sigma_i^2/\Delta\tau$ given by eq. 6 (recall that $k_i \ln \Lambda$ is a constant for each direction i). When doing so, we combine the velocity dispersions measured at $R_{1/4}$, $R_{1/2}$ and $R_{3/4}$ in all of our fiducial runs and in *all* simulation outputs for which $\sigma_i^2 < 0.2 \times \sigma_{DM}^2$ is satisfied (recall that we expect departures from eq. 6 unless $\sigma_i \ll \sigma_{DM}$). Note that the fitting is carried out separately for the vertical and radial velocity components.

Figure 3 plots the standard deviation in $k_i \ln \Lambda$ as a function of

Table 2. Best-fit parameters for eq. 6 and 16. The rms deviation from the best-fit model is quoted as $\sigma_{k_i \ln \Lambda}$

Component	$\sigma_{k_i \ln \Lambda}$	α	β	$k_i \ln \Lambda$
σ_z	0.082	-0.356	0.0	18.80
σ_R	0.132	-0.331	0.0	35.97

α for $\beta = 0$ (solid lines; the value anticipated by LO) and $\beta = 1$ (dashed lines). Note that the minimum variance is largely independent of β (we have verified this result for the range $0 \leq \beta \leq 3$), but depends strongly on α . The best-fit values are $\alpha_z = -0.356$ for σ_z and $\alpha_R = -0.331$ for σ_R ; the corresponding best-fit values of $k_i \ln \Lambda$ are 18.80 and 35.97 for σ_z and σ_R , respectively. This suggests that, at fixed DM particle mass, collisional heating depends sensitively on the local density of DM but not on its velocity dispersion. This is likely because the density of DM exhibits strong radial gradients across the extent of the disk, whereas its velocity dispersion does not.

The solid black lines plotted in Figure 2 corresponds to eq. 6 plotted using these best-fit values of α and $k_i \ln \Lambda$, and $\beta = 0$. For comparison, the dotted lines in the left- and right-most panels show extrapolations of the $R = R_{1/2}$ result (middle panels) assuming $\alpha = 0$, i.e. the value anticipated by LO85. Although $\alpha = 0$ describes our numerical simulations rather well, the data prefers a somewhat weaker dependence on the local density of DM.

Note too that our simple empirical model works well provided $\sigma_i \ll \sigma_{DM}$. As $\sigma_i \rightarrow \sigma_{DM}$, however, a systematic departure from this relation is evident in Figure 2. This is seen most clearly for the two largest values of m_{DM} (i.e. $10^{7.5} M_\odot$, red, and $10^8 M_\odot$, blue), for which heating rates are seen to saturate, approaching an asymptotic constant $\sigma_i \approx \sigma_{DM}$ (horizontal grey lines). This behaviour can be captured quite accurately by the formula

$$\sigma_i^2 = \sigma_{DM}^2 \left[1 - \exp\left(-\frac{(t+t_0)}{t_{vir}}\right) \right], \quad (16)$$

where t_{vir} is given by eq. 8, and t_0 is chosen so that $\sigma_i(t=0) = \sigma_{i,0}$, i.e.

$$\frac{t_0}{t_{vir}} = \ln\left(1 + \frac{\sigma_{i,0}^2}{\sigma_{DM}^2 + \sigma_{i,0}^2}\right). \quad (17)$$

Equation 16 has useful features: 1) it depends only on parameters already introduced in eq. 6; 2) it predicts $\sigma_i \rightarrow \sigma_{DM}$ as $\tau \rightarrow \infty$, and 3) in the limit $\sigma_{i,0} \rightarrow 0$ it returns the heating rates predicted by the LO85 model, eq. 6 (this can be easily verified by taking the second-order Taylor expansion of eq. 16 and substituting in the definitions of τ_{max} and t_0 , i.e. eq. 8 and eq. 17, respectively). Note that eq. 16 also ensures that the effects of collisional heating are the same regardless of the initial velocity dispersion of stellar particles. The black dashed lines in Figure 2 correspond to eq. 16 for the same best-fit parameters described above (and provided in Table 2).

4.3 Evolution of the vertical and radial stellar velocity dispersion

The simple empirical model described above (eq. 16) successfully captures the collisional heating rates inferred from all of our numerical simulations. This is summarized in Figure 4, where we plot the evolution of the vertical (σ_z^2 , upper panels) and radial (σ_R^2 , lower

Figure 3. Standard deviation in $k_i \ln \Lambda$ obtained for our fiducial simulations plotted as a function of the local density exponent, α . Different line styles correspond to different (fixed) values of β , as indicated in the legend. The parameters α and β determine the degree to which the evolution of the stellar velocity dispersion due to collisional heating depends on the local value of the DM density and velocity dispersion, respectively (see eq. 6; $\alpha = \beta = 0$ are the values anticipated by LO85). Blue lines corresponds to the vertical velocity dispersion, σ_z , and orange lines to the radial dispersion, σ_R .

panels) stellar velocity dispersion measured at several characteristic radii for a range of galaxy/halo models. All models correspond to a DM halo with $V_{200} = 200 \text{ km s}^{-1}$ and $\lambda_{DM} = 0.03$ and a disk mass fraction of $f_* = 0.01$, but other properties of the halo or disk are varied as described below.

The left-most panels show results for different m_{DM} (coloured lines) and μ (line styles); in all cases, velocity dispersions have been measured at the initial half-stellar mass radius, $R_{1/2}$. When expressed in scaled units, all models evolve approximately self-similarly, suggesting that *heating rates are driven primarily by the mass resolution of the DM component, and are largely insensitive to the stellar particle mass*, although at arbitrarily late times the dispersion is significantly lower for $\mu = 1$ (see Appendix B and Ludlow et al. 2019a for a more detailed discussion).

In the middle panels we plot results for three values of the halo's concentration parameter: $c = 7$ (dotted-dashed), $c = 15$ (dashed) and our fiducial value $c = 10$ (solid). Results are shown for $\mu = 5$ and for two different DM particle masses – $m_{DM} = 10^7 M_\odot$, green, and $m_{DM} = 10^8 M_\odot$, blue. Note that for $c = 7$ and 15 we have adjusted f_j (see Table 1 for details) to ensure that the stellar mass profile remains unchanged – $R_{1/2} = 4.14 \text{ kpc}$ is therefore independent of concentration, but the local density and velocity dispersion of DM particles at that radius are not. For our fiducial model, $c = 10$, we plot results at the initial value of $R_{1/2}$; for the other concentrations, we plot results at the radius that has the same local DM density but *not* the same stellar density or DM velocity dispersion (the latter are indicated by horizontal grey lines).

Finally, the right-most panels of Figure 4 compare results for three different initial disk scale-heights. Results are shown at the initial radius $R_{1/2}$. Our fiducial models ($z_d = 0.05 R_d$) are shown using solid lines; initial scale-heights two and four times larger are shown using dashed and dotted lines, respectively. As with the middle panels, in this case we plot results for $\mu = 5$, and for two different DM parti-

Figure 4. Vertical (upper panels) and radial (lower panels) dimensionless velocity dispersion, σ_i^2/V_{200}^2 , as a function of (dimensionless) time, $\tau \equiv t/t_c$ (see text for details), for variations of our fiducial model. Left panels adopt $f_j = 1$, but vary the dark matter particle mass and $\mu = m_{\text{DM}}/m_*$ as indicated; results are plotted at the initial half-stellar mass radius, $R_{1/2}$, which is the same for all models. Middle panels vary the concentration of the dark matter halo, with f_j adjusted in order to keep the stellar mass profile fixed (see Table 1). In this case, results for our fiducial model, i.e. $c = 10$, are plotted at the initial radius $R_{1/2}$, but for the other values of c we plot results at the radius corresponding to the same DM density (note that the DM velocity dispersion and stellar density differ at these radii for the three models plotted). The right-hand panels vary the disk’s initial scale height, z_d , and are plotted at the initial value of $R_{1/2}$ (which is the same for all three values of z_0). The horizontal grey lines in each panel mark the local one-dimensional velocity dispersion of DM particles at the corresponding radii. The solid black lines (shown only in the left- and right-hand panels, for clarity) show the best-fitting eq. 6, for which $\sigma_i \propto t$. Expressed in scaled units, the stellar velocity dispersion evolves approximately self-similarly, and are better described by eq. 16 (purple dot-dashed curves).

cle masses. Note that varying the disk’s initial scale-height leads to different initial values for the velocity dispersion at $R_{1/2}$ (Table 1), but the differences are small compared to the integrated effects of collisional heating.

The solid black lines in the left and right hand panels (not plotted in the middle panels for clarity) of Figure 4 correspond to eq. 6, plotted using the best-fit value of α (and $\beta = 0$) determined above for our fiducial models (Table 2); these curves describe the simulation results reasonably well in the regime $\sigma_i \ll \sigma_{\text{DM}}$ (horizontal lines indicate σ_{DM}^2 at $R_{1/2}$). However, as hinted at in Figure 2, systematic departures are noticeable at late times, and are better captured by eq. 16, shown as dot-dashed purple lines for the same parameters.

Overall, these results suggest that eq. 16 provides an accurate account of the collisional heating rates measured in all of our simulations, accommodating a wide variation in the masses of both DM and stellar particles, the local density and velocity dispersion of DM particles, and the initial height (and velocity dispersion) of disk stars. We next turn our attention to the evolution of the vertical scale-height of disks.

4.4 The impact of collisional heating on the vertical structure of thin disks

4.4.1 Thick versus thin disks

In Section 2.3 we outlined the relation between the vertical velocity dispersion of disk stars and their scale height. The latter is determined by the balance between the vertical “pressure” gradients exerted by disk stars and the combined vertical compressive forces of the DM halo and disk. In the thin disk limit, where the latter are dominated by either the disk (self-gravitating, SG) or the halo (non-self-gravitating, NSG), scale heights are given by eqs. 10 or 11, respectively. The actual scale heights will be the smaller of the two, or in cases where disk and halo make comparable contributions to the vertical acceleration profile, smaller than both. As in Benítez-Llambay et al. (2018), we find that a useful approximation for the thin-disk vertical half-mass scale height is given by the square harmonic mean of the SG and NSG cases, i.e

$$z_{1/2}^{\text{thin}} = \left(\frac{1}{z_{\text{SG}}^2} + \frac{1}{2z_{\text{SG}}z_{\text{NSG}}} + \frac{1}{z_{\text{NSG}}^2} \right)^{-1/2}, \quad (18)$$

where we have used the super-script ‘thin’ to indicate that the thin-disk approximation has been used.

In Figure 5 we plot z_{SG} (dot-dashed lines), z_{NSG} (short dashed

Figure 5. The vertical half-mass scale height of stellar particles (normalized by r_{200}) plotted as a function of their vertical velocity dispersion (normalized by V_{200}). Results are shown for the cylindrical radii corresponding to the initial values of $R_{1/4}$, $R_{1/2}$ and $R_{3/4}$ (left to right, respectively). Black dots show results obtained from our fiducial simulations (limited to $\mu = 5$, but shown for all V_{200} and m_{DM}). The various curves correspond to different theoretical prescriptions for calculating the vertical scale heights of disks: z_{NSG} (dashed lines) and z_{SG} (dot-dashed) are the non-self-gravitating and self-gravitating disk scale heights calculated in the thin disk limit, respectively; the dashed orange line (labelled $z_{1/2}^{\text{thin}}$) is their squared harmonic mean (eq. 18). The thick green line corresponds to the half-mass scale height of a “thick” non-self-gravitating disk, $z_{\text{NSG}}^{\text{thick}}$ (see text for details), and the blue line is an extrapolation between $z_{1/2}^{\text{thin}}$ (at low σ_z) and $z_{\text{NSG}}^{\text{thick}}$ (at high σ_z ; see eq. 21).

line) and $z_{1/2}^{\text{thin}}$ (dashed orange lines) as a function of the vertical stellar velocity dispersion at radii corresponding to the initial values of $R_{1/4}$, $R_{1/2}$ and $R_{3/4}$ (left-to-right panels, respectively). Results obtained from our fiducial set of simulations ($c = 10$ and $\mu = 5$, but for all values of V_{200} and m_{DM}) are underlaid as black points (each simulation output is shown as a single point). Note that, provided the vertical velocity dispersion remains small ($\sigma_z^2/V_{200}^2 \lesssim 0.1$, or so), eq. 18 accurately describes the σ_z -dependence of the measured half-mass scale-heights. This is shown explicitly in Figure 6, where we plot the time-dependence of $z_{1/2}$, measured at the initial value of $R_{1/4}$, for one of our fiducial models ($V_{200} = 200 \text{ km s}^{-1}$, $\mu = 5$), with m_{DM} spanning $10^6 M_\odot$ to $10^8 M_\odot$ (different sets of curves). Results obtained when assuming the *thin* disk approximation, $z_{1/2}^{\text{thin}}$, are again shown using dashed orange lines. This simple model works remarkably well *provided the disk remains thin*; If it does not, then the model does not, which is most readily apparent for the lowest-resolution run in which disks are maximally heated; in this case $z_{1/2}^{\text{thin}}$ systematically under-predicts the measured scale height.

These extreme instances of collisional disk thickening can nevertheless be accommodated by renouncing the thin disk approximation and directly solving eq. 12 in the NSG limit. In general, this must be done numerically, but admits an analytic solution for a Hernquist DM potential. In this case, eq. 12 becomes

$$\frac{1}{\rho_\star} \frac{\partial \rho_\star}{\partial z'} = - \frac{1}{(1-f_\star) v_z^2} \frac{z'}{(\sqrt{R'^2 + z'^2} + a')^2 \sqrt{R'^2 + z'^2}}, \quad (19)$$

where $v_z = \sigma_z/V_{200}$ and the primes indicate quantities normalized by r_{200} (i.e. $x' = x/r_{200}$). This can be easily integrated, yielding

$$\ln \left[\frac{\rho_\star(R', z')}{\rho_\star(R', 0)} \right] = \frac{1}{(1-f_\star) v_z^2} \left[\frac{1}{\sqrt{R'^2 + z'^2} + a'} - \frac{1}{R' + a'} \right]. \quad (20)$$

Note that in the limit $z' \rightarrow \infty$, eq. 20 approaches a constant density, and the vertical mass profile diverges; this is clearly nonphysical. The vertical scale-height, $z_{\text{NSG}}^{\text{thick}}$, can nonetheless be estimated robustly. To do so, we integrate eq. 20 along the z' -direction to obtain the mass enclosed by an infinitesimally-thin cylinder of radius $R_{1/2}$ and height $z'_{\text{max}} = 1$ (i.e. we integrate to a maximum vertical height

Figure 6. Evolution of the vertical half-mass scale height, $z_{1/2}$, measured at the radius $R_{1/4}$ enclosing one-quarter of the initial stellar mass. Results are shown for one of our fiducial models ($V_{200} = 200 \text{ km s}^{-1}$, $\mu = 5$) and for a range of dark matter particle masses (increasing from $m_{\text{DM}} = 10^6 M_\odot$ to $10^8 M_\odot$ in equally-spaced steps of $\Delta \log m_{\text{DM}} = 0.5$). Black lines show the simulation results; orange dashed lines show the evolution expected for *thin* stellar disks ($z_{1/2}^{\text{thin}}$; eq. 18) and blue lines correspond to $z_{1/2}^{\text{thick}}$ (see text and eq. 21 for details). In both cases, the vertical velocity dispersion is calculated from eq. 16 using the best-fit parameters provided in Table 2.

of $z = r_{200}$), and then determine the value of z' that encloses half of the integrated mass⁴. The results are shown in Figure 5 as thick green lines. The thick blue lines labelled $z_{1/2}^{\text{thick}}$ show a simple inter-

⁴ Note that, for the stellar disks and DM haloes considered in this paper, the vertical scale-heights obtained this way are largely insensitive to the initial upper-limit of integration, z'_{max} . For example, adopting $z'_{\text{max}} = 0.5, 2.0$, or 10 yield scale-heights that are virtually indistinguishable from those plotted in Figure 5.

pole between $z_{1/2}^{\text{thin}}$ (at low σ_z) and $z_{\text{NSG}}^{\text{thick}}$ (at high σ_z), specifically,

$$z_{1/2}^{\text{thick}} = z_{1/2}^{\text{thin}} - (z_{\text{NSG}}^{\text{thick}} - z_{\text{NSG}}). \quad (21)$$

Equation 21 is also plotted in Figure 6 using thick blue lines, which in poorly resolved systems provides a clear improvement over scale-heights estimated using the thin disk approximation.

We show in Appendix D that the impact of spurious heating on the vertical scale-height is suppressed in simulations that adopt gravitational softening lengths $\varepsilon \gtrsim z_d$. Nevertheless, the relation between $z_{1/2}$ and σ_z depicted in Figure 5 remains the same, suggesting that, regardless of ε , vertical scale heights of simulated disks can be inferred from their vertical velocity dispersion profiles as described above.

4.4.2 Evolution of the vertical half-mass scale height

The upper-panels of Figure 7 plot the evolution of the normalized vertical half-mass scale heights (i.e. $z_{1/2}/r_{200}$) for the same models used for Figure 4. The left-most panel shows results for a range of m_{DM} and μ (see legend); in this case, scale heights are measured at the initial radius $R_{1/2}$. The middle panel shows results for runs that vary the halo concentration (with the disk's spin parameter adjusted to ensure the stellar mass profile remains the same between models). In this case, results are plotted at $R_{1/2}$ for our fiducial model ($c = 10$), but at radii corresponding to the same DM density for the other models (which correspond to different DM velocity dispersion and stellar density). In the right-most panel we plot results measured at $R_{1/2}$ but for disks with varying initial scale heights ($z_d/R_d = 0.05, 0.1$ and 0.2 , as indicated). In all panels, simulations are shown as coloured lines; dot-dashed purple curves show the half-mass scale height, $z_{1/2}^{\text{thick}}$ (eq. 21), calculated as described above (note that the time-dependence of the scale height is driven solely by the predicted evolution of the vertical velocity dispersion at $R_{1/2}$, i.e. by eq. 16).

The lower panels of Figure 7 plot the evolution of $z_{1/2}$ for the same models as in the upper-left panel, but at initial radii corresponding to $R_{1/4}$ (left), $R_{1/2}$ (middle) and $R_{3/4}$ (right; each of these characteristic radii are shown as horizontal dashed lines in the respective panel). In this case, quantities are expressed in physical units to better emphasize the impact of collisional heating on the thickness of simulated disks. As with the upper panels, dot-dashed purple lines show the predicted scale-height evolution, $z_{1/2}^{\text{thick}}$. In all cases, the simple empirical model described above for the vertical scale-heights of disks – combined with the evolution of the vertical velocity dispersion profiles predicted by eq. 16 – provides an accurate description of our numerical results.

5 APPLICATION TO COSMOLOGICAL SIMULATIONS OF GALAXY FORMATION

The results presented above suggest that simulated disk galaxies in the presence of coarse-grained dark matter haloes are susceptible to spurious collisional heating by dark matter particles. This echoes the findings of Sellwood (2013), who showed that the relaxation of stellar disks due to star-star encounters also poses a threat to simulations of galaxy formation.

Collisional heating affects the kinematics of disk particles and their vertical structure at essentially all galacto-centric radii (unlike the collisional relaxation of DM haloes, which is primarily limited to their central regions). Heating rates depend primarily on the mass, local density and characteristic velocity of DM particles, and can be

accurately described by a simple semi-empirical model which has its roots in the analytic theory developed by Chandrasekhar (1960) and extended by Lacey & Ostriker (1985). Below we discuss the implementation of our model and its implications for the interpretation of disk galaxy structure in cosmological simulations.

5.1 Implementation of the empirical disk heating model

5.1.1 Velocity dispersions

Our empirical model for collisional disk heating is simple to implement. The time-dependence of the stellar velocity dispersion at any cylindrical radius R is given by eq. 16. In practice, heating rates depend on galacto-centric radius through the radial dependence of the local density and velocity dispersion of DM particles (eq. 13 and 14 for the models discussed in this paper, respectively), which can be determined from numerical simulations or calculated using empirical models that describe the structure of DM haloes as a function of their mass (e.g. Ludlow et al. 2014; Correa et al. 2015; Diemer & Kravtsov 2015; Ludlow et al. 2016; Diemer & Joyce 2019).

The best-fit parameters for eq. 16 were determined in Section 4.2 and are provided in Table 2, for reference. Only one parameter remains to be specified: $\sigma_{i,0}$, the *initial* components of the stellar velocity dispersion. To gauge the potential importance of spurious collisional heating we can set $\sigma_{i,0}$ to zero and compare the predicted values of $\sigma_i(R, t)$ to values of interest. The solid black line in the upper-left panel of Figure 8, for example, shows the vertical velocity dispersion (normalized by V_{200}) due to collisional heating at the characteristic disk scale radius, R_d , after $\Delta t = 13.6$ Gyr (results are similar for σ_R). The results are plotted as a function of the number of DM particles N_{200} within the virial radius r_{200} . We assume a Hernquist profile for the DM halo with an NFW-equivalent concentration of $c = 10$, but show the impact of varying c between 5 and 15 using a shaded region. For the disk, we assume $R_d = 0.2 \times r_{-2}$ where $r_{-2} = r_{200}/c$ is the scale radius of the corresponding NFW halo. Colored lines show results for the same set-up, but for various non-zero values of the initial stellar velocity dispersion, $\sigma_{z,0}$. For $c = 10$, the vertical velocity dispersion at R_d reaches 90 per cent of its maximum value of $\Delta\sigma_z \approx \sigma_{\text{DM}} \approx 0.52 V_{200}$ after $\Delta t = 13.6$ Gyr for $N_{200} \lesssim 3.26 \times 10^4$, although the precise value depends on concentration. Note that this is a limiting case, and collisional heating can be substantial for even larger N_{200} . For example, $\Delta\sigma_z(R_d) \approx 0.1 \times V_{200}$ for $N_{200} \approx 2.96 \times 10^6$ (assuming $c = 10$).

The upper-middle panel of Figure 8 shows the vertical velocity dispersion of disk stars (normalized by V_{200}) after $\Delta t = 13.6$ Gyr at $0.25 R_d$ (dot-dashed red line), R_d (black line) and $4 R_d$ (dashed blue line) within DM haloes of varying N_{200} . As in the left-hand panel, we assume $R_d = 0.2 \times r_{-2}$ and $c = 10$ (the grey shaded region again indicates the effect of varying c from 5 to 15, but is shown only for the $R = R_d$ curve). Note that σ_z reaches different maximum values at different radii and for different values of N_{200} . This is because the maximum stellar velocity dispersion is determined by the local velocity dispersion of DM, which has a weak radial gradient. Note too that, for a given N_{200} , collisional heating is less problematic at large R where the density of DM particles is lower. This suggests that collisional heating may alter the morphologies of simulated galaxies, a possibility we will explore in future work.

The upper right-hand panel of Figure 8 compares the vertical velocity dispersion at R_d after $\Delta t = 2$ Gyr (red dot-dashed line), 6 Gyr (dashed blue line) and 13.6 Gyr (black line). As above, we assume a Hernquist DM halo with NFW-equivalent concentration $c = 10$ (the shaded region has the same meaning as in the other panels), and plot

Figure 7. *Upper panels:* Same as Figure 4, but for the evolution of the vertical half-mass scale height, $z_{1/2}$. *Lower panels:* Evolution of the half-mass scale height at the initial radii $R_{1/4}$ (left), $R_{1/2}$ (middle) and $R_{3/4}$ (right; indicated by horizontal dashed lines) for our fiducial models. Different colours correspond to different dark matter particle masses and different line styles to different $\mu = m_{\text{DM}}/m_*$. In all panels, the purple dot-dashed curves correspond to $z_{1/2}^{\text{thick}}$ (see eq. 21 and Section 4.4 for details) calculated assuming σ_z evolves according to eq. 16.

results as a function of N_{200} . Fewer particles are required to suppress collisional heating over shorter timescales, as expected since it is an integrated effect. For example, the vertical velocity dispersion (measured at R_d) of an initially thin stellar disk will not exceed ≈ 10 per cent of V_{200} after 2 Gyr provided $N_{200} \gtrsim 4.29 \times 10^5$, whereas the same degree of heating will be reached for $N_{200} \lesssim 1.28 \times 10^6$ or $\lesssim 2.96 \times 10^6$ for integration times of 6 and 13.6 Gyr, respectively.

Although the results plotted in Figure 8 strictly apply to the vertical velocity dispersion of stellar particles, similar results are obtained for the radial velocity dispersion.

5.1.2 Vertical scale height

As discussed in Section 4.4, the evolution of σ_z determines the evolution of the vertical scale height of simulated disks. In our analysis, we characterized vertical scale heights by means of the vertical half-mass height, $z_{1/2}$. A useful approximation for $z_{1/2}$ (at least over the range of radii and galaxy/halo properties studied in this paper) is given by eq. 21, which provides a means of modelling $z_{1/2}$ in terms of σ_z , R and the structural properties of the disk and halo.

In the lower panels of Figure 8 we plot $z_{1/2}^{\text{thick}}$ (normalized by r_{200}) for the same set of models displayed in the upper panels for σ_z . Note that, even in the most extreme cases of collisional heating, the scale heights of disks do not exceed a few per cent of their host’s virial radius, r_{200} . There are, however, a few additional results in Figure 8 worth highlighting. One: the *minimum* vertical scale height of sim-

ulated disks are determined by the initial vertical velocity dispersion of their stars (lower left-hand panel). Two: our model predicts a outward flaring of stellar disks as a result of numerical heating; i.e., if allowed to reach its maximum thickness, the scale height will increase with R , at least over the radial range $R_d/4 \leq R \leq 4 \times R_d$ (lower middle panel; see also the lower-left panel of Figure 1). And three: vertical scale-heights grow monotonically with time, before plateauing at a maximum physical scale height that is determined by the local velocity dispersion of halo particles (lower right panel).

5.2 Implications for existing and future cosmological simulations

Our model can therefore be used to assess the importance of collisional disk heating for cosmological simulations of galaxy formation. In Figure 9 we plot, as a function of the DM particle mass, the halo masses below which disk heating is expected to become important. We assume a Hernquist (1990) profile for the DM halo with mass-dependent NFW-equivalent concentrations taken from the analytic model of Ludlow et al. (2016). The various lines correspond to different amounts of collisional heating affecting an initially perfectly-thin (i.e. $\sigma_{z,0} = 0$) stellar disk, and are plotted at $R = R_d$ (we assume $R_d = 0.2 \times r_{-2}$; for several curves we show the impact of varying R from 0.5 to $2 \times R_d$ using shaded regions). The dot-dashed green lines correspond to $\Delta\sigma_z = 0.03 \times V_{200}$, the solid red lines to $\Delta\sigma_z = 0.1 \times V_{200}$, and the dashed blue lines to $\Delta\sigma_z = 0.3 \times V_{200}$; dotted black lines indicate a maximally-heated disk, $\Delta\sigma_z = \sigma_{\text{DM}}$. Solid

Figure 8. The vertical stellar velocity dispersion (squared, and normalized by V_{200}^2 ; top panels) and the vertical half-mass disk scale-height (normalized by r_{200} ; bottom panels) as a function of the number of DM particles N_{200} within the virial radius, r_{200} . Different columns show the effect of varying different model parameters. Panels on the left show the impact of varying $\sigma_{z,0}$, the initial stellar vertical velocity dispersion. The black line corresponds to the limiting case, $\sigma_{z,0} = 0$; red (dot-dashed), blue (dashed) and green (dotted) lines to $\sigma_{z,0}/V_{200} = 0.05, 0.1$ and 0.2 , respectively. Each of these curves measure the integrated effect of collisional heating measured at the disk scale radius, R_d (approximated here as $R_d = 0.2 \times r_{-2}$, where r_{-2} is the scale radius of an NFW halo) after $\Delta t = 13.6$ Gyr in a DM halo with concentration parameter $c = 10$. The shaded region shows the effect of varying the halo concentration between 5 and 15 (shown only for $\sigma_{z,0} = 0$). The middle panels show the impact of collisional heating at different radii. Black lines correspond to $R = R_d = 0.2 \times r_{-2}$ (the shaded region again shows the impact of varying c from 5 to 15), whereas blue (dashed) and red (dot-dashed) lines correspond to $R = R_d/4$ and $R = 4 \times R_d$, respectively (each curve assumes $\sigma_{z,0} = 0$ and $\Delta t = 13.6$ Gyr). The right-most panels show σ_z and $z_{1/2}$ measured at R_d for different integration times: $\Delta t = 2, 6$, and 13.6 Gyr ($\sigma_{z,0} = 0$).

black lines (in order of increasing thickness) correspond to fixed physical degrees of heating: $\Delta\sigma_z = 1 \text{ km s}^{-1}$, 3 km s^{-1} , 10 km s^{-1} , and 30 km s^{-1} respectively. The left- and right-hand panels of Figure 9 correspond to integration times of $\Delta t = 10$ Gyr and 2 Gyr respectively.

For the dark matter particle masses adopted for many recent cosmological simulations, $m_{\text{DM}} \approx 10^4 M_\odot$ to $\approx 10^8 M_\odot$, collisional heating is expected to be important over a broad range of halo masses. More specifically, the locations at which a vertical line in Figure 9 intersects the curves corresponding to a fixed (fractional or absolute) amount of heating mark the halo masses below which we expect collisional heating to be important for that particle mass. Clearly the properties of galaxies in many simulations may be unduly affected by collisional heating, particularly their structure, morphology and kinematics.

For example, all recent large-volume cosmological hydrodynamical simulations (i.e. $V \gtrsim 100^3 \text{ Mpc}^3$) used DM particle masses $m_{\text{DM}} \gtrsim 10^6 M_\odot$ (this includes the EAGLE simulations, Illustris, TNG300, TNG100, TNG50, Horizon-AGN, New Horizon and Magneticum, but also Mufasa and Simba runs; see Table 3 for simulation details and references). We therefore expect collisional heating in these runs, integrated over ≈ 10 Gyr, to increase the vertical velocity dispersion of disk stars at $R = R_d$ to $\gtrsim 0.1 V_{200}$ ($0.03 V_{200}$) for halo masses $M_{200} \lesssim 1.6 \times 10^{12} M_\odot$ ($1.4 \times 10^{13} M_\odot$), corresponding to haloes resolved with fewer than $N_{200} = 1.6 \times 10^6$ (1.4×10^7) DM particles. These mass scales are reduced somewhat for shorter integration times, but remain problematic. For example, for $\Delta t =$

2 Gyr we find $\Delta\sigma_z \gtrsim 0.1 V_{200}$ ($0.03 V_{200}$) for $M_{200} \lesssim 3.8 \times 10^{11} M_\odot$ ($3.3 \times 10^{12} M_\odot$), corresponding to $N_{200} = 3.8 \times 10^5$ (3.3×10^6).

Results become substantially worse as the DM particle mass increases, and a few of the runs mentioned above are therefore *more* vulnerable to collisional heating than just mentioned. For $m_{\text{DM}} = 10^7 M_\odot$ – which is comparable to, or even better than the mass resolution of TNG300, TNG100, Horizon-AGN, Magneticum, and the intermediate-resolution EAGLE simulation – collisional heating is likely to strongly affect the structure and kinematics of simulated disks (i.e. $\Delta\sigma_z \gtrsim 0.1 V_{200}$) in haloes with masses less than about $M_{200} \lesssim 1.2 \times 10^{13} M_\odot$ (or equivalently, $N_{200} \lesssim 1.2 \times 10^6$). Thus, virtually all star-forming main sequence galaxies will be spuriously heated in these runs. More specifically, galaxies hosted by haloes with $M_{200} \approx 10^{13} M_\odot$ ($\approx 10^{12} M_\odot$) will experience a spurious increase in their vertical velocity dispersion at $R \approx R_d$ of $\approx 40 \text{ km s}^{-1}$ ($\approx 62 \text{ km s}^{-1}$) over roughly $\Delta t = 10$ Gyr; their radial velocity dispersion will increase by $\Delta\sigma_R \gtrsim 58 \text{ km s}^{-1}$ ($\gtrsim 84 \text{ km s}^{-1}$) and their half-mass scale-heights by $\Delta z_{1/2} \gtrsim 750 \text{ pc}$ ($\gtrsim 1.5 \text{ kpc}$) over the same timescale. Collisional heating is also problematic over much shorter timescales: For example, for the same halo masses $\Delta\sigma_z \gtrsim 18 \text{ km s}^{-1}$ ($\gtrsim 30 \text{ km s}^{-1}$), $\Delta\sigma_R \gtrsim 26 \text{ km s}^{-1}$ ($\gtrsim 43 \text{ km s}^{-1}$) and $\Delta z_{1/2} \gtrsim 196 \text{ pc}$ ($\gtrsim 515 \text{ pc}$) for $\Delta t = 2$ Gyr.

Increasing the mass resolution of DM particles in cosmological simulations is the best way to suppress the spurious collisional heating of galaxy disks. For DM particle masses $m_{\text{DM}} \approx 10^5 M_\odot$ (which is comparable to the highest uniform-resolution cosmological simulation published to date, i.e. the $V = (12.5 h^{-1} \text{ Mpc})^3$ Simba

Figure 9. The mass of haloes, M_{200} , below which simulated disk galaxies are expected to be unduly affected by collisional heating. Results are plotted as a function of the dark matter particle mass, m_{DM} . The dot-dashed green lines corresponds to a spurious increase in the vertical velocity dispersion of disk stars by $\Delta\sigma_z = 0.03 \times V_{200}$ over $\Delta t = 10$ Gyr (left) and 2 Gyr (right), the solid red lines to $\Delta\sigma_z = 0.1 \times V_{200}$, the dashed blue line to $\Delta\sigma_z = 0.3 \times V_{200}$, and the dotted black line to the maximum possible value $\Delta\sigma_z = \sigma_{\text{DM}}$ (note that we assume $\sigma_{z,0} = 0$ for each curve). All results are calculated at the disk’s characteristic radius, R_d , with shaded regions indicating the broader radial range $R = (0.5 - 2) R_d$. In order of increasing thickness, solid black lines correspond to the halo masses below which collisional heating increases the vertical velocity dispersion of disk stars by $\Delta\sigma_z = 1, 3, 10$ and 30 km s^{-1} (in these cases, results are plotted only at $R = R_d$ for clarity). We assume $R_d = 0.2 \times r_{-2}$, where $r_{-2} = r_{200}/c$ is the characteristic radius of the NFW dark matter halo; the latter is prescribed using the analytic model of Ludlow et al. (2016).

run), spurious disk heating over a timescale of 10 Gyr is below the level of $\Delta\sigma_z \approx 0.03 \times V_{200}$ ($\approx 6 \text{ km s}^{-1}$) for $M_{200} \gtrsim 1.8 \times 10^{12} M_\odot$ ($N_{200} \approx 1.8 \times 10^7$) and below $\approx 0.1 \times V_{200}$ ($\approx 10 \text{ km s}^{-1}$) for $M_{200} \gtrsim 2.1 \times 10^{11} M_\odot$ ($N_{200} \gtrsim 2.1 \times 10^6$). For a DM particle mass of $m_{\text{DM}} \approx 5 \times 10^5$ (comparable to the value adopted for the TNG50 simulation), we expect, over ≈ 10 Gyr, $\Delta\sigma_z \approx 0.03 \times V_{200}$ ($\approx 10 \text{ km s}^{-1}$) for $M_{200} \approx 7.4 \times 10^{12} M_\odot$ ($N_{200} \approx 1.5 \times 10^7$) and $\Delta\sigma_z \approx 0.1 \times V_{200}$ ($\approx 16 \text{ km s}^{-1}$) for $M_{200} \approx 8.7 \times 10^{11} M_\odot$ ($N_{200} \approx 1.7 \times 10^6$). This may explain why Pillepich et al. (2019) reported poor convergence in the vertical scale-heights of disks in TNG50 with increasing mass resolution.

The most recent generation of zoom simulations, which sacrifice galaxy statistics in favour of resolution, typically adopt DM particle masses in the range $10^4 \lesssim m_{\text{DM}}/M_\odot \lesssim \text{a few} \times 10^5$ (see Table 3 for several examples). These runs often target Milky Way-mass galaxies ($M \approx 10^{12} M_\odot$) and, although it remains challenging to suppress collisional heating below $\Delta\sigma_z \lesssim 0.03 V_{200}$ ($\approx 4.9 \text{ km s}^{-1}$) over 10 Gyr in these halos (which would require $m_{\text{DM}} \lesssim 5.2 \times 10^4 M_\odot$), they typically succeed in suppressing it below $\Delta\sigma_z \lesssim 0.1 V_{200}$ ($\approx 16 \text{ km s}^{-1}$), which requires $m_{\text{DM}} \lesssim 5.9 \times 10^5 M_\odot$.

These are, of course, upper limits on spurious disk heating as galaxies may not experience secular evolution at all times, but they will experience spurious DM heating at all times, even if they also occasionally merge with other galaxies.

Our results can therefore be used to determine the range of DM particle masses for which spurious collisional heating will be unimportant in DM haloes of a given virial mass, and thus can be used as a guide for the selection of numerical parameters for simulations of galaxy formation and evolution. As discussed above, suppressing numerical relaxation in simulations will always favor higher (DM) mass resolution. The optimal particle mass for a particular simula-

tion will therefore represent a compromise between the simulation volume (or equivalently the number of resolved objects) and high resolution. Figure 10 helps visualize the situation. Here we plot, for a number of recent state-of-the-art cosmological and zoom simulations, the DM particle mass versus the average number of resolved DM haloes with virial masses above $M_{200} \geq 5 \times 10^9 M_\odot$ (which roughly corresponds to the halo mass above which galaxy occupation reaches ≈ 100 per cent; Benitez-Llambay & Frenk 2020). Vertical grey lines mark fixed volume increments, and the horizontal dashed line corresponds to particle masses for which $5 \times 10^9 M_\odot$ haloes are resolved with 20 particles.

The brown shaded regions in Figure 10 indicate the DM particle masses below which collisional heating (integrated over $\Delta t = 10$ Gyr) does not exceed $0.1 \times V_{200}$ in haloes of different mass. Darker shades correspond to decreasing halo masses, with values ranging from $M_{200} = 10^{13} M_\odot$ to $10^8 M_\odot$ in steps of 1 dex, as indicated. Note that the majority of zoom simulations targeting Milky Way-mass galaxies or local group analogues (e.g. Auriga, Eris, Apostle, FIRE, Latte, Elvis, CLUES, Hestia) are unlikely to be strongly impacted by spurious collisional heating, although both the progenitors of the galaxies that these simulations target and their present-day satellites may strongly be. However, most simulations of large volumes, e.g. $\gtrsim (100^3 \text{ Mpc}^3)$, which typically adopt DM particle masses $m_{\text{DM}} \gtrsim 10^6 M_\odot$, are likely to fall victim to the numerical heating effects discussed in this paper.

Finally, we stress that the magnitude of the heating one may wish to suppress will depend on the subgrid physics adopted for the run, and on the scientific goals of the simulation programme. Simulations that properly resolve the cold phase of the inter-stellar medium, for example, are expected to result in some galaxies having cold, molecular gas disks with $\sigma_z \ll 10 \text{ km s}^{-1}$, giving rise to stellar disks that, at

Figure 10. The DM particle mass versus the average number of resolved haloes with virial mass $M_{200} > 5 \times 10^9 M_\odot$ for a number of recent cosmological and zoom hydrodynamical simulations. Halo abundances are estimated using the Tinker et al. (2008) mass function; the value $M_{200} = 5 \times 10^9 M_\odot$ roughly corresponds to the halo mass above which galaxy occupation reaches 100 per cent (Benítez-Llambay & Frenk 2020). Vertical grey lines mark fixed volumes, increasing by factors of 8 from $(2 \text{ Mpc})^3$ to $(4096 \text{ Mpc})^3$; the horizontal dashed line indicates the minimum DM particle mass required to resolve $M_{200} = 5 \times 10^9 M_\odot$ haloes with at least 20 particles. The brown shaded regions highlight the DM particle masses below which spurious collisional heating (at $R = R_d$, integrated over $\Delta t = 10 \text{ Gyr}$) remains below $\Delta\sigma_z \lesssim 0.1 V_{200}$; darker shades correspond to haloes of decreasing virial mass, M_{200} , as labelled. As in previous figures, we assume $R_d = 0.2 \times r_{-2}$, where $r_{-2} = r_{200}/c$ is the NFW-equivalent scale radius of the Hernquist (1990) DM halo. The majority of state-of-the-art zoom simulations, and a few of the highest-resolution, smallest-volume cosmological runs, are in principle able to resolve cool stellar disks (e.g. $\Delta\sigma_z \lesssim 0.1 V_{200}$) in Milky Way-mass haloes, but most large volume cosmological runs cannot.

least initially, have very low vertical velocity dispersions and small scale heights. In these cases, DM particle masses should be chosen so that $\Delta\sigma_z \approx$ a few km s^{-1} in order to ensure that spurious collisional heating does not alter the kinematics of stellar particles. Simulations that oppose artificial fragmentation by preventing gas cooling below $\approx 10^4 \text{ K}$ have a natural kinematic floor of $\approx 10 \text{ km s}^{-1}$, which provides a sensible target above which collisional heating should be suppressed. The solid black curves in Figure 9 can be used to guide the selection of particle masses that adhere to these restrictions. For example, the vertical heating of disk stars will not exceed $\Delta\sigma_z \approx 3 \text{ km s}^{-1}$ over 10 Gyr in a simulated dwarf galaxy of mass $\approx 10^{11} M_\odot$ provided $m_{\text{DM}} \lesssim 7.1 \times 10^3 M_\odot$ (corresponding to $N_{200} \gtrsim 1.4 \times 10^7$); the heating will not exceed $\Delta\sigma_z \approx 10 \text{ km s}^{-1}$ over 10 Gyr provided $m_{\text{DM}} \lesssim 8.0 \times 10^4 M_\odot$ (or $N_{200} \gtrsim 1.3 \times 10^6$).

Some large-volume simulations are of course not intended for studies of galaxy structure, but rather for studies of the structure of the inter- or circum-galactic medium, the role of environmental effects in galaxy evolution, the properties of galaxy groups or clusters, or the imprint of baryons on cosmology and large-scale structure. It seems unlikely that spurious collisional disk heating is detrimental to simulations aimed at addressing these topics, but the effect should nonetheless be recognized as one of their important limitations.

In Table 3, we tabulate the halo masses below which collisional heating (specifically, $\Delta\sigma_z/V_{200} = 0.1$ over $\Delta t = 10 \text{ Gyr}$) may become important for the cosmological hydrodynamical simulations plotted in Figure 10.

6 SUMMARY AND CONCLUSIONS

Spurious collisional heating of simulated stellar disks by coarse-grained DM particles affects the vertical and radial kinematics of disk stars, and as a consequence, the structure of their stellar distribution. The effects are irreversible, and depend primarily on the number of DM particles and the structural properties of the disk's DM halo. A rigorous theoretical description of disk heating was outlined by Lacey & Ostriker (1985), who calculated (among other quantities) the rate at which orthogonal components of the stellar velocity dispersion of disk stars increases with time as a result of scattering off point-mass perturbers (black holes in their case, which were at the time viable candidates for DM). Their analytic model provides a surprisingly accurate description of our simulations results (see Figure 2), but cannot account for the strong radial gradients in the dark matter density across stellar disks that are embedded within realistic DM haloes, nor the asymptotic velocity dispersion of stellar particles set by either energy equipartition or the local escape velocity of the halo (eq. 7). We provided a simple empirical amendment to their equations that accurately describes the results of our simulations (see eq. 16).

Collisional heating is largely independent of stellar particle mass, but may be exacerbated in simulations that adopt stellar-to-DM particle mass ratios that differ substantially from unity (see Appendix B and Ludlow et al. 2019a). As a result, attempts to artificially increase the baryon-to-DM mass resolution (i.e. to decrease the baryon-to-DM particle mass ratio) in simulations will only benefit runs that reach sufficiently high DM-mass resolution to ensure that collisional heating due to DM particles can be safely ignored. If it cannot, the effect can instead be suppressed by increasing the resolution of the DM component relative to that of the stars.

Our results have implications for the reliability with which simulations can be used to interpret the vertical, radial and kinematic gradients in the ages and metallicities of stars. For example, in the solar neighbourhood the velocity dispersion of stars increases with age and with decreasing metallicity (e.g. Wielen 1977; Quillen & Garnett 2001); at fixed cylindrical radius R , the scale height of stars increases with age (e.g. Casagrande et al. 2016). It is plausible that such trends arise as a result of secular evolution, and reflect the cumulative effects orbital scattering (due to, for example, molecular clouds or globular clusters), or dynamical heating due to accretion events or disk instabilities. Disks in cosmological simulations, if not sufficiently well-resolved, may however exhibit similar age- or metallicity-dependent gradients as a result of *spurious* collisional heating (although see Navarro et al. 2018 and Grand et al. 2016 for examples of simulations in which they are not spurious). Studies of metallicity and age gradients based on simulated disk galaxies should suppress these numerical effects.

In simulations, collisions with DM particles randomly perturb stellar particle orbits, which may disturb ordered flows and disrupt the coherent motions reminiscent of rotationally-supported disks or tidal streams. The spatial distribution and kinematics of stars in simulated galaxies serve as proxies for their morphology, and are often used to decompose them into distinct kinematic components. The possible importance of relaxation and its effects on the structure and morphology of simulated galaxies should be given serious consideration, particularly in studies based on uniform-resolution cosmological simulations, in which galaxies with different stellar masses are resolved with different numbers of dark matter particles. We will

address the implications of collisional heating for the morphological transformation of galaxies in a follow-up paper.

Because galaxy disks are dynamically cold relative to their surrounding dark matter haloes, collisions between dark matter and stellar particles will, over time, increase the velocity dispersion of stars in an attempt to establish energy equipartition between the two components. This is true regardless of the DM-to-stellar particle mass ratio, μ (equipartition is satisfied when $\sigma_* m_* = \sigma_{\text{DM}} m_{\text{DM}}$), although it will be accelerated when $\mu > 1$. Spurious collisional heating of galaxies will therefore artificially increase their sizes, as the increased kinetic energy of stellar particles drives them onto higher energy orbits within the halo potential (see also Ludlow et al. 2019a).

The highlights of our work can be summarized as follows:

- (i) Spurious collisional heating due to DM halo particles affects the velocity dispersion and scale heights of disk stars at all galacto-centric radii (Figure 1). The magnitude of the effect depends on the mass of DM particles (or on the number of particles at fixed halo mass), but also on their density and velocity dispersion (Figure 2; the former dominates due to its much stronger radial gradients). As a result, heating is often more substantial in the denser, colder centres of DM haloes than in their outskirts, an effect that may drive morphological changes in disk galaxies. The effects of collisional heating on the (radial and vertical) velocity dispersion of disk stars (Figure 4), and on their scale-heights (Figure 4), can be described by eqs. 16 and 21, respectively.
- (ii) The effects of collisional heating are approximately independent of the mass of stellar particles, suggesting that perturbations to their orbits are due primarily to shot noise in the DM particle distribution rather than the gradual exchange of energy between DM and stellar particles of unequal mass (see e.g. Figure 1 or 4, but also the discussion in Appendix B and Ludlow et al. 2019a). Because the total mass of the DM halo greatly exceeds that of the stars, other galaxy components, such as bulges or stellar haloes, contribute little to the heating of disk stars (see Appendix C). Collisional heating is independent of the gravitational softening length (ϵ) provided the vertical scale-height (z_d) of a simulated galaxy is resolved with at least one softening length, i.e. $\epsilon \lesssim z_d$ (Appendix D).
- (iii) Our results provide a tool to help interpret results from current cosmological simulations (see Figures 9 and 10, Table 3, and the discussion in section 5.2), and to guide the selection of DM particle masses and gravitational softening lengths for future high-resolution simulations of galaxy formation. For example, to ensure that the vertical collisional heating of stars in the disk of a typical Milky Way-mass galaxy (i.e. $M \approx 10^{12} M_\odot$) does not exceed $\Delta\sigma_z \approx 5 \text{ km s}^{-1}$ (roughly the minimum scale set by a gas particle temperature floor of 10^4 K) over $\Delta t = 10 \text{ Gyr}$, the DM particle mass must be $m_{\text{DM}} \lesssim 5.5 \times 10^4 M_\odot$ ($N_{200} \gtrsim 1.8 \times 10^7$). Maintaining $\Delta\sigma_z \lesssim 0.1 V_{200} \approx 16 \text{ km s}^{-1}$ (again for $\Delta t = 10 \text{ Gyr}$) requires $m_{\text{DM}} \lesssim 5.9 \times 10^5 M_\odot$ ($N_{200} \gtrsim 1.7 \times 10^6$). For a typical dwarf galaxy, $M_{200} \approx 10^{11} M_\odot$, DM particle masses of $m_{\text{DM}} \lesssim 2.0 \times 10^4 M_\odot$ and $4.5 \times 10^4 M_\odot$ (or equivalently, $N_{200} \gtrsim 5 \times 10^6$ and $N_{200} \gtrsim 2.2 \times 10^6$, respectively) ensure collisional heating does not exceed $\Delta\sigma_z = 5 \text{ km s}^{-1}$ and $0.1 V_{200}$, respectively. The results are based on heating rates measured at the characteristic disk scale length, which we take to be $\approx 1.4 \text{ kpc}$ and 3.8 kpc for the dwarf and Milky Way-mass galaxies, respectively (i.e. $0.2 \times r_{-2}$ for typical values of the DM halo's concentrations). Softening lengths should be smaller than the vertical scale-height of a thin, molecular disk, i.e. $\epsilon \lesssim 100 \text{ pc}$.

In closing, we note a number of limitations of our study that could be improved upon in future work. One is that by design our idealized

Table 3. A summary of the halo masses in recent state-of-the-art simulations below which we expect collisional vertical disk heating to be of order $0.1 \times V_{200}$ over a timescale of $\Delta t = 10^{10}$ yr (taken from Figure 9). The first three columns provide the simulation name, the journal reference and the type of simulation (i.e. whether a uniform mass-resolution cosmological simulation or zoom). Next we list the DM particle mass (m_{DM}), and the virial mass and corresponding DM particle number (M_{200} and N_{200} , respectively) below which spurious heating of initially-cold stellar disks results in $\Delta\sigma_z/V_{200} = 0.1$; the corresponding amount of physical heating (i.e. in units of kms^{-1}) is also provided, as are the corresponding changes in vertical disk scale height, $\Delta z_{1/2}$, calculated assuming $f_* = 0.01$ (both in physical units and as a fraction of r_{200}). For zoom simulations that vary the mass resolution (either for an individual galaxy, or a sample of galaxies), we quote values for the highest-resolution run published.

Simulation	Reference	Type	m_{DM} [$M_\odot$]	M_{200} [$M_\odot$]	N_{200} [10^6]	$\Delta\sigma_z$ [V_{200}]	$\Delta\sigma_z$ [kms^{-1}]	$\Delta z_{1/2}$ [r_{200}]	$\Delta z_{1/2}$ [kpc]
Horizon-AGN	Dubois, et al (2014)	Cosmo.	8×10^7	7.5×10^{13}	0.9	0.10	69.0	2.6×10^{-3}	1.76
New Horizon	Dubois et al. (2020)	Cosmo.	1.2×10^6	1.9×10^{12}	1.6	0.10	20.2	1.4×10^{-3}	0.27
Mufasa-1	Davé et al. (2016)	Cosmo.	9.6×10^7	8.8×10^{13}	0.9	0.10	72.6	2.8×10^{-3}	1.99
Mufasa-2	"	Cosmo.	1.2×10^7	1.4×10^{13}	1.2	0.10	39.7	1.9×10^{-3}	0.75
Mufasa-3	"	Cosmo.	1.5×10^6	2.3×10^{12}	1.5	0.10	21.6	1.3×10^{-3}	0.30
Simba-1	Davé et al. (2019)	Cosmo.	9.6×10^7	8.8×10^{13}	0.9	0.10	72.6	2.8×10^{-3}	1.99
Simba-2	"	Cosmo.	1.2×10^7	1.4×10^{13}	1.2	0.10	39.7	1.9×10^{-3}	0.75
Simba-3	"	Cosmo.	1.5×10^6	2.3×10^{12}	1.5	0.10	21.6	1.4×10^{-3}	0.30
Simba-4	"	Cosmo.	1.9×10^5	3.6×10^{11}	1.9	0.10	11.7	1.1×10^{-2}	0.12
Magneticum	Dolag et al. (2016)	Cosmo.	1.3×10^{10}	6.1×10^{15}	0.5	0.10	298.1	5.9×10^{-3}	17.5
Ilustris-1	Vogelsberger et al. (2014)	Cosmo.	6.3×10^6	8.1×10^{12}	1.3	0.10	32.8	1.7×10^{-3}	0.56
Ilustris-2	"	Cosmo.	5.0×10^7	5.0×10^{13}	1.0	0.10	60.1	2.4×10^{-3}	1.42
Ilustris-3	"	Cosmo.	4.0×10^8	3.0×10^{14}	0.8	0.10	109.4	3.4×10^{-3}	3.72
TNG300	Springel et al. (2018)	Cosmo.	5.9×10^7	5.8×10^{13}	1.0	0.10	62.9	2.5×10^{-3}	1.58
TNG100	"	Cosmo.	7.5×10^6	9.5×10^{12}	1.3	0.10	34.5	1.8×10^{-3}	0.63
TNG50	Pillepich et al. (2019)	Cosmo.	4.5×10^5	7.9×10^{11}	1.8	0.10	15.0	1.2×10^{-3}	0.18
EAGLE-MR	Schaye et al. (2015)	Cosmo.	9.7×10^6	1.2×10^{13}	1.2	0.10	37.3	1.9×10^{-3}	0.69
EAGLE-HR	"	Cosmo.	1.2×10^6	1.9×10^{12}	1.6	0.10	20.3	1.4×10^{-3}	0.27
BAHAMAS	McCarthy et al. (2017)	Cosmo.	5.5×10^9	2.9×10^{15}	0.5	0.10	231.2	5.1×10^{-3}	11.82
Massive Black-II	Khandai et al. (2015)	Cosmo.	1.6×10^6	2.4×10^{12}	1.5	0.10	21.8	1.4×10^{-3}	0.30
Romulus25	Tremmel et al. (2017)	Cosmo.	3.4×10^5	6.1×10^{11}	1.8	0.10	13.8	1.1×10^{-3}	0.15
Eris	Guedes et al. (2011)	zoom	9.8×10^4	2.0×10^{11}	2.0	0.10	9.6	9.7×10^{-4}	0.09
Auriga	Grand et al. (2017)	zoom	3.0×10^5	5.5×10^{11}	1.8	0.10	13.4	1.1×10^{-3}	0.15
Apostle (L1)	Sawala et al. (2016)	zoom	5.0×10^4	1.1×10^{11}	2.2	0.10	7.8	8.8×10^{-4}	0.07
CLUES	Libeskind et al. (2010)	zoom	2.9×10^5	5.3×10^{11}	1.8	0.10	13.2	1.1×10^{-3}	0.15
HESTIA	Libeskind et al. (2020)	zoom	1.5×10^5	2.9×10^{11}	2.0	0.10	10.9	9.9×10^{-4}	0.11
NIHAO-UHD	Buck et al. (2020)	zoom	5.1×10^5	8.8×10^{11}	1.7	0.10	15.7	1.2×10^{-3}	0.19
Latte	Wetzel et al. (2016)	zoom	3.5×10^4	7.9×10^{10}	2.3	0.10	7.0	8.6×10^{-4}	0.06
FIRE-II (m12i)	Hopkins et al. (2018)	zoom	4.3×10^4	9.6×10^{10}	2.2	0.10	7.4	8.7×10^{-4}	0.07
Elvis	Garrison-Kimmel et al. (2014)	zoom	1.9×10^5	3.6×10^{11}	1.9	0.10	11.7	1.1×10^{-3}	0.13
Hydrangea	Bahé et al. (2017)	zoom	9.7×10^6	1.2×10^{13}	1.2	0.10	37.3	1.9×10^{-3}	0.69
The 300 Project	Cui et al. (2018)	zoom	1.9×10^9	1.1×10^{15}	0.6	0.10	169.5	4.5×10^{-3}	7.58

disks do not account for a number of important physical processes that may also give rise to secular disk heating. Among them are accretion-driven heating, or orbital scattering by instabilities, globular clusters, giant molecular clouds or spiral density waves; spurious heating of disks by star particles in stellar haloes or bulges are also ignored (but see Appendix C). Nevertheless, collisional heating is an integrated effect, and we expect relaxation rates estimated from eq. 16 to yield sensible *lower limits* to the spurious heating due to coarse-grained DM particles. Perhaps most importantly, our study neglects the time-dependent evolution of the DM potential (which introduces a time-dependent DM density and characteristic velocity), as well as the fact that most simulated disk galaxies actively form stars, thus replenishing a thin disk with a low velocity dispersion stellar component. These issues should be the focus of future work.

ACKNOWLEDGEMENTS

ADL and DO acknowledge financial supported from the Australian Research Council through their Future Fellowship scheme (project

numbers FT160100250, FT190100083, respectively). This work is partly funded by Vici grant 639.043.409 from the Dutch Research Council (NWO). This work has benefited from the following public PYTHON packages: SCIPY (Virtanen et al. 2020), NUMPY (Harris et al. 2020), MATPLOTLIB (Hunter 2007) and IPYTHON (Pérez & Granger 2007).

DATA AVAILABILITY

The simulations used in this paper can be made available upon request. Theoretical results are reproducible using the equations provided in the paper.

REFERENCES

- Bahé Y. M., et al., 2017, MNRAS, 470, 4186
- Benítez-Llambay A., Frenk C., 2020, MNRAS, 498, 4887
- Benítez-Llambay A., Navarro J. F., Frenk C. S., Ludlow A. D., 2018, MNRAS, 473, 1019
- Binney J., Knebe A., 2002, MNRAS, 333, 378

- Binney J., Tremaine S., 2008, *Galactic Dynamics: Second Edition*. Princeton University Press
- Buck T., Obreja A., Macciò A. V., Minchev I., Dutton A. A., Ostriker J. P., 2020, *MNRAS*, 491, 3461
- Casagrande L., et al., 2016, *MNRAS*, 455, 987
- Chandrasekhar S., 1960, *Principles of stellar dynamics*
- Clauwens B., Schaye J., Franx M., Bower R. G., 2018, *MNRAS*, 478, 3994
- Correa C. A., Wyithe J. S. B., Schaye J., Duffy A. R., 2015, *MNRAS*, 452, 1217
- Correa C. A., Schaye J., Clauwens B., Bower R. G., Crain R. A., Schaller M., Theuns T., Thob A. C. R., 2017, *MNRAS*, 472, L45
- Cui W., et al., 2018, *MNRAS*, 480, 2898
- Davé R., Thompson R., Hopkins P. F., 2016, *MNRAS*, 462, 3265
- Davé R., Anglés-Alcázar D., Narayanan D., Li Q., Rafieferantsoa M. H., Appleby S., 2019, *MNRAS*, 486, 2827
- Diemer B., Joyce M., 2019, *ApJ*, 871, 168
- Diemer B., Kravtsov A. V., 2015, *ApJ*, 799, 108
- Dolag K., Komatsu E., Sunyaev R., 2016, *MNRAS*, 463, 1797
- Du M., Ho L. C., Debattista V. P., Pillepich A., Nelson D., Zhao D., Hernquist L., 2020, *ApJ*, 895, 139
- Dubois, et al 2014, *MNRAS*, 444, 1453
- Dubois Y., Peirani S., Pichon C., Devriendt J., Gavazzi R., Welker C., Volonteri M., 2016, *MNRAS*, 463, 3948
- Dubois Y., et al., 2020, arXiv e-prints, p. arXiv:2009.10578
- Furlong M., et al., 2017, *MNRAS*, 465, 722
- Garrison-Kimmel S., Boylan-Kolchin M., Bullock J. S., Lee K., 2014, *MNRAS*, 438, 2578
- Grand R. J. J., Springel V., Gómez F. A., Marinacci F., Pakmor R., Campbell D. J. R., Jenkins A., 2016, *MNRAS*, 459, 199
- Grand R. J. J., et al., 2017, *MNRAS*, 467, 179
- Guedes J., Callegari S., Madau P., Mayer L., 2011, *ApJ*, 742, 76
- Harris C. R., et al., 2020, *Nature*, 585, 357–362
- Henon M., 1973, in Contopoulos G., Henon M., Lynden-Bell D., eds, *Saas-Fee Advanced Course 3: Dynamical Structure and Evolution of Stellar Systems*. p. 183
- Hernquist L., 1990, *ApJ*, 356, 359
- Hopkins P. F., et al., 2018, *MNRAS*, 480, 800
- Huang K.-H., et al., 2017, *ApJ*, 838, 6
- Hunter J. D., 2007, *Computing in Science & Engineering*, 9, 90
- Khandai N., Di Matteo T., Croft R., Wilkins S., Feng Y., Tucker E., DeGraf C., Liu M.-S., 2015, *MNRAS*, 450, 1349
- Kravtsov A. V., 2013, *ApJ Lett.*, 764, L31
- Lacey C. G., Ostriker J. P., 1985, *ApJ*, 299, 633
- Libeskind N. I., Yepes G., Knebe A., Gottlöber S., Hoffman Y., Knollmann S. R., 2010, *MNRAS*, 401, 1889
- Libeskind N. I., et al., 2020, *MNRAS*, 498, 2968
- Ludlow A. D., Navarro J. F., Angulo R. E., Boylan-Kolchin M., Springel V., Frenk C., White S. D. M., 2014, *MNRAS*, 441, 378
- Ludlow A. D., Bose S., Angulo R. E., Wang L., Hellwing W. A., Navarro J. F., Cole S., Frenk C. S., 2016, *MNRAS*, 460, 1214
- Ludlow A. D., et al., 2017, *Phys. Rev. Lett.*, 118, 161103
- Ludlow A. D., Schaye J., Schaller M., Richings J., 2019a, *MNRAS*, 488, L123
- Ludlow A. D., Schaye J., Bower R., 2019b, *MNRAS*, 488, 3663
- Ludlow A. D., Schaye J., Schaller M., Bower R., 2020, *MNRAS*, 493, 2926
- McCarthy I. G., Schaye J., Bird S., Le Brun A. M. C., 2017, *MNRAS*, 465, 2936
- Navarro J. F., Frenk C. S., White S. D. M., 1996, *ApJ*, 462, 563
- Navarro J. F., et al., 2018, *MNRAS*, 476, 3648
- Nelson D., et al., 2018, *MNRAS*, 475, 624
- Pérez F., Granger B. E., 2007, *Computing in Science and Engineering*, 9, 21
- Pillepich A., et al., 2019, arXiv e-prints,
- Posti L., Fall S. M., 2021, arXiv e-prints, p. arXiv:2102.11282
- Power C., Navarro J. F., Jenkins A., Frenk C. S., White S. D. M., Springel V., Stadel J., Quinn T., 2003, *MNRAS*, 338, 14
- Quillen A. C., Garnett D. R., 2001, in Funes J. G., Corsini E. M., eds, *Astronomical Society of the Pacific Conference Series Vol. 230, Galaxy Disks and Disk Galaxies*. pp 87–88
- Revaz Y., Jablonka P., 2018, *A&A*, 616, A96
- Sawala et al. 2016, *MNRAS*, 457, 1931
- Schaye J., et al., 2010, *MNRAS*, 402, 1536
- Schaye et al. 2015, *MNRAS*, 446, 521
- Sellwood J. A., 2013, *ApJ Lett.*, 769, L24
- Springel V., 2005, *MNRAS*, 364, 1105
- Springel V., Di Matteo T., Hernquist L., 2005, *MNRAS*, 361, 776
- Springel V., et al., 2018, *MNRAS*, 475, 676
- Thob et al 2019, *MNRAS*, 485, 972
- Tinker J., Kravtsov A. V., Klypin A., Abazajian K. a., 2008, *ApJ*, 688, 709
- Trayford J. W., et al., 2017, *MNRAS*, 470, 771
- Trayford J. W., Frenk C. S., Theuns T., Schaye J., Correa C., 2019, *MNRAS*, 483, 744
- Tremmel M., Karcher M., Governato F., Volonteri M., Quinn T. R., Pontzen A., Anderson L., Bellovary J., 2017, *MNRAS*, 470, 1121
- Virtanen P., et al., 2020, *Nature Methods*, 17, 261
- Vogelsberger M., et al., 2014, *MNRAS*, 444, 1518
- Wetzel A. R., Hopkins P. F., Kim J.-h., Faucher-Giguère C.-A., Kereš D., Quataert E., 2016, *ApJ Lett.*, 827, L23
- Wielen R., 1977, *A&A*, 60, 263
- Yurin D., Springel V., 2014, *MNRAS*, 444, 62
- van de Sande, et al 2019, *MNRAS*, 484, 869

APPENDIX A: JUSTIFICATION FOR SEVERAL ASSUMPTIONS AND CHOICES USED IN THIS PAPER

A1 $f_{\star} = 0.01$

Because the collisional heating rate anticipated by eq 16 depends only on properties of the DM halo, it should be relatively insensitive to the stellar distribution, and in particular (within reason) to the stellar mass fraction. All of the simulations analyzed in this paper adopted a stellar to *total* mass fraction of $f_{\star} \equiv M_{\star}/M_{200} = 0.01$. This value is relatively low, particularly for the interesting case of haloes with masses $\approx 10^{12} M_{\odot}$, for which $f_{\star} \gtrsim 0.03$. We nevertheless adopt $f_{\star} = 0.01$ since, for each of the halo models we consider, it ensures that the disk is gravitationally sub-dominant at all radii, and has a negligible impact on the dynamics and spatial distribution of halo particles. For studies such as ours, larger values of $f_{\star}$ are undesirable for a couple of reasons: massive disks 1) may lead to a contraction of the inner halo, and 2) are vulnerable to instabilities, which rapidly alter the disk structure. Both of these effects complicate the interpretation of the simulation results and the development of the empirical model described in Section 4.2. Collisional heating of more massive disks will be studied in future work.

The first point above is shown explicitly in Figure A1, where we plot the circular velocity profiles due to DM (blue) and stars (red) for a few example halo/galaxy models. The left-hand panel shows a DM halo with concentration $c = 10$ (our fiducial choice) and stellar disks with $f_{\star} = 0.01$ (also our fiducial choice, shown using solid red lines) and $f_{\star} = 0.02$ (dashed red lines). In both cases, thin and thick lines, respectively, distinguish the initial profiles (shown here using the analytic profiles used to construct each model) from those measured in the simulations after $\Delta t = 9$ Gyr. Note that the more massive disk significantly alters the mass profile of the DM halo on spatial scales $\lesssim R_{1/2}$ (outsized circles mark the location of the *initial* disk half-mass radius). This may lead to difficulties interpreting the underlying heating rates anticipated by eq. 16 since the local density of dark matter is no longer independent of the properties of the disk.

The right-hand panel shows analogous results for our fiducial disk (i.e. $f_{\star} = 0.01$ and f_j adjusted to ensure a fixed stellar mass profile) but for DM haloes with $c = 7$ (dashed lines), 10 (solid) and 15 (dot-

Figure A1. Circular velocity profiles due to DM (blue curves) and stars (red) for models adopting $V_{200} = 200 \text{ km s}^{-1}$ and $m_{\text{DM}} = 10^{7.5} M_{\odot}$. The left-hand panel corresponds to a DM halo with concentration $c = 10$ and spin parameter $\lambda_{\text{DM}} = 0.03$. Two sets of red curves correspond to circular velocity profiles of disk stars (both disks have $f_j = 1$) for $f_* = 0.01$ (lighter colour, solid lines) or $f_* = 0.02$ (darker colour, dashed lines). Thin lines show the initial profiles for each component and thick lines the corresponding profiles after $\Delta t = 9 \text{ Gyr}$ of evolution. Note that $f_* = 0.02$ leads to a significant contraction of the DM halo within the initial stellar half-mass radius (shown as outsized circles), whereas $f_* = 0.01$ leaves the halo’s mass profile unaltered. The right-hand panel, similarly, shows the circular velocity profiles due to stars ($f_* = 0.01$; red curves) and the surrounding DM halo for varying concentration parameters ($c = 7, 10$ and 15 ; note that for each value of c we have adjusted f_j to ensure that the stellar mass profile remains fixed). As on the left, the initial profiles are plotted using thin lines and the final profiles using thick lines. Regardless of the DM halo’s concentration, the $f_* = 0.01$ stellar disk does not affect its mass profile.

dashed). In none of these cases does the presence of the stellar disk alter the distribution of dark matter.

A2 Limitations due to collisional relaxation of the DM halo

In cosmological DM-only simulations, the radius at which the local collisional relaxation time is of order the age of the universe, t_{H} , i.e. where $\kappa(r) \equiv t_{\text{relax}}(r)/t_{\text{H}} \approx 1$, marks the well-known “convergence radius” of a simulated halo. Specifically, circular velocity profiles of haloes converge (to better than $\lesssim 10$ per cent) to those of arbitrarily-high mass and force resolution provided $\kappa \simeq 0.6$ (for *individual* haloes); see Power et al. 2003) or $\kappa \simeq 0.18$ (for *median* mass profiles; see Ludlow et al. 2019b). For radii $r \lesssim r_{\text{conv}}$, the collisional self-heating of dark matter particles thermalizes their velocity distribution and suppresses the internal density. It is desirable to test the validity of eq. 16 at radii deemed “converged” by similar arguments in order to ensure the local DM density does not develop an (unknown and unwanted) time dependence. But what value of κ is relevant for our idealized runs? What determines the innermost resolved radius?

Figure A2 provides some clues. Here we plot the measured convergence radii for stars (connected squares) and DM (circles) in the two lowest-resolution realisations of our fiducial galaxy/halo model (i.e. $V_{200} = 200 \text{ km s}^{-1}$, $f_* = 0.01$, $\mu = 5$). We calculate r_{conv} by locating the radius beyond which the $V_c(r)$ profiles converge to $\lesssim 10$ per cent relative to the *initial* profiles. The convergence radius grows with time, as expected, and can be described reasonably well assuming $\kappa_* = 1.7$ (for stars) and $\kappa_{\text{DM}} = 0.08$ for DM (note that we have assumed $t_{\text{relax}} \equiv N_{\text{DM}}/(8 \ln N_{\text{DM}})$, such that relaxation is driven by the DM particles; t_{H} is the current age of the universe). These results depend slightly on μ (not shown), but primarily on N_{DM} . For the purposes of this paper it suffices to say that, for DM, $r_{\text{conv}} \lesssim R_{1/4}$ even at the lowest (mass) resolution studied (horizontal dotted lines mark the initial values of $R_{1/4}$ and $R_{1/2}$). We therefore expect the

impact of collisional *self*-heating of the DM halo to be negligible at $r \gtrsim R_{1/4}$ in all models, which is the *minimum* radius we considered in our analysis. Hence, our analysis is robust to collisional heating in the DM component.

Figure A2 also reveals that the convergence radius of stars exceeds that of the DM by nearly a factor of 5, but it is premature (and unlikely correct) to assume that this is a generally-valid result. For example, it is *not* due to mass segregation: it is approximately valid for $\mu = 1$ as well. It is more likely that this is a general manifestation of energy diffusion associated with equipartition: in our model, stars form a dynamically-cold disk in the centre of a comparatively *hot* halo: stars, on average, *gain* energy through collisions with DM particles, causing them to more readily flow from the central regions of haloes outward. It is therefore likely that the convergence radii of simulated galaxies depend not only on the distribution and number density of DM particles, but also on μ , and on the initial phase-space distribution of stellar particles (i.e. on their initial segregation with respect to the DM, which will depend on the initial scale height, angular momentum, mass fraction, etc). We conclude that the convergence radii of galaxies generally exceed those of their DM haloes.

APPENDIX B: ASYMPTOTIC DEPENDENCE OF VERTICAL AND RADIAL HEATING RATES ON DARK MATTER-TO-STELLAR PARTICLE MASS RATIO, μ

The collisional heating of stellar disks reported in this paper is largely insensitive to the dark matter-to-stellar particle mass ratio, $\mu = m_{\text{DM}}/m_*$: at fixed m_{DM} , similar results were obtained for all values of m_* considered. As discussed in Section 2.2, complete equipartition of energy should drive the stellar velocity dispersion asymptotically to $\sigma_i \approx \sqrt{\mu} \sigma_{\text{DM}}$; in practice, the virial theorem imposes an asymptotic limit of $\sigma_i \approx \sqrt{2} \sigma_{\text{DM}}$. The latter limit is applicable to runs with $\mu \geq 2$, the former for those with $\mu < 2$ (see eq. 7).

Figure A2. Convergence radii (r_{conv} ; where simulated $V_c(r)$ profiles first deviate by $\lesssim 10$ per cent from their initial profile) for our fiducial model ($V_{200} = 200 \text{ km s}^{-1}$, $f_* = 0.01$, $\mu = 5$) run with $m_{\text{DM}} = 10^{7.5} M_\odot$ (green) and $m_{\text{DM}} = 10^8 M_\odot$ (blue). Results are shown separately for the stellar disk (connected squares) and for the DM halo (circles). Dashed and solid lines show the radii at which the relaxation time of the stars or DM halo (in units of the current age of the universe) are $\kappa_* = 1.7$ and $\kappa_{\text{DM}} = 0.08$, respectively (see text for details). The horizontal dotted lines mark the initial characteristic radii $R_{1/4}$ and $R_{1/2}$. Note that the convergence radius of the DM halo does not exceed $R_{1/4}$, even in our lowest-resolution run (corresponding to $m_{\text{DM}} = 10^8 M_\odot$ for the fiducial runs shown here).

Whether full energy equipartition will be reached, however, depends on a number of factors, including μ , the *total* relative masses of each component (i.e. $f_*^{-1} - 1$) as well as the number of particles in each, their initial kinematic and spatial segregation, and the duration of the simulation, Δt .

The main results of our study, presented in Section 4, were based on simulations carried out for a total of $\Delta t = 9.6 \text{ Gyr}$. In Figure A3, we present results for three of our fiducial runs (i.e. $V_{200} = 200 \text{ km s}^{-1}$, $\lambda_{\text{DM}} = 0.03$, $f_* = 0.01$, $c = 10$), extended to $\Delta t = 100 \text{ Gyr}$. Upper and lower panels plot σ_z^2 and σ_R^2 , respectively (measured at the radius $R_{1/4}$ that encloses a quarter of the initial stellar mass). The DM particle mass is $m_{\text{DM}} = 10^8 M_\odot$ and $\mu = 1$ (blue solid lines), 5 (red dot-dashed) and 25 (green dashed). The relatively high DM particle mass is chosen in order to hasten collisional heating.

The kinematics of disk stars undergo an initial phase of very rapid heating, lasting roughly $\Delta t \approx 10 \text{ Gyr}$. At later times the effects of collisional heating taper off, and the stellar velocity dispersion obtained from the various models approach the asymptotic values expected from eq. 7, namely $\sigma_i \approx \sigma_{\text{DM}}$ for $\mu = 1$ and $\sigma_i \approx \sqrt{2} \sigma_{\text{DM}}$ for $\mu = 5$ and 25. These results confirm our expectations for the asymptotic velocity dispersion based on energy equipartition and the virial theorem.

The collisional heating of our galaxy disks is driven by two processes: potential fluctuations due to shot noise in the particle distribution, and by the gradual exchange of energy between particles of unequal mass. The former is present in all simulations, regardless of μ ; the latter only affects runs for which $\mu \neq 1$. Both processes give rise to the spurious growth of galaxy sizes in the EAGLE simulations

Figure A3. The evolution of σ_z^2 (upper panel) and σ_R^2 (lower panel) measured at $R_{1/4}$ for our fiducial model (see Section 3.1) with $m_{\text{DM}} = 10^8 M_\odot$ and for three values of μ . Results are shown for $\Delta t = 100 \text{ Gyr}$ to emphasize the asymptotic behaviour of the velocity dispersion of stellar particles. Horizontal lines mark two characteristic values: σ_{DM}^2 , the 1-dimensional dark matter velocity dispersion (squared); and $v_{\text{esc}}^2/V_{200}^2 = 2 \times \sigma_{\text{DM}}^2$, the local escape velocity for a system in virial equilibrium. Note that a simulated stellar system in energy equipartition with its dark matter halo would have $\sigma_*^2 = \mu \times \sigma_{\text{DM}}^2$, which can only be achieved for $\mu \leq 2$; for larger μ , the transfer of energy from dark matter to stellar particles leads to their slow evaporation from the system, such that, regardless of μ , the maximum local velocity dispersion of stars is approximately $\sqrt{2} \times \sigma_{\text{DM}}$.

reported by Ludlow et al. (2019a), but it is the former that dominates the disk heating effect presented in this paper.

APPENDIX C: COLLISIONAL HEATING DUE TO STELLAR HALOES AND BULGES

In Section 4 we established that spurious collisional heating of thin stellar disks by coarse-grained DM haloes is cause for concern in simulations of galaxy formation. Realistic simulations, however, will produce galaxies with a wide range of morphologies, even when limited to the disk galaxy population: disks are often partnered with bulges or stellar haloes, for example. Are star particles in these components also able to collisionally excite stellar disks? What about collisions between disk stars?

We investigate this in Figure C1, where we plot the evolution of the vertical and radial velocity dispersion (upper and middle panels, respectively) and the half-mass scale height (lower panels) obtained

Figure C1. Contribution to collisional heating due to dark matter and stellar haloes. Top panels plot the evolution of the vertical velocity dispersion, σ_z^2/V_{200}^2 , middle panels the radial velocity dispersion, σ_R^2/V_{200}^2 , and bottom panels the vertical half-mass scale height, $z_{1/2}/r_{200}$. All quantities are measured at the *initial* stellar half-mass radius of the disk, $R_{1/2}$. Different colours correspond to different galaxy/halo configurations: black to a live dark matter halo and a live thin disk; yellow to a live DM halo, live disk and live stellar halo; blue to a static DM halo, live disk and live stellar halo; green to a static dark matter halo and live disk. All models correspond to our fiducial halo ($V_{200} = 200 \text{ km s}^{-1}$, $\lambda = 0.03$) and disk ($f_{\text{disk}} = 0.01$), but for the *stellar* halo we consider mass fractions of $f_{\text{halo}} = 0.01$ (left panels) and $f_{\text{halo}} = 0.03$ (right panels). Different line-styles correspond to DM haloes sampled with particles of mass $m_{\text{DM}} = 10^7 M_{\odot}$ (solid) and $10^8 M_{\odot}$ (dash-dotted). Note that, in all cases, the heating and structural evolution of the disk is driven primarily by the presence of a coarse-grained DM halo.

from a number of supplementary simulations. All models adopt a Hernquist profile for the DM (the NFW-equivalent halo has a concentration $c = 10$ and $V_{200} = 200 \text{ km s}^{-1}$) and $\lambda_{\text{DM}} = 0.03$. As for the other models considered in this paper, the disk contains a fraction $f_{\star} = 0.01$ of the system’s total mass, and the disk-to-halo specific angular momentum ratio is $f_j = 1$. Different line styles correspond to different DM particle masses, as indicated; the stellar particle masses are chosen so that $\mu = 5$ in all cases. Lines of different colour denote different simulations, as follows. Black lines correspond to our fiducial model: a coarse-grained DM halo and a thin, rotationally-supported stellar disk with parameters described above. Yellow lines are the same, but with an additional stellar bulge/halo component with mass fractions equal to $f_{\text{halo}} = 0.01$ (left) and 0.03 (right). Note that the spheroidal stellar component is also modelled as a Hernquist sphere with a characteristic scale radius equal to one-tenth that of the DM halo’s. Comparing these curves, we conclude

that collisional heating by stellar haloes or bulges is negligible compared to that due to the DM halo.

In order to verify these results, and to test the impact of collisions between stars *in the disk*, we have repeated these two sets of simulations, but after replacing the “live” DM halo with a fixed Hernquist potential of equivalent mass and characteristic size. The results are plotted as blue and green lines for the cases with and without a “live” stellar halo, respectively. In this case, the heating rate drops considerably, and remains largely independent of the presence or absence of a coarse-grained stellar halo/bulge. We have verified that these results apply to a relatively broad range of stellar spheroidal structures, ranging from compact bulges to extended stellar haloes.

Finally, note that spurious collisional heating is still present, albeit much reduced, even when the DM halo is modelled using a fixed analytic potential. This suggests that collisional heating due to star-star encounters also affects simulations of galaxy structure (in agreement with Sellwood 2013).

APPENDIX D: THE IMPACT OF THE GRAVITATIONAL SOFTENING LENGTH

The simulations presented in this paper and all previous Appendices used a gravitational softening length equal to the initial scale-height of the stellar disk, which is a fixed fraction of the disk scale radius, i.e. $\epsilon = z_d = 0.05 R_d$. As discussed in Section 3.1, this has the advantage of preserving the scale-invariance of our runs, since the softening length scales self-consistently with the size of the disk in models with widely varying V_{200} and R_d . However, as discussed by Ludlow et al. (2020), the softening length adopted in hydrodynamical simulations of galaxy formation has a non-trivial relationship to galaxy structure: if it is too large, it suppresses the small-scale clustering of stellar and DM particles, resulting in lower central densities, larger galaxies, and DM haloes with softened “cores”; if it is too *small*, it exacerbates the collisional relaxation of stellar structures, escalating the spurious growth of galaxies sizes. Similar softening-dependent effects may infiltrate the results present in this work.

We explore the softening dependence of collisional disk heating for a couple of our fiducial models in Figure D1 ($V_{200} = 200 \text{ km s}^{-1}$, $c = 10$, $\lambda_{\text{DM}} = 0.03$; $f_\star = 0.01$, $f_{\text{disk}} = 0.01$, $\mu = 5$). We focus on the radial profiles of the vertical velocity dispersions ($\sigma_z(R)$, upper panels), radial velocity dispersion ($\sigma_R(R)$, middle) and vertical half-mass scale-height ($z_{1/2}$, bottom). Results are plotted at $t = 0$ (left columns) and after $\Delta t = 2 \text{ Gyr}$ (middle) and $\Delta t = 9 \text{ Gyr}$ (right). Line styles indicate the mass of DM particles (solid for $m_{\text{DM}} = 10^7 M_\odot$; dashed for $m_{\text{DM}} = 10^8 M_\odot$), and different coloured lines correspond to runs carried out with different softening lengths, ranging from $\epsilon/z_d = 0.25$ to 4 (physical values of ϵ are marked using arrows in each panel).

The plot elicits a few comments. First, despite the wide range of softening lengths considered, the consequences of collisional heating are apparent in all simulations, and at all radii. Nevertheless, the integrated effects are suppressed in runs in which the softening is largest, particularly in the galaxy’s central regions in runs with $\epsilon > z_d$ (see Ludlow et al. 2020 and Pillepich et al. 2019 for similar results based on cosmological simulations).

Importantly, however, the results are largely independent of ϵ provided $\epsilon \lesssim z_d$, i.e. provided softening length does not exceed the scale height of the disk. This implies that – in cosmological simulations, which typically adopt softening lengths that are fixed in physical or comoving coordinates – collisional disk heating may be suppressed in low-mass, poorly-resolved galaxies, whose characteristic sizes can be comparable to the softening length. Clearly such systems can, however, not be considered spatially resolved in the first place.

Despite the tendency for collisional heating to be suppressed when $\epsilon \gtrsim z_0$, the relation between $z_{1/2}$ and σ_z appears largely independent of the softening length. As a result, the scale heights of simulated disks can still be inferred from their vertical velocity dispersion profiles. We show this explicitly in Figure D2 (which is analogous to Figure 5 in the main body of the paper), where we plot the vertical scale height of stellar particles (normalized by r_{200}) versus their vertical velocity dispersion (normalized by V_{200}) at different galacto-centric radii ($R_{1/4}$, $R_{1/2}$ and $R_{3/4}$ from left to right, respectively; R_f is the cylindrical radius enclosing a fraction f of the galaxy’s initial stellar mass). The results, which correspond to the same set of runs plotted in Figure D1, clearly show that, despite the softening dependence of $z_{1/2}$ and σ_z , the relation between these two quantities (at fixed R_f) is independent of softening.

Figure D1. The effect of gravitational softening on the radial profiles of the vertical (top panels) and radial (middle panels) velocity dispersion of stellar particles for our fiducial galaxy/halo models. Bottom panels show the radial dependence of the vertical half-mass scale height. Panels on the left correspond to $t = 0$, i.e. the initial conditions of our simulations, middle panels to $\Delta t = 2$ Gyr, and right-most panels to $\Delta t = 9$ Gyr (the initial profiles in the highest-resolution run are shown for comparison using a thin black line in all panels). In all cases, profiles are plotted to the innermost radius that encloses at least 10 stellar particles. All runs adopt $V_{200} = 200 \text{ km s}^{-1}$, $f_j = 0.01$, and $f_{\text{disk}} = 0.01$, but were repeated for $m_{\text{DM}} = 10^7 M_\odot$ (solid curves) and $m_{\text{DM}} = 10^8 M_\odot$ (dashed curves). Different colours correspond to different softening lengths ϵ (indicated by arrows), which vary from $\epsilon = z_0/4$ to $\epsilon = 4 \times z_0$ (our fiducial value is $\epsilon = z_0$). Vertical dotted lines mark the initial characteristic radii $R_{1/4}$, $R_{1/2}$ and $R_{3/4}$, from left to right, respectively. Measured scale heights, $z_{1/2}(R)$, enclose half of the stellar mass relative to the disk mid-plane. Collisional heating rates are suppressed when ϵ is large, but are independent of softening provided $\epsilon/z_d \lesssim 1$.

Figure D2. Half-mass vertical scale height of stellar particles (normalized by r_{200}) plotted as a function of their vertical velocity dispersion (squared, and normalized by V_{200}^2). From left to right, different panels correspond to measurements made at cylindrical radii corresponding to $R_{1/4}$, $R_{1/2}$ and $R_{3/4}$, respectively, where R_f is the radius enclosing a fraction f of the galaxy's initial stellar mass. Simulation results are plotted as individual points for the same fiducial models shown in Figure D1; points are color-coded by gravitational softening length, ϵ , with different symbols used for runs with different DM particle mass, as indicated. The dashed orange line shows the square harmonic mean of the vertical scale heights expected for thin self gravitating (SG) and non-self gravitating (NSG) disks, i.e. eq. 18; the solid blue line shows the $z_{1/2} - \sigma_z$ relation for thick disks, i.e. eq. 21. Note that although both $z_{1/2}$ and σ_z depend on ϵ (most noticeably when $\epsilon \gtrsim z_0$; Figure D1), the relation between them does not. The vertical scale heights of simulated disks can therefore be estimated from their vertical velocity dispersion profiles independently of the gravitational softening length.