



Universiteit  
Leiden  
The Netherlands

## Optical and infrared phase curves of the lava planet K2-141 b

Zieba, S.; Zilinskas, M.; Kreidberg, L.; Cowan, N.; Nguyen, G.; Miguel, Y.; ... ; Carone, L.

### Citation

Zieba, S., Zilinskas, M., Kreidberg, L., Cowan, N., Nguyen, G., Miguel, Y., ... Carone, L. (2021). Optical and infrared phase curves of the lava planet K2-141 b. *Epsc Abstracts*, EPSC2021-476. doi:10.5194/epsc2021-476

Version: Publisher's Version

License: [Creative Commons CC BY 4.0 license](#)

Downloaded from: <https://hdl.handle.net/1887/3275512>

**Note:** To cite this publication please use the final published version (if applicable).



## Optical and Infrared Phase Curves of the Lava Planet K2-141 b

**Sebastian Zieba**<sup>1,2</sup>, Mantas Zilinskas<sup>2</sup>, Laura Kreidberg<sup>1</sup>, Nicolas Cowan<sup>3,4,5</sup>, Giang Nguyen<sup>6</sup>, Yamila Miguel<sup>2</sup>, Raymond Pierrehumbert<sup>7</sup>, Roxana Lupu<sup>8</sup>, Lisa Dang<sup>3,4</sup>, Mark Hammond<sup>9</sup>, Luca Malavolta<sup>10</sup>, and Ludmila Carone<sup>1</sup>

<sup>1</sup>Max-Planck-Institut für Astronomie, Königstuhl 17, 69117 Heidelberg, Germany (zieba@mpia.de)

<sup>2</sup>Leiden Observatory, Leiden University, Niels Bohrweg 2, 2333CA Leiden, The Netherlands

<sup>3</sup>Department of Physics, McGill University, 3600 rue University, Montréal, QC H3A 2T8, Canada

<sup>4</sup>McGill Space Institute; Institute for Research on Exoplanets; Centre for Research in Astrophysics of Quebec

<sup>5</sup>Department of Earth & Planetary Sciences, McGill University, 3450 rue University, Montréal, QC H3A 0E8, Canada

<sup>6</sup>Centre for Research in Earth and Space Sciences, York University, 4700 Keele st., Toronto, Ontario, M3J 1P3, Canada

<sup>7</sup>Department of Physics (Atmospheric, Oceanic and Planetary Physics), University of Oxford, Parks Rd, Oxford, OX1 3PU, UK

<sup>8</sup>BAER Institute, NASA Ames Research Center, Naval Air Station, Moffett Field, CA 94035, USA

<sup>9</sup>Department of the Geophysical Sciences, University of Chicago, USA

<sup>10</sup>Dipartimento di Fisica e Astronomia "Galileo Galilei", Università di Padova, Padua, Italy

K2-141 b is a transiting, small (1.5 RE) Ultra-Short-Period (USP) planet orbiting its star every 6.7 hours discovered by the Kepler space telescope. The planet's high surface temperature of more than 2000 K makes it an excellent target for atmospheric studies by the observation of its thermal emission. We present 65 hours of continuous photometric observations of K2-141 b collected with Spitzer's IRAC Channel 2 at 4.5 microns spanning 10 full phases of the orbit. Our best fit model of the Spitzer data shows no significant offset of the thermal hotspot and is inconsistent with the observed offset of the well-studied USP planet 55 Cnc e at a 3.7 sigma level. We measure an eclipse depth of  $142 \pm 40$  ppm and an amplitude variation of  $120 \pm 40$  ppm in the infrared. The joint analysis of the observations collected in the two photometric bands favors a non-zero geometric albedo with  $A_g = 0.26 \pm 0.07$  and a tentative temperature gradient. With a dayside temperature of  $2141 - 361 + 352$  K and a night-side temperature of  $1077 - 623 + 473$  K we also find no evidence of heat redistribution on the planet. We compare the observations to a 1D rock vapor model and a 1D circulation toy model and argue that the data are best explained by a thin rock vapor atmosphere with a thermal inversion.