



Universiteit
Leiden
The Netherlands

Food insecurity, dietary quality and health in the Netherlands

Velde, L.A. van der

Citation

Velde, L. A. van der. (2022, March 1). *Food insecurity, dietary quality and health in the Netherlands*. Retrieved from <https://hdl.handle.net/1887/3277065>

Version: Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/3277065>

Note: To cite this publication please use the final published version (if applicable).

CHAPTER 4

Needs and perceptions regarding healthy eating among people at risk of food insecurity: a qualitative analysis

Laura A. van der Velde, Linde A. Schuilenburg, Jyothi K. Thrivikraman, Mattijs E. Numans, Jessica C. Kiefte-de Jong

Published in the *International Journal for Equity in Health*.
2019 Nov 27;18(1):184.

Abstract

Background: Healthy eating behavior is an essential determinant of overall health. This behavior is generally poor among people at risk of experiencing food insecurity, which may be caused by many factors including perceived higher costs of healthy foods, financial stress, inadequate nutritional knowledge, and inadequate skills required for healthy food preparation. Few studies have examined how these factors influence eating behavior among people at risk of experiencing food insecurity. We therefore aimed to gain a better understanding of the needs and perceptions regarding healthy eating in this target group.

Methods: We conducted a qualitative exploration grounded in data using inductive analyses with 10 participants at risk of experiencing food insecurity. The analysis using an inductive approach identified four core factors influencing eating behavior: Health related topics; Social and cultural influences; Influences by the physical environment; and Financial influences.

Results: Overall, participants showed adequate nutrition knowledge. However, eating behavior was strongly influenced by both social factors (e.g., child food preferences and cultural food habits), and physical environmental factors (e.g., temptations in the local food environment). Perceived barriers for healthy eating behavior included poor mental health, financial stress, and high food prices. Participants had a generally conscious attitude towards their financial situation, reflected in their strategies to cope with a limited budget. Food insecurity was mostly mentioned in reference to the past or to others and not to participants' own current experiences. Participants were familiar with several existing resources to reduce food-related financial strain (e.g., debt assistance) and generally had a positive attitude towards these resources. An exception was the Food Bank, of which the food parcel content was not well appreciated. Proposed interventions to reduce food-related financial strain included distributing free meals, facilitating social contacts, increasing healthy food supply in the neighborhood, and lowering prices of healthy foods.

Conclusion: The insights from this study increase understanding of factors influencing eating behavior of people at risk of food insecurity. Therefore, this study could inform future development of potential interventions aiming at helping people at risk of experiencing food insecurity to improve healthy eating, thereby decreasing the risk of diet-related diseases.

Background

Healthy eating behavior is an essential determinant of overall health. Previous literature extensively shows that people with lower socioeconomic status (SES) generally exhibit less healthy eating behaviors (1) and have increased risk of obesity and related illnesses (2, 3). The same holds for people experiencing food insecurity (4-6), which is an inadequate physical and economic access to adequate food that meets dietary needs and food preferences (7). The concept of food insecurity is closely related to lower SES, although this is a complex relationship and people with lower SES do not always experience food insecurity and vice versa (8). However, it is evident that food insecurity is more common among people with lower SES and therefore people with lower SES or living in disadvantaged neighborhoods have an increased risk of experiencing food insecurity (9).

Thus far, knowledge on food insecurity in Europe is limited (10). A previous study among Dutch Food Bank recipients found a food insecurity prevalence of almost 73 percent (11). Our recent study has shown that approximately one quarter of families living in disadvantaged neighborhoods in The Netherlands experienced food insecurity (12). Results of this study further showed that general health, diet quality, and weight were suboptimal, especially among food insecure participants. A possible intervention for reducing food insecurity is the Food Bank, but despite the high prevalence of food insecurity it was hardly used (12). The Dutch Food Bank is a non-governmental organization that distributes donated food to offer temporal food aid to people in need (13). This is done through providing food parcels, meant to supplement the usual diet, to eligible persons. Eligibility is based on household size-adjusted monthly disposable income. The food parcel content largely depends on donated foods and therefore varies per time and location of Food Bank. Recent research indicated that the parcel content was generally not in line with nutritional guidelines, which may contribute to suboptimal dietary intake among people eligible for Food Bank use (14).

Various factors may contribute to the generally suboptimal eating behavior among people at risk of experiencing food insecurity, including stress (15-17), inadequate knowledge and skills regarding healthy eating and food preparation (18), and higher costs of healthy foods (19). These higher costs might be an even more prominent issue than previously, since the Dutch Government recently increased taxes of all basic necessities such as foods (including foods that are considered healthy like fruit

and vegetables) from 6 to 9 percent (20). This price increase may lead to less healthy eating behavior, as previous research shows that pricing affects food choices (21, 22).

Much uncertainty still exists about contributing factors to suboptimal eating behavior among people at risk of experiencing food insecurity. Improving insight is essential for developing targeted interventions to support this population, focused on improving healthy eating behavior and thereby decreasing diet-related disease risk. Therefore, we aimed to gain a better understanding of the needs and perceptions regarding healthy eating behavior of people at risk of experiencing food insecurity living in disadvantaged neighborhoods in the Netherlands.

Methods

Rationale and study sample

Participants were selected from a sample of 242 participants included in a cross-sectional study on food insecurity in disadvantaged neighborhoods in The Hague, The Netherlands (12). These neighborhoods were selected based on predefined criteria used by the Dutch Government to identify disadvantaged neighborhoods in the Netherlands (23). Participants lived in or near the preselected disadvantaged neighborhoods and had at least one child below the age of 18 years living at home. A detailed description of the methods and results of this study are described elsewhere (12). Participants who provided valid contact information were invited to take part in an interview. None of the participants that agreed to participate dropped out of the study. Reasons for refusing to participate included being too busy, thinking an interview of approximately 60 minutes was too long, and being or going on holiday. A convenience sample, taking into account the diversity of the study sample, of a total of 10 participants (either fathers or mothers, one parent per household) were interviewed. After those 10 interviews, thematic saturation was reached. Interviews were conducted between April and July 2018. Sociodemographic characteristics, food insecurity status and diet quality scores of the participants were previously assessed (12). Food insecurity status was assessed using the 18-item United States Department of Agriculture (USDA) Household Food Security Survey Module. Affirmative responses to the questions (described in **Additional Table 1**) were summed and resulted in a continuum of food insecurity status ranging from 0-18, categorized as 'food secure' (0-2 affirmative responses), and 'food insecure' (≥ 3 affirmative responses), according to the USDA standards (24, 25). Dietary intake was assessed using the Dutch Healthy

Diet Food Frequency Questionnaire (DHD-FFQ) (26). Based on this dietary intake data we constructed a food group-based 6-component diet quality score (**Additional Table 2**). Each component score reflected the adherence to the dietary guidelines of the concerning food group. Component scores were summed to obtain the total diet quality score (range 0-60), with higher scores indicating a better diet quality. Written informed consent was obtained from all participants. Participants received a financial compensation of 10 euros for their effort and any travel expenses were refunded. The study was approved by the Medical Ethics Committee of Leiden University Medical Centre (P17.164).

Study design

Face-to-face open interviews were conducted, guided by a topic list (**Additional Table 3**). The topic list was created at the start of the study based on issues raised in the previous study (12) and consisted of topics to discuss and open ended example questions for each topic to guide the interviewer. These topics and example questions were discussed within the research team. The interviews started with general questions concerning participants' background, family, and living conditions to make the participant feel at ease, followed by questions focusing on perceptions regarding healthy eating, including knowledge; skills; external, social, and cultural influences; health; finances; stress; environmental factors; opinions about eating on a low budget; existing resources; and Food Bank use. Interviewees were also free to introduce other topics that were of interest to them. The topic list was merely used as guidance during the interviews and was re-evaluated after each interview and if appropriate adjusted or complemented with new topics that emerged during the interview. During the interviews, two members of the study team were present; one of them conducted the interview and the other observed. All interviews were audio-recorded with participants' permission using a digital voice recorder and transcribed verbatim. Participants were interviewed at a time and place that was most convenient to them. Interviews were held for 22 to 76 minutes with an average interview time of 47 minutes.

Analysis

We used a general inductive approach to analyze the data (27). Segments of the interview texts in the transcripts were coded using open coding, i.e., codes were built and modified throughout the coding process. Some text segments were assigned to

more than one code category and text segments that were not relevant for the study objectives were not included in any category. During the process, some of the codes were merged with other codes that had a similar meaning, resulting in 79 codes. One researcher coded the interviews. A second researcher coded two randomly selected interviews to check inter-rater reliability (IRR) (28), calculated as:

$$IRR = \frac{\text{number of agreements}}{\text{number of agreements} + \text{disagreements}}$$

We found an IRR of 93%.

Codes were grouped into subthemes, which were then grouped into main themes (29). Four main themes were identified that comprised the allocated codes for all transcripts. No new themes emerged towards the end of the study, suggesting thematic saturation was reached.

The software Atlas.ti version 7.5.6 (Scientific Software Development, Berlin) was used to assist the coding process and extraction of quotes and themes. The quotes presented in this paper were chosen based on their illustration of the described theme or clarifying role of the common or uncommon viewpoints.

Results

Two males and eight females were interviewed, aged between 35 and 55 years (**Table 1**). Most participants had an income below the basic needs budget and were lower educated. Six participants were single parents and half of the participants had a paid job. Participants had a Moroccan, Colombian, Surinamese, Curacao, or Polish migration background. Participants were all either overweight or obese, based on their self-reported height and weight. Seven participants were classified as food insecure. The four main themes related to healthy eating behavior and the corresponding subthemes that were identified in the analyses are described below and depicted in **Figure 1**.

Table 1 Sociodemographic characteristics of the participants (n=10)

Participant number	Age category in years	Sex	Educational level	Household income	Employment status	Marital status	Migration background	Food Bank use	BMI1 category	Diet quality score	Food security status
1	45-50	Male	ISCED 2	Below basic needs budget	Currently paid job	Two parent household	Moroccan	No	Overweight	36/ 60	Food insecure
2	40-45	Female	ISCED 2	Below basic needs budget	Currently no paid job	Single parent	Colombian	No	Overweight	31/ 60	Food insecure
3	45-50	Female	ISCED 2	Below basic needs budget	Currently no paid job	Single parent	Surinamese	No	Obese Class I (moderately obese)	29/ 60	Food insecure
4	40-45	Female	ISCED 5	Below basic needs budget	Currently paid job	Single parent	Surinamese	No	Overweight	33/ 60	Food secure
5	40-45	Female	ISCED 2	Below basic needs budget	Currently paid job	Single parent	Curacao	No	Obese Class I (moderately obese)	41/ 60	Food insecure
6	40-45	Male	ISCED 1	Below basic needs budget	Currently no paid job	Two parent household	Moroccan	Yes	Obese Class I (moderately obese)	35/ 60	Food insecure
7	35-40	Female	ISCED 4	Below basic needs budget	Currently no paid job	Two parent household	Polish	No	Obese Class I (moderately obese)	31/ 60	Food secure
8	50-55	Female	ISCED 1	Below basic needs budget	Currently no paid job	Single parent	Moroccan	No	Obese Class III (Very severely obese)	46/ 60	Food insecure
9	45-50	Female	ISCED 7	Above basic needs budget	Currently paid job	Two parent household	Surinamese	No	Overweight	32/ 60	Food secure
10	35-40	Female	ISCED 3	Above basic needs budget	Currently paid job	Single parent	Surinamese	No	Overweight	43/ 60	Food insecure

¹BMI: Body Mass Index




Figure 1. Main themes and their corresponding subthemes

Theme 1. Health related topics

Perceived healthy and unhealthy eating

Overall, participants demonstrated relatively good nutrition knowledge; adequate fresh fruit and vegetable intakes were perceived as essential components of a healthy diet. Snacks, fast-food, fatty foods, sugar, and overeating were considered unhealthy. Brown bread consumption was generally considered healthy, in contrast to white bread. Some participants indicated that bread consumption could lead to becoming overweight. Participants had conflicting opinions about whether meat consumption was healthy. Some participants considered meat as an essential component of a healthy diet, whereas others considered meat to be very unhealthy.

A frequently mentioned strategy to improve dietary intake was to replace sugar-containing beverages with water. Another strategy to improve dietary intake, and control intakes of unfavorable meal constituents like salt, was home cooking (e.g.,

making pizza from scratch). Barriers for healthy eating included feeling rushed and pressed for time or tired (e.g., after a working day).

“Hurry hurry, you know. For example, if you have to go somewhere, for example they have extra lessons in the mosque. Then I notice, quickly baking chips with a minced-meat hot dog and stuff. [...] Sometimes you have those empty moments. And then you bake a minced-meat hot dog.”

Participant 1

Some participants indicated that healthy cooking and home cooking were difficult and laborious compared to unhealthy cooking and takeaway foods, whereas in the opinion of others healthy cooking was not difficult at all, because healthier cooking techniques (like steaming and oven cooking) were considered easier than less healthy techniques (like frying). Some misconceptions about dietary advice were present, e.g. stating coconut oil as being specifically beneficial for health, while saturated fats like coconut oil are usually not recommended in international and national dietary guidelines (30, 31). Participants mentioned mostly consulting social media or acquaintances for information regarding healthy eating.

Physical and mental health and disease

Most participants clearly linked a healthier diet to chronic disease prevention for themselves and their children.

“If children eat healthy, they are not ill. Have fewer problems with everything. With concentration too.”

Participant 7

Participant 6 really regretted his unhealthy eating pattern in the past, which in his opinion had led to diabetes, and he wanted to prevent that from happening to his children:

“An example of me. I have always eaten unhealthy and now I have it [disease]. Custard, ice cream, chocolate... [...] I should not have done that. But you never knew in advance that you could become a diabetic. If my parents had said that, I would not have done it. But they did not say much. [...] They never said: ‘that is good and that is bad’. [...] It is a pity, but... I did not get it from them.”

Participant 6

Another participant became more aware of her lifestyle after being warned by her physician to lose weight in order to prevent cardiovascular diseases and diabetes. One participant mentioned experiencing poorer mental and physical health because of an unhealthy diet and overeating. Contrariwise, poor mental health was seen as a cause of unfavorable eating behavior. Participants explained they lacked energy to prioritize healthy eating or cooking when feeling unwell, worried, stressed or depressed.

“Everyone has a difficult situation and you are not in the mood, yes then it is easy to get a bag of fries and throw them in [the frying pan] and everyone has fries. Because it requires fewer actions and if you do not feel mentally well, then washing the dishes is really too much. Going to a supermarket uh, getting out of bed even, is just too much.”

Participant 3

Broader health concepts

Besides a healthy diet, a healthy weight was considered an important aspect of overall health. Many participants mentioned healthy eating and physical activity as ways to obtain or maintain a healthy weight. One participant felt these factors were interrelated:

“But I think that if you start exercising, that you, that diet is going to change automatically a little bit.”

Participant 2

Some participants mentioned having the intention to exercise more often but not (yet) actually had changed their physical activity level, for example because it was perceived too hard to make time or set one’s mind to it. Costs were not discussed as a barrier for physical activity.

Theme 2. Social and cultural influences

Influences by children

Children played a major role in food choices and food purchases. Participants indicated finding it difficult not to give in to their child’s unhealthy food wishes. Various reasons were indicated for giving in: participants felt sorry for their children if they would not give in, they found it hard to repeatedly reject their child, or they

wanted to compensate their lack of time for their child (e.g., due to a busy work schedule) by buying food that the child liked:

"I work a lot. Night shifts, day shifts and evening shifts. She [child] is alone, I am there with my aunt, but then I felt guilty and then when I left, she started to cry. When I came back, I had cookies for her, 'mommy has brought you cake'. [...] You know, or I went to get her at the babysitter and then she said: 'I missed you, you should not go to work anymore'. 'That's okay, mommy will buy a cake for you okay?'"

Participant 10

Child food preferences also influenced food purchases and dinner choices. Parents mentioned several strategies to broaden their children's exposure to and taste for healthy food including repeated exposure to disliked foods so children could get used to the taste and cooking preferred dishes in a healthier way, such as a homemade pizza rather than store bought or hiding vegetables within a (favorite) dish.

*"It's weird, but they [children] do not want vegetables. But yes, if you for example make chili con carne or for example sauce for spaghetti, then you just throw it through that zucchini. But that is how they eat it. *laughing* So yes, that's how you do it."*

Participant 1

Setting a good example for their child was mentioned as a motivation for healthy eating by some participants. Further, school food regulations positively influenced child-eating behavior at school and sometimes also translated into healthier eating behaviors at home. For example, at some schools, unhealthy snacks or drinks were not allowed in class, which also made the children and parents reconsider consuming these products at home. Most participants had a positive attitude towards these school food regulations as they considered it a helpful contribution to adopting healthier eating behavior.

Influences by culture, family and friends

Besides child influences, extended family and friends also influenced eating and food purchasing behaviors. Eating with friends was generally more associated with having a nice time than with healthy eating. Attempts to adopt healthier cooking styles were sometimes hindered by other family members, e.g., when they disliked the lower-

salt meals. Eating at family gatherings mostly negatively influenced dietary intake, as family gatherings were often accompanied by unhealthy eating, overeating and sometimes setting bad examples:

“Well, uh, not really influence but they [family] try to force trough their vision or their will and I find that difficult. For example, if I go to my mother, well that she uh thinks he [child] should eat peppers, well, I don’t agree with that. [...] After a day at Grandma’s, he [child] goes home and then he ate chocolate, he ate crisps, he ate cake, he ate candy, he ate dinner and preferably ate three other things as well and then also coke and ice cream. Yes, I just think that, I’m really annoyed by that. Really that is just such a frustration.”

Participant 3

One participant even decided to limit family visits to reduce her child’s exposure to unhealthy eating habits of the family. Another mentioned strategy was to bring healthy products to these gatherings themselves. Positive influences were also mentioned, as friends and family sometimes served as an exemplary role for healthy behavior or provided guidance about child upbringing:

“But the bigger she [child] grew, the more rebellious she became, and I say, ‘no, this is not going to happen’. Then I went to talk to my aunt, and she coached me a bit and told me I should be strong. No remains no. That’s how I started to learn.”

Participant 10

Participants’ cultural background also influenced their eating behavior, which was reflected in food customs (e.g., providing and consuming large quantities of food at social gatherings) and food choices (e.g., purchasing and cooking traditional foods, mostly indicated to be unhealthy, fatty or sugary foods).

Theme 3. Influences by the physical environment

Presence of food outlets

Participants lived in or near a disadvantaged neighborhood in The Hague. The presence of sufficient food shops and other facilities in these neighborhoods was appreciated:

"Advantages are uhm, yes you can get almost everything here, also from your own culture the groceries. Everything is close by."

Participant 3

The abundance of supermarkets, small food shops (e.g., Turkish shops) and the market were mentioned in this regard. The market was seen as a place to buy large quantities of cheap fruit and vegetables, although some mentioned that these products did not last long enough as they were not fresh. A downside of the abundance of food outlets in the neighborhood was mentioned to be the food outlets offering unhealthy foods, as participants felt that the presence of these food outlets tempted them into making unhealthy food choices. The food supply at the supermarket checkouts was also considered unhealthy and tempting. Resisting these temptations was especially difficult for children.

"I also want to leave this neighborhood. Because [...] you cannot blame [name child] because he walks out and it already starts, that Bulgarian there, the fries shop there. I mean, in the morning at around a quarter past eight, he already has fried chicken. Yes, you go with your child to the market to get watermelon, he is twice in the fight at the Kentucky. And then he looks at me like that again [...] and then, yes you have to disappoint him. And as a mother you also get tired of that no, no, no [...]. So, uh sometimes we have a little fight about this too. [...] I just want to live somewhere that if you walk out the first ten minutes you will not come across a single snack something. [...] this is really too bad for a child."

Participant 3

The school food environment was mostly viewed as healthy by the participants, which is not surprising as most schools adhered to healthy school food regulations. However, as long as the food outlets surrounding the schools offered unhealthy foods, children were tempted to buy those unhealthy foods during the breaks or after school.

Livability of the neighborhood

Participants had a mostly positive attitude towards their neighborhoods, for example because of the closeness of shops and facilities, social support of the neighbors, perceived safety, openness towards each other and towards different cultures, and

multicultural influences in the neighborhood. Some negative aspects about the neighborhoods were mentioned as well, for example noise pollution, dirty streets and perceived lack of safety of the neighborhood resulting in restricting the child's outdoor activities.

Theme 4. Financial influences

Coping

Most participants had an income below the basic needs limit and prices were considered important for food purchasing. Various strategies were used to cope with a limited budget, such as careful budgeting and planning, budget-friendly cooking, buying secondhand items and buying cheap groceries or groceries on sale. Supermarkets where specific products were the cheapest at that moment were consciously selected, and some participants went to the market around closing time when products were sold for dumping prices. Advantages of planning grocery shopping in advance were firstly preventing buying unnecessary things and thereby saving money, and secondly sticking to healthy eating intentions. Some participants indicated specific financially induced adaptations in their food purchasing behavior, such as limiting outdoor eating to save money and switching from premium brands to cheaper alternatives of the same products, although the budget products sometimes were perceived less tasty or induced feelings of shame:

"Yes, I used to be ashamed to buy cheap products [...]. I really thought those people would think that I don't have money. That's how I thought. Some colleagues also said: 'you should not be ashamed, even if all your groceries are premium brands, it's all the same'. It's just another package, just look, it's all the same. I used to buy Cornflakes of 3 euros while I could also get Cornflakes of 1 euro."

Participant 10

Non-basic needs like a holiday with the family or visiting family abroad were important motivators for saving money.

Financial perception

Healthy foods (e.g., fruit and vegetables) were perceived to be generally more expensive compared to less healthy foods (e.g., sweets and snacks), making choosing unhealthy options tempting.

“Well then you go and look and the healthy things are actually really expensive. Yes then you are inclined, [...] we better take a sausage roll, you almost want to say that.”

Participant 3

Some participants felt discontented about that and indicated that lowering healthy food prices would be a great help in achieving healthier eating behavior in the population.

“But the worst help there is are all those sweets in the shops. Those are cheap and the ones that you need are expensive. That is the worst thing they can have. And then some people think: ‘Yes, that is cheap?’ That is why we have a lot of children with obesity here, too many children. Children from 4 years and older, some children are only 5, all teeth are rotten. Wherever you go, [for] 50 cents you have a bag full of candy. You are not going to have a bag full of vegetables for 50 cents. You do not have that. So if you turn that mentality around, it would be better.”

Participant 5

However, it was mentioned that using the right strategies (e.g., coping strategies for dealing with a limited budget like buying frozen vegetables) it was possible to buy healthy foods despite having a limited budget. Participants generally felt in control over their grocery shopping behavior and felt this was not greatly influenced by external factors. Participants demonstrated a conscious attitude towards their financial situation, as reflected in their coping strategies for dealing with a limited budget, knowingly buying products that were a bit more expensive if they lasted longer, and prioritizing basic needs over luxury needs.

Financial stress

Despite their generally low incomes, participants overall felt relatively comfortable with their financial situation. As described above, various coping strategies were applied to cope with a limited budget and financial stress. Besides, some participants indicated that money was not the most important thing in their lives. For example, health was considered much more important.

“For me, money is not everything. For me it is that I can get up every day, that I can breathe every day, that I thank my god. Every day of my life because

not everyone can do that and I think that's the best you can do as a person, especially when you get up. Because we cannot buy that, not with any money."

Participant 5

However, as also indicated in the theme about mental health, financial stress was a barrier for healthy eating behavior, as participant 8 indicated about the time when she was in debt:

"I did not really buy healthy food then, I just bought what was cheap. I only want to live because you are in the cramp, it's not possible, it's difficult."

Participant 8

Regarding basic needs like food and clothes, participants clearly prioritized their children over themselves. For example, participants mentioned to rather skip a meal themselves than that the child would be short on something.

"I do not care because I prefer [caring for] them [children] rather than myself. I can do with a few slices of bread and peanut butter and then I go to sleep. But they can't."

Participant 6

Food insecurity was mostly mentioned in reference to the past or to others and not to the participants' own current experiences, i.e., mentioning past experiences of having insufficient money for food due to debts, or knowing others that were unable to afford sufficient food. Interestingly, participant 1 was classified as food insecure according to the previous questionnaire, but during the interview he specifically mentioned not to worry about going hungry:

"So, you always have to pay close attention and put everything in order when it comes to finances. For the rest just happy. I mean, my family also. I mean, I'm not worried about, for example, that I'm going to starve, not that."

Participant 1

He made a clear link with the quantity aspect of food security for himself and his family:

"Healthy eating for me and my family means ensuring that there always is food. Yes. That is first of all healthy, that you have to eat. And secondly, yes,

that you pay attention to your diet.”

Participant 1

Existing and proposed solutions to reduce food-related financial strain

Participants were familiar with several existing resources to reduce financial strain or improve eating behavior, like several foundations, allowances, debt assistance, dieticians, the Food Bank, and local initiatives. They generally had a positive attitude towards these resources, which were perceived as a welcome helping hand, although some indicated that they would rather not need it. Conceptually the Food Bank was appreciated, but the actual content of the food parcels distributed by the Food Banks was criticized. Participants mentioned that the distributed products were not suitable for preparing a meal and were sometimes rotten or past the expiry date. If bread was provided it was sometimes stale. Suggested improvements for the content of the food parcels were to provide more fresh products like fruit, vegetables, potatoes and other products that can be used to prepare a proper meal. It was further deemed desirable that social contacts would be promoted and facilitated by Food Banks or other organizations, for example by facilitating getting together for a coffee and conversation.

“The only thing they [Food Banks] don’t have is social contacts.”

Participant 6

Other proposed solutions to reduce financial strain and improve dietary habits were providing free meals for those in need, increasing healthy food supply in the neighborhood (specifically limiting unhealthy snacks at supermarket checkouts and decreasing the number of fast-food outlets) and lowering healthy food prices.

“What would help me? To eat healthier? If the store prices of those things drop a little, that would be super helpful. Not just for me but for many people.”

Participant 5

Barriers for using resources included feeling ashamed, thinking not to belong to the target group, not being eligible for the desired resources, finding it too difficult to register for resources or not knowing where to find the right information. Further, dietary advice provided by dieticians was mentioned to be insufficiently suitable for different cultural backgrounds:

“For dietary advice, it’s just hard in such a neighborhood as this because you have different cultures. [...] I also experienced that at the dietician, yes okay I do get the dietician, but I don’t eat all that. And you can’t expect that if it is in your roots not to eat certain things that you just change it.”

Participant 3

Several participants felt that resources like Food Banks and allowances were often misused by people who did not need it and that people who actually needed help not always asked for or accepted help.

Discussion and conclusions

The current study aimed to provide better insight in the needs and perceptions regarding healthy eating among parents living in disadvantaged neighborhoods in the Netherlands at risk of experiencing food insecurity. Overall, participants showed relatively adequate nutrition knowledge and awareness of the importance of healthy eating behavior for optimal mental and physical health. Nevertheless, participants indicated various social, environmental and financial barriers to healthy eating behavior.

Comparison with previous literature

Consistent with previous research (32), participants acknowledged the importance of healthful eating for chronic disease prevention and overall health. Weight maintenance and child weight maintenance through healthful eating and physical activity was a recurring topic. This finding is in contrast with a previous study (33) that found that participants recognized the importance of improving health habits for themselves but not for their children. Our participants were clearly highly aware of the importance of child weight control, but nevertheless child overweight was a common concern among participants.

Some studies confirm the association between lower nutrition knowledge and lower SES (18, 34) and low (but not very low) food security (35), whereas others indicate adequate nutrition knowledge in these groups (36, 37), which is in line with our findings. Nevertheless, participants generally had a suboptimal diet quality and physical activity level, suggesting that a lack of knowledge was not the driving factor influencing eating behavior. This is in line with various psychological theories related

to health behavior, all consisting of multiple constructs indicating that a variety of factors influence the eventual health behavior (38).

Participants voiced several social, environmental and financial barriers to healthy eating behavior. Social barriers included unhealthy foods offered at social gatherings, bad exemplary roles of others, lacking social support for adopting healthier eating habits, and cultural customs that were associated with overeating and unhealthy food products. Social and family relations are shown to influence eating behavior (39). Especially children were noted to play an important role in family food habits (39), which is in line with the views of our participants. Therefore, it is important to consider child influences when developing interventions to improve eating behavior among families at risk of food insecurity. In line with previous studies (34, 40), lack of time to prepare or cook a meal was another perceived barrier for healthy eating.

Environmental barriers for a healthy eating and lifestyle behavior included an unfavorable food environment (e.g., an abundance of fast-food outlets). A systematic review on environmental factors and obesogenic dietary intakes showed that the food environment (i.e. less access to supermarkets or greater access to takeaway outlets) was consistently associated with higher overweight prevalence, and mixed results were found for the association between the food environment and dietary behaviors (41). Living in a disadvantaged neighborhood may act as a barrier for healthy eating behavior through increased access to takeaway outlets, thereby increasing the ease of making unhealthy choices (41). Further, perceived lack of safety was mentioned as a barrier to outdoor activities like physical activity and child outdoor play. Previous research among low-SES women also indicated unsafe neighborhood environments as barrier for physical activity (42). Also in line with this study (42), despite the generally low income of this study population and of our participants, costs were not discussed as a barrier for physical activity.

Financial considerations were mentioned as a barrier for healthy eating in two ways. Firstly, some believed that healthy foods were too expensive. Strikingly, this perception will probably only intensify because of the recent national tax increase, which came into force on January 2019 (20). As the interviews were conducted before January 2019, we were not able to assess the impact of the tax increase on price perceptions and eating behavior in our study. Therefore, future studies should focus on the effects of the tax increase on eating behavior, especially in low-SES groups. The perception that healthy foods are expensive is in line with previous studies indicating financial

considerations as important barriers for health behavior among low-SES groups (33, 40, 43-45), although participants were resourceful in finding ways to save money and get healthy foods. Secondly, in line with previous studies (46-48), financial stress and poor mental health were associated with poorer eating behavior. Interestingly, while most participants had low incomes and 7 participants were previously classified as food insecure (12), participants had an overall positive attitude towards their financial situation and barely mentioned personally experiencing food insecurity at the present. Participants did mention experiencing food insecurity in reference to the past or to others. This might be due to feelings of discomfort or shame when disclosing personal experiences with food insecurity during an interview (49).

To improve healthy eating behavior among people at risk of food insecurity, participants perceived that changes were needed at the governmental and community and social level. Suggested changes at the governmental level included improving existing resources, for example improving the quality and healthfulness of the Food Bank parcel content. Opposite to the perceptions of our participants, most participants of another Dutch study were satisfied with the food parcels and perceived them as healthy (50), even though their content did not conform to Dutch nutritional guidelines (14). Another proposed governmental intervention was decreasing healthy food prices. Previous studies consistently show that food taxation and subsidies can effectively improve population dietary behavior (22), suggesting that subsidizing healthy foods might be a very promising intervention. This makes the recent decision of the Dutch government to increase food taxes (20) highly undesirable. Suggested changes at the community and social level included promoting and facilitating social contacts in the neighborhood as this was currently lacking according to some participants. The importance of eating in a social context was also highlighted in a previous study among charity-run soup kitchen users (36). Facilitating social contacts could for example be done at Food Banks by providing a suitable location for social interaction. This might also reduce shame and stigmatization associated with Food Bank use, as this was indicated as a barrier for Food Bank use in previous literature (51, 52) and in our study.

Methodological considerations

This study deepens the understanding of needs and perceptions of parents at risk of experiencing food insecurity. Our qualitative, open interview approach enabled identifying important themes regarding healthy eating behavior in this difficult to reach target population. Our analyses confirmed some of the themes that were expected to play a role in healthy eating behavior based on our previous study and

the literature (e.g., family influences) and deepened knowledge on these topics. Additionally, some less anticipated themes emerged during the interviews (e.g., influence of the food environment and importance of social contacts). Our results may not be representative for a national sample of people at risk of food insecurity because we only recruited participants from the current study on food insecurity in disadvantaged neighborhoods in The Hague, The Netherlands (12). Also, participants volunteered to be interviewed which may have led to a sample with a larger-than-usual interest in nutrition. However, the included participants varied in terms of migration background and other characteristics. Also, thematic saturation for all themes was reached, suggesting that the sample size was sufficient for the aims of our study.

Implications

Nutrition knowledge and motivation to improve healthy eating behavior were relatively high among participating parents at risk of food insecurity, yet they indicated various social, environmental and financial barriers to healthy eating behavior. Therefore, interventions aimed at improving eating behavior in this unique population should not merely focus on nutrition education but take into account a wider range of social, environmental and financial factors. Because our study population consisted specifically of families with young children living in or near disadvantaged neighborhoods, the identified themes, barriers and interventions may not be generalizable to other populations at risk of food insecurity. Therefore, future studies are needed to confirm the needs and perceptions regarding healthy eating behavior in other populations at risk of experiencing food insecurity, e.g., young or elderly populations, childless people, and people with other migration backgrounds. Suggested interventions to improve eating behavior and reduce food-related financial strain that were identified in our study include facilitating social contacts (thereby potentially enhancing social support for both financial and food-related issues), improving existing recourses (e.g., Food Bank parcel content), culture-specific dietary advice, parenting training focused on handling child food choice influences, and improving the neighborhood food environment. Also, financial and mental issues should be addressed prior to focusing on improving eating behavior. Further, possibilities for subsidizing healthy foods or taxing unhealthy foods in the Netherlands should be explored as a potentially promising intervention to improve eating behavior.

List of abbreviations

BMI Body Mass Index

IRR Inter Rater Reliability

-

References

1. Darmon N, Drewnowski A. Does social class predict diet quality? *The American journal of clinical nutrition*. 2008;87(5):1107-17.
2. Everson SA, Maty SC, Lynch JW, Kaplan GA. Epidemiologic evidence for the relation between socioeconomic status and depression, obesity, and diabetes. *Journal of psychosomatic research*. 2002;53(4):891-5.
3. McLaren L. Socioeconomic status and obesity. *Epidemiologic reviews*. 2007;29(1):29-48.
4. Hanson KL, Connor LM. Food insecurity and dietary quality in US adults and children: a systematic review. *The American journal of clinical nutrition*. 2014;100(2):684-92.
5. Moradi S, Mirzababaei A, Dadfarma A, Rezaei S, Mohammadi H, Jannat B, et al. Food insecurity and adult weight abnormality risk: a systematic review and meta-analysis. *European journal of nutrition*. 2019;58(1):45-61.
6. Seligman HK, Laraia BA, Kushel MB. Food insecurity is associated with chronic disease among low-income NHANES participants. *The Journal of nutrition*. 2009;140(2):304-10.
7. (FAO) FaOotUN. Rome Declaration on World Food Security and World Food Summit Plan of Action. Rome1996.
8. Foley W, Ward P, Carter P, Coveney J, Tsourtos G, Taylor A. An ecological analysis of factors associated with food insecurity in South Australia, 2002–7. *Public health nutrition*. 2010;13(2):215-21.
9. Gundersen C, Kreider B, Pepper J. The Economics of Food Insecurity in the United States. *Applied Economic Perspectives and Policy*. 2011;33(3):281-303.
10. Borch A, Kjærnes U. Food security and food insecurity in Europe: An analysis of the academic discourse (1975–2013). *Appetite*. 2016;103:137-47.
11. Neter JE, Dijkstra SC, Visser M, Brouwer IA. Food insecurity among Dutch food bank recipients: a cross-sectional study. *BMJ open*. 2014;4(5):e004657.
12. van der Velde LA, Nyns CJ, Engel MD, Neter JE, van der Meer IM, Numans ME, et al. Exploring food insecurity and obesity in Dutch families: a cross-sectional mediation analysis. *BMC Public Health*. 2020;20(1):569.
13. Voedselbanken Nederland (Dutch Food Bank). Feiten en Cijfers Voedselbanken Nederland 2019 (Facts and Figures Dutch Food Bank 2019) [Available from: <https://www.voedselbankennederland.nl/>].
14. Neter JE, Dijkstra SC, Visser M, Brouwer IA. Dutch food bank parcels do not meet nutritional guidelines for a healthy diet. *British Journal of Nutrition*. 2016;116(3):526-33.
15. Laitinen J, Ek E, Sovio U. Stress-related eating and drinking behavior and body mass index and predictors of this behavior. *Preventive medicine*. 2002;34(1):29-39.
16. Torres SJ, Nowson CA. Relationship between stress, eating behavior, and obesity. *Nutrition*. 2007;23(11-12):887-94.
17. Zellner DA, Loaiza S, Gonzalez Z, Pita J, Morales J, Pecora D, et al. Food selection changes under stress. *Physiology behavior*. 2006;87(4):789-93.
18. Parmenter K, Waller J, Wardle J. Demographic variation in nutrition knowledge in England. *Health education research*. 2000;15(2):163-74.

19. Rao M, Afshin A, Singh G, Mozaffarian D. Do healthier foods and diet patterns cost more than less healthy options? A systematic review and meta-analysis. *BMJ open*. 2013;3(12):e004277.
20. Snel M. Kamerstuk 31532 nr 191 Voedingsbeleid Brief van de staatssecretaris van financiën. 2018.
21. French SA, Story M, Jeffery RW. Environmental influences on eating and physical activity. *Annual review of public health*. 2001;22(1):309-35.
22. Niebylski ML, Redburn KA, Duhaney T, Campbell NR. Healthy food subsidies and unhealthy food taxation: A systematic review of the evidence. *Nutrition*. 2015;31(6):787-95.
23. Vogelaar CP. Brief van de minister voor wonen, wijken en integratie 2007 [Available from: <https://zoek.officielebekendmakingen.nl/kst-30995-1.html>].
24. Economic Research Service. U.S. Household Food Security Survey Module: three-stage design, with screeners. In: USDA, editor. 2012.
25. United States Department of Agriculture (USDA). Definitions of Food Security 2017 [Available from: <https://www.ers.usda.gov/topics/food-nutrition-assistance/food-security-in-the-us/definitions-of-food-security>].
26. van Lee L, Feskens EJ, Meijboom S, van Huysduynen EJH, van't Veer P, de Vries JH, et al. Evaluation of a screener to assess diet quality in the Netherlands. *British Journal of Nutrition*. 2016;115(3):517-26.
27. Thomas DR. A general inductive approach for analyzing qualitative evaluation data. *American journal of evaluation*. 2006;27(2):237-46.
28. McAlister A, Lee D, Ehler K, Kajfez R, Faber C, Kennedy M, editors. Qualitative coding: An approach to assess inter-rater reliability. ASEE Annual Conference & Exposition, Columbus, Ohio <https://peer.asee.org/28777>; 2017.
29. Saldana J. The coding manual for qualitative researchers An introduction to codes and coding. 3. London: Sage Publications Ltd; 2009.
30. Netherlands Nutrition Center. Richtlijnen Schijf van Vijf (Wheel of Five guidelines). In: (Voedingscentrum) NNC, editor. The Hague 2016.
31. Herforth A, Arimond M, Álvarez-Sánchez C, Coates J, Christianson K, Muehlhoff E. A Global Review of Food-Based Dietary Guidelines. *Advances in Nutrition*. 2019.
32. Evans AE, Wilson DK, Buck J, Torbett H, Williams J. Outcome expectations, barriers, and strategies for healthful eating: a perspective from adolescents from low-income families. *Family Community Health*. 2006;29(1):17-27.
33. Davis AM, Befort C, Steiger K, Simpson S, Mijares M. The nutrition needs of low-income families regarding living healthier lifestyles: Findings from a qualitative study. *Journal of Child Health Care*. 2013;17(1):53-61.
34. Skuland SE. Healthy Eating and Barriers Related to Social Class. The case of vegetable and fish consumption in Norway. *Appetite*. 2015;92:217-26.
35. Fitzgerald N, Hromi-Fiedler A, Segura-Pérez S, Pérez-Escamilla R. Food insecurity is related to increased risk of type 2 diabetes among Latinas. *Ethnicity & disease*. 2011;21(3):328.
36. Wicks R, Trevena LJ, Quine S. Experiences of food insecurity among urban soup kitchen consumers: insights for improving nutrition and well-being. *Journal of the American Dietetic Association*. 2006;106(6):921-4.

37. Evans A, Banks K, Jennings R, Nehme E, Nemec C, Sharma S, et al. Increasing access to healthful foods: a qualitative study with residents of low-income communities. 2015;12(1):S5.
38. Linke SE, Robinson CJ, Pekmezi D. Applying psychological theories to promote healthy lifestyles. *American Journal of Lifestyle Medicine*. 2014;8(1):4-14.
39. Coveney J. What does research on families and food tell us? Implications for nutrition and dietetic practice. *Nutrition Dietetics: The Journal of the Dietitians Association of Australia*. 2002;59(2):113-20.
40. Eikenberry N, Smith C. Healthful eating: perceptions, motivations, barriers, and promoters in low-income Minnesota communities. *Journal of the American Dietetic Association*. 2004;104(7):1158-61.
41. Giskes K, van Lenthe F, Avendano-Pabon M, Brug J. A systematic review of environmental factors and obesogenic dietary intakes among adults: are we getting closer to understanding obesogenic environments? *Obesity reviews*. 2011;12(5):e95-e106.
42. Ball K, Salmon J, Giles-Corti B, Crawford D. How can socio-economic differences in physical activity among women be explained? A qualitative study. *Women & health*. 2006;43(1):93-113.
43. Kamphuis CB, van Lenthe FJ, Giskes K, Brug J, Mackenbach JP. Perceived environmental determinants of physical activity and fruit and vegetable consumption among high and low socioeconomic groups in the Netherlands. *Health & place*. 2007;13(2):493-503.
44. Dammann KW, Smith C. Factors affecting low-income women's food choices and the perceived impact of dietary intake and socioeconomic status on their health and weight. *Journal of nutrition education behavior*. 2009;41(4):242-53.
45. Williams L, Ball K, Crawford D. Why do some socioeconomically disadvantaged women eat better than others? An investigation of the personal, social and environmental correlates of fruit and vegetable consumption. *Appetite*. 2010;55(3):441-6.
46. Bratanova B, Loughnan S, Klein O, Claassen A, Wood R. Poverty, inequality, and increased consumption of high calorie food: Experimental evidence for a causal link. *Appetite*. 2016;100:162-71.
47. Sarlio-Lähteenkorva S, Lahelma E, Roos E. Mental health and food habits among employed women and men. *Appetite*. 2004;42(2):151-6.
48. O'Neil A, Quirk SE, Housden S, Brennan SL, Williams LJ, Pasco JA, et al. Relationship Between Diet and Mental Health in Children and Adolescents: A Systematic Review. *American Journal of Public Health (AJPH)*. 2014;104(10):e31-e42.
49. Lachance L, Sean Martin M, Kaduri P, Godoy-Paiz P, Ginieniewicz J, Tarasuk V, et al. Food insecurity, diet quality, and mental health in culturally diverse adolescents. *Ethnicity and Inequalities in Health Social Care*. 2014;7(1):14-22.
50. Neter JE, Dijkstra S, Dekkers A, Ocke M, Visser M, Brouwer I. Dutch food bank recipients have poorer dietary intakes than the general and low-socioeconomic status Dutch adult population. *European journal of nutrition*. 2017;1-12.
51. Hoogland H, Berg J. Ervaringen van schaamte en psychologisch lijden door voedselbankklanten. *Journal of Social Intervention: Theory Practice*. 2016;25(1).
52. van der Horst H, Pascucci S, Bol W. The "dark side" of food banks? Exploring emotional responses of food bank receivers in the Netherlands. *British Food Journal*. 2014;116(9):1506-20.

Additional material Chapter 4

Additional Table 1. Food insecurity status assessment

Statement/ question
I (or other family members) worried whether my (or our) food would run out before I (or we) got money to buy more. ¹
The food that I (or we) bought just didn't last, and I (or we) didn't have money to get more. ¹
I (or we) couldn't afford to eat balanced meals. ¹
In the last 12 months, did you (or other adults in your household) ever cut the size of your meals or skip meals because there wasn't enough money for food? ²
How often did this happen in the last 12 months? ³
In the last 12 months, did you ever eat less than you felt you should because there wasn't enough money for food? ²
In the last 12 months, were you ever hungry but didn't eat because there wasn't enough money for food? ²
In the last 12 months, did you lose weight because there wasn't enough money for food? ²
In the last 12 months, did you (or other adults in your household) ever not eat for a whole day because there wasn't enough money for food? ²
How often did this happen in the last 12 months? ³
I (or we) relied on only a few kinds of low-cost food to feed my (or our) child/children because I was (or we were) running out of money to buy food. ¹
I (or we) couldn't feed my (or our) child/children a balanced meal, because I (or we) couldn't afford that. ¹
My (or our) child was/children were not eating enough because I (or we) just couldn't afford enough food. ¹
In the last 12 months, did you ever cut the size of your child's/ any of the children's meals because there wasn't enough money for food? ²
In the last 12 months, did your child/ children ever skip meals because there wasn't enough money for food? ²
How often did this happen in the last 12 months? ³
In the last 12 months, was your child/were your children ever hungry but you just couldn't afford more food? ²
In the last 12 months, did your child/ any of the children ever not eat for a whole day because there wasn't enough money for food? ²

¹Answer options: Often true/ Sometimes true/ Never true/ I don't know

²Answer options: Yes/ No/ I don't know

³Answer options: Almost every month/ Some months but not every month/ Only 1 or 2 months/ I don't know

Additional Table 2. Diet quality score components, dietary guidelines and scoring per component

Component	Recommendations by the Health Council of the Netherlands ¹ and/ or the Netherlands Nutrition Centre ²	% contribution to component score	Units	Score		
				0	5	10
Vegetables	Eat at least 200 grams of vegetables daily	100	Grams/ d	0	<i>Continuous</i>	≥200
Fruit	Eat at least 200 grams of fruit daily	100	Pieces/ d	0	<i>Continuous</i>	≥ 2
Fish	Eat one serving of fish weekly, preferably oily fish	50	Servings/ w	0	<1	≥ 1
		50	No fish consumed		Lean or both lean and fatty fish	Mostly fatty fish
Bread	Replace refined cereal products by whole-grain products	50		Mostly white bread	Both white and brown/ whole-grain bread	Mostly brown/ whole-grain bread
	Women: 4-5 brown/ whole-grain sandwiches daily	50	Sandwiches/ d	0	<i>Continuous</i>	≥ 4
	Men: 6-8 brown/ whole-grain sandwiches daily		Sandwiches/ d	0	<i>Continuous</i>	≥ 6
Oils and fats	Replace butter, hard margarines and cooking fats by soft margarines, liquid cooking fats, and vegetable oils	50		Butter, hard margarines	Both butter, hard margarines and oils margarines and soft margarines	Oils and soft margarines
		50		Butter on bread or bread is not buttered at all	Semi-skimmed butter or hard margarine on bread	Diet margarine on bread
Sweet and savory snacks	For products outside the Wheel of Five: choose an item from the daily selection no more than three to five times per day, and something from the weekly selection no more than three times a week	25	Sweet snacks (larger serving)/ w	≥ 3	<1 to 2	Not consumed
		25	Sweet snacks (small serving)/ d	> 3	<i>Continuous</i>	Not consumed
		25	Savory snacks (larger serving)/ w	≥ 3	1 to 2	Not consumed
		25	Savory snacks (small serving)/ d	> 3	<i>Continuous</i>	Not consumed

¹Health Council of the Netherlands. *Guidelines for a healthy diet 2015* (Richtlijnen Goede Voeding 2015). The Hague 2015.² Netherlands Nutrition Center. *Wheel of Five guidelines* (Richtlijnen Schijf van Vijf). The Hague 2016.

Additional Table 3. Topic list and example questions

<i>Topic</i>	<i>Example question</i>
General/ introductory topics	
Birthplace and culture	When and where were you born?
	For how long have you been living in the Netherlands
Household composition	What does your family look like?
	Who lives at your home?
Living conditions	Where do you live/ which neighborhood?
	What do you think of the neighborhood where you live?
	What kind of house do you have?
Specific topics	
Healthy eating	What is healthy eating for you?
	What do you think of healthy eating?
Skills	What do you think about cooking a healthy meal?
	What do you find easy or difficult when cooking a healthy meal?
Influences on eating and food purchasing	What influences how you eat or what kind of foods you buy?
Healthy lifestyle	What can you tell about health and nutrition?
	What impact does your health have on what you eat?
	How does your weight affect what you eat?
	How do you think about exercise and health?
Eating in a social context	Do other people influence what you eat? Who are they?
	How do they (reference to previous question) affect what you eat?
	What can you tell about eating and coming together with people, for example on parties or social gatherings?
Neighborhood	How does the neighborhood where you live affect what you eat?
	What kind of temptations (for you/ for the children) are there in your neighborhood?
Cultural influences on eating	What influence does your culture have on your eating habits?
Family	What does healthy eating mean for your children (for you as a parent)?
	How important is this (reference to previous question)?
	How do you ensure that your children eat healthy?
	What would you like to teach your children about food and health?
Upbringing	What do you find easy or difficult when raising your child?
	What barriers do you experience when raising your child?
Financial status	Would you describe your own financial status?
	How do you influence that in daily life?

Stress/ financial stress	How does stress affect what you eat or what food you buy? What impact do your finances have on how stressed you feel?
Food costs	What role does money play in what you eat or what food you buy? How do you take into account food costs? How do you take into account offers?
Priorities	What do you find important or what do you pay attention to when spending your money? What do you find important or what do you pay attention to when buying groceries?
Nutrition and health	Does eating have an impact on your health? How do you notice that (reference to previous question)? How do you describe your own health? Does your health status influence where or how you buy your groceries? Does your physical health prevent you from for example going to the market to buy groceries?
Help	What could help you to eat healthier? What can people with a limited budget help to have sufficient and healthy food to eat? How can the municipality help? Which help from the municipality / which foundations do you know? What can for example the school or supermarket do to make healthy eating easier? What do you think of receiving vouchers to get fruit and vegetables at the market? What do you think of the Food Bank?
Rearrangement of the neighborhood	If you could rearrange the neighborhood you live in, how would you do that? What would you like to remove from the neighborhood or add to the neighborhood?
Closing topics	
Future	How would you like your future to look like?
Unaddressed topics	Is there anything else you would like to address in this interview?

