



Universiteit
Leiden

The Netherlands

Advanced echocardiography in characterization and management of patients with secondary mitral regurgitation

Namazi, F.

Citation

Namazi, F. (2022, May 10). *Advanced echocardiography in characterization and management of patients with secondary mitral regurgitation*. Retrieved from <https://hdl.handle.net/1887/3303481>

Version: Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/3303481>

Note: To cite this publication please use the final published version (if applicable).


Part I

Epidemiology in secondary mitral regurgitation



Chapter two

Sex differences in prognosis of significant secondary mitral regurgitation



Farnaz Namazi, Pieter van der Bijl, N. Mai Vo, Suzanne E., van Wijngaarden, Nina Ajmone Marsan, Victoria Delgado, Jeroen J. Bax.

ESC Heart Fail. 2021 Aug 6. doi: 10.1002/ehf2.13503.

Abstract

Aims

Secondary mitral regurgitation (MR) is more frequent in men than in women. However, little is known about differences in prognosis between men and women with secondary MR. The objective of this study is to investigate the sex distribution of secondary MR and the prognostic differences between sexes.

Methods

Patients with significant secondary MR, of both ischemic and non-ischemic etiology, were identified through the departmental electronic patient files and retrospectively analyzed. The primary endpoint was all-cause mortality.

Results

A total of 698 patients (mean age 66 ± 11 years) with significant secondary MR were included: 471 (67%) men and 227 (33%) women. Ischemic heart failure was significantly more common in men (61%), whereas non-ischemic heart failure was more prevalent in women (63%). Women had significantly smaller left ventricular (LV) volumes when compared to men and more preserved LV systolic function when assessed with LV global longitudinal strain (GLS; $8.5 \pm 4.1\%$ vs. $7.5 \pm 3.6\%$; $P = 0.004$). Women more often underwent surgical mitral valve repair (34%) when compared to men (26%), whilst no differences were observed for transcatheter mitral valve repair. During a median follow-up of 57 [interquartile range 29-110] months, 373 (53%) patients died. Women showed significantly lower mortality rates at 1-, 2- and 5-year follow-up (9%, 16% and 33% vs. 10%, 20% and 42%) when compared to men ($P = 0.001$).

Conclusions

Significant secondary MR is more frequently observed in men as compared to women and is associated with worse prognosis.

Introduction

Mitral regurgitation (MR) is one of the most common valvular diseases with a growing incidence and significant MR is associated with poor prognosis ^{1,2}. The frequency of significant MR in men and women is comparable ³. However, when evaluating the etiology of MR, significant differences are observed: secondary MR is more prevalent in men, whereas primary MR is more frequently present in women ³⁻⁵. Previous studies have shown differences between men and women undergoing surgical mitral valve repair for primary or secondary MR, with women having a higher mortality risks ^{4,6,7}. In patients undergoing transcatheter edge-to-edge mitral valve repair with the MitraClip device, no sex differences were observed ⁸. However, these studies were mostly performed in patients with primary MR. A recent publication evaluating the sex differences in patients with secondary MR undergoing surgical mitral valve repair showed higher mortality risk in women as compared to men ⁹. The factors underlying these differences remain elusive, particularly among patients with secondary MR ¹⁰. Accordingly, the objective of this study is to investigate the sex distribution and long-term prognosis of patients with significant secondary MR.

Methods

Patient population

Patients with significant (moderate and severe) secondary MR were identified from 1999-2018 through the departmental echocardiographic database of the Leiden University Medical Center (Leiden, The Netherlands). Patients with previous mitral valve intervention were excluded. Baseline demographic, clinical and echocardiographic characteristics were prospectively collected through the departmental clinical database (EPD-Vision 11.8.4.0; Leiden University Medical Center, Leiden, The Netherlands) and were analyzed retrospectively. The institutional review board approved this retrospective study of clinically acquired data and waived the need for written patient informed consent.

Clinical variables included heart failure etiology (i.e. ischemic vs. non-ischemic), New York Heart Association (NYHA) functional class, comorbidities and medication use. Body surface area was calculated according to the Du Bois formula ¹¹. Ischemic heart failure was defined based on coronary artery disease confirmed by coronary angiography, prior coronary revascularization with percutaneous coronary intervention and/or coronary artery bypass grafting (CABG). Mitral valve intervention included surgical mitral valve repair or replacement and transcatheter edge-to-edge mitral valve repair with the MitraClip device.

Echocardiography

Transthoracic echocardiography was performed with commercially available systems (General Electric Vingmed Ultrasound, Milwaukee, USA), and images were digitally stored for offline analysis (EchoPAC 201.0.0, General Electric Vingmed Ultrasound, Milwaukee, USA). Image acquisition was performed with patients in hemodynamic stable conditions at rest in the left lateral decubitus position. Using 3.5 MHz or M5S transducers two-dimensional images, M-mode and Doppler data were acquired from parasternal, apical and subcostal views. From the apical 2- and 4-chamber views, left ventricular (LV) volumes (end-diastolic and end-systolic) were measured and the LV ejection fraction (EF) was quantified using the Simpson's biplane method¹². LV volumes were indexed for body surface area. MR severity was assessed using a multiparametric approach according to current guidelines and graded as moderate (grade 2), moderate-to-severe (grade 3) and severe (grade 4)¹³⁻¹⁵. From standard 2-dimensional transthoracic echocardiography, LV global longitudinal strain (GLS) was measured using apical 4-chamber, 2-chamber and long-axis views of the LV and processed offline using commercially available software (EchoPAC 201.0.0, General Electric Vingmed Ultrasound, Milwaukee, USA)¹⁶. LV GLS is a measure of global shortening of the myocardium and is conventionally expressed as negative values. However, in this study we have treated this variable as absolute value and therefore, a higher LV GLS value represents better LV systolic function.

Follow-up

Patients were followed-up for the occurrence of the primary endpoint of all-cause mortality. The follow-up started from the date of the first echocardiogram showing significant secondary MR. Data on survival were collected from the departmental cardiology information system (EPD-Vision 11.8.4.0; Leiden University Medical Center, Leiden, The Netherlands) which is linked to the governmental death registry database. Follow-up was complete for all patients.

Statistical analysis

Continuous data are presented as mean \pm standard deviation (when normally distributed) or as median with interquartile range (when not normally distributed). An independent sample Student t-test or Mann-Whitney U test (when appropriate) was used for the comparison of continuous data. Categorical data are presented as absolute numbers and percentages and a χ^2 -test was used for the comparison between groups. To estimate the cumulative survival rates, a Kaplan-Meier analysis was performed and the log-rank test was used to compare the cumulative survival rates between men and women. Based on the Kaplan-Meier analysis, a post hoc landmark analysis was

performed at 36 months of follow-up to evaluate early vs. late outcomes between men and women. Independent associates for all-cause mortality were evaluated using Cox proportional hazards regression analysis. The hazard ratio (HR) and 95% confidence intervals (CI) were calculated and reported. Variables with a p-value <0.05 were considered statistically significant and were included in the multivariable model. All statistical analyses were performed using SPSS for Windows, version 23.0 (IBM, Armonk, NY, USA), with a two-tailed p-value <0.05 being considered statistically significant.

Results

Patient population

A total of 698 patients (mean age 66 ± 11 years) with significant secondary MR were included. Ischemic heart failure was present in 53% of the total population. The majority of the patients presented with heart failure symptoms NYHA functional class III (60%). The mean LVEF was $29\pm 11\%$ and the mean LV GLS was $7.8\pm 3.8\%$. The majority of the patients (82%) had moderate-to-severe or severe MR. Tables 1 and 2 summarize the baseline clinical and echocardiographic characteristics for the total population and the differences between sexes.

There were 471 (67%) men (mean age 67 ± 10 years) and 227 (33%) women (mean age 65 ± 13). No differences were observed in the prevalence of atrial fibrillation or other cardiovascular risk factors (i.e. hypertension and/or diabetes mellitus). In terms of heart failure etiology, men more frequently had ischemic heart failure (61%) whereas women more frequently had non-ischemic heart failure (63%). Although women had more severe heart failure symptoms as compared to men, the difference did not reach statistical significance. In terms of echocardiographic characteristics, women had significantly smaller LV volumes when compared to men, but no significant difference was observed in LVEF. However, LV systolic function was slightly better in women than in men when assessed with LV GLS (LV GLS $7.5\pm 3.6\%$ in men vs. $8.5\pm 4.1\%$ in women; $P=0.004$). No differences were observed in MR grade between men and women.

Table 1. Clinical characteristics.

	Total population (n= 698)	Men (n= 471)	Women (n= 227)	P-value
Age (years)	66±11	67±10	65±13	0.051
BSA (m ²)	1.92±0.21	1.99±0.19	1.78±0.19	<0.001
Atrial fibrillation, n (%)	289 (41)	206 (44)	83 (37)	0.071
Hypertension, n (%)	275 (39)	175 (37)	100 (44)	0.081
Diabetes mellitus, n (%)	167 (24)	120 (26)	47 (21)	0.166
eGFR (mL/min/1.73m ²)	62±26	63±25	61±27	0.417
Heart failure etiology, n (%)				
Ischemic	370 (53)	287 (61)	83 (37)	<0.001
Non-ischemic	328 (47)	184 (39)	144 (63)	<0.001
NYHA class, n (%)				
I	34 (5)	26 (6)	8 (4)	0.251
II	170 (24)	116 (25)	54 (24)	0.809
III	415 (60)	282 (60)	133 (59)	0.747
IV	79 (11)	47 (10)	32 (14)	0.108
Medication, n (%)				
Beta-blockers	492 (71)	326 (69)	166 (73)	0.288
Diuretics	580 (83)	387 (82)	193 (85)	0.346
ACEi/ARB	565 (81)	390 (83)	175 (77)	0.072
MRA	304 (44)	193 (41)	111 (49)	0.048

Continuous data are presented as mean ± SD or median [interquartile range]. Categorical data are presented as numbers and percentages.

ACEi, angiotensin-converting enzyme inhibitor; ARB, angiotensin receptor blocker; BSA, body surface area; eGFR, estimated glomerular filtration rate; MRA, mineralocorticoid receptor antagonists; NYHA, New York Heart Association.

Table 2. Echocardiographic characteristics.

	Total population (n= 698)	Men (n= 471)	Women (n= 227)	P-value
LVEDVi (mL/m ²)	101 [76-129]	104 [81-132]	92 [68-122]	<0.001
LVESVi (mL/m ²)	73 [51-97]	77 [53-98]	66 [44-92]	<0.001
LVEF (%)	29±11	29±11	30±12	0.180
LV GLS (%) *	7.8±3.8	7.5±3.6	8.5±4.1	0.004
MR grade, n (%)				
Moderate	125 (18)	87 (19)	38 (17)	0.576
Moderate-to-severe	305 (44)	210 (45)	95 (42)	0.495
Severe	268 (38)	174 (37)	94 (41)	0.256

Continuous data are presented as mean ± SD or median [interquartile range]. Categorical data are presented as numbers and percentages. *LV GLS feasible in N= 660 patients.

LVEF, left ventricular ejection fraction; LVEDVi, indexed left ventricular end-diastolic volume; LVESVi, indexed left ventricular end-systolic volume; LV GLS, left ventricular global longitudinal strain; MR, mitral regurgitation.

Follow-up

During follow-up, two-thirds of the patients (64%) received cardiac resynchronization therapy, and 308 patients (44%) received mitral valve intervention: 28% underwent surgical mitral valve repair and 16% received transcatheter edge-to-edge mitral valve repair. Men more frequently received cardiac resynchronization therapy when compared to women. In terms of invasive mitral valve treatment, women more often underwent surgical mitral valve repair (34%), whilst no differences were observed in transcatheter edge-to-edge mitral valve repair. A summary of device implantation and mitral valve intervention at follow-up and differences between men and women is shown in Table 3.

Table 3. Intervention during follow-up.

	Total population (n= 698)	Men (n= 471)	Women (n= 227)	p-value
Device therapy, n (%)				
Cardiac resynchronization therapy	414 (64)	294 (68)	120 (55)	0.001
MV intervention, n (%)				
Medical therapy only	387 (55)	279 (59)	108 (48)	0.004
Mitral valve repair	198 (28)	122 (26)	76 (34)	0.037
Mitral valve replacement	3 (0.4)	1 (0.2)	2 (0.9)	0.206
MitraClip	110 (16)	69 (15)	41 (18)	0.246
Concomitant procedure, n (%) *				
CABG	60 (9)	46 (10)	14 (6)	0.112
Tricuspid valve annuloplasty	131 (19)	83 (18)	46 (21)	0.264
LV reconstruction	17 (2)	12 (3)	5 (2)	0.782
Cardiac support device (CoreCap)	63 (9)	39 (8)	24 (11)	0.322
MAZE	30 (4)	21 (5)	9 (4)	0.763

Continuous data are presented as mean \pm SD or median [interquartile range]. Categorical data are presented as numbers and percentages. * Concomitant procedures with mitral valve treatment.

CABG, coronary artery bypass graft

Survival analysis

During a median follow-up of 57 [interquartile range 29-110] months, 373 (53%) patients died. Women showed significantly lower mortality rates at 1-, 2- and 5-years follow-up as compared to men (women 9%, 16% and 33% vs. men 10%, 20% and 42%, respectively; $P= 0.001$, Figure 1, panel A). Based on the Kaplan-Meier curves, an additional landmark analysis was performed at 36 months of follow-up demonstrating

that the differences in survival were significant between men and women after this time point ($P < 0.001$, Figure 1, panel B). In patients with ischemic heart failure, there was no significant difference in outcome between men and women ($P = 0.179$, Figure 2, Panel A). On the contrary, in patients with non-ischemic heart failure, a significant difference in outcome between men and women was seen with women having a better outcome than men ($P = 0.017$, Figure 2, Panel B).




Figure 1. Kaplan-Meier curves for all-cause mortality. Panel A demonstrates time to all-cause mortality according to sex: women (green) and men (red). Panel B demonstrates the landmark analysis at 36 months of follow-up with time-to-event curves for all-cause mortality according to sex.




Figure 2. Kaplan-Meier curves for all-cause mortality in patients with ischemic and non-ischemic heart failure. Panel A demonstrates time to all-cause mortality according to sex in patients with ischemic heart failure: women (green) and men (red). Panel B demonstrates time to all-cause mortality according to sex in patients with non-ischemic heart failure

Considering patients receiving medical therapy only, women had lower mortality rates than men (5-year estimated rates 77% in women vs. 62% in men; $P=0.001$). On univariable analysis, age, male sex, chronic kidney disease, diabetes mellitus, ischemic heart failure, LVEF and LV GLS were significantly associated with all-cause mortality (Table 4). On multivariable analysis, after correcting for various clinical and echocardiographic parameters, male sex remained independently associated with all-cause mortality (HR 1.423; 95% CI, 1.109-1.826; $P=0.006$).

Table 4. Uni- and multivariable Cox regression analyses to identify associates of all-cause mortality.

Variable	Univariate analysis			Multivariate analysis		
	HR	95% CI	p-value	HR	95% CI	p-value
Age	1.030	1.019-1.040	<0.001	1.020	1.009-1.032	0.001
Male sex	1.474	1.175-1.850	0.001	1.423	1.109-1.826	0.006
BSA (m ²)	1.050	0.652-1.689	0.842			
eGFR (mL/min/1.73m ²)	0.977	0.972-0.981	<0.001	0.980	0.975-0.985	<0.001
Hypertension	0.940	0.762-1.160	0.566			
Diabetes mellitus	1.429	1.132-1.805	0.003	1.309	1.021-1.678	0.034
Atrial fibrillation	1.145	0.932-1.406	0.197			
Ischemic aetiology	1.291	1.052-1.585	0.015	1.071	0.855-1.342	0.550
NYHA classification \geq II	1.026	0.621-1.694	0.920			
Beta-blockers	0.845	0.681-1.048	0.125			
CRT	1.092	0.859-1.390	0.471			
MV intervention *	1.051	0.852-1.298	0.642			
LVEF (%)	0.988	0.977-0.998	0.025	0.999	0.985-1.013	0.864
LV GLS (%)	1.082	1.047-1.118	<0.001	1.066	1.024-1.109	0.002

BSA, body surface area; CI, confidence interval; CRT, cardiac resynchronization therapy; EDV, end-diastolic volume; eGFR, estimated glomerular filtration rate; HR, hazard ratio; LVEF, left ventricular ejection fraction; LV GLS, left ventricular global longitudinal strain; MV, mitral valve; NYHA, New York Heart Association.

* Combined surgical mitral valve repair, mitral valve replacement and percutaneous edge-to-edge mitral valve repair

Discussion

The present study shows that significant secondary MR is more frequent in men as compared to women. Ischemic etiology is the most frequent underlying mechanism in men whereas non-ischemic cardiomyopathy is more frequently observed in women. Men had a worse prognosis, compared to women.

Sex differences in secondary mitral regurgitation

Important differences in prevalence, pathophysiology and prognosis of various cardiovascular diseases have been described between men and women ¹⁷. In mitral valve pathology two different mechanisms of MR can be distinguished; primary MR (i.e. the result of pathology of the MV apparatus itself) and secondary MR (i.e. the result of changes in the LV geometry due to ischemic or non-ischemic heart failure) ¹⁸. A large echocardiography-based registry by Monteagudo Ruiz *et al.* ³ described the sex differences among 3309 patients with moderate and severe MR and showed that secondary MR was more prevalent among men vs. women (39.2% vs. 21.8%, respectively). However, the study did not elaborate further on the differences between men and women with significant secondary MR. The randomized clinical trial from the Cardiothoracic Surgical Trials Network (CTSN) comparing mitral valve repair vs. replacement in 251 patients with ischemic secondary MR showed that 38% were women while the remaining 62% were men ⁹. In the present study ischemic etiology of secondary MR was observed in 61% of men vs. 37% of women. When analyzing the clinical and echocardiographic characteristics of the patients enrolled in the CTNS trial ⁹, men were comparable to women in terms of age, body mass index, frequency of atrial fibrillation, history of myocardial infarction and renal dysfunction; however, men more frequently had a history of ventricular arrhythmias while women presented more frequently with diabetes mellitus and hypertension. Furthermore, men had larger LV dimensions and effective regurgitant orifice areas when compared to women. However, when the effective orifice regurgitant orifice area was corrected for the LV end-diastolic volume, women had a larger ratio than men suggesting more severe MR in women than in men. These results are similar to the data reported in this study. However, it should be noted that the present study population is larger and includes both etiologies of secondary MR (ischemic and non-ischemic). Although other studies have compared sex-differences in MR^{19, 20}, this is the first and largest study focusing specifically on secondary MR.

Sex differences in outcomes in patients with secondary MR

Although men with significant secondary MR seem to receive interventional treatment more often than women,^{1, 21-23} men seem to have a worse prognosis. A study by Estevez-Loureiro *et al.* ⁸ investigated the effect of gender on results after transcatheter edge-to-edge MV repair with the Mitraclip in 173 patients (64 women vs. 109 men) and showed no differences between the sexes in MR reduction, heart failure hospitalization and all-cause mortality. Similarly, data from the German TRANscatheter Mitral Valve Interventions (TRAMI) registry, including 501 men and 327 women showed no differences in prognosis after successful transcatheter MV repair with the MitraClip ²⁰.

Although, those studies consisted mainly of patients with secondary MR, a significant proportion of patients had other etiologies of MR and sub-analyses concerning only patients with secondary MR were not performed. In contrast, in the randomized CTNS trial, Giustino *et al.* ⁹ showed that among patients with secondary MR undergoing MV repair or replacement, women had a significantly higher risk for mortality when compared to men (27.1% vs. 17.4%, respectively; $P= 0.03$). The reasons for the worse outcome in women after MV surgery remain unclear. However, it is important to note that both men and women showed similar LV reverse remodeling at 2 years follow-up while women had smaller baseline LV volumes and larger effective regurgitant orifice areas when corrected for LV end-diastolic volume as compared to men, suggesting that women had more severe MR than men at baseline and, after correcting the volume overload with MV repair or replacement, their LVs benefited less. Data on more sensitive measures of LV systolic function such as LV GLS or tissue characterization with cardiac magnetic resonance were not available, but it can be speculated that women have stiffer, more fibrotic LVs than men.

In the present study, including ischemic and non-ischemic secondary MR, male gender was independently associated with worse outcome. However, the differences between men and women became evident 3 years after diagnosis of significant secondary MR (Figure 1) which could indicate progression of the underlying mechanism of secondary MR (progression of LV remodelling). Although no statistically significant difference was seen between men and women with ischemic heart failure (possibly due to the low number of women with ischemic heart failure in the present study), a trend of worse survival is seen in men with ischemic heart failure (Figure 2, Panel A). In men, the etiology of secondary MR was ischemic heart failure which is known to respond less well to heart failure therapies and may further progress over time, leading to worse outcome ²⁴⁻²⁶.

Study limitations

This is a single center retrospective study, which limits the generalizability of the results. In addition, this is a tertiary referral center where the patients are referred for specific treatments which may have introduced some selection bias. Measures for heart failure, such as NT-proBNP unfortunately were not systematically available and therefore could not be provided for the present study. Cardiac mortality was not systematically documented in our centre and due to the retrospective design of the study these data could not be acquired. However, to the best of our knowledge, this is the first and largest registry evaluating gender differences and long-term outcome in heart failure patients with secondary MR. The present study did not evaluate changes

over time in LV systolic function and dimensions which could shed more light into the association between heart failure progression and outcomes.

Conclusion

Significant secondary MR is more frequently observed in men as compared to women and is associated with worse prognosis.

References

1. Iung B, Delgado V, Rosenhek R, Price S, Prendergast B, Wendler O, De Bonis M, Tribouilloy C, Evangelista A, Bogachev-Prokophiev A, Apor A, Ince H, Laroche C, Popescu BA, Pierard L, Haude M, Hindricks G, Ruschitzka F, Windecker S, Bax JJ, Maggioni A, Vahanian A. Contemporary Presentation and Management of Valvular Heart Disease: The EURObservational Research Programme Valvular Heart Disease II Survey. *Circulation* 2019.
2. Yadgir S, Johnson CO, Aboyans V, Adebayo OM, Adedoyin RA, Afari-deh M, Alahdab F, Alashi A, Alipour V, Arabloo J, Azari S, Barthelemy CM, Benziger CP, Berman AE, Bijani A, Carrero JJ, Carvalho F, Daryani A, Duraes AR, Esteghamati A, Farid TA, Farzadfar F, Fernandes E, Filip I, Gad MM, Hamidi S, Hay SI, Ilesanmi OS, Irvani SSN, Jurisson M, Kasaeian A, Kengne AP, Khan AR, Kisa A, Kisa S, Kolte D, Manafi N, Manafi A, Mensah GA, Mirrakhimov EM, Mohammad Y, Mokdad AH, Negoii RI, Nguyen HLT, Nguyen TH, Nixon MR, Otto CM, Patel S, Pilgrim T, Radfar A, Rawaf DL, Rawaf S, Rawasia WF, Rezapour A, Roever L, Saad AM, Saadatagah S, Senthilkumaran S, Sliwa K, Tesfay BE, Tran BX, Ullah I, Vaduganathan M, Vasankari TJ, Wolfe CDA, Yonemoto N, Roth GA. Global, Regional, and National Burden of Calcific Aortic Valve and Degenerative Mitral Valve Diseases, 1990-2017. *Circulation* 2020.
3. Monteagudo Ruiz JM, Galderisi M, Buonauro A, Badano L, Aruta P, Swaans MJ, Sanchis L, Saraste A, Monaghan M, Theodoropoulos KC, Papitsas M, Liel-Cohen N, Kobal S, Bervar M, Berlot B, Filipatos G, Ikonomidis I, Katsanos S, Tanner FC, Cassani D, Faletra FF, Leo LA, Martinez A, Matabuena J, Grande-Trillo A, Alonso-Rodriguez D, Mesa D, Gonzalez-Alujas T, Sitges M, Carrasco-Chinchilla F, Li CH, Fernandez-Golfin C, Zamorano JL. Overview of mitral regurgitation in Europe: results from the European Registry of mitral regurgitation (EuMiClip). *Eur Heart J Cardiovasc Imaging* 2018;**19**(5):503-507.
4. Vakamudi S, Jellis C, Mick S, Wu Y, Gillinov AM, Mihaljevic T, Cosgrove DM, Svensson L, Cho L. Sex Differences in the Etiology of Surgical Mitral Valve Disease. *Circulation* 2018;**138**(16):1749-1751.
5. Seeburger J, Eifert S, Pfannmuller B, Garbade J, Vollroth M, Misfeld M, Borger M, Mohr FW. Gender differences in mitral valve surgery. *Thorac Cardiovasc Surg* 2013;**61**(1):42-6.
6. Kisilitsina ON, Zareba KM, Bonow RO, Andrei AC, Kruse J, Puthumana J, Akhter N, Chris Malaisrie S, McCarthy PM, Rigolin VH. Is mitral valve disease treated differently in men and women? *Eur J Prev Cardiol* 2019;**26**(13):1433-1443.
7. Vassileva CM, Stelle LM, Markwell S, Boley T, Hazelrigg S. Sex differences in procedure selection and outcomes of patients undergoing mitral valve surgery. *Heart Surg Forum* 2011;**14**(5):E276-82.
8. Estevez-Loureiro R, Settergren M, Winter R, Jacobsen P, Dall'Ara G, Sondergaard L, Cheung G, Pighi M, Ghione M, Ihlemann N, Moat NE, Price S, Streit Rosenberg T, Di Mario C, Franzen O. Effect of

- gender on results of percutaneous edge-to-edge mitral valve repair with MitraClip system. *Am J Cardiol* 2015;**116**(2):275-9.
9. Giustino G, Overbey J, Taylor D, Ailawadi G, Kirkwood K, DeRose J, Gillinov MA, Dagenais F, Mayer ML, Moskowitz A, Bagiella E, Miller M, Grayburn P, Smith PK, Gelijns A, O'Gara P, Acker M, Lala A, Hung J. Sex-Based Differences in Outcomes After Mitral Valve Surgery for Severe Ischemic Mitral Regurgitation: From the Cardiothoracic Surgical Trials Network. *JACC Heart Fail* 2019;**7**(6):481-490.
 10. Lampert BC, Lindenfeld J, Abraham WT. Too Different or Too Late?: Gender Differences in Outcomes After Mitral Valve Surgery. *JACC Heart Fail* 2019;**7**(6):491-492.
 11. Du BOIS D, Du BOIS EF. CLINICAL CALORIMETRY: TENTH PAPER A FORMULA TO ESTIMATE THE APPROXIMATE SURFACE AREA IF HEIGHT AND WEIGHT BE KNOWN. *JAMA Internal Medicine* 1916;**XVII**(6_2):863-871.
 12. Lang RM, Badano LP, Mor-Avi V, Afilalo J, Armstrong A, Ernande L, Flachskampf FA, Foster E, Goldstein SA, Kuznetsova T, Lancellotti P, Muraru D, Picard MH, Rietzschel ER, Rudski L, Spencer KT, Tsang W, Voigt JU. Recommendations for cardiac chamber quantification by echocardiography in adults: an update from the American Society of Echocardiography and the European Association of Cardiovascular Imaging. *Eur Heart J Cardiovasc Imaging* 2015;**16**(3):233-70.
 13. Foster E, Wasserman HS, Gray W, Homma S, Di Tullio MR, Rodriguez L, Stewart WJ, Whitlow P, Block P, Martin R, Merlino J, Herrmann HC, Wiegers SE, Silvestry FE, Hamilton A, Zunamon A, Kraybill K, Gerber IL, Weeks SG, Zhang Y, Feldman T. Quantitative assessment of severity of mitral regurgitation by serial echocardiography in a multicenter clinical trial of percutaneous mitral valve repair. *Am J Cardiol* 2007;**100**(10):1577-83.
 14. Grayburn PA, Carabello B, Hung J, Gillam LD, Liang D, Mack MJ, McCarthy PM, Miller DC, Trento A, Siegel RJ. Defining "severe" secondary mitral regurgitation: emphasizing an integrated approach. *J Am Coll Cardiol* 2014;**64**(25):2792-801.
 15. Lancellotti P, Tribouilloy C, Hagendorff A, Popescu BA, Edvardsen T, Pierard LA, Badano L, Zamorano JL. Recommendations for the echocardiographic assessment of native valvular regurgitation: an executive summary from the European Association of Cardiovascular Imaging. *Eur Heart J Cardiovasc Imaging* 2013;**14**(7):611-44.
 16. Negishi K, Negishi T, Kurosawa K, Hristova K, Popescu BA, Vinereanu D, Yuda S, Marwick TH. Practical guidance in echocardiographic assessment of global longitudinal strain. *JACC Cardiovasc Imaging* 2015;**8**(4):489-492.
 17. Regitz-Zagrosek V, Oertelt-Prigione S, Prescott E, Franconi F, Gerds E, Foryst-Ludwig A, Maas AH, Kautzky-Willer A, Knappe-Wegner D, Kintscher U, Ladwig KH, Schenck-Gustafsson K, Stangl V. Gender in cardiovascular diseases: impact on clinical manifestations, management, and outcomes. *Eur Heart J* 2016;**37**(1):24-34.
 18. Enriquez-Sarano M, Akins CW, Vahanian A. Mitral regurgitation. *Lancet* 2009;**373**(9672):1382-94.
 19. Dziadzko V, Clavel MA, Dziadzko M, Medina-Inojosa JR, Michelena

- H, Maalouf J, Nkomo V, Thapa P, Enriquez-Sarano M. Outcome and undertreatment of mitral regurgitation: a community cohort study. *Lancet* 2018;**391**(10124):960-969.
20. Werner N, Puls M, Baldus S, Lubos E, Bekeredjian R, Sievert H, Schofer J, Kuck KH, Möllmann H, Hehrlein C, Nickenig G, Boekstegers P, Ouarrak T, Senges J, Zahn R. Gender-related differences in patients undergoing transcatheter mitral valve interventions in clinical practice: 1-year results from the German TRAMI registry. *Catheter Cardiovasc Interv* 2020;**95**(4):819-829.
 21. Acker MA, Parides MK, Perrault LP, Moskowitz AJ, Gelijns AC, Voisine P, Smith PK, Hung JW, Blackstone EH, Puskas JD, Argenziano M, Gammie JS, Mack M, Ascheim DD, Bagiella E, Moquete EG, Ferguson TB, Horvath KA, Geller NL, Miller MA, Woo YJ, D'Alessandro DA, Ailawadi G, Dagenais F, Gardner TJ, O'Gara PT, Michler RE, Kron IL. Mitral-valve repair versus replacement for severe ischemic mitral regurgitation. *N Engl J Med* 2014;**370**(1):23-32.
 22. Obadia JF, Messika-Zeitoun D, Leurent G, lung B, Bonnet G, Piriou N, Lefevre T, Piot C, Rouleau F, Carrie D, Nejari M, Ohlmann P, Leclercq F, Saint Etienne C, Teiger E, Leroux L, Karam N, Michel N, Gilard M, Donal E, Trochu JN, Cormier B, Armoiry X, Boutitie F, Maucort-Boulch D, Barnel C, Samson G, Guerin P, Vahanian A, Mewton N. Percutaneous Repair or Medical Treatment for Secondary Mitral Regurgitation. *N Engl J Med* 2018.
 23. Stone GW, Lindenfeld J, Abraham WT, Kar S, Lim DS, Mishell JM, Whisenant B, Grayburn PA, Rinaldi M, Kapadia SR, Rajagopal V, Sarembock IJ, Brieke A, Marx SO, Cohen DJ, Weissman NJ, Mack MJ. Transcatheter Mitral-Valve Repair in Patients with Heart Failure. *N Engl J Med* 2018.
 24. Cleland J, Freemantle N, Ghio S, Fruhwald F, Shankar A, Marijanowski M, Verboven Y, Tavazzi L. Predicting the long-term effects of cardiac resynchronization therapy on mortality from baseline variables and the early response a report from the CARE-HF (Cardiac Resynchronization in Heart Failure) Trial. *J Am Coll Cardiol* 2008;**52**(6):438-45.
 25. Ferreira JP, Duarte K, McMurray JJV, Pitt B, van Veldhuisen DJ, Vincent J, Ahmad T, Tromp J, Rossignol P, Zannad F. Data-Driven Approach to Identify Subgroups of Heart Failure With Reduced Ejection Fraction Patients With Different Prognoses and Aldosterone Antagonist Response Patterns. *Circ Heart Fail* 2018;**11**(7):e004926.
 26. Kloosterman M, van Stipdonk AMW, ter Horst I, Rienstra M, Van Gelder IC, Vos MA, Prinzen FW, Meine M, Vernooij K, Maass AH. Association between heart failure aetiology and magnitude of echocardiographic remodelling and outcome of cardiac resynchronization therapy. *ESC Heart Failure* 2020;**7**(2):645-653.

