



Universiteit
Leiden
The Netherlands

Reproductive and sexual health care in oncology: current practice and challenges

Krouwel, E.M.

Citation

Krouwel, E. M. (2022, May 12). *Reproductive and sexual health care in oncology: current practice and challenges*. Retrieved from <https://hdl.handle.net/1887/3303552>

Version: Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/3303552>

Note: To cite this publication please use the final published version (if applicable).



Chapter 10

An Educational Need regarding Treatment-related Infertility and Fertility Preservation; A National Survey among Members of the Dutch Society for Medical Oncologists

E.M. Krouwel, E.M.L. Birkhoff, M.P.J. Nicolai, S. Osanto, H. Putter,
R.C.M. Pelger, H.W. Elzevier

J Cancer Educ 2021 Oct. Online ahead of print.

INTRODUCTION

Cancer treatments are associated with a variety of undesirable side effects; of which one of specific concern to young men and women is the effect on their endocrine health and future reproductive ability [38]. Due to the increasing 5-year, and overall survival over the past few decades, consideration for physical and psychological consequences becomes progressively prioritized [11]. Loss of fertility is a devastating side effect for young cancer survivors with severe emotional impact [2], and specifically resulting from treatment with chemotherapy or radiation therapy [16, 27, 32]. Moreover, the prospect of facing treatment-induced infertility for women of reproductive age is proven to affect their cancer treatment decisions in up to 41% [22, 34].

Cytostatic cancer drugs are designed to target dividing cells; implying that in addition to inhibiting cancer cell growth, proliferating primordial follicles which enfold oocytes are conjointly harmed [6]. As for men, infertility and persistent azoospermia is a common long-term adverse effect [7]. Alkylating drugs are feared most for their effect to fertility, by inducing both impaired fertility and early menopause [33]. Although the exact risk for cancer- and treatment related infertility depends on the chemotherapy agent, the chemotherapeutic regime and the age of the patient, it should not be underestimated considering the long-term impact. Being the physician prescribing cytostatic drugs, the medical oncologist is responsible for informing about the risk of infertility before commencing a gonad toxic treatment, referral to a reproductive specialist for fertility preservation (FP) and discussing alternative treatment strategies if applicable. Nonetheless, various studies suggest the information provision regarding fertility issues is often experienced as inadequate by patients. Furthermore, it is suggested that cancer care physicians do not possess sufficient knowledge regarding fertility risks and options for FP [36, 25, 8, 13]. As a result, information is not timely provided or in some cases is not provided at all [3, 4, 23]. At the time of diagnosis, fertility issues are often outweighed by the focus on survival. A Dutch observational study, showed in 2011, a total of 9.8% of female patients were referred for FP counselling. However, the absolute numbers of patients receiving FP counselling increased over time [5]. And indeed, informing cancer patients of reproductive age about possible reduced fertility and referral to a reproductive specialist in a timely manner is recommended by national [18, 29], European [31] and international guidelines [24]. Fertility counselling performed by a fertility specialist prior to cancer treatment, in comparison to the oncologist alone, is associated with better psychological health. Those patients who undertook counselling and proceeded with FP reported reduced regret, compared with those who did not proceed to FP [22]. Patients who felt fertility concerns have not been given full consideration at the time of diagnosis have been shown to cope with psychological distress, expressed by uncertainty and concern, as well as higher levels of depression and cancer or fertility-related trauma during survivorship [22].

Over the last decade, we have experienced a surge of scientific reports on aspects of altered fertility in young adults with cancer; these particularly include the growing number of available preservation options [1] and the devastating impact of the loss of fertility to cancer survivors

[9, 30, 37]. In addition, various studies investigated practice regarding (referral for) FP counselling by physicians involved with oncology patients. An overview of quantitative studies among oncology care providers published in the past ten years regarding knowledge level, the discussion of fertility, referral to fertility specialists and barriers regarding this discussion is provided in Supplement 1. Pediatric studies have been excluded in this overview.

To date, published quantitative surveys have suggested there may be a deficiency in medical oncologists' knowledge about FP options and that the provision of information to patients about FP may be suboptimal. The purpose of the hereby presented nationwide study was to evaluate Dutch medical oncologists' practice patterns, knowledge, educational need, attitudes and barriers regarding treatment-related infertility and FP among men and women of reproductive age.

METHODS

Study design and cohort identification

A questionnaire was used for collecting data in a cross-sectional postal survey. The sample consisted of 433 members of the NVMO (Dutch Society for Medical Oncology) with several areas of expertise. Our sampling strategy aimed for representation with regard to different tumors, employment setting, level of education, years of oncology experience, type of hospital, age and gender.

Instrument design and development

The questionnaire was developed by the authors. The content of the questionnaire was evaluated by 4 oncologists working in Leiden University Medical Center through an anonymous pilot study and modified using their feedback. The final version comprised a demographic sheet, including professional background, experience in oncology practice, gender and age. Furthermore, Likert-scale items measured practices, attitudes, content of sexual and fertility counselling, responsibility, need for education and barriers regarding the discussing of sexual function and fertility issues. In addition, a list was made of most common medications measuring knowledge about their possible side effects to sexual function, to future reproductive ability and regarding teratogenicity. Data concerning the discussing of sexuality issues and knowledge about medication were processed separately [19].

Survey administration

The questionnaires were sent to all medical oncologists who were member of the NVMO, members not practicing in the Netherlands have been excluded. After the initial mailing was finished, reminders were sent to non-responders after 6 and 12 months. An information letter concerning the study and a post-paid return envelope were added, as well as an opt-out possibility. Data were collected anonymously in order to prevent a self-reporting bias.

Supplement 1. Overview of quantitative surveys among care providers regarding knowledge, practice, referral and barriers about the discussing of fertility and FP with cancer patients of childbearing age; 1) conducted/published in the past 10 years, 2) excluding qualitative studies and studies concerning pediatric care providers.

Study	Year of survey	Country	Study population	Type of care providers	Eligible participants	Final participants	Completion rate (%) _a	Female (a) and male (b) patients	Knowledge of FP (%)	Discussing fertility and referral (%) _b	Two most mentioned barriers (%)
Krouwel <i>et al.</i> [current survey, 2021]	2013-2014	The Netherlands	Dutch Society for Medical Oncology	Medical oncologists	392	120	30.6%	a, b	47.5% sufficient knowledge 76% wants to improve knowledge	Discussing fertility: 68.3% often/always. Referral: 44.6% men, 28.9% women	Prognosis is poor 53%, unlikely patient will survive 43.1%
Vesali <i>et al.</i> [49]	2015-2016	Iran	Attendees congresses	Radiation oncologists and hematologists	131	103	78.6%	a, b	Knowledge GnRH/sperm cryopreservation 2.77 and 2.64	Information provision FP 13.6%. Referral 71.8%.	N.A.
Post <i>et al.</i> [38]	2016	United States	Nationwide	Radiation Oncologists	N.A.	352	N.A.	b	N.A.	Fertility counselling always recommended by 52%	N.A.
Sallem <i>et al.</i> [44]	2012-2013	France	Nationwide	Oncologists	N.A.	102	N.A.	a, b	14% knowledgeable regarding FP	Discussing fertility by 46%, referral 22%	Poor prognosis 54%, urgent treatment 51%
Melo <i>et al.</i> [33]	2013-2015	Portugal	Clinical institutions, Portuguese Oncology Society	Doctors who assist female cancer patients	N.A.	111	N.A.	a	N.A.	Fertility risks discussed 65.7%, preservation 59.3%. Referral 7.38%.	Time, knowledge.
Louwe <i>et al.</i> [30]	Unknown	The Netherlands	Seven of eight regions (85% of the country).	Hematologists, oncologists, radiotherapists, surgeons, gynecologists	406	280	69%	a	Knowledge treatment effect 55-68% chemotherapy 55%, radiation 68%. FP knowledge 13-36%.	79% usually or always discuss fertility issues.	Lack of available reproductive specialists 11%, insufficient time 10%

Supplement 1. Overview of quantitative surveys among care providers regarding knowledge, practice, referral and barriers about the discussing of fertility and FP with cancer patients of childbearing age; 1) conducted/published in the past 10 years, 2) excluding qualitative studies and studies concerning pediatric care providers. (*continued*)

Study	Year of survey	Country	Study population	Type of care providers	Eligible participants	Final participants	Completion rate (%) _a	Female (a) and male (b) patients	Knowledge of FP (%)	Discussing fertility and referral (%) _b	Two most mentioned barriers (%)
Kcimm-Malpass <i>et al.</i> [23]	2014	United States	NCI-designated cancer centers	Nurses	94	52	55.3%	a, b	adds uncertainty and concern, alongside higher levels of depression (not specific to fertility; CES-D) and cancer or fertility-related traumas	Rarely/never discuss FP by 73.1%. Referral N.A.	Lack of available information, patient cannot afford FP
Chung <i>et al.</i> [13]	2016	China (Hong Kong)	Clinicians working in Hong Kong	Oncology; hematology, gynecology; pediatricians, surgery physicians	467	167	36.5%	a, b	45.6% reported familiarity with FP.	Percentage of discussing N.A. 68.3% never referred for FP.	No time before treatment 60.6%, risk of cancer recurrence 53.8%
Takeuchi <i>et al.</i> [47]	Unknown	Japan	Nationwide	Physicians involved with cancer patients	412	180	43.7%	a, b	N.A.	Discussing fertility 42.7% sometimes, always. Referral N.A.	Insufficient knowledge
Shah <i>et al.</i> [46]	Unknown	United States	Members Society of Gynecologic Oncology	Gynecologic oncologists	1087	152	14%	a	N.A.	Assess fertility status 68% always. Referral easy according 64.9%.	Predictors: number of young women, cancer stage.
Rosenberg <i>et al.</i> [40]	2015	United States	Oncologists involved in endocrine studies.	Medical oncologists caring for patients with breast cancer	301	93	31%	a	N.A.	Discussing fertility 98%, referral 97%	Cost/insurance 47%; patient does not want to discuss FP 27%.

Supplement 1. Overview of quantitative surveys among care providers regarding knowledge, practice, referral and barriers about the discussing of fertility and FP with cancer patients of childbearing age; 1) conducted/published in the past 10 years, 2) excluding qualitative studies and studies concerning pediatric care providers. (continued)

Study	Year of survey	Country	Study population	Type of care providers	Eligible participants	Final participants	Completion rate (%) ^a	Female (a) and male (b) patients	Knowledge of FP (%)	Discussing fertility and referral (%) ^b	Two most mentioned barriers (%)
Micaux <i>et al.</i> [34]	2015	Sweden	National	Physicians in oncology/hematology	821	329	55%	a, b	High knowledge male risk fertility 59%, female risk fertility/early menopause 65%	Discussing fertility: 74% female, 70% male; discussing FP 57% female, 63% male	Poor prognosis 78%, patient anxious or overwhelmed by having cancer 54%
Krouwel <i>et al.</i> [26]	2012	The Netherlands	Members Dutch Oncology Nursing Society	Oncology nurses	N.A.	421	N.A.	a, b	Sufficient knowledge of FP options: 31.1%.	Discussion of FP in more than half of the cases 42.6%	Lack of knowledge 25.2%, poor prognosis 16.4%
Ghazeei <i>et al.</i> [18]	2012-2013	Lebanon	Lebanese Society Medical Oncology, practitioners two medical centers	Oncologists, clinical practitioners (i.e. students, doctors and nurses)	81 oncologists, all practitioners	53 oncologists, 88 practitioners	65.4% for oncologists	a, b	92.3% of oncologists estimated sperm cryopreservation should be done.	73.1% of oncologists routinely discussed, 39.6% regularly refers to fertility specialist	N.A.
Buske <i>et al.</i> [11]	2011-2012	Germany	Oncologists	Oncologists	Unknown.	120	Unknown.	a, b	49.6% knowledge sperm preservation, 37.3% well informed about FP measures.	Discussing possible fertility impairment by 65.8%, FP by 65.3%. Referral by 96.6%	Poor prognosis 62.7%, over 35 years 47%
Biglia <i>et al.</i> [8]	Not reported.	Italy	Breast surgeons, oncologists	Representatives of all Italians regions	Unknown.	181	Unknown.	a	N.A.	91% discusses fertility, 60% referral	N.A.
Louwe <i>et al.</i> [31]	Not reported.	The Netherlands	2 comprehensive cancer centers	Gynecologists, oncologists, surgeons, radiotherapists, hematologists	206	96	46.6%	a	N.A.	62% took action to protect the ovarian function	Poor prognosis or need for immediate therapy 62%, costs

Supplement 1. Overview of quantitative surveys among care providers regarding knowledge, practice, referral and barriers about the discussing of fertility and FP with cancer patients of childbearing age; 1) conducted/published in the past 10 years, 2) excluding qualitative studies and studies concerning pediatric care providers. (*continued*)

Study	Year of survey	Country	Study population	Type of care providers	Eligible participants	Final participants	Completion rate (%) ^a	Female (a) and male (b) patients	Knowledge of FP (%)	Discussing fertility and referral (%) ^b	Two most mentioned barriers (%)
Adams <i>et al.</i> [1]	Not reported.	United Kingdom	Members doctors.net. of GMC registered doctors.	Medical and clinical oncologists	1488	100	N.A.	a, b	Sperm preservation 64% sufficient; ovarian preservation 82% insufficient.	Providing informed consent 97%, 45% routinely referral fertility specialist	Perception FP poor success rates 81%, lack of knowledge 63%
Duffy <i>et al.</i> [16]	Not reported.	United States of America	National; sample American Medical Association.	Oncologists, hematologists, radiation oncologists, gynecologists	1088	344	31.6%	a, b	±50% moderate/high knowledge confidence in fertility/ FP options.	Not reported.	Not reported.
King <i>et al.</i> [24]	Not reported.	United Kingdom	Cancer institutes, networks, associations.	Nurses, nurse specialists, oncology/surgery residents, surgeons, oncologists.	N.A.	306	N.A.	a	Knowledge local preservation options variable.	Always discussed 48%, most of the times discussed 34%.	Age (77%), disease status (37.9%)
Arafa <i>et al.</i> [3]	Not reported.	Saudi Arabia	3 regions; Eastern, Jeddah and Riyadh.	Medical, radiation and surgical oncologists.	180	103	57.2%	a, b	Familiarity with ICSI 49.5%.	Routinely discussing cryopreservation 41.7% Referral 19.5%.	Type of cancer (92%), age of patient (87%)

N.A.: Not available

^a Eligible participants/final participants (%)

^b Discussing of FP depending on how questions have been phrased.

Analysis

Data analysis was performed using SPSS (Release 22; SPSS Inc., USA). Frequency distribution was used to calculate demographic information. Respondents were recoded regarding age (set at median age 47: 46 years or under vs. over 46), experience (0-10 years vs. 11 years or more), knowledge (none to some vs. sufficient to a lot) and residents vs. qualified specialists. Observed differences between demographic information and specific answers were identified using the Pearson's chi-square test, linear-by-linear association, paired T-test and independent sample T-test. *P*-values < 0.05 were considered statistically significant.

Ethics

In The Netherlands, studies that do not involve patients or interventions are not subject to approval from an ethical board. The ethical board was consulted for a comparable previous study, as the study did not concern information recorded by the investigator in such a manner that subjects could be identified, and as it did not compromise the study participants' integrity, the Committee declared that no official ethical approval was needed.

RESULTS

Demographics

Of the 433 invited participants 209 responses were received, resulting in a response-rate of 48.3%. Of the 209 responses, 26 oncologists reported to be retired, 6 physicians were not medical oncologists and 9 were returned to sender. Forty-eight oncologists reported they were not willing to participate due to a lack of time (*n*=35), a lack of interest (*n*=2), a lack of experience (*n*=2), the length of the instrument (*n*=4) or other reasons (*n*=5). Out of 392 eligible participants, 120 oncologists completed the questionnaire (30.6%). Majority of the respondents reported breast cancer (73.3%) and colorectal cancer (65.8%) as area of expertise (multiple areas of expertise possible). For extensive information on the respondents characteristics see Table 1.

Table 1. Demographic characteristics

Demographic characteristics (n=120)	n (%)
Age (years)	
Mean 45.8	119 (99.2)
Median 47 (range 30-64)	
Gender	
Male	56 (46.7)
Female	63 (52.5)
Unknown	1 (0.8)

Table 1. Demographic characteristics (*continued*)

Demographic characteristics (n=120)	n (%)
Oncology experience (years)	
1–2	19 (15.8)
3–5	27 (22.5)
6–10	13 (10.8)
11–15	19 (15.8)
>15	40 (33.3)
Unknown	2 (1.7)
Function	
Oncologist	74 (61.7)
Oncologist and haematologist	15 (12.5)
Haematologist	12 (10)
Resident medical oncology	19 (15.8)
Hospital type	
University hospital	40 (33.3)
Top clinical teaching hospital	27 (22.5)
District general hospital	47 (39.2)
Categorical cancer hospital	3 (2.5)
University hospital and district general hospital	2 (1.7)
Unknown	1 (0.8)
Areas of interest^a	
Breast cancer	88 (73.3)
Colorectal cancer	79 (65.8)
Palliative medicine	57 (47.5)
Urological cancer	55 (45.8)
Gynaecological cancer	53 (44.2)
Haematology	37 (30.8)
Lymphoma	32 (26.7)
Neuroendocrine cancer	15 (12.5)
Head and neck cancer	14 (11.7)
Melanoma	9 (7.5)
Sarcomas	8 (6.7)
Lung cancer	3 (2.5)
Experimental research (phase I-II)	3 (2.5)
Gastro-intestinal	2 (1.7)
Hepatic and biliary tract cancer	1 (0.8)

a) Multiple areas of interest possible.

Practice

When it comes to discussing the impact of cancer treatment to fertility, almost seventy percent of respondents (n=82, 68.3%) stated to discuss fertility often or always, 20% (n=24) declared to discuss it in more than half of the cases, 5.8% (n=7) in half of the cases, 3.3% (n=4) in less than half of the cases and 2.5% (n=3) hardly never/never. Among oncologists working in a district general hospital, it was less usual to discuss fertility. In district general hospitals 84.1% discussed fertility in more than half of the cases vs. 90.3% in all others hospitals ($p=0.033$, linear-by-linear). No significant differences were observed regarding gender, all different specialties, residents vs. oncologists, age through 46 years or older, experience through 10 years or more, level of knowledge and availability of a department protocol.

When cancer treatment has the prospect to cure, almost all respondents (n=114, 98.3%) discuss fertility. Yet, if cancer treatment has no prospect of cure, only half of the respondents (n=61, 52.6%) discuss fertility. When treatment is at palliative stage, a quarter of the respondents stated to discuss fertility (23.3%, n=27).

Fertility is discussed with women until the age of 44 on average (range 35-100 year, SD 8.2, n=114) and on average with men until the age of 53 (range 37-100, SD 10.6, n=107).

Topics that were reported to be discussed with women, are a desire to start a family (n=120, 100%), menopausal symptoms (n=105, 87.5%), fear for premature termination of pregnancy (n=15, 12.5%), fear for congenital abnormalities (n=51, 42.5%) and heredity (n=79, 65.8%). With men, frequently discussed topics were a desire to start a family (n=117, 97.5%), erectile dysfunction (n=60, 50%), ability to ejaculate (n=24, 20%), fear for congenital abnormalities (n=39, 32.5%) and heredity (n=63, 52.5%).

Knowledge

To the question: 'How much knowledge do you possess regarding FP for cancer patients?', 47.5% of the respondents (n=57) answered sufficient knowledge, 41.7% (n=50) had some knowledge, 7.5% (n=9) said they did not have much knowledge and 3.3% (n=4) reported they possessed a lot of knowledge. Oncologists estimated their knowledge significantly higher in comparison to residents (linear-by-linear association $p=0.004$).

Three-quarters of the oncologists (n=86, 75.4%) would like to improve their knowledge towards fertility issues and management of fertility issues. Residents significantly more often wish to improve their knowledge ($p=0.041$). Experience however, is not of significant influence to the wish for more knowledge ($p=0.081$). Almost three-quarters (n=84, 74.3%) of the respondents believe there is too little attention for fertility issues and management of fertility issues during residency. Respondents estimated that initial cryopreservation of semen would cost 693,15 euro (range 30-15000 euro; SD 1801; n=75) with an annual cost for cryopreservation of 103,07 euro (range 0-500 euro; SD 124.8; n=75).

Local practice

Approximately forty percent (n=45, 38.5%) of the respondents reported a protocol or a standard at their current workplace stating the routine discussing of fertility, 41.9% (n=49) did not have such a protocol and 19.7% (n=23) was not aware of such a protocol. During multidisciplinary oncology meetings, according to 37.1% (n=43) fertility is regularly discussed. Seventy-three oncologists (62.9%) reported fertility is not routinely discussed in multidisciplinary meetings. Half of the oncologists reported there is sufficient patient information available in their department regarding fertility (n=56, 48.3%), a quarter (25%, n=29) reported there is not sufficient information and the remaining quarter (26.7%, n=31) did not know whether patient information is available. Seventy percent often or always registered fertility consultation in the patient's file (n=82), 12 percent (n=14) does so in more than half of the cases and 21 oncologists (18%) reported to register in half of the cases or less. Eighty-four percent (n=101) never/hardly never prescribed a gonadotropin-releasing hormone agonist (GnRH-A) before starting chemotherapy, for protection of the ovaria. Twelve oncologists (10%) did so in less than half of the cases, 2.5% (n=3) in half of the cases and 3.3% (n=4) did so in more than half of the cases or almost always.

Responsibility

According to 36.4% (n=43), responsibility for the discussion of fertility was assigned to their department, 53 oncologists (44.9%) stated there were no agreements regarding responsibility and 18.6% (n=22) did not know if there were agreements. Majority of the participants 93.2% (n=110, question answered by n=118) was convinced that it is the responsibility of the oncologist to discuss fertility with patients of reproductive age, 5.8% (n=7) disagreed to this responsibility. One oncologist did not know whether it is within the treatment responsibility to discuss fertility. In addition, 78% (n=92) believed that there is also a responsibility for the oncology nurse, 17% (n=20) did not believe that it is the oncology nurses' responsibility to address fertility and 5.1% (n=6) did not know if it should be oncology nurses' responsibility.

Barriers

The respondents were given a list of 30 possible barriers for discussing fertility, in order for them to indicate to which extent they agreed. The three barriers most agreed upon by the respondents were: 'Prognosis is poor' (53%), 'Unlikely patient will survive treatment' (43.1%) and 'High chance on fertility recovery after treatment' (28.7%). The three the barriers most disagreed upon by respondents were: 'Patient cannot afford treatment' (91.3%), 'Patient is single' (90.6%) and 'No contact information available of fertility specialist' (88.8%). A complete overview of barriers can be found in Table 2.

Table 2. Barriers towards discussing fertility

I would tend not to discuss fertility with a patient because:	Agree	Neither agree nor disagree	Disagree
	n (%)	n (%)	n (%)
Patient cannot afford treatment	1 (0.9)	9 (7.8)	105 (91.3)
Patient is single	2 (1.7)	9 (7.7)	106 (90.6)
No contact information available of fertility specialist	2 (1.7)	11 (9.5)	103 (88.8)
Patient is a teenager	3 (2.8)	4 (3.7)	101 (93.5)
This may raise fear and discomfort	3 (2.5)	20 (17)	95 (80.5)
Semen cryopreservation is not adolescent friendly	3 (2.7)	20 (17.7)	90 (79.6)
Uncomfortable to discuss fertility	4 (3.5)	23 (20.4)	86 (76.2)
Patient already has a child/children	5 (4.3)	22 (18.8)	90 (76.9)
Semen cryopreservation is expensive for patient	5 (4.5)	28 (25.2)	78 (70.3)
Hereditary tumor (risk of passing to child)	7 (6)	17 (14.5)	93 (79.5)
Pregnancy during or after chemotherapy may induce malformation of child	8 (6.9)	26 (22.4)	82 (70.7)
I do not possess enough knowledge regarding fertility preservation options	8 (6.8)	27 (23.1)	82 (70.1)
Patient is homosexual	10 (9)	19 (17.1)	82 (73.9)
Fear of possible malignant transformation of ovarian tissue	11 (9.7)	18 (15.9)	84 (74.3)
Fertility treatment may influence success of cancer treatment	12 (10.3)	27 (23)	79 (66.7)
Treatment delay	12 (10.3)	27 (23.1)	78 (66.7)
Patient does not bring up the subject	13 (11)	25 (21.2)	80 (67.8)
Possibility of reintroducing cancer or metastasis by ovarian tissue cryopreservation	13 (11.4)	26 (22.8)	75 (65.8)
Culture or religion of patient does not support assisted fertility	13 (11.5)	32 (28.3)	68 (60.2)
Curation has priority	19 (16.2)	33 (28.2)	65 (55.6)
Difficult to predict risk for deteriorated fertility	19 (16.4)	33 (28.4)	64 (55.2)
Ovarian tissue cryopreservation is experimental	20 (17)	36 (30.5)	62 (52.5)
Patient is treated before with chemo and/or radiotherapy	23 (19.8)	30 (25.9)	63 (54.3)
Lack of time during consult	25 (21.4)	22 (18.8)	70 (59.8)
Patient is HIV positive	28 (25.7)	26 (23.9)	55 (50.4)
Hormonal treatment may increase risk of recurrence	28 (25)	31 (27.7)	53 (47.3)
Patient is not able to ejaculate, therefore cryopreservation is not possible	31 (27.4)	22 (19.5)	60 (53.1)
High chance on rapid recovery of reproductive function after treatment	33 (28.7)	31 (27)	51 (44.3)
Unlikely patient will survive treatment	50 (43.1)	25 (21.6)	41 (35.3)
Prognosis is poor	62 (53)	27 (23.1)	28 (23.9)

For ease of presentation, results in response categories 'Strongly agree' and 'agree' have been merged, as have 'strongly disagree' and 'disagree'. Total number of respondents may differ per barrier, as some respondents skipped barriers.

Referral to fertility specialist

When asked which percentage of all patients of reproductive age has been referred to a fertility specialist, on average 44.6% of men (range 0-100%; SD 37.1), and on average 28.9% of women (range 0-100%; SD 31.4), is referred. The percentage of women being referred is significantly

lower in comparison to men ($p < 0.001$, paired T-test). The percentage of men being referred was more often by oncologists of 47 years and older ($p = 0.028$, ind. sample T test) and by residents ($p = 0.001$, ind. sample T test). There were no significant differences in gender or hospital type of the respondents in reference to the percentage of referred patients.

Ethics

The participating oncologists were asked to give their opinion upon post-mortem use of preserved material for assistant reproduction for the partner. Half of the oncologists ($n = 56$) believed this is acceptable, 39 oncologists (35.5%) believed this is not acceptable and 15 oncologists (13.6%) were not aware of the existence of this possibility.

DISCUSSION

Main findings

Considering it is crucial that medical oncologists address the impact of cancer treatment to fertility with patients of childbearing age, this survey intended to represent current practice and knowledge among medical oncologists practicing in the Netherlands. Main findings of our study demonstrate an accountable attitude with regards to fertility issues among responding oncologists, yet about two-third of the participants stated to often or always discuss the impact to fertility. Practice behavior is mainly influenced by patients' prognosis, type of hospital and fertility recovery chances. On average, 44.6% of reproductive men and 28.9% of reproductive women are referred to a fertility specialist. Half of our respondents said to possess sufficient knowledge concerning FP. Three-quarter of the oncologists believed too little training is paid to the subject during residency and expressed a wish for additional education on fertility issues and preservation options.

Comparison to literature and interpretation of findings

In the past decade, several international surveys have been performed amongst oncology care providers regarding the provision of FP (Supplement 1). Response-rates of previous surveys differ from 14 to 78.6 percent (mean 47.3%). Much of what is known about fertility and cancer is the result of studies conducted in the US, the UK and other countries. Two studies have been performed in the Netherlands, reflecting on the practice of physicians from several different cancer specialties, not solely medical oncologists. Our findings indicate that responsibility for fertility concerns is acknowledged by oncologists, however in practice the discussing of fertility concerns may vary. Other surveys among oncologists across the world show greatly varying results, with discussing percentages ranging from 13.6 to 98 percent and referral percentages from 15 to 97 percent (Supplement 1). Although the counselling percentages vary in countries

and regions, it is clear that we are facing a generic concern probably applicable to all cancer care institutes in a greater or lesser extent.

In comparison to previous surveys worldwide, the surveyed oncologists score average on discussing fertility concerns in practice. As for referral to a fertility specialist, scores of the Dutch oncologists are slightly lower than average. However, most of the reviewed surveys were conducted in single centers, selected populations (e.g. among oncologists who had previously enrolled women on premenopausal studies) or demarcated areas, and often only investigated FP regarding female cancer or breast cancer patients as seen in Supplement 1. Therefore the questioned populations may be biased, as local practices may differ significantly. Differences in several types of practices are empowered by our finding that oncologists employed in a district general hospital were less likely to discuss fertility issues. This proven variety in practice between district general hospitals and top clinical- and university hospitals may be explained by limited access to fertility departments which are usually located in university or top clinical teaching hospitals. This phenomenon was previously observed by Hariton *et al.*, showing the association between the establishment of a oncofertility clinic and increased consultations for FP [17]. In addition, a recent Dutch survey showed the relevance of available reproductive specialists with specific expertise with regards to women with cancer, as a lack of available specialists was reported a major barrier against discussing fertility issues with patients [25]. Furthermore, Louwe *et al.* revealed a positive correlation between the number of FP options available and the number of information sources available in regards to confidence in the physicians' knowledge [25]. Negative correlation was shown between the frequency of discussion fertility issues and a lack of reproductive specialists in the geographic region, which is very similar to our results.

Barriers most mentioned by our respondents, were a poor prognosis and unlikelihood the patient will survive the treatment. In comparison to literature, these are often mentioned barriers towards discussing fertility issues by clinicians working in oncology departments. By way of comparison, in a Swedish survey the barrier 'poor prognosis' was mentioned by 78% [28], in a German survey by 62.7% [8], in a Dutch pilot survey by 62% [26], in a Canadian survey by 66.4% [39], in a UK survey by 78.6% [15] and in an American survey by 30% [14]. Besides medical reasons, one out of five oncologists stated lack of time during the consult as a barrier towards discussing fertility.

In addition we asked the clinicians to estimate costs of semen cryopreservation. Estimations of the costs of semen cryopreservation were variable, on average the estimated costs were fairly overestimated. Specifying, costs for initial cryopreservation were estimated €693,15, actual costs are €119,82, with additional costs per sample of €62,81 (reimbursements 2018). Annual costs for cryopreservation were estimated €103,07, actual annual costs are €60,12 (reimbursements 2018). As the costs are fairly overestimated (specifically the initial costs), patients may be informed incorrectly by their clinicians. In some cases, this may result in the decision to withhold from cryopreservation, an undesirable consequence.

Clearly, there is a reported lack of training for fertility issues and their management during residency. Consequently, a wish for additional education is expressed by the majority of the respondents, implying a major role for the development of training courses and implementation of the subject fertility issues during residency. By these means, early referral by oncologists before initiation of chemotherapy and radiotherapy will be enabled, as this is a key factor for success in (female) FP [35].

Strengths and limitations

The completion rate of 30.6% is lower than the average response rate of physicians surveys [10], also on the lower limits in comparison to physicians surveys performed in the Netherlands with response rates running from 28-55% [21, 20, 12]. The completion rate may be explained by the length of the questionnaire, the content of the questionnaire (assessing treatment side-effect knowledge, which may be embarrassing if unfamiliar with this knowledge) as well as the sensitivity of the topics sexuality and fertility. Yet, a non-response bias may have occurred. Oncologists with affinity for the subject may have been more inclined to answer than oncologists who are less committed to fertility concerns. Demographic characteristics of non-responders have not been made available, consequently non-response calculations could not be made. A non-validated questionnaire has been used as a validated instrument was not available. Nevertheless, a pilot study has been conducted to check for validity and reliability. Subdivision by area of specialization resulted in small numbers of medical oncologists in each specialization group. Accordingly, it was not possible to do sub analyses for every separate area of specialization.

Clinical implications

Awareness and sufficient knowledge among medical oncologists regarding possible toxic effects to endocrine and reproductive health is of critical importance for young men and women with cancer. Due to a lack of knowledge, referral possibilities and counselling barriers the ability to start or complete a family after treatment may not be extensively discussed. Subsequently many men and women of reproductive age with cancer could be missing the opportunity to investigate their FP options. We recommend expansion of education of fertility treatment risks and preservation options starting in medical school, continued during residency training and updates when practicing as a medical oncologist. A culture of shared decision making should be pursued, through the development of clear fertility referral pathways including psychosocial support to improve care for men and women of childbearing age facing a cancer treatment.

CONCLUSION

In conclusion, the results suggest that medical oncologists take responsible attitudes towards fertility preservation in oncology practice. Self-reported knowledge regarding fertility preserva-

tion is strongly varying and the majority expressed a wish for additional education. Practice attitudes remain under influence of factors like poor prognosis, a lack of knowledge, treatment-delay and local availability of fertility specialists. Efforts to develop educational training on treatment fertility risks, communication skills and acquaintance with fertility preservation options are highly recommended. Improvement of awareness regarding fertility preservation and in addition availability of fertility specialists in district general hospitals may increase referral of young cancer patients for fertility preservation. Timely referral to discuss preserving options for endocrine and reproductive health is crucial, before irreversible damage to the gonads is done.

REFERENCES

1. Ajala, Tosin, Junaid Rafi, Peter Larsen-Disney, and Richard Howell. 2010. Fertility Preservation for Cancer Patients: A Review. *Obstetrics and Gynecology International* 2010:160386. doi:10.1155/2010/160386.
2. Armuand, G. M., L. Wettergren, K. A. Rodriguez-Wallberg, and C. Lampic. 2014. Desire for children, difficulties achieving a pregnancy, and infertility distress 3 to 7 years after cancer diagnosis. *Support Care Cancer* 22 (10):2805-2812. doi:10.1007/s00520-014-2279-z.
3. Armuand, G.M., K.A. Rodriguez-Wallberg, L. Wettergren, J. Ahlgren, G. Enblad, M. Hoglund, and C. Lampic. 2012. Sex differences in fertility-related information received by young adult cancer survivors. *J. Clin. Oncol* 30 (17):2147-2153. doi:JCO.2011.40.6470 [pii];10.1200/JCO.2011.40.6470 [doi].
4. Balthazar, U., A. M. Deal, M. A. Fritz, L. A. Kondapalli, J. Y. Kim, and J. E. Mersereau. 2012. The current fertility preservation consultation model: are we adequately informing cancer patients of their options? *Hum Reprod* 27 (8):2413-2419. doi:10.1093/humrep/des188.
5. Bastings, L., O. Baysal, C.C. Beerendonk, D.D. Braat, and W.L. Nelen. 2014. Referral for fertility preservation counselling in female cancer patients. *Hum. Reprod* 29 (10):2228-2237. doi:deu186 [pii];10.1093/humrep/deu186 [doi].
6. Blumenfeld, Zeev. 2012. Chemotherapy and fertility. *Best Practice & Research Clinical Obstetrics & Gynaecology* 26 (3):379-390. doi:https://doi.org/10.1016/j.bpobgyn.2011.11.008.
7. Brydoy, M., S. D. Fossa, O. Dahl, and T. Bjoro. 2007. Gonadal dysfunction and fertility problems in cancer survivors. *Acta Oncol* 46 (4):480-489. doi:10.1080/02841860601166958.
8. Buske, D., A. Sender, D. Richter, E. Brahler, and K. Geue. 2016. Patient-Physician Communication and Knowledge Regarding Fertility Issues from German Oncologists' Perspective-a Quantitative Survey. *J Cancer Educ* 31 (1):115-122. doi:10.1007/s13187-015-0841-0.
9. Carter, J., L. Raviv, L. Applegarth, J.S. Ford, L. Josephs, E. Grill, C. Sklar, Y. Sonoda, R.E. Baser, and R.R. Barakat. 2010. A cross-sectional study of the psychosexual impact of cancer-related infertility in women: third-party reproductive assistance. *J. Cancer Surviv* 4 (3):236-246. doi:10.1007/s11764-010-0121-2 [doi].
10. Cook, Julia V., Heather O. Dickinson, and Martin P. Eccles. 2009. Response rates in postal surveys of healthcare professionals between 1996 and 2005: An observational study. *BMC Health Services Research* 9 (1):160. doi:10.1186/1472-6963-9-160.
11. Demark-Wahnefried, W., N. M. Aziz, J. H. Rowland, and B. M. Pinto. 2005. Riding the crest of the teachable moment: promoting long-term health after the diagnosis of cancer. *J Clin Oncol* 23 (24):5814-5830. doi:10.1200/jco.2005.01.230.
12. Dikmans, R. E., E. M. Krouwel, M. Ghasemi, T. C. van de Grift, M. B. Bouman, Mjpf Ritt, H. W. Elzevier, and M. G. Mullender. 2018. Discussing sexuality in the field of plastic and reconstructive surgery: a national survey of current practice in the Netherlands. *Eur J Plast Surg* 41 (6):707-714. doi:10.1007/s00238-018-1452-x.
13. Duffy, C., S.M. Allen, C. Dube, and K. Dickersin. 2012. Oncologists' confidence in knowledge of fertility issues for young women with cancer. *J. Cancer Educ* 27 (2):369-376. doi:10.1007/s13187-011-0304-1 [doi].
14. Forman, E. J., C. K. Anders, and M. A. Behera. 2010. A nationwide survey of oncologists regarding treatment-related infertility and fertility preservation in female cancer patients. *Fertil Steril* 94 (5):1652-1656. doi:10.1016/j.fertnstert.2009.10.008.
15. Gilbert, E., A. Adams, H. Mehanna, B. Harrison, and G.M. Hartshorne. 2011. Who should be offered sperm banking for fertility preservation? A survey of UK oncologists and haematologists. *Ann. Oncol* 22 (5):1209-1214. doi:mdq579 [pii];10.1093/annonc/mdq579 [doi].

16. Green, D. M., T. Kawashima, M. Stovall, W. Leisenring, C. A. Sklar, A. C. Mertens, S. S. Donaldson, J. Byrne, and L. L. Robison. 2010. Fertility of male survivors of childhood cancer: a report from the Childhood Cancer Survivor Study. *J Clin Oncol* 28 (2):332-339. doi:10.1200/jco.2009.24.9037.
17. Hariton, Eduardo, Pietro Bortoletto, Eden R. Cardozo, Ephraim P. Hochberg, and Mary E. Sabatini. 2016. The Role of Oncofertility Clinics in Facilitating Access to Reproductive Specialists. *Journal of patient experience* 3 (4):131-136. doi:10.1177/2374373516685960.
18. IKR. 2010. Cryopreservatie van zaadcellen bij oncologiepatiënten; version 2.0.
19. Krouwel, E. M., L. F. Albers, M. P. J. Nicolai, H. Putter, S. Osanto, R. C. M. Pelger, and H. W. Elzevier. 2019. Discussing Sexual Health in the Medical Oncologist's Practice: Exploring Current Practice and Challenges. *J Cancer Educ*. doi:10.1007/s13187-019-01559-6.
20. Krouwel, E. M., M. P. Nicolai, G. J. van der Wielen, H. Putter, A. D. Krol, R. C. Pelger, L. Incrocci, and H. W. Elzevier. 2015. Sexual Concerns after (Pelvic) Radiotherapy: Is There Any Role for the Radiation Oncologist? *J Sex Med* 12 (9):1927-1939. doi:10.1111/jsm.12969.
21. Krouwel, E.M., Hagen, J.H., Nicolai, M.P.J., Vahrmeijer, A.L., Putter, H., Pelger, R.C.M., Elzevier, H.W., and 2015. Management of Sexual Side Effects in the Surgical Oncology Practice: A Nationwide Survey of Dutch Surgical Oncologists. In *Eur J Surg Onc*.
22. Logan, S., J. Perz, J. M. Ussher, M. Peate, and A. Anazodo. 2019. Systematic review of fertility-related psychological distress in cancer patients: Informing on an improved model of care. *Psychooncology* 28 (1):22-30. doi:10.1002/pon.4927.
23. Logan, S., J. Perz, J. Ussher, M. Peate, and A. Anazodo. 2017. Clinician provision of oncofertility support in cancer patients of a reproductive age: A systematic review. *Psychooncology*. doi:10.1002/pon.4518.
24. Loren, A. W., P. B. Mangu, L. N. Beck, L. Brennan, A. J. Magdalinski, A. H. Partridge, G. Quinn, W. H. Wallace, and K. Oktay. 2013. Fertility preservation for patients with cancer: American Society of Clinical Oncology clinical practice guideline update. *J Clin Oncol* 31 (19):2500-2510. doi:10.1200/jco.2013.49.2678.
25. Louwe, L. A., A. M. Stiggelbout, A. Overbeek, Cgjm Hilders, M. H. van den Berg, E. Wendel, E. van Dulmen-den Broeder, and M. M. Ter Kuile. 2018. Factors associated with frequency of discussion of or referral for counselling about fertility issues in female cancer patients. *Eur J Cancer Care (Engl)* 27 (1). doi:10.1111/ecc.12602.
26. Louwe, L. A., M. M. ter Kuile, C. G. Hilders, E. Jenninga, S. M. Tiemessen, A. A. Peters, J. W. Nortier, and A. M. Stiggelbout. 2013. Oncologists' practice and attitudes regarding fertility preservation in female cancer patients: a pilot study in the Netherlands. *J Psychosom Obstet Gynaecol* 34 (3):129-132. doi:10.3109/0167482x.2013.821977.
27. Meirow, D., and D. Nugent. 2001. The effects of radiotherapy and chemotherapy on female reproduction. *Hum Reprod Update* 7 (6):535-543.
28. Micaux Obol, C., G. M. Armuand, K. A. Rodriguez-Wallberg, J. Ahlgren, P. Ljungman, L. Wettergren, and C. Lampic. 2017. Oncologists and hematologists' perceptions of fertility-related communication - a nationwide survey. *Acta Oncol* 56 (8):1103-1110. doi:10.1080/0284186x.2017.1310394.
29. NVOG. 2016. Fertilitieitsbehoud bij vrouwen met kanker, version 1.0. *National guideline IKNL*.
30. Partridge, A.H., S. Gelber, J. Peppercorn, E. Sampson, K. Knudsen, M. Laufer, R. Rosenberg, M. Przystypny, A. Rein, and E.P. Winer. 2004. Web-based survey of fertility issues in young women with breast cancer. *J. Clin. Oncol* 22 (20):4174-4183. doi:22/20/4174 [pii];10.1200/JCO.2004.01.159 [doi].
31. Peccatori, F. A., H. A. Azim, Jr., R. Orecchia, H. J. Hoekstra, N. Pavlidis, V. Kesic, and G. Pentheroudakis. 2013. Cancer, pregnancy and fertility: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. *Ann Oncol* 24 Suppl 6:vi160-170. doi:10.1093/annonc/mdt199.

32. Reinmuth, S., C. Hohmann, R. Rendtorff, M. Balcerek, S. Holzhausen, A. Muller, G. Henze, T. Keil, and A. Borgmann-Staudt. 2013. Impact of chemotherapy and radiotherapy in childhood on fertility in adulthood: the FeCt-survey of childhood cancer survivors in Germany. *J Cancer Res Clin Oncol* 139 (12):2071-2078. doi:10.1007/s00432-013-1527-9.
33. Rosendahl, Mikkel, Claus Yding Andersen, Nina la Cour Freiesleben, Anders Juul, Kristine Løssl, and Anders Nyboe Andersen. 2010. Dynamics and mechanisms of chemotherapy-induced ovarian follicular depletion in women of fertile age. *Fertil Steril* 94 (1):156-166. doi:https://doi.org/10.1016/j.fertnstert.2009.02.043.
34. Ruddy, Kathryn J., Shari I. Gelber, Rulla M. Tamimi, Elizabeth S. Ginsburg, Lidia Schapira, Steven E. Come, Virginia F. Borges, Meghan E. Meyer, and Ann H. Partridge. 2014. Prospective Study of Fertility Concerns and Preservation Strategies in Young Women With Breast Cancer. *Journal of Clinical Oncology* 32 (11):1151-1156. doi:10.1200/JCO.2013.52.8877.
35. Salama, M., and T. K. Woodruff. 2017. Anticancer treatments and female fertility: clinical concerns and role of oncologists in oncofertility practice. *Expert Rev Anticancer Ther* 17 (8):687-692. doi:10.1080/14737140.2017.1335199.
36. Sallem, A., J. Shore, I. Ray-Coquard, L. Ferreux, M. Bourdon, C. Maignien, C. Patrat, J. P. Wolf, and K. Pocate-Cheriet. 2018. Fertility preservation in women with cancer: a national study about French oncologists awareness, experience, and feelings. *J Assist Reprod Genet* 35 (10):1843-1850. doi:10.1007/s10815-018-1251-0.
37. Schover, L.R., L.A. Rybicki, B.A. Martin, and K.A. Bringelsen. 1999. Having children after cancer. A pilot survey of survivors' attitudes and experiences. *Cancer* 86 (4):697-709. doi:10.1002/(SICI)1097-0142(19990815)86:4<697::AID-CNCR20>3.0.CO;2-J [pii].
38. Vassilakopoulou, M., E. Boostandoost, G. Papaxoinis, T. de La Morthe Rouge, D. Khayat, and A. Psyrri. 2016. Anticancer treatment and fertility: Effect of therapeutic modalities on reproductive system and functions. *Crit Rev Oncol Hematol* 97:328-334. doi:10.1016/j.critrevonc.2015.08.002.
39. Yee, S., E. Fuller-Thomson, A. Lau, and E. M. Greenblatt. 2012. Fertility preservation practices among Ontario oncologists. *J Cancer Educ* 27 (2):362-368. doi:10.1007/s13187-011-0301-4.

