



Universiteit  
Leiden  
The Netherlands

## Measuring shared decision making in oncology: an informed approach

Bomhof-Roordink, H.

### Citation

Bomhof-Roordink, H. (2022, June 7). *Measuring shared decision making in oncology: an informed approach*. Retrieved from <https://hdl.handle.net/1887/3307663>


Version: Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/3307663>

**Note:** To cite this publication please use the final published version (if applicable).





---

GENERAL INTRODUCTION

Shared decision making (SDM) between patient and healthcare professional about treatment options is becoming 'the new normal' in the Netherlands, envisioned Bruno Bruins, the former Dutch minister of Health in 2019.<sup>1</sup> The Dutch Federation of Medical Specialists considers that SDM should become a habit<sup>2</sup> and, consequently, it may become the new normal. 'The new normal' has gained a completely different meaning in the Netherlands since 2020. It refers to the behaviours asked from each individual to slow the spread of COVID-19.<sup>3</sup> For example, here and elsewhere people have been asked to avoid physical contact and to wash their hands frequently.<sup>4, 5</sup> The recommended behaviours have repeatedly been communicated by the government and are quite easy in themselves, but still adherence has been low. Evidently, commitment to new behaviours is not easy, even for simple behaviours. In contrast, SDM between patient and healthcare professional involves two or more individuals who need to commit to complex behaviours during and outside the clinical consultation. Communication about these behaviour changes by healthcare professional organisations, among others is challenging, and they may not easily become routine for patients and healthcare professionals.

In 1972, Veatch described the contractual model in which there is true sharing of ethical authority and responsibility between patient and physician, next to sharing of decision making. Ten years later, ethically valid informed consent was stated to involve a process of SDM.<sup>6</sup> In the 1990's several journals published papers on SDM,<sup>7-9</sup> and Charles and colleagues presented the first SDM model in 1997 (see Box 1).<sup>10</sup> The BMJ embraced patient partnership with a contribution by Charles and colleagues<sup>11</sup> and by illustrating it as a tangoing couple on their cover in 1999, upon Charles et al. revisiting their SDM model.<sup>12</sup> In 2006, Makoul & Clayman identified 31 separate concepts used to explicate SDM, from 161 different definitions.<sup>13</sup> A year later, Moumjid et al. concluded that while clear SDM definitions were available, they were poorly cited and that the term SDM was being used inconsistently.<sup>14</sup> Over the following years, the number of publications on SDM increased rapidly.<sup>15</sup>

In 2011 the Salzburg statement called upon patients and healthcare professionals 'to work together to be coproducers of health', with specific tasks for each of them.<sup>16</sup> To date, a range of implementation activities have been undertaken to support SDM, such as: training of healthcare professionals,<sup>17, 18</sup> development of pocket cards for healthcare professionals,<sup>19</sup> and development of patient decision aids.<sup>20, 21</sup> Dutch national campaigns have been launched ('3 goede vragen',<sup>22, 23</sup> 'consultkaart',<sup>24</sup> 'begin een goed gesprek'<sup>25</sup>) to create awareness about SDM, informed by e.g., the AskShareKnow,<sup>26, 27</sup> the Ask 3 questions campaigns,<sup>28</sup> and Option Grids.<sup>29</sup> SDM has even been established by Dutch law; the Dutch Medical Treatment Agreement Act (Wet op de geneeskundige behandelovereenkomst (WGBO)) which regulates the rights and obligations of patients, was adapted recently (January 1, 2020) and now includes reference to SDM.<sup>30</sup>

**Box 1.** First SDM model by Charles et al.<sup>10, 12</sup>

1. At a minimum, both the physician and patient are involved in the treatment decision-making process.
2. Both the physician and patient share information with each other.
3. Both the physician and the patient take steps to participate in the decision-making process by expressing treatment preferences.
4. A treatment decision is made, and both the physician and patient agree on the treatment to implement.

**SDM measurement challenges**

While many SDM implementation activities have been launched, measurement difficulties remain.<sup>31-33</sup> In 2011, Scholl et al. identified 28 SDM measurement instruments from the literature and concluded that further psychometric testing was needed, since validity had often not been sufficiently investigated.<sup>34</sup> Moreover, these and more recent measurement instruments only assess healthcare professionals behaviour, or include patient and healthcare professional behaviour in one item. This makes it impossible to assess the patients' role, while their responsibilities have been clearly emphasized since the first SDM models.<sup>10, 12</sup> Measurement of behaviours outside consultations is also lacking, while SDM extends to the world outside the consultation room.<sup>35</sup>

In previous research, patients and healthcare professionals have been involved in the development of SDM measurement instruments to a limited extent only, even though this is recommended.<sup>36</sup> This lack of involvement may partly explain poor correlations between SDM assessments from different viewpoints,<sup>33, 37-40</sup> including an independent observer (e.g., OPTION-5<sup>41</sup>), the patient (e.g., SDM-Q-9,<sup>42</sup>) or the healthcare professional (e.g., SDM-Q-Doc<sup>43</sup>). Patient and healthcare professional involvement will likely improve the content validity of the measurement instruments and for questionnaires, their feasibility and acceptability.

Last but not least, for most existing measurement instruments, the developers apparently have assumed a reflective model, as they assessed factor structure and/or internal consistency. They have thereby neglected the formative nature of the SDM construct. That is, SDM in itself may not be something already present, in contrast to e.g., intelligence.<sup>44, 45</sup> SDM is formed by the behaviours of patients and healthcare professionals, both during and outside consultations. What these behaviours entail, may vary per context. Together the items of a measurement instrument form the construct, while for e.g., intelligence, the items reflect the construct. A consequence of assuming a formative measurement model is that another approach is needed to inform item selection and to determine the validity of a measurement instrument.

**Aim and outline**

We aimed to develop and validate questionnaires to assess the SDM process in oncology from both the patient and the physician viewpoint. We chose the participant perspective and decided to develop questionnaires instead of a coding scheme to be completed by an independent observer, since questionnaires are far more easy to use in research. To guide our development and validation process, we used the original COnsensus-based

Standards for the selection of health Measurement INstruments (COSMIN) checklist<sup>46, 47</sup> and wrote two reviews: one on published SDM measurement instruments and one on published SDM models. Next, we used several consecutive studies to develop, test, and validate the questionnaires. We chose to develop the questionnaires specifically for oncology, since cancer patients often face preference-sensitive decisions,<sup>48</sup> a decision type for which SDM is considered to be the appropriate approach.<sup>49</sup> Cancer patients' treatment preferences vary<sup>50, 51</sup> and often differ from physicians' treatment preferences.<sup>52, 53</sup> Survival, for example, may be weighed differently by patients and physicians.<sup>52</sup> To ensure that treatment is in line with individual patients' preferences, cancer patients' involvement in decision making is of utmost importance. Fortunately, most cancer patients prefer an active or collaborative role in treatment decision making.<sup>54-56</sup>

In chapter 2, we present an overview of existing SDM measurement instruments and an assessment of the level of evidence for 10 measurement properties. This assessment was informed by the methodological quality of the respective validation study or studies, and by the psychometric quality of the measurement properties. In chapter 3, we present an overview of models defining SDM between a patient and a healthcare professional, the components making up the models, who is seen as responsible for the occurrence of the SDM components, the inclusion of the components over time, and we present a frequency map of SDM components per healthcare setting. In chapter 4, views of stakeholders are integrated into a model of SDM in oncology. Chapter 5 describes the development and first testing of the iSHAREpatient and iSHAREphysician. These questionnaires aim to measure SDM in oncology, from the viewpoint of the patient and of the physician, respectively. In chapter 6 we demonstrate construct validity of the iSHAREpatient and iSHAREphysician, test-retest agreement of the iSHAREpatient, and agreement between scores on the iSHAREpatient and iSHAREphysician. In chapter 7 the main findings are summarized and discussed, including strengths and limitations, practice implications, suggestions for future research and concluding remarks.

## REFERENCES

1. Stijnman S. Interview: Minister Bruno Bruins. Samen beslissen wordt het nieuwe normaal. *Wisselstof* September 2019. Available from: <https://www.wisselstof.nl/magazine/60/nieuw-magazine/176/samen-beslissen-wordt-het-nieuwe-normaa>. Date last accessed: 07-09-2021.
2. Federatie Medisch Specialisten. Visiedocument Samen Beslissen. Mei 2019. Available from: [https://www.demedischspecialist.nl/sites/default/files/FMS\\_Visiedoc-SamenBeslissen%282019%29\\_v03.pdf](https://www.demedischspecialist.nl/sites/default/files/FMS_Visiedoc-SamenBeslissen%282019%29_v03.pdf). Date last accessed: 07-09-2021.
3. Algemeen Nederlands Persbureau. RIVM: het nieuwe normaal moeten we normaal gaan vinden. *Nederlands Dagblad*. Available from: <https://www.nd.nl/nieuws/varia/972447/rivm-het-nieuwe-normaal-moeten-we-normaal-gaan-vinden>. Date last accessed: 07-09-2021.
4. Backer JA, Mollema L, Vos ER, Klinkenberg D, van der Klis FR, de Melker HE, et al. Impact of physical distancing measures against COVID-19 on contacts and mixing patterns: repeated cross-sectional surveys, the Netherlands, 2016-17, April 2020 and June 2020. *Euro Surveill*. 2021;26.
5. Jalloh MF, Nur AA, Nur SA, Winters M, Bedson J, Pedi D, et al. Behaviour adoption approaches during public health emergencies: implications for the COVID-19 pandemic and beyond. *BMJ Glob Health*. 2021;6.
6. President's Commission for the Study of Ethical Problems in Medicine and Biomedical and Behavioral Research (1982). *Making Health Care Decisions. The ethical and legal implications of informed consent in the patient-practitioner relationship*. Washington, D.C. <https://pubmed.ncbi.nlm.nih.gov/12041401/>.
7. Brock DW. The ideal of shared decision making between physicians and patients. *Kennedy Inst Ethics J*. 1991;1:28-47.
8. Mitchell P. Shared decision-making poses problems for UK. *Lancet*. 1997;349:1306.
9. Emanuel EJ, Emanuel LL. Four models of the physician-patient relationship. *JAMA*. 1992;267:2221-6.
10. Charles C, Gafni A, Whelan T. Shared decision-making in the medical encounter: what does it mean? (or it takes at least two to tango). *Soc Sci Med*. 1997;44:681-92.
11. Charles C, Whelan T, Gafni A. What do we mean by partnership in making decisions about treatment? *BMJ*. 1999;319:780-2.
12. Charles C, Gafni A, Whelan T. Decision-making in the physician-patient encounter: revisiting the shared treatment decision-making model. *Soc Sci Med*. 1999;49:651-61.
13. Makoul G, Clayman ML. An integrative model of shared decision making in medical encounters. *Patient Educ Couns*. 2006;60:301-12.
14. Moumjid N, Gafni A, Bremond A, Carrere MO. Shared decision making in the medical encounter: are we all talking about the same thing? *Med Decis Making*. 2007;27:539-46.
15. Lu C, Li X, Yang K. Trends in Shared Decision-Making Studies From 2009 to 2018: A Bibliometric Analysis. *Front Public Health*. 2019;7:384.
16. Salzburg Global Seminar, Salzburg statement on shared decision making. *BMJ*. 2011;342:d1745.
17. Legare F, Adekpedjou R, Stacey D, Turcotte S, Kryworuchko J, Graham ID, et al. Interventions for increasing the use of shared decision making by healthcare professionals. *Cochrane Database Syst Rev*. 2018;7:CD006732.
18. Henselmans I, van Laarhoven HWM, de Haes H, Tokat M, Engelhardt EG, van Maarschalkerweerd PEA, et al. Training for Medical Oncologists on Shared Decision-Making About Palliative Chemotherapy: A Randomized Controlled Trial. *Oncologist*. 2019;24:259-65.
19. Nederlandse Federatie van Kankerpatiënten organisaties. *Tools bij Samen Beslissen*. Available from: <https://nfk.nl/themas/samen-beslissen/tools-bij-samen-beslissen>. Date last accessed: 07-09-2021.
20. Stacey D, Legare F, Lewis K, Barry MJ, Bennett CL, Eden KB, et al. Decision aids for people facing health treatment or screening decisions. *Cochrane Database Syst Rev*. 2017;4:CD001431.

21. van den Berg M, van der Meij E, Bos AME, Boshuizen MCS, Determann D, van Eekeren R, et al. Development and testing of a tailored online fertility preservation decision aid for female cancer patients. *Cancer Med.* 2021;10:1576-88.
22. 3 goede vragen. Available from: <https://3goedevragen.nl/>. Date last accessed: 07-09-2021.
23. Garvelink MM, Jillissen M, Knops A, Kremer JAM, Hermens R, Meinders MJ. Implementation of the three good questions-A feasibility study in Dutch hospital departments. *Health Expect.* 2019;22:1272-84.
24. Samen beslissen met behulp van Consultkaarten. Available from: <https://consultaart.nl/>. Date last accessed: 07-09-2021.
25. Begin een goed gesprek. Available from: <https://beginengoesprek.nl/>. Date last accessed: 07-09-2021.
26. Shepherd HL, Barratt A, Trevena LJ, McGeechan K, Carey K, Epstein RM, et al. Three questions that patients can ask to improve the quality of information physicians give about treatment options: a cross-over trial. *Patient Educ Couns.* 2011;84:379-85.
27. Shepherd HL, Barratt A, Jones A, Bateson D, Carey K, Trevena LJ, et al. Can consumers learn to ask three questions to improve shared decision making? A feasibility study of the ASK (AskShareKnow) Patient-Clinician Communication Model((R)) intervention in a primary health-care setting. *Health Expect.* 2016;19:1160-8.
28. Lloyd A, Joseph-Williams N, Edwards A, Rix A, Elwyn G. Patchy 'coherence': using normalization process theory to evaluate a multi-faceted shared decision making implementation program (MAGIC). *Implement Sci.* 2013;8:102.
29. Seal RP, Kynaston J, Elwyn G, Smith PE. Using an Option Grid in shared decision making. *Pract Neurol.* 2014;14:54-6.
30. Ubbink DT GP, Gosens T, Brand PLP. Meer 'samen beslissen' nodig door aangescherpte Wgbo. *Ned Tijdschr Geneeskd* 2021;165:D5775.
31. Gerwing J, Gulbrandsen P. Contextualizing decisions: Stepping out of the SDM track. *Patient Educ Couns.* 2019;102:815-6.
32. Blumenthal-Barby J, Opel DJ, Dickert NW, Kramer DB, Tucker Edmonds B, Ladin K, et al. Potential Unintended Consequences Of Recent Shared Decision Making Policy Initiatives. *Health Aff (Millwood).* 2019;38:1876-81.
33. Scholl I, Kriston L, Dirmaier J, Harter M. Comparing the nine-item Shared Decision-Making Questionnaire to the OPTION Scale - an attempt to establish convergent validity. *Health Expect.* 2015;18:137-50.
34. Scholl I, Koelewijn-van Loon M, Sepucha K, Elwyn G, Legare F, Harter M, et al. Measurement of shared decision making - a review of instruments. *Z Evid Fortbild Qual Gesundheitswes.* 2011;105:313-24.
35. Clayman ML, Gulbrandsen P, Morris MA. A patient in the clinic; a person in the world. Why shared decision making needs to center on the person rather than the medical encounter. *Patient Educ Couns.* 2017;100:600-4.
36. de Vet HCW, Terwee CB, Mokkink LB, Knol DL. *Measurement in Medicine.* Cambridge University Press. Cambridge. 2011.
37. Kriston L, Harter M, Scholl I. A latent variable framework for modeling dyadic measures in research on shared decision-making. *Z Evid Fortbild Qual Gesundheitswes.* 2012;106:253-63.
38. Geessink NH, Ofstad EH, Olde Rikkert MGM, van Goor H, Kasper J, Schoon Y. Shared decision-making in older patients with colorectal or pancreatic cancer: Determinants of patients' and observers' perceptions. *Patient Educ Couns.* 2018;101:1767-74.
39. Calderon C, Jimenez-Fonseca P, Ferrando PJ, Jara C, Lorenzo-Seva U, Beato C, et al. Psychometric properties of the Shared Decision-Making Questionnaire (SDM-Q-9) in oncology practice. *Int J Clin Health Psychol.* 2018;18:143-51.
40. Kasper J, Heesen C, Kopke S, Fulcher G, Geiger F. Patients' and observers' perceptions of involvement differ. Validation study on inter-relating measures for shared decision making. *PLoS One.* 2011;6:e26255.



41. Elwyn G, Tsulukidze M, Edwards A, Legare F, Newcombe R. Using a 'talk' model of shared decision making to propose an observation-based measure: Observer OPTION 5 Item. *Patient Educ Couns.* 2013;93:265-71.
42. Kriston L, Scholl I, Holzel L, Simon D, Loh A, Harter M. The 9-item Shared Decision Making Questionnaire (SDM-Q-9). Development and psychometric properties in a primary care sample. *Patient Educ Couns.* 2010;80:94-9.
43. Scholl I, Kriston L, Dirmaier J, Buchholz A, Harter M. Development and psychometric properties of the Shared Decision Making Questionnaire--physician version (SDM-Q-Doc). *Patient Educ Couns.* 2012;88:284-90.
44. Wollschlager D. Short communication: Where is SDM at home? putting theoretical constraints on the way shared decision making is measured. *Z Evid Fortbild Qual Gesundheitswes.* 2012;106:272-4.
45. Dowie J. Shared decision making is a Preference-sensitive Formative Construct: the Implications. *Eur J Pers Cent Healthc.* 2019;7:506-17.
46. Mokkink LB, Terwee CB, Patrick DL, Alonso J, Stratford PW, Knol DL, et al. The COSMIN checklist for assessing the methodological quality of studies on measurement properties of health status measurement instruments: an international Delphi study. *Qual Life Res.* 2010;19:539-49.
47. Terwee CB, Mokkink LB, Knol DL, Ostelo RWJG, Bouter LM, de Vet HCW. Rating the methodological quality in systematic reviews of studies on measurement properties: a scoring system for the COSMIN checklist. *Qual Life Res.* 2012;21:651-7.
48. Politi MC, Studts JL, Hayslip JW. Shared decision making in oncology practice: what do oncologists need to know? *Oncologist.* 2012;17:91-100.
49. Stiggelbout AM, Pieterse AH, De Haes JC. Shared decision making: Concepts, evidence, and practice. *Patient Educ Couns.* 2015;98:1172-9.
50. Schmidt K, Damm K, Vogel A, Golpon H, Manns MP, Welte T, et al. Therapy preferences of patients with lung and colon cancer: a discrete choice experiment. *Patient Prefer Adherence.* 2017;11:1647-56.
51. Post CCB, Mens JWM, Haverkort MAD, Koppe F, Jurgenliemk-Schulz IM, Snyers A, et al. Patients' and clinicians' preferences in adjuvant treatment for high-risk endometrial cancer: Implications for shared decision making. *Gynecol Oncol.* 2021.
52. Davidson PM, Jiwa M, Goldsmith AJ, McGrath SJ, Digiacomio M, Phillips JL, et al. Decisions for lung cancer chemotherapy: the influence of physician and patient factors. *Support Care Cancer.* 2011;19:1261-6.
53. Pieterse AH, Stiggelbout AM, Baas-Thijssen MC, van de Velde CJ, Marijnen CA. Benefit from preoperative radiotherapy in rectal cancer treatment: disease-free patients' and oncologists' preferences. *Br J Cancer.* 2007;97:717-24.
54. Moth E, McLachlan SA, Veillard AS, Muljadi N, Hudson M, Stockler MR, et al. Patients' preferred and perceived roles in making decisions about adjuvant chemotherapy for non-small-cell lung cancer. *Lung Cancer.* 2016;95:8-14.
55. Hamelinck VC, Bastiaannet E, Pieterse AH, van de Velde CJH, Liefers GJ, Stiggelbout AM. Preferred and Perceived Participation of Younger and Older Patients in Decision Making About Treatment for Early Breast Cancer: A Prospective Study. *Clin Breast Cancer.* 2018;18:e245-e53.
56. Roydhouse JK, Gutman R, Wilson IB, Kehl KL, Keating NL. Patient and proxy reports regarding the experience of treatment decision-making in cancer care. *Psychooncology.* 2020;29:1943-50.