



Universiteit
Leiden
The Netherlands

Reduced psychosocial functioning in subacromial pain syndrome is associated with persistence of complaints after 4 years

Overbeek, C.L.; Gademan, M.G.J.; Kolk, A.; Visser, C.P.J.; Zwaal, P. van der; Nagels, J.; Nelissen, R.G.H.H.

Citation

Overbeek, C. L., Gademan, M. G. J., Kolk, A., Visser, C. P. J., Zwaal, P. van der, Nagels, J., & Nelissen, R. G. H. H. (2021). Reduced psychosocial functioning in subacromial pain syndrome is associated with persistence of complaints after 4 years. *Journal Of Shoulder And Elbow Surgery*, 30(2), 223-228. doi:10.1016/j.jse.2020.08.039

Version: Publisher's Version

License: [Creative Commons CC BY 4.0 license](https://creativecommons.org/licenses/by/4.0/)

Downloaded from: <https://hdl.handle.net/1887/3196062>

Note: To cite this publication please use the final published version (if applicable).



ELSEVIER



SHOULDER

Reduced psychosocial functioning in subacromial pain syndrome is associated with persistence of complaints after 4 years

Celeste L. Overbeek, MD^{a,b,*}, Maaïke G.J. Gademan, MSc^{a,c}, Arjen Kolk, MD^{a,b}, Cornelis P.J. Visser, MD, PhD^d, Peer van der Zwaal, MD, PhD^e, Jochem Nagels, MD^{a,b}, Rob G.H.H. Nelissen, MD, PhD^{a,b}

^aDepartment of Orthopaedics, Leiden University Medical Center, Leiden, The Netherlands

^bLaboratory for Kinematics and Neuromechanics, Department of Orthopaedics and Rehabilitation, Leiden University Medical Center, Leiden, The Netherlands

^cDepartment of Clinical Epidemiology, Leiden University Medical Center, Leiden, The Netherlands

^dDepartment of Orthopaedics, Alrijne Hospital, Leiden, The Netherlands

^eDepartment of Orthopaedics, Haaglanden Medical Center, The Hague, The Netherlands

Background: Patients with subacromial pain syndrome (SAPS) frequently present with coexisting psychosocial problems; however, whether this also associates with long-term outcome is currently unknown. We assessed whether psychosocial functioning in patients with SAPS is associated with persistence of complaints after 4 years of routine care.

Methods: In a longitudinal study, 34 patients with SAPS were selected after clinical and radiologic evaluation and assessed at baseline and after 4 years. For the assessment of psychosocial functioning, the RAND-36 questionnaire domains of social functioning, role limitations due to emotional problems, mental health, vitality, and general health were evaluated. Complaint persistence at follow-up was assessed by (1) an anchor question (reduced, persistent, or increased symptoms), (2) change in pain (change in visual analog scale score), and (3) change in quality of life (change in Western Ontario Rotator Cuff index score).

Results: Lower baseline mental health (odds ratio [OR], 0.92; 95% confidence interval [CI], 0.85-0.98; $P = .013$), vitality (OR, 0.90; 95% CI, 0.83-0.98; $P = .011$), and general health (OR, 0.93; 95% CI, 0.88-0.98; $P = .009$) were associated with persistent complaints as reported by the anchor question, change in visual analog scale score, and change in Western Ontario Rotator Cuff index score.

Conclusions: Evaluating psychosocial functioning parallel to physical complaints is currently not standard procedure in the treatment of SAPS. In this study, we showed that factors related to psychosocial functioning are associated with long-term persistence of complaints in SAPS. Future studies may investigate whether a multimodal treatment with assessment of psychosocial functioning may facilitate pain relief and recovery in SAPS.

Level of evidence: Level I; Prospective Design; Prognosis Study

© 2020 The Author(s). This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Keywords: Subacromial impingement; subacromial pain syndrome; psychosocial health; RAND-36; longitudinal; depression; prognostic; multimodal

This study was approved by the Medical Ethical Committee of the Leiden University Medical Center (no. P09.227). All participants provided written informed consent.

*Reprint requests: Celeste L. Overbeek, MD, Laboratory for Kinematics and Neuromechanics, Department of Orthopaedics and

Rehabilitation, Leiden University Medical Center, Room B0-57, Albinusdreef 2, 2333 ZA Leiden, Postbus 9600, 2300, RC Leiden, The Netherlands.

E-mail address: c.l.overbeek@lumc.nl (C.L. Overbeek).

Shoulder pain affects 70% of the general population at some point in life.²⁶ Most frequently, shoulder pain is attributed to irritation of subacromial tissues and is accordingly called subacromial pain (or impingement) syndrome.³⁷ Subacromial pain syndrome (SAPS) is associated with significant individual and socioeconomic consequences as it compromises the ability to perform daily activities owing to related pain and disability.^{5,12,23,26,35,37} Current treatments focus primarily on symptom relief of the shoulder. Such treatments include physical therapy to improve neuromuscular control and subacromial clearance, as well as the use of subacromial corticosteroid injections to reduce subacromial inflammation. Failure to respond to these usually reliable treatments suggests a nonorganic chronic condition in up to 40% of all patients with SAPS.^{11,26,32,37}

In various musculoskeletal pain disorders, a close relation between pain and psychosocial functioning (eg, depression, anxiety, or social support) is documented.^{18,25} Impaired psychosocial functioning is frequently viewed to be a result of pain; however, it may also enhance the perception of pain.¹³ In the knee and hip, it has repetitively been observed that patients who have depressive feelings tend to respond poorly to interventions targeting the painful joint.^{3,34} Furthermore, treating concurrent depression in patients with chronic low-back pain has been shown to result in pain relief in nearly 25% of patients.³³ These studies show the importance of psychosocial functioning in the treatment of musculoskeletal pain.²¹

Regarding research on SAPS, there has been a focus on peripheral pathology of the shoulder whereas psychosocial factors have received less attention.^{4,7,19,22,24,27,28} It has been established that patients with SAPS frequently present with coexisting psychosocial problems; however, whether this also associates with long-term outcome is unknown.^{7,19,28} Therefore, we assessed whether psychosocial functioning in patients with SAPS is associated with persistence of complaints after 4 years of routine care.

Materials and methods

For this longitudinal prognostic analysis, eligible patients were recruited between April 2010 and December 2012 at the Leiden University Medical Center, Haaglanden Medical Center, and Alrijne Hospital, under a previously registered and published study protocol (trial register no. NTR2283).⁸ Consecutive patients with SAPS were selected through physical examination, shoulder radiographs, and magnetic resonance arthrography by dedicated shoulder surgeons. The inclusion criteria were patients who were aged 35-60 years, who had unilateral shoulder complaints for >3 months, and who received a clinical diagnosis of SAPS based on a positive Hawkins test and Neer impingement test with lidocaine.⁸ The exclusion criteria were insufficient language skills, inflammatory glenohumeral (GH) arthritis, clinical signs of GH or acromioclavicular osteoarthritis, previous shoulder surgery, fracture or dislocation, cervical radiculopathy, GH instability,

decreased passive GH mobility (eg, frozen shoulder), and presence of electronic implants (eg, pacemaker). Additionally, patients were excluded in case other specific conditions were diagnosed on radiographs or magnetic resonance arthrography such as calcific tendinitis, full-thickness rotator cuff tear, and labral or ligament pathology.⁸ Patients who provided written informed consent were included and contacted for a follow-up visit between June 2014 and September 2015 (ie, 3-4 years later).

Psychosocial functioning

At baseline and follow-up, psychosocial functioning was assessed by means of the RAND-36 questionnaire.^{1,15,40} The RAND-36 questionnaire is a widely used and validated survey for the evaluation of health-related quality of life in 8 domains: physical functioning, social functioning, role limitations due to physical problems, role limitations due to emotional problems, mental health, vitality, bodily pain, and general health.^{1,15,40} The reliability of these individual domains ranges between an α of 0.78 and an α of 0.93.²⁹ Each domain is separately scored by standardization of scales, aggregation of scale scores, and transformation to summary scores, ranging between 0 and 100. Higher scores represent better function.^{1,15,40} The RAND-36 scores on the domains of social functioning, role limitations due to emotional problems, mental health, vitality, and general health were evaluated to assess psychosocial functioning.⁴⁰

Persistence of complaints

At follow-up, an anchor question was used to assess whether patients experienced persistent, reduced, or increased complaints compared with the first visit.

At baseline and follow-up, a visual analog scale (VAS) for pain during movement was assessed using a 100-mm VAS scale (on which 0 indicated no pain and 100 indicated maximal pain). Changes in pain over time were expressed as Δ VAS score.

At baseline and follow-up, the Western Ontario Rotator Cuff (WORC) index was used to assess quality of life through 5 domains on a scale from 0 (worst possible) to 100 (best possible).^{17,20} Changes in WORC scores were expressed as Δ WORC index score.

Statistical analysis

We first determined whether there were baseline differences in psychosocial functioning (RAND-36 score) and other characteristics (eg, age, sex, or treatment) between patients who had increased, persistent, and reduced complaints at follow-up by use of the independent-samples *t* test, Wilcoxon rank sum test, or χ^2 test depending on the type and distribution of data. Subsequently, logistic and linear regression analyses were performed to assess whether baseline psychosocial functioning (RAND-36 score) was associated with complaint persistence (anchor question, Δ VAS score, and Δ WORC index score). For this, the dependent variables (Δ VAS score and Δ WORC index score) were checked for a normal distribution. Additionally, we determined from scatter plots that the relations between the RAND-36 domains and the Δ VAS score and Δ WORC index score were linear and the residual errors had a normal distribution. The

analyses were also performed with the inclusion of age and sex to assess the influence of these factors on the estimated group differences.³⁸ Results from logistic and linear regression analyses were presented as odds ratios (ORs) and unstandardized β values, respectively.

The data were stored in a Microsoft Access 2010 database (version 14.0.7195.5000; Microsoft, Redmond, WA, USA) with SP2 MSO (Service Pack 2, Microsoft Office, version 14.0.7214.5000; Microsoft). For statistical analyses, SPSS software (version 20; IBM, Armonk, NY, USA) was used. A 2-sided P value $\leq .05$ was considered statistically significant.

Results

Thirty-four patients fulfilled our eligibility criteria. After a follow-up period of 3.8 years (standard deviation [SD], 0.5 years), 3 patients declined to participate, 1 patient had died, and 2 patients could not be contacted. The 6 patients lost to follow-up were aged 53 years (SD, 4.6 years), with 50% ($n = 3$) being women; the median complaint duration was 12

months (25th quartile–75th quartile, 12–30 months) at baseline. The eventual study cohort of 28 patients (82%) had a mean age of 50 years (SD, 6.5 years), with 61% ($n = 17$) being women; the median complaint duration was 17 months (25th quartile–75th quartile, 12–26 months). All these patients were treated conservatively with physical therapy ($n = 21$, 75%) and/or subacromial infiltrations ($n = 17$, 61%). Two patients were treated operatively after failed conservative management by Neer acromioplasty ($n = 1$) and distal clavicle resection ($n = 1$).

Compared with the first visit, none of the patients reported increased complaints on the anchor question at follow-up, whereas 9 patients (32%) reported persistent complaints and 19 (68%) reported reduced complaints. There were no differences in the received treatment between the 2 subgroups (Table I). Patients with persistent complaints at follow-up had a significantly longer median duration of complaints at presentation (30 months; quartiles, 20–69 months) than patients with reduced complaints at follow-up (12 months; quartiles, 9–24 months). There were no baseline differences in physical functioning and

Table I Baseline characteristics of patients with persistent or reduced shoulder complaints at follow-up

Patient characteristics	SAPS complaints at follow-up		Group difference	
	Persistent ($n = 9$)	Reduced ($n = 19$)	95% CI or statistic*	P value
Demographic characteristics				
Age, mean (SD), yr	47 (4.7)	52 (6.8)	-0.3 to 10	.063
Female, n (%)	6 (67)	11 (58)	0.2	.657
Right-side dominance, n (%)	9 (100)	15 (79)	2.2	.137
Dominant side affected, n (%)	7 (78)	10 (53)	1.6	.203
Anamnestic duration of complaints, median (quartiles), mo	30 (20 to 69)	12 (9.3 to 24)	-2.6	.009 [†]
Therapy, n (%)				
Physiotherapy	7 (78)	14 (74)	0.1	.815
Subacromial infiltration	7 (78)	10 (53)	1.6	.203
Operation	1 (11)	1 (5.3)	0.3	.575
Pain and disability				
VAS score during movement, median (quartiles)	44 (25 to 62)	32 (18 to 62)	-0.4	.681
WORC score, mean (SD)	59 (15)	58 (19)	-16 to 14	.864
RAND-36 score				
Physical functioning, median (quartiles)	75 (90 to 100)	80 (75 to 90)	-2.6 to 20	.127
Social functioning, median (quartiles)	75 (56 to 81)	88 (75 to 100)	-1.7	.099
Role limitations due to physical problems, median (quartiles)	0 (0 to 63)	75 (50 to 100)	-2.3	.023 [†]
Role limitations due to emotional problems, median (quartiles)	100 (67 to 100)	100 (100 to 100)	-0.1	.918
Mental health, mean (SD)	65 (14)	82 (12)	5.9 to 27	.003 [†]
Vitality, median (quartiles)	50 (45 to 60)	70 (60 to 80)	-2.9	.004 [†]
Bodily pain, mean (SD)	56 (15)	58 (16)	-11 to 14	.793
General health, median (quartiles)	45 (25 to 63)	75 (65 to 85)	-2.7	.008 [†]

SAPS, subacromial pain syndrome; CI, confidence interval; SD, standard deviation; VAS, visual analog scale; WORC, Western Ontario Rotator Cuff.

Complaint persistence was assessed with an anchor question at 4 years' follow-up. Depending on the type and distribution of data (histograms), data were analyzed with the independent-samples t test, Wilcoxon rank sum test, or χ^2 test.

* χ^2 Value or z score.

[†] Statistically significant ($P < .05$).

pain (RAND-36, VAS, or WORC score) between patients with persistent complaints and those with reduced complaints at follow-up (Table I). The patients who reported persistent complaints at follow-up did have lower baseline scores on the psychosocial functioning domains of the RAND-36 (ie, role limitations due to physical problems, mental health, vitality, and general health) than patients who reported reduced complaints.

Lower baseline levels of mental health (OR, 0.92; 95% confidence interval [CI], 0.85 to 0.98; $P = .013$), vitality (OR, 0.90; 95% CI, 0.83 to 0.98; $P = .011$), and general health (OR, 0.93; 95% CI, 0.88 to 0.98; $P = .009$) were associated with persistence of complaints as indicated on the anchor question (Table II). In accordance, baseline mental health (β , -0.90 ; 95% CI, -1.65 to -0.16), vitality (β , -1.03 ; 95% CI, -1.72 to -0.33), and general health (β , -0.79 ; 95% CI, -1.27 to -0.31) were negatively associated with Δ VAS score (Table II). Moreover, baseline mental health (β , 0.84; 95% CI, 0.19 to 1.48), vitality (β , 0.88; 95% CI, 0.26 to 1.50), and general health (β , 0.57; 95% CI, 0.11 to 1.02) were associated with Δ WORC index score. The association between baseline psychosocial functioning and persistence of complaints at follow-up was not affected by inclusion of sex and age in the analysis, as described in Supplementary Table S1.

Discussion

In this longitudinal study, we showed that lower baseline levels of mental health, vitality, and general health (RAND-36) are associated with persistence of complaints after approximately 4 years of routine care in patients with SAPS. The baseline scores of patients with persistent complaints in these RAND-36 domains were 12, 17, and 24 points lower, respectively, than those in the Dutch

population and comparable to those obtained from patients with depression.^{1,16,30}

In SAPS, few studies have been conducted on the association between shoulder complaints and psychosocial functioning.^{4,27} In confirmation with our study, depressive symptoms have been associated with recurring shoulder symptoms in a community-based sample from the general population.¹⁴ Furthermore, pain self-efficacy, expectations, and pain catastrophizing have been associated with chronic shoulder symptoms.^{4,32,36} In this longitudinal study, we showed that there is an association between lower levels of mental health, vitality, and general health and long-term persistence of complaints in SAPS, which emphasizes the role of psychosocial factors in SAPS.

Our study has some limitations. First, the calculations were performed on a small sample size. Because of limited power, we could not control for various factors such as the type of treatment received. However, because there were no differences in received treatment between the 2 subgroups with persistent vs. reduced complaints, we do not think that this introduced bias. Second, we did not assess all psychosocial constructs that may determine the course of complaints. Baseline psychosocial variables such as anxiety or catastrophizing may have an effect on long-term pain and function as well and should be evaluated in future studies.^{25,39} Third, there was a longer baseline duration of complaints in patients with persistent complaints at follow-up than in those with reduced complaints at follow-up. We therefore could not clarify cause-and-effect relationships, but this was also out of the scope of our study.

In SAPS, complaints become chronic in up to 40% of all patients.^{11,26,32,37} Clinicians generally focus on anatomic deformities and damaged subacromial tissues, whereas complaints do not correspond with magnetic resonance imaging pathologies in nearly half of all patients.^{9,10} Conditions external to the shoulder—for

Table II Baseline psychosocial functioning associated complaints at follow-up

RAND-36	Persistent complaints at follow-up		Δ VAS score		Δ WORC index score	
	OR (95% CI)	P value	β (95% CI)	P value	β (95% CI)	P value
Social functioning	0.97 (0.93 to 1.01)	.152	-0.39 (-0.99 to 0.20)	.183	0.08 (-0.45 to 0.61)	.755
Role limitations due to emotional problems	1.00 (0.98 to 1.02)	.843	-0.11 (-0.46 to 0.24)	.513	-0.07 (-0.38 to 0.24)	.627
Mental health	0.92 (0.85 to 0.98)	.013*	-0.90 (-1.65 to -0.16)	.020*	0.84 (0.19 to 1.48)	.013*
Vitality	0.90 (0.83 to 0.98)	.011*	-1.03 (-1.72 to -0.33)	.006*	0.88 (0.26 to 1.50)	.008*
General health	0.93 (0.88 to 0.98)	.009*	-0.79 (-1.27 to -0.31)	.002*	0.57 (0.11 to 1.02)	.017*

VAS, visual analog scale; WORC, Western Ontario Rotator Cuff; OR, odds ratio; CI, confidence interval.

Logistic regression was performed with complaint persistence as the dependent variable (with reduced complaints as the reference value), whereas linear regression was performed with the following dependent variables: change in VAS score for pain during movement over time (Δ VAS score) and change in WORC index score over time (Δ WORC score). The independent variables are the RAND-36 domains related to psychosocial functioning.

* Statistically significant ($P < .05$).

example, psychosocial functioning—may contribute to the perception of pain as well, and studies in other pain conditions have shown that addressing these factors results in quicker relief of pain and recovery.^{2,6,9,31,33} Future studies may investigate whether such a multimodal treatment may also benefit patients with SAPS.³³

Conclusion

In this prospective longitudinal study with nearly 4 years' follow-up, we showed that reduced psychosocial functioning in patients with SAPS is associated with long-term persistence of complaints. In other musculoskeletal pain conditions, it has been suggested that addressing coexisting psychosocial problems may enhance treatment outcome.^{2,6,31,33} Future studies may investigate whether a multimodal treatment with assessment of psychosocial functioning may also facilitate pain relief and recovery in SAPS.

Disclaimer

The institution of one or more of the authors has received funding from the Dutch Arthritis Foundation (grant no. 13-1-303).

The authors, their immediate families, and any research foundations with which they are affiliated have not received any financial payments or other benefits from any commercial entity related to the subject of this article.

Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.jse.2020.08.039>.

References

- Aaronson NK, Muller M, Cohen PD, Essink-Bot ML, Fekkes M, Sanderman R, et al. Translation, validation, and norming of the Dutch language version of the SF-36 Health Survey in community and chronic disease populations. *J Clin Epidemiol* 1998;51:1055-68.
- Bair MJ, Robinson RL, Katon W, Kroenke K. Depression and pain comorbidity: a literature review. *Arch Intern Med* 2003;163:2433-45. <https://doi.org/10.1001/archinte.163.20.2433>
- Brander V, Gondek S, Martin E, Stulberg SD. Pain and depression influence outcome 5 years after knee replacement surgery. *Clin Orthop Relat Res* 2007;464:21-6. <https://doi.org/10.1097/BLO.0b013e318126c032>
- Chester R, Jerosch-Herold C, Lewis J, Shepstone L. Psychological factors are associated with the outcome of physiotherapy for people with shoulder pain: a multicentre longitudinal cohort study. *Br J Sports Med* 2018;52:269-75. <https://doi.org/10.1136/bjsports-2016-096084>
- Chipchase LS, O'Connor DA, Costi JJ, Krishnan J. Shoulder impingement syndrome: preoperative health status. *J Shoulder Elbow Surg* 2000;9:12-5.
- Chou R, Qaseem A, Snow V, Casey D, Cross JT Jr, Shekelle P, et al. Diagnosis and treatment of low back pain: a joint clinical practice guideline from the American College of Physicians and the American Pain Society. *Ann Intern Med* 2007;147:478-91. <https://doi.org/10.7326/0003-4819-147-7-200710020-00006>
- De Baets L, Matheve T, Meeus M, Struyf F, Timmermans A. The influence of cognitions, emotions and behavioral factors on treatment outcomes in musculoskeletal shoulder pain: a systematic review. *Clin Rehabil* 2019;33:980-91. <https://doi.org/10.1177/0269215519831056>
- de Witte PB, Nagels J, van Arkel ER, Visser CP, Nelissen RG, de Groot JH. Study protocol subacromial impingement syndrome: the identification of pathophysiologic mechanisms (SISTIM). *BMC Musculoskelet Disord* 2011;12:282. <https://doi.org/10.1186/1471-2474-12-282>
- de Witte PB, Overbeek CL, Navas A, Nagels J, Reijnen M, Nelissen RG. Heterogeneous MR arthrography findings in patients with subacromial impingement syndrome—diagnostic subgroups? *J Electromyogr Kinesiol* 2016;29:64-73. <https://doi.org/10.1016/j.jelekin.2015.06.006>
- Diercks R, Bron C, Dorrestijn O, Meskers C, Naber R, de Ruiter T, et al. Guideline for diagnosis and treatment of subacromial pain syndrome: a multidisciplinary review by the Dutch Orthopaedic Association. *Acta Orthop* 2014;85:314-22. <https://doi.org/10.3109/17453674.2014.920991>
- Dorrestijn O, Stevens M, Winters JC, van der Meer K, Diercks RL. Conservative or surgical treatment for subacromial impingement syndrome? A systematic review. *J Shoulder Elbow Surg* 2009;18:652-60. <https://doi.org/10.1016/j.jse.2009.01.010>
- Feleus A, Bierma-Zeinstra SM, Miedema HS, Bernsen RM, Verhaar JA, Koes BW. Incidence of non-traumatic complaints of arm, neck and shoulder in general practice. *Man Ther* 2008;13:426-33. <https://doi.org/10.1016/j.math.2007.05.010>
- Gamsa A, Vikis-Freibergs V. Psychological events are both risk factors in, and consequences of, chronic pain. *Pain* 1991;44:271-7.
- Gill TK, Shanahan EM, Taylor AW, Buchbinder R, Hill CL. Shoulder pain in the community: an examination of associative factors using a longitudinal cohort study. *Arthritis Care Res (Hoboken)* 2013;65:2000-7. <https://doi.org/10.1002/acr.22082>
- Hays RD, Morales LS. The RAND-36 measure of health-related quality of life. *Ann Med* 2001;33:350-7.
- Hays RD, Sherbourne CD, Mazel RM. The RAND 36-Item Health Survey 1.0. *Health Econ* 1993;2:217-27.
- Holtby R, Razmjou H. Measurement properties of the Western Ontario rotator cuff outcome measure: a preliminary report. *J Shoulder Elbow Surg* 2005;14:506-10. <https://doi.org/10.1016/j.jse.2005.02.017>
- Jayakumar P, Overbeek CL, Lamb S, Williams M, Funes CJ, Gwilym S, et al. What factors are associated with disability after upper extremity injuries? A systematic review. *Clin Orthop Relat Res* 2018;476:2190-215. <https://doi.org/10.1097/CORR.0000000000000427>
- Kennedy P, Joshi R, Dhawan A. The effect of psychosocial factors on outcomes in patients with rotator cuff tears: a systematic review. *Arthroscopy* 2019;35:2698-706. <https://doi.org/10.1016/j.arthro.2019.03.043>
- Kirkley A, Griffin S, Dainty K. Scoring systems for the functional assessment of the shoulder. *Arthroscopy* 2003;19:1109-20. <https://doi.org/10.1016/j.arthro.2003.10.030>
- Kroenke K, Bair MJ, Damush TM, Wu J, Hoke S, Sutherland J, et al. Optimized antidepressant therapy and pain self-management in primary care patients with depression and musculoskeletal pain: a randomized controlled trial. *JAMA* 2009;301:2099-110. <https://doi.org/10.1001/jama.2009.723>
- Kromer TO, Sieben JM, de Bie RA, Bastiaenen CH. Influence of fear-avoidance beliefs on disability in patients with subacromial shoulder

- pain in primary care: a secondary analysis. *Phys Ther* 2014;94:1775-84. <https://doi.org/10.2522/ptj.20130587>
23. Kuijpers T, van Tulder MW, van der Heijden GJ, Bouter LM, van der Windt DA. Costs of shoulder pain in primary care consultants: a prospective cohort study in The Netherlands. *BMC Musculoskelet Disord* 2006;7:83. <https://doi.org/10.1186/1471-2474-7-83>
 24. Lentz TA, Barabas JA, Day T, Bishop MD, George SZ. The relationship of pain intensity, physical impairment, and pain-related fear to function in patients with shoulder pathology. *J Orthop Sports Phys Ther* 2009;39:270-7. <https://doi.org/10.2519/jospt.2009.2879>
 25. Linton SJ, Shaw WS. Impact of psychological factors in the experience of pain. *Phys Ther* 2011;91:700-11. <https://doi.org/10.2522/ptj.20100330>
 26. Luime JJ, Koes BW, Hendriksen IJ, Burdorf A, Verhagen AP, Miedema HS, et al. Prevalence and incidence of shoulder pain in the general population; a systematic review. *Scand J Rheumatol* 2004;33:73-81. <https://doi.org/10.1080/03009740310004667>
 27. Martinez-Calderon J, Meeus M, Struyf F, Miguel Morales-Asencio J, Gijon-Nogueron G, Luque-Suarez A. The role of psychological factors in the perpetuation of pain intensity and disability in people with chronic shoulder pain: a systematic review. *BMJ Open* 2018;8:e020703. <https://doi.org/10.1136/bmjopen-2017-020703>
 28. Martinez-Calderon J, Struyf F, Meeus M, Luque-Suarez A. The association between pain beliefs and pain intensity and/or disability in people with shoulder pain: a systematic review. *Musculoskelet Sci Pract* 2018;37:29-57. <https://doi.org/10.1016/j.msksp.2018.06.010>
 29. McHorney CA, Ware JE Jr, Lu JF, Sherbourne CD. The MOS 36-item Short-Form Health Survey (SF-36): III. Tests of data quality, scaling assumptions, and reliability across diverse patient groups. *Med Care* 1994;32:40-66.
 30. McHorney CA, Ware JE Jr, Raczek AE. The MOS 36-Item Short-Form Health Survey (SF-36): II. Psychometric and clinical tests of validity in measuring physical and mental health constructs. *Med Care* 1993;31:247-63.
 31. Patten SB. Long-term medical conditions and major depression in the Canadian population. *Can J Psychiatry* 1999;44:151-7.
 32. Reilingh ML, Kuijpers T, Tanja-Harfterkamp AM, van der Windt DA. Course and prognosis of shoulder symptoms in general practice. *Rheumatology (Oxford)* 2008;47:724-30. <https://doi.org/10.1093/rheumatology/ken044>
 33. Rej S, Dew MA, Karp JF. Treating concurrent chronic low back pain and depression with low-dose venlafaxine: an initial identification of "easy-to-use" clinical predictors of early response. *Pain Med* 2014;15:1154-62. <https://doi.org/10.1111/pme.12456>
 34. Rolfson O, Dahlberg LE, Nilsson JA, Malchau H, Garellick G. Variables determining outcome in total hip replacement surgery. *J Bone Joint Surg Br* 2009;91:157-61. <https://doi.org/10.1302/0301-620X.91B2.20765>
 35. Umer M, Qadir I, Azam M. Subacromial impingement syndrome. *Orthop Rev (Pavia)* 2012;4:e18. <https://doi.org/10.4081/or.2012.e18>
 36. Vahakari M, Leppilahti J, Hyvonen P, Ristiniemi J, Paivansalo M, Jalovaara P. Acromial shape in asymptomatic subjects: a study of 305 shoulders in different age groups. *Acta Radiol* 2010;51:202-6. <https://doi.org/10.3109/02841850903476556>
 37. van der Windt DA, Koes BW, Boeke AJ, Deville W, De Jong BA, Bouter LM. Shoulder disorders in general practice: prognostic indicators of outcome. *Br J Gen Pract* 1996;46:519-23.
 38. VanderZee KI, Sanderman R, Heyink JW, de Haes H. Psychometric qualities of the RAND 36-Item Health Survey 1.0: a multidimensional measure of general health status. *Int J Behav Med* 1996;3:104-22.
 39. Vranceanu AM, Jupiter JB, Mudgal CS, Ring D. Predictors of pain intensity and disability after minor hand surgery. *J Hand Surg Am* 2010;35:956-60. <https://doi.org/10.1016/j.jhssa.2010.02.001>
 40. Ware JE Jr, Sherbourne CD. The MOS 36-item short-form health survey (SF-36). I. Conceptual framework and item selection. *Med Care* 1992;30:473-83.