



Universiteit
Leiden
The Netherlands

Treatment strategies for metastatic soft tissue sarcomas

Wiltink, L.M.; Haas, R.L.M.; Gelderblom, H.; Sande, M.A.J. van de

Citation

Wiltink, L. M., Haas, R. L. M., Gelderblom, H., & Sande, M. A. J. van de. (2021). Treatment strategies for metastatic soft tissue sarcomas. *Cancers*, 13(7).
doi:10.3390/cancers13071722

Version: Publisher's Version

License: [Creative Commons CC BY 4.0 license](#)

Downloaded from: <https://hdl.handle.net/1887/3196192>

Note: To cite this publication please use the final published version (if applicable).

Treatment Strategies for Metastatic Soft Tissue Sarcomas

Lisette M. Wiltink ^{1,*}, Rick L. M. Haas ^{1,2} , Hans Gelderblom ³  and Michiel A. J. van de Sande ⁴ 

¹ Department of Radiation Oncology, Leiden University Medical Center, 2300 RC Leiden, The Netherlands; r.haas@nki.nl

² Department of Radiation Oncology, Netherlands Cancer Institute, 1006 BE Amsterdam, The Netherlands

³ Department of Medical Oncology, Leiden University Medical Center, 2300 RC Leiden, The Netherlands; A.J.Gelderblom@lumc.nl

⁴ Department of Orthopedic Surgery, Leiden University Medical Center, 2300 RC Leiden, The Netherlands; M.A.J.van_de_Sande@lumc.nl

* Correspondence: l.m.wiltink@lumc.nl

Soft tissue sarcomas (STS) are a diverse group of rare tumors of mesenchymal origin with different clinical, histologic and molecular characteristics [1]. While most patients have localized disease at diagnosis, about 10% present with synchronous metastatic disease [2]. Although myxoid and dedifferentiated liposarcomas metastasize to limb, soft tissue, bone, retroperitoneum and the pelvis, the most common site for metastasis of other sarcomas is the lungs. About 50% of high-grade STS patients eventually develop pulmonary metastases. Within 70% of these patients, the lungs remain the only organ with metastatic disease [3]. Patients with metastasized STS generally have a poor prognosis, with a median survival of 12 months [4]. However, for example, in the PICASSO III study, an improved median overall survival of 16–17 months is reported after systemic treatment [5].

The current ESMO clinical practice guideline recommends mainly systemic treatment for advanced and metastatic STS [6]. Anthracyclines (e.g., doxorubicin) are advised as first-line treatment. Combined treatment with ifosfamide can lead to higher tumor response rates and a prolonged progression free survival [7]. Clearly the anticipated side effects of this systemic therapy should be balanced with the expected benefits, in an effort to decrease the anticipated side effects of adding ifosfamide to anthracyclines. Dacarbazine could be considered, instead of ifosfamide, for leiomyosarcomas and solitary fibrous tumors [8,9]. Doxorubicin, in combination with the anti-PDGFRα biological olaratumab, has been investigated, but failed to show a survival benefit as compared to doxorubicin alone [10]. For several metastatic STS subtypes, such as angiosarcomas, leiomyosarcomas and dermatofibrosarcomas protuberans, histotype tailored chemotherapeutic regimes are recommended [6]. For other STS subtypes, histotype tailored chemotherapy was less effective than standard chemotherapy [11]. Further-line therapies, such as molecular targeted agents for selected subtypes, can be considered for patients with a progressive disease [6,12].

Surgery is generally only advised for metachronous lung metastasis, with a disease-free interval for more than one year, if the lesions are located in the lungs alone, and if all lesions could be removed [6,13]. The benefit of (neo)adjuvant chemotherapy in a metastasectomy setting is not yet clear, as a randomized study on neoadjuvant chemotherapy before lung metastasectomy has been initiated (EORTC 62933), but was prematurely stopped [14]. In a retrospective cohort comparing metastasectomy or chemotherapy, followed by metastasectomy, no difference in disease specific or progression free survival was found; however, an improved progression free survival was reported in patients with a good (complete) histological response following chemotherapy [15]. Radiological response was found to be a prognostic factor for improved progression free survival and disease specific survival [15].

In very selected cases, surgery, stereotactic radiotherapy, or ablation of metastasis could be considered without chemotherapy [6]. Recently, an improved survival, from 10.5



Citation: Wiltink, L.M.; Haas, R.L.M.; Gelderblom, H.; van de Sande, M.A.J. Treatment Strategies for Metastatic Soft Tissue Sarcomas. *Cancers* **2021**, *13*, 1722. <https://doi.org/10.3390/cancers13071722>

Received: 18 February 2021

Accepted: 31 March 2021

Published: 6 April 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

to 38.9 months, was found in soft tissue sarcoma patients after pulmonary metastasectomy alone. In multivariate analysis, metastasectomy was found to be the most important prognostic factor for survival [3]. In a cohort comparing patients with and without metastasectomy, a significantly improved 10 year overall survival of 17% versus 3% was found after surgery, leading to a median overall survival of 3.3 years compared to 0.9 years without surgery [16]. Furthermore, others studies have shown a better overall survival after metastasectomy [17,18]. Moreover, Wigge et al reported that repeated resections from different localizations was a strong predictor for improved survival [18]. Resection of oligometastatic hepatic metastasis can lead to an overall survival of 90.3% at one year, and, if a radical resection of the lesion could be achieved, extrahepatic disease should not be considered as a contraindication for resection [19]. Radiotherapy could also be used to locally control metastatic disease. Several cohorts show excellent outcomes, with minimal toxicity after stereotactic radiation therapy [20–23]. At a one year follow-up, a local control rate of 94% and an overall survival of 76% are reported [21]. Brachytherapy of metastasis with (125)I seeds is reported to improve progression free survival after a failure of first-line chemotherapy, from 3.6 to 7 months, compared to chemotherapy alone [24]. A few studies have been published about interventional radiology treatment options, such as radio frequent ablation. In one of these, ablation is shown to be effective for local metastatic control in leiomyosarcomas, with a local control rate at one year of 95.2% [25].

All studies focusing on local control of metastatic disease and survival are retrospective, which may lead to selection bias, as, typically, only selected patients are offered local therapy for metastases. Since STSs are relatively rare, it is difficult to design prospective patient cohorts, and even more difficult to include enough patient numbers in randomized controlled trials, in order to compare different treatment options in a homogeneous patient population. There is also reluctance from patients and physicians to randomize between local therapy and no local therapy, as there is almost always a center preference. For that reason, these trials are likely to fail due to poor recruitment. With the relatively poor prognosis of STS, quality of life during and after treatment is even regarded as more important; yet, in almost none of the studies is quality of life after treatment reported.

Finally, another treatment option not yet discussed, is best supportive care, which always remains the alternative for patients unfit or unwilling to undergo local or systemic treatments. Short-course palliative radiotherapy could be part of this strategy by relieving pain caused by metastasis, with minimal toxicity [26].

In conclusion, palliative systemic treatment remains the standard treatment in metastatic disease, but, accumulating evidence shows improved overall survival after management of oligometastatic disease with surgery, radiotherapy, or interventional radiological ablation techniques. Traditionally, metastasectomy is the treatment of choice to achieve local control, but similar local control rates are reported after stereotactic radiotherapy techniques. It is likely that only a subset of STS patients will benefit from local metastatic treatment. Typically, these decisions can only be made in a multidisciplinary sarcoma tumor board in a reference center, as STS includes over 80 subtypes [6]. Specific treatment for each metastasized subtype will not be available in the near future. Well-designed studies should be encouraged to support subgroup analysis, as should quality of life outcome measures to help develop the best treatment strategy for each STS subtype.

Funding: This research received no external funding.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Fletcher, C.D. *Pathology of Soft Tissue Sarcomas*; Churchill Livingstone: Edinburgh, UK, 1990.
2. Komdeur, R.; Hoekstra, H.J.; van den Berg, E.; Molenaar, W.M.; Pras, E.; de Vries, E.G.; van der Graaf, W.T. Metastasis in soft tissue sarcomas: Prognostic criteria and treatment perspectives. *Cancer Metastasis Rev.* **2002**, *21*, 167–183. [[CrossRef](#)] [[PubMed](#)]
3. Kawamoto, T.; Hara, H.; Morishita, M.; Fukase, N.; Kawakami, Y.; Takemori, T.; Fujiwara, S.; Kitayama, K.; Yahiro, S.; Miyamoto, T.; et al. Prognostic influence of the treatment approach for pulmonary metastasis in patients with soft tissue sarcoma. *Clin. Exp. Metastasis* **2020**, *37*, 509–517. [[CrossRef](#)] [[PubMed](#)]

4. Italiano, A.; Mathoulin-Pelissier, S.; Cesne, A.L.; Terrier, P.; Bonvalot, S.; Collin, F.; Michels, J.J.; Blay, J.Y.; Coindre, J.M.; Bui, B. Trends in survival for patients with metastatic soft-tissue sarcoma. *Cancer* **2011**, *117*, 1049–1054. [[CrossRef](#)] [[PubMed](#)]
5. Ryan, C.W.; Merimsky, O.; Agulnik, M.; Blay, J.Y.; Schuetz, S.M.; Van Tine, B.A.; Jones, R.L.; Elias, A.D.; Choy, E.; Alcindor, T.; et al. PICASSO III: A Phase III, Placebo-Controlled Study of Doxorubicin With or Without Palifosfamide in Patients With Metastatic Soft Tissue Sarcoma. *J. Clin. Oncol.* **2016**, *34*, 3898–3905. [[CrossRef](#)] [[PubMed](#)]
6. Casali, P.G.; Abecassis, N.; Aro, H.T.; Bauer, S.; Biagini, R.; Bielack, S.; Bonvalot, S.; Boukovinas, I.; Bovee, J.; Brodowicz, T.; et al. Soft tissue and visceral sarcomas: ESMO-EURACAN Clinical Practice Guidelines for diagnosis, treatment and follow-up. *Ann. Oncol.* **2018**, *29*, iv51–iv67. [[CrossRef](#)]
7. Judson, I.; Verweij, J.; Gelderblom, H.; Hartmann, J.T.; Schöffski, P.; Blay, J.Y.; Kerst, J.M.; Sufliarsky, J.; Whelan, J.; Hohenberger, P.; et al. Doxorubicin alone versus intensified doxorubicin plus ifosfamide for first-line treatment of advanced or metastatic soft-tissue sarcoma: A randomised controlled phase 3 trial. *Lancet Oncol.* **2014**, *15*, 415–423. [[CrossRef](#)]
8. D’Ambrosio, L.; Touati, N.; Blay, J.Y.; Grignani, G.; Flippot, R.; Czarnecka, A.M.; Piperno-Neumann, S.; Martin-Broto, J.; Sanfilippo, R.; Katz, D.; et al. Doxorubicin plus dacarbazine, doxorubicin plus ifosfamide, or doxorubicin alone as a first-line treatment for advanced leiomyosarcoma: A propensity score matching analysis from the European Organization for Research and Treatment of Cancer Soft Tissue and Bone Sarcoma Group. *Cancer* **2020**, *126*, 2637–2647. [[CrossRef](#)]
9. Stacchiotti, S.; Tortoreto, M.; Bozzi, F.; Tamborini, E.; Morosi, C.; Messina, A.; Libertini, M.; Palassini, E.; Cominetti, D.; Negri, T.; et al. Dacarbazine in solitary fibrous tumor: A case series analysis and preclinical evidence vis-a-vis temozolomide and antiangiogenics. *Clin. Cancer Res.* **2013**, *19*, 5192–5201. [[CrossRef](#)] [[PubMed](#)]
10. Tap, W.D.; Wagner, A.J.; Schöffski, P.; Martin-Broto, J.; Krarup-Hansen, A.; Ganjoo, K.N.; Yen, C.C.; Abdul Razak, A.R.; Spira, A.; Kawai, A.; et al. Effect of Doxorubicin Plus Olaratumab vs Doxorubicin Plus Placebo on Survival in Patients With Advanced Soft Tissue Sarcomas: The ANNOUNCE Randomized Clinical Trial. *JAMA* **2020**, *323*, 1266–1276. [[CrossRef](#)] [[PubMed](#)]
11. Gronchi, A.; Ferrari, S.; Quagliuolo, V.; Broto, J.M.; Pousa, A.L.; Grignani, G.; Basso, U.; Blay, J.Y.; Tendero, O.; Beveridge, R.D.; et al. Histotype-tailored neoadjuvant chemotherapy versus standard chemotherapy in patients with high-risk soft-tissue sarcomas (ISG-ST5 1001): An international, open-label, randomised, controlled, phase 3, multicentre trial. *Lancet Oncol.* **2017**, *18*, 812–822. [[CrossRef](#)]
12. van der Velden, D.L.; Hoes, L.R.; van der Wijngaart, H.; van Berge Henegouwen, J.M.; van Werkhoven, E.; Roepman, P.; Schilsky, R.L.; de Leng, W.W.J.; Huitema, A.D.R.; Nuijen, B.; et al. The Drug Rediscovery protocol facilitates the expanded use of existing anticancer drugs. *Nature* **2019**, *574*, 127–131. [[CrossRef](#)] [[PubMed](#)]
13. Blackmon, S.H.; Shah, N.; Roth, J.A.; Correa, A.M.; Vaporciyan, A.A.; Rice, D.C.; Hofstetter, W.; Walsh, G.L.; Benjamin, R.; Pollock, R.; et al. Resection of pulmonary and extrapulmonary sarcomatous metastases is associated with long-term survival. *Ann. Thorac Surg.* **2009**, *88*, 877–884. [[CrossRef](#)] [[PubMed](#)]
14. ClinicalTrials.gov. Surgery With or Without Combination Chemotherapy in Treating Patients With Lung Metastases From Soft Tissue Sarcoma. Available online: <https://clinicaltrials.gov/ct2/show/study/NCT00002764> (accessed on 4 February 2021).
15. Ohnstad, H.O.; Bruland, O.S.; Taksdal, I.; Bjerkeheggen, B.; Nenadovic, M.; Sæter, G.; Jørgensen, L.H.; Hall, K.S. Response to preoperative chemotherapy in patients undergoing resection of pulmonary metastasis from soft tissue sarcoma—a predictor of outcome? *Acta Oncol.* **2014**, *53*, 1180–1187. [[CrossRef](#)] [[PubMed](#)]
16. Smolle, M.A.; van Praag, V.M.; Posch, F.; Bergovec, M.; Leitner, L.; Friesenbichler, J.; Heregger, R.; Riedl, J.M.; Pichler, M.; Gerger, A.; et al. Surgery for metachronous metastasis of soft tissue sarcoma—A magnitude of benefit analysis using propensity score methods. *Eur. J. Surg. Oncol.* **2019**, *45*, 242–248. [[CrossRef](#)]
17. Krishnan, C.K.; Kim, H.S.; Park, J.W.; Han, I. Outcome after Surgery for Extremity Soft Tissue Sarcoma in Patients Presenting With Metastasis at Diagnosis. *Am. J. Clin. Oncol.* **2018**, *41*, 681–686. [[CrossRef](#)] [[PubMed](#)]
18. Wigge, S.; Heißner, K.; Steger, V.; Ladurner, R.; Traub, F.; Sipos, B.; Bösmüller, H.; Kanz, L.; Mayer, F.; Kopp, H.G. Impact of surgery in patients with metastatic soft tissue sarcoma: A monocentric retrospective analysis. *J. Surg. Oncol.* **2018**, *118*, 167–176. [[CrossRef](#)] [[PubMed](#)]
19. Marudanayagam, R.; Sandhu, B.; Perera, M.T.; Bramhall, S.R.; Mayer, D.; Buckels, J.A.; Mirza, D.F. Liver resection for metastatic soft tissue sarcoma: An analysis of prognostic factors. *Eur. J. Surg. Oncol.* **2011**, *37*, 87–92. [[CrossRef](#)]
20. Baumann, B.C.; Bernstein, K.A.; DeLaney, T.F.; Simone, C.B., 2nd; Kolker, J.D.; Choy, E.; Levin, W.P.; Weber, K.L.; Muniappan, A.; Berman, A.T.; et al. Multi-institutional analysis of stereotactic body radiotherapy for sarcoma pulmonary metastases: High rates of local control with favorable toxicity. *J. Surg. Oncol.* **2020**, *122*, 877–883. [[CrossRef](#)] [[PubMed](#)]
21. Baumann, B.C.; Nagda, S.N.; Kolker, J.D.; Levin, W.P.; Weber, K.L.; Berman, A.T.; Staddon, A.; Hartner, L.; Hahn, S.M.; Glatstein, E.; et al. Efficacy and safety of stereotactic body radiation therapy for the treatment of pulmonary metastases from sarcoma: A potential alternative to resection. *J. Surg. Oncol.* **2016**, *114*, 65–69. [[CrossRef](#)]
22. Lindsay, A.D.; Haupt, E.E.; Chan, C.M.; Spiguel, A.R.; Scarborough, M.T.; Zlotecki, R.A.; Gibbs, P.C. Treatment of Sarcoma Lung Metastases with Stereotactic Body Radiotherapy. *Sarcoma* **2018**, *2018*, e9132359. [[CrossRef](#)]
23. Loi, M.; Duijm, M.; Baker, S.; Rossi, L.; Grunhagen, D.; Verhoef, C.; Nuytens, J. Stereotactic body radiotherapy for oligometastatic soft tissue sarcoma. *Radiol. Med.* **2018**, *123*, 871–878. [[CrossRef](#)] [[PubMed](#)]
24. Mo, Z.; Zhang, T.; Zhang, Y.; Xiang, Z.; Yan, H.; Zhong, Z.; Gao, F.; Zhang, F. Feasibility and clinical value of CT-guided (125)I brachytherapy for metastatic soft tissue sarcoma after first-line chemotherapy failure. *Eur. Radiol.* **2018**, *28*, 1194–1203. [[CrossRef](#)] [[PubMed](#)]

-
25. Gravel, G.; Yevich, S.; Tselikas, L.; Mir, O.; Teriitehau, C.; De Baère, T.; Deschamps, F. Percutaneous thermal ablation: A new treatment line in the multidisciplinary management of metastatic leiomyosarcoma? *Eur. J. Surg. Oncol.* **2017**, *43*, 181–187. [[CrossRef](#)] [[PubMed](#)]
 26. Janssen, S.; Bolm, L.; Käsmann, L.; Bartscht, T.; Rades, D. Palliative Radiation Therapy for Spinal Cord Compression from Metastatic Soft Tissue Sarcoma. *In Vivo* **2016**, *30*, 529–531. [[PubMed](#)]