



Universiteit  
Leiden  
The Netherlands

## **Characterization of dynamic changes in Matrix Gla Protein (MGP) gene expression as function of genetic risk alleles, osteoarthritis relevant stimuli, and the vitamin K inhibitor warfarin**

Houtman, E.; Almeida, R.C. de; Tuerlings, M.; Suchiman, H.E.D.; Broekhuis, D.; Nelissen, R.G.H.H.; ... ; Meulenbelt, I.

### **Citation**

Houtman, E., Almeida, R. C. de, Tuerlings, M., Suchiman, H. E. D., Broekhuis, D., Nelissen, R. G. H. H., ... Meulenbelt, I. (2021). Characterization of dynamic changes in Matrix Gla Protein (MGP) gene expression as function of genetic risk alleles, osteoarthritis relevant stimuli, and the vitamin K inhibitor warfarin. *Osteoarthritis And Cartilage*, 29(8), 1193-1202. doi:10.1016/j.joca.2021.05.001

Version: Publisher's Version

License: [Creative Commons CC BY-NC-ND 4.0 license](https://creativecommons.org/licenses/by-nc-nd/4.0/)

Downloaded from: <https://hdl.handle.net/1887/3206810>

**Note:** To cite this publication please use the final published version (if applicable).

## Characterization of dynamic changes in Matrix Gla Protein (MGP) gene expression as function of genetic risk alleles, osteoarthritis relevant stimuli, and the vitamin K inhibitor warfarin



E. Houtman †, R. Coutinho de Almeida †, M. Tuerlings †, H.E.D. Suchiman †, D. Broekhuis ‡, R.G.H.H. Nelissen ‡, Y.F.M. Ramos †, J.B.J. van Meurs §, I. Meulenbelt †\*

† Molecular Epidemiology, Department of Biomedical Data Sciences, Leiden University Medical Center, Leiden, the Netherlands

‡ Department of Orthopaedics, Leiden University Medical Center, Leiden, the Netherlands

§ Department of Internal Medicine, Erasmus MC, University Medical Center, Rotterdam, the Netherlands

### ARTICLE INFO

#### Article history:

Received 25 November 2020

Accepted 5 May 2021

#### Keywords:

Osteoarthritis  
Articular cartilage  
Subchondral bone  
Matrix Gla protein  
Warfarin  
Vitamin K  
Genetic risk

### SUMMARY

**Objective:** We here aimed to characterize changes of Matrix Gla Protein (*MGP*) expression in relation to its recently identified OA risk allele rs1800801-T in OA cartilage, subchondral bone and human *ex vivo* osteochondral explants subjected to OA related stimuli. Given that *MGP* function depends on vitamin K bioavailability, we studied the effect of frequently prescribed vitamin K antagonist warfarin.

**Methods:** Differential (allelic) mRNA expression of *MGP* was analyzed using RNA-sequencing data of human OA cartilage and subchondral bone. Human osteochondral explants were used to study exposures to interleukin one beta (IL-1 $\beta$ ; inflammation), triiodothyronine (T3; Hypertrophy), warfarin, or 65% mechanical stress (65%MS) as function of rs1800801 genotypes.

**Results:** We confirmed that the *MGP* risk allele rs1800801-T was associated with lower expression and that *MGP* was significantly upregulated in lesioned as compared to preserved OA tissues, mainly in risk allele carriers, in both cartilage and subchondral bone. Moreover, *MGP* expression was downregulated in response to OA like triggers in cartilage and subchondral bone and this effect might be reduced in carriers of the rs1800801-T risk allele. Finally, warfarin treatment in cartilage increased *COL10A1* and reduced *SOX9* and *MMP3* expression and in subchondral bone reduced *COL1A1* and *POSTN* expression.

**Discussion & conclusions:** Our data highlights that the genetic risk allele lowers *MGP* expression and upon OA relevant triggers may hamper adequate dynamic changes in *MGP* expression, mainly in cartilage. The determined direct negative effect of warfarin on human explant cultures functionally underscores the previously found association between vitamin K deficiency and OA.

© 2021 The Authors. Published by Elsevier Ltd on behalf of Osteoarthritis Research Society International. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

### Introduction

Osteoarthritis (OA) is the most common degenerative disease of joints and its incidence is rising with increasing obesity and age, resulting in a high social and economic burden on society. Interacting risk factors for OA include obesity, age, sex, abnormal loading and genetic factors. The genetic component of OA is estimated to be

in the range of 40–60%<sup>1,2</sup>. For that matter, large-scale genome wide association studies (GWAS) have identified strong, in other words highly significant and reproducible, OA risk genes involved in the aetiology of OA, whereas follow-up studies have shown that risk single nucleotide polymorphisms (SNPs) frequently modulate pathology due to altering transcription of the genes in *cis* both in bone and cartilage<sup>3–6</sup>.

In this regard Matrix Gla protein (*MGP*) via rs4764133<sup>7</sup> with proxy SNPs rs1800801 and rs4236<sup>8</sup>, was previously identified as strong OA risk gene for hand OA with the OA conferring allele associated with lower expression of *MGP* relative to the non-risk allele<sup>7</sup> in a range of joint tissues but its effect was most profound in cartilage and subchondral bone<sup>7,9</sup>. On the other hand, these studies

\* Address correspondence and reprint requests to: I. Meulenbelt, Department of Medical Statistics and Bioinformatics, Section of Molecular Epidemiology, Leiden University Medical Center, LUMC Post-zone S-05-P, PO Box 9600, 2300 RC, Leiden, the Netherlands.

E-mail address: [i.meulenbelt@lumc.nl](mailto:i.meulenbelt@lumc.nl) (I. Meulenbelt).

could not identify significant differential expression of *MGP* in OA pathophysiology in macroscopically lesioned OA compared to preserved cartilage<sup>7</sup> nor in macroscopically preserved cartilage compared to healthy cartilage<sup>9</sup>. These differential expression analyses were, however, determined in a relatively small sample size.

*MGP* regulates extracellular calcium levels via high affinity to its  $\gamma$ -carboxyglutamic acid (Gla) residues. Low *MGP* levels results in higher calcification of cartilage tissue and a reduced bone mineral density<sup>10–12</sup>. As the OA risk allele (rs1800801) has been associated with a reduced *MGP* gene expression<sup>12</sup> and with increased vascular calcification<sup>13</sup>, this would suggest increased cartilage calcification in carriers of the OA risk allele. The latter was further justified by recapitulating downregulation of *MGP* in cartilage chondrocytes resulting in pro-catabolic (*ADAMTS4*, *MMP13*), as well as pro-hypertrophic (*COL10A1*, *VEGFA*) mRNA signalling<sup>9</sup>. The *MGP* protein is produced by the cell in inactive form and is dependent on vitamin K for activation, via carboxylation (c-*MGP*). As such, low vitamin K levels have been hypothesized to play a role in OA pathogenesis<sup>14,15</sup>. Similarly, vitamin K antagonists such as warfarin, that are frequently prescribed for the prevention of thromboembolic events in patients with atrial fibrillation<sup>16</sup>, have been suggested to predispose to OA<sup>17</sup>. Nonetheless, the direct effect of warfarin on human articular cartilage tissue homeostasis has not been assessed.

Here we set out to explore *MGP* gene expression in relation to the OA risk allele rs1800801-T, in a large RNA-sequencing dataset containing both macroscopically preserved and lesioned cartilage<sup>18</sup> and subchondral bone<sup>19</sup> as well as in our recently established full thickness human *ex vivo* osteochondral explant model<sup>20</sup>. The latter allowing us to study the effect of the OA risk allele on the dynamic *MGP* response to different OA related stimuli, such as inflammation (Interleukin one beta (IL-1 $\beta$ )), hypertrophy (Triiodothyronine (T3)) and 65% mechanical stress (65%MS). Moreover, we used the human *ex vivo* explant model to study the direct effect of vitamin K antagonist, warfarin, on articular cartilage and subchondral bone homeostasis.

## Material and methods

### Sample description

Human material was obtained from the Research in Articular Osteoarthritis Cartilage (RAAK) biobank as previously described in detail<sup>21</sup>. The RAAK study is approved by the medical ethics committee of the Leiden University Medical Center (P08.239/P19.013). In this study, RNA-sequencing data was included of paired macroscopically preserved and lesioned OA cartilage of  $N = 35$  participants<sup>18</sup> and subchondral bone of  $N = 24$  participants<sup>19</sup> for which sample characteristics have previously been described. In total 136 osteochondral explants were harvested from the macroscopically preserved condyle knee joints of  $N = 18$  participants and divided over the different experiment (Supplementary Fig. 1). Multiple osteochondral explants containing both cartilage and bone (diameter of 8 mm) were extracted per participant and washed in sterile PBS before taking into culture. Donor characteristics of osteochondral explants are described in Supplementary Table 1 and study design is described in Supplementary Fig. 1. For additional details on neo-cartilage deposition, RNA and DNA isolations, Taq-Man genotyping, RNA sequencing data of cartilage and subchondral bone, unconfined dynamic (cyclic) compression, expression quantitative trait loci (eQTL), Allelic Expression Imbalance (AEI) and data analysis, see the Supplementary Methods.

### Treatment of osteochondral explants

Explants were cultured in 24 wells plates (Greiner CELLSTAR; Sigma) supplemented with 1.5 ml CDM in a 5% (v/v) CO<sub>2</sub> incubator

at 37°C. Three days after extraction, explants were treated with either IL-1 $\beta$  (10 ng/ml), triiodothyronine (T3, 10 nM; Sigma) or warfarin (50  $\mu$ M; Sigma), depicted in Fig. 1(A). Six days after extraction, dynamic unconfined compression was applied to explant tissues using the Mach-1 mechanical testing system (Biomomentum Inc., Laval, QC, Canada) on four subsequent days [Fig. 1(B)]. Mechanical stress was applied at a strain of 65% of cartilage height and at a frequency of 1 Hz, mimicking walking speed [Fig. 1(C)]. Cartilage and bone were separated using a scalpel, snap-frozen in liquid nitrogen, and stored at  $-80^{\circ}\text{C}$  for RNA isolation.

### Reverse transcription and Real-Time PCR


Real-Time PCR for gene expression was performed with QuantStudio 6 Real-Time PCR system (Applied Biosystems) using Fast Start Sybr Green Master mix (Roche Applied Science). Primer sequences (Table 1) used were tested for linear amplification and missing datapoints for genes are summarized in Supplementary Tables 2 and 3. Details on normalization can be found in the Supplementary Methods.

### Statistical analyses

Differential *MGP* expression analyses between preserved and lesioned OA cartilage and bone including false discovery rates (FDR) as multiple testing correction for the genome wide analyses were reproduced from Coutinho<sup>18</sup> and Tuerlings<sup>19</sup>, respectively. Description on their study design and sample numbers are in Supplementary Methods. To assess allelic expression imbalance (AEI) we applied our previously published methodology in R<sup>22</sup> to RNA sequence data of *MGP* in the larger dataset of cartilage<sup>18</sup> and a dataset of bone<sup>19</sup> which is further outlined in Supplementary Methods.

To test expression quantitative trait analyses (eQTL) and differential expression of *MGP* in genotype strata in the current manuscript, we used the variance stabilizing transformation (VST) normalized *MGP* expression levels of these RNA sequencing datasets and used generalized estimating equations (GEE)<sup>23</sup> to effectively adjust for dependencies of genotypes among donors by adding a random effect for sample donor. Details of the models applied are outlined in Supplementary Methods.

*MGP* expression by RT-qPCR in the *in vitro* 3D-neo cartilage formation was estimated using a generalized linear mixed model (GLMM) using *MGP* levels ( $-\Delta\text{CT}$ ) as dependent variable and time as repeated measure:  $MGP\ level \sim Time + (1|Donor)$ . In the osteochondral explant models fold changes (FC) of RT-qPCR expression were determined by calculating the log<sub>2</sub> of the  $-\Delta\Delta\text{CT}$  for each sample ( $2^{-\Delta\Delta\text{CT}}$ ) where  $FC > 1$  is upregulation and  $FC < 1$  is downregulation of treated samples compared to control samples. The reported *P*-values were determined by applying GEE to  $-\Delta\text{CT}$  values to effectively adjust for dependencies among donors of the explants by adding a random effect for sample donor as we did not have perfect pairs for each analysis. We followed a linear GEE model, with *MGP* level as dependent variable, treatment as factor and exchangeable working matrix:  $MGP\ level \sim Treatment + (1|Donor)$ <sup>24</sup>. Differences in effect sizes between strata was determined by performing unpaired student's *t*-test on the fold changes corrected for control samples. Warfarin treated osteochondral explants samples were paired hence a paired sample *t*-test was performed to determine between-group differences and *p*-values. Except for AEI, Statistical analyses were performed in SPSS statistics 23 (IBM). Outliers were investigated using Grubbs's test and normal distribution was determined using Shapiro–Wilk test and visually

**Fig. 1**

**Schematic representation of different perturbations applied to osteochondral explants.** Osteochondral explants were punched from the still macroscopically preserved looking knee condyle area and taken into culture. **[A]** Explants were subjected to treatment with IL-1 $\beta$  (10 ng/ml), triiodothyronine (T3; 10 nM) or warfarin (50  $\mu$ M). **[B]** Explants received mechanical stresses at a strain of 65% for 10 min per day on four subsequent days. On day 13, cartilage and bone was separated, snap frozen and stored at  $-80^{\circ}\text{C}$ . **[C]** Schematic representation of the dynamic (cyclic) compression applied to osteochondral explants. Legend: N=Newton; h=height.

Osteoarthritis  
and Cartilage

inspecting Q–Q plots. The boxplots represent 25th, 50th and 75th percentiles, and whiskers extend to the 95%CI.

## Results

### Expression patterns of MGP in previous established RNA sequencing datasets of preserved and lesioned OA cartilage and subchondral bone

We used our previously established RNA sequencing dataset of macroscopically preserved and lesioned OA cartilage samples ( $N = 35$  pairs<sup>18</sup>) and subchondral bone ( $N = 24$  pairs<sup>19</sup>), to examine differential MGP expression with OA tissue status and with the OA risk SNPs (see [Supplementary Methods and Figures](#)). An increased expression of MGP in lesioned compared to preserved OA cartilage was observed (FC = 1.45, 95%CI [1.24; 1.61],  $P$ -value =  $1.78 \times 10^{-3}$ ) and this increase of MGP was genome wide significant (FDR = 0.021). Similarly, MGP was upregulated in lesioned compared to preserved OA subchondral bone (FC = 1.53, 95%CI [1.22; 1.64],  $P$ -value = 0.023), but this was not genome wide significant (FDR = 0.12). Together, these results show a robust upregulation of MGP expression with ongoing OA pathophysiology.

Here we studied whether the MGP differential expression between preserved and lesioned OA tissues was affected by MGP risk allele carriership. As shown in [Fig. 2\(A\)](#), the MGP upregulation occurs particularly among risk allele carriers rs1800801-T in lesioned compared to preserved OA cartilage independent of age


and sex of donors (OR = 2.70, 95%CI [1.16; 6.29],  $P$ -value = 0.021). Notably however, the overall MGP expression remains lower among risk allele carriers rs1800801-T as compared to carriers of the reference allele rs1800801-C. The same effect was observed in subchondral bone, where MGP was found to be upregulated in lesioned compared to preserved tissue only in risk allele carriers rs1800801-T (OR = 3.04, 95%CI [1.24; 7.45],  $P$ -value = 0.015) independent of age and sex of donors ([Fig. 2\(B\)](#)).

Next, we attempted to replicate the previously shown AEI of MGP in association with the OA risk SNP rs1800801<sup>7</sup> in heterozygous individuals in this larger RNA sequencing dataset of preserved and lesioned OA cartilage<sup>18</sup> and a novel dataset of OA subchondral bone<sup>19</sup>. Additionally, we explored whether the effect size in AEI differed in these tissues between preserved and lesioned areas. As shown in [Supplementary Fig. 2\(A\)](#) we confirmed AEI expression of MGP in preserved OA cartilage with the risk-conferring allele rs1800801-T associated to a reduced MGP expression of 10% (95%CI [2.24; 18.64]) relative to the reference allele rs1800801-C. In lesioned OA cartilage the AEI was very comparable with rs1800801-T associated to a reduced MGP expression of 11% (95%CI [2.25; 19.49]) relative to the reference allele rs1800801-C. In subchondral bone, genotype of rs1800801 could not be called thus we used its proxy SNP rs4236 ( $r^2 = 0.93$  with rs1800801), which was also investigated previously<sup>7</sup>. As shown in [Supplementary Fig. 2\(B\)](#), we confirmed AEI of MGP in preserved OA subchondral bone with the risk-conferring allele rs4236-C associated to a reduced MGP expression of 10% (95%CI [5.70; 14.52]) relative to the reference

| Gene name | Forward 5'-3'            | Reverse 5'-3'            |
|-----------|--------------------------|--------------------------|
| SDHA      | TGGAGCTGCAGAACCTGATG     | TGTAGTCTTCCTGGCATGC      |
| MGP       | CGCCCCAGATTGATAAGTA      | TCTCCTTTGACCCTCACTGC     |
| SOX9      | CCCCAACAGATCGCCTACAG     | CTGGAGTTCTGGTGGTCGGT     |
| ACAN      | AGAGACTCACACAGTCGAAACAGC | CTATGTTACAGTGCTCGCCAGTG  |
| COL2A1    | CTACCCCAATCCAGCAAACGT    | AGGTGATGTTCTGGGAGCCTT    |
| RUNX2     | CTGTGGTACTGTGATGGCG      | AGGTAGTACTTGGGGAGGA      |
| ALPL      | CAAAGGCTTCTTCTGTGGTG     | CCTGCTTGCTTTTCTTCA       |
| COL1A1    | GTGCTAAAGGTGCCAATGGT     | ACCAGGTTACCGCTGTAC       |
| COL10A1   | GGCAACAGCATTATGACCA      | TGAGATCGATGATGGCACTCC    |
| MMP3      | GAGGCATCCACACCTAGGTT     | TCAGAAATGGTGCATCGATT     |
| MMP13     | TTGAGCTGGACTCATTGTCG     | GGAGCCTCTCAGTCATGGAG     |
| ADAMTS5   | TGGCTCACGAAATCGGACAT     | GGCTTATCTTCTGTGGAAAC     |
| COMP      | ACAATGACGGAGTCCTGAC      | TCGCATCAAAGTCGCTCTG      |
| OMD       | GGACACAACAAATTAAGCAAGC   | TGGTGTAATGTAGTGGGTCA     |
| BGLAP     | CCCTCTGCTYGGACACAAA      | CACACTCTCGCCCTATTGG      |
| OGN       | TGATGAAATGCCACGTGTC      | TTTGTAAGGGTGGTACAGCA     |
| SPP1      | GCCAGTTGCAGCTTCTCA       | AAAAGCAAATCACTGCAATTCTCA |
| TNFRSF11B | TTGATGAAAGCTTACCCGGGA    | TCTGGTCACTGGGTTTGCATC    |
| BMP2      | TCCATGTGGACGCTCTTCA      | AGCAGCAACGCTAGAAGACA     |
| POSTN     | TACACTTGTGGACCTGT        | TTAAGGAGCGCTGATCCA       |

**Table 1** Primer sequence used to determine gene expression levels in real-time PCR

Osteoarthritis  
and Cartilage



**Fig. 2**

**MGP expression as function of the transcript and OA risk SNP rs1800801.** [A] Variance stabilizing transformation (VST) normalized *MGP* expression levels extracted from the RNA sequencing dataset in preserved and lesioned OA cartilage stratified for rs1800801 genotype CC ( $n_{\text{Preserved}} = 19$  vs  $n_{\text{Lesioned}} = 16$ ) and CT + TT ( $n_{\text{Preserved}} = 36$  vs  $n_{\text{Lesioned}} = 27$ ). [B] Variance stabilizing transformation (VST) normalized *MGP* expression levels extracted from the RNA sequencing dataset in preserved and lesioned OA subchondral bone stratified for rs1800801 genotype CC ( $n_{\text{Preserved}} = 5$  vs  $n_{\text{Lesioned}} = 5$ ) and CT + TT ( $n_{\text{Preserved}} = 9$  vs  $n_{\text{Lesioned}} = 9$ ). The boxplots represent the median with the 25th, 50th and 75th percentiles, and whiskers reflect the total range. Independent samples are depicted by black dots in each graph. To adjust for donor variation, *P*-values were estimated by performing logistic generalized estimation equations, with tissue status as dependent variable and *MGP* level, age and sex as covariate:  $Tissue\ status \sim MGP\ level + age + sex + (1|Donor)$ . \* $P \leq 0.05$ .

Osteoarthritis  
and Cartilage

allele rs4236-T. In lesioned OA subchondral bone AEI was very comparable, with rs4236-C associated to a reduced *MGP* expression of 12% (95%CI [8.80; 14.55]) relative to the reference allele rs4236-T.


Finally, we analysed *MGP* expression levels among genotype carriers of one or two of the OA risk alleles rs1800801-T (eQTL) and confirmed that also overall *MGP* expression in cartilage is reduced in a dose responsive manner with *MGP* risk alleles,

independent of donor, age, sex, and OA status, i.e., preserved or lesioned (OR = 0.73, 95%CI [0.64; 0.84], *P*-value =  $4.00 \times 10^{-6}$ ; Supplementary Fig. 3(A)). In subchondral bone we observed a similar pattern, however this was not significant (Supplementary Fig. 3(B)). Together these data confirm that innate lower *MGP* expression levels confer risk to OA, though its effect seems more pronounced in articular cartilage.

### MGP expression patterns in human *in vitro* and *ex vivo* models and as function of OA related cues

First, we investigated expression of *MGP* during neo-cartilage formation using a human *in vitro* 3D pellet culture with primary chondrocytes. As shown in Fig. 3, *MGP* is expressed in primary chondrocytes (day-0) and increases during cartilage extracellular matrix (ECM) deposition until day-14, suggesting that *MGP* expression can be considered a marker of neo-cartilage formation.

Next, we explored dynamic changes in *MGP* expression in cartilage and subchondral bone in an established human *ex vivo* osteochondral explant model<sup>20</sup> as function of OA related stimuli being inflammation (IL-1 $\beta$ ), hypertrophy (T3), and 65% mechanical stress (65%MS). As shown in Fig. 4, we observed in cartilage a consistent and significant downregulation of *MGP* expression after treatment with IL-1 $\beta$  (FC = 0.03, 95%CI [0.02; 0.06], *P*-value =  $4.40 \times 10^{-7}$ ), T3 (FC = 0.80, 95%CI [0.56; 0.97], *P*-value = 0.046), as well as with mechanical stress (FC = 0.65, 95%CI [0.45; 0.85], *P* = 0.002). Notable, in Fig. 4, is an outlier in the mechanical stress group, however removing this datapoint did not influence our result (FC = 0.67, 95%CI [0.47; 0.87], *P*-value = 0.046). In subchondral bone we were not able to isolate RNA for IL-1 $\beta$  treated samples and only observed a significant downregulation of *MGP* expression after treatment with T3 (FC = 0.81, 95%CI [0.52; 1.10], *P*-value = 0.015).



**Fig. 3**

**Gene expression of *MGP* in an *in vitro* 3D model of neo-cartilage.** Gene expression of *MGP* in an *in vitro* 3D chondrocyte pellet model of neo-cartilage formation (*N* = 3 donors; day 0: *n* = 2, all other time points: *n* = 3). Data is depicted as mean expression ( $-\Delta\text{CT}$ )  $\pm$  standard error of the mean (SEM) and each dot represents a sample of two combined biological duplicates. Statistical analysis was performed by generalized linear mixed model (GLMM) using *MGP* levels as dependent variable and time as repeated measure:  $MGP \text{ level} \sim \text{Time} + (1|\text{Donor})$ .

Osteoarthritis  
and Cartilage

### Changes in *MGP* expression in the *ex vivo* OA models as function of the transcript and OA risk SNP rs1800801

Since general *MGP* expression was identified to change between preserved and lesioned OA cartilage and subchondral bone, and in a osteochondral explant model to several OA related stimuli (Fig. 4), we next explored whether the OA risk allele rs1800801-T modified these effects. Hereto we investigated the observed dynamic downregulation of *MGP*, upon inducing hypertrophy (T3 exposure; Fig. 5(A) and (C)) and mechanical stress (65%MS; Fig. 5(B) and (D)) in our *ex vivo* cartilage explant model stratified by rs1800801 genotypes. For IL-1 $\beta$  treatment, donor numbers were too low to explore the effect of genotype.

In cartilage (Fig. 5(A) and (B)), we observed that downregulation of *MGP* occurred particularly among carriers of the reference allele rs1800801-C for hypertrophy (T3) (FC = 0.69, 95%CI [0.49; 0.89]) and for mechanical stress (FC = 0.26, 95%CI [0.14; 0.38]) as compared to carriers of the risk allele rs1800801-T for hypertrophy (FC = 0.92, 95%CI [0.57; 1.27]) and for mechanical stress (FC = 0.85, 95%CI [0.61; 1.09]). Also in the data shown in Fig. 5(B) is the previously identified outlier in the mechanical stress group which did not influence our result upon removal (FC = 0.29, 95%CI [0.17–0.41], *P*-value = 0.045). As shown in Supplementary Table 4, the difference in response (FC) among carriers of the reference allele rs1800801-C relative to carriers of the OA risk allele rs1800801-T is significant for mechanical stress (FC = 0.34, 95% CI [0.28–0.38], *P*-value =  $2.8 \times 10^{-3}$ ). Similarly in bone [Fig. 5(C)–(D)], we observed that downregulation of *MGP* expression in subchondral bone upon hypertrophy induction (T3) was more pronounced among carriers of the reference allele rs1800801-C (FC = 0.50, 95%CI [0.09; 0.91] as compared to the carriers of the risk allele rs1800801-T (FC = 0.82, 95%CI [0.37; 1.27]). This difference, however, did not reach statistical significance (Supplementary Table 4). For mechanical stress no effects were observed in subchondral bone. Together these data suggest that particularly in cartilage the OA risk allele rs1800801-T may have a different response in *MGP* expression upon OA relevant cues.

### Treatment of osteochondral explants with warfarin

Since the activation of *MGP* is dependent on vitamin K and innate lower *MGP* expression confers risk to OA, we next investigated the direct effect of the vitamin K antagonist warfarin on articular chondrocyte and subchondral bone signalling. Hereto, *ex vivo* osteochondral explants (*n* = 15 pairs for cartilage and *n* = 13 pairs for subchondral bone) were treated with warfarin. The effect of this reduced vitamin K bioavailability on the cartilage homeostasis was determined by measuring chondroprotective genes (*SOX9*, *COL2A1* and *ACAN*), genes involved in early/late cartilage hypertrophy (*RUNX2*, *ALPL*, *COL1A1*, *COL10A1* and *MGP*) and catabolic genes (*MMP3*, *MMP13* and *ADAMTS5*). As shown in Fig. 6(A), warfarin exposure to cartilage reduced expression of *SOX9* (FC = 0.87, 95%CI [0.77; 0.97], *P*-value = 0.023) and *MMP3* (FC = 0.56, 95%CI [0.43; 0.69], *P*-value =  $1.02 \times 10^{-5}$ ), while increasing *COL10A1* (FC = 2.26, 95%CI [1.14; 3.38], *P*-value = 0.045). In addition, *RUNX2* (FC = 1.43, 95%CI [0.96; 1.90], *P*-value = 0.094), a master transcriptional regulator of chondrocyte maturation, and *ALPL* (FC = 6.21, 95%CI [1.36; 11.06], *P*-value = 0.059) show a trend towards upregulation in response to warfarin treatment. In subchondral bone, genes involved in matrix formation (*COL10A1*, *RUNX2*, *ALPL*, *COL1A1*, *OMD*, *BGLAP* and *OGN*) and remodelling (*MGP*, *SPP1*, *TNFRSF11B*, *BMP2* and *POSTN*) were measured. As shown in Fig. 6(B), warfarin exposure to subchondral bone significantly reduced expression of the bone formation marker *COL1A1* (FC = 0.81, 95%CI [0.59; 1.03], *P*-value = 0.046) and




Fig. 4

**Gene expression of *MGP* in response to three different OA relevant cues in cartilage of osteochondral explants.** Gene expression of *MGP* ( $-\Delta\text{CT}$ ) in an *ex vivo* osteochondral explant model in articular cartilage (**A**, **B**, **C**) and subchondral bone (**D**, **E**). *MGP* expression, represented by the housekeeping gene corrected value ( $-\Delta\text{CT}$ ), in articular cartilage upon perturbation with **[A]** IL-1 $\beta$  ( $n_{\text{Control}} = 6$  vs  $n_{\text{Treated}} = 6$ ) **[B]** T3 ( $n_{\text{Control}} = 21$  vs  $n_{\text{Treated}} = 21$ ) and **[C]** posttraumatic OA after 65% MS ( $n_{\text{Control}} = 30$  vs  $n_{\text{Treated}} = 23$ ). *MGP* expression in subchondral bone upon perturbation with **[D]** T3 ( $n_{\text{Control}} = 16$  vs  $n_{\text{Treated}} = 16$ ) and **[E]** posttraumatic OA after 65% MS ( $n_{\text{Control}} = 23$  vs  $n_{\text{Treated}} = 19$ ). The boxplots represent the median with 25th, 50th and 75th percentiles, and whiskers reflect the total range.  $-\Delta\text{CT}$  of each independent sample is depicted by black dots in the graphs. To adjust for donor variation *P*-values were determined by performing linear generalized estimation equations, with *MGP* levels as dependent variable and treatment as factor:  $MGP \text{ level} \sim \text{Treatment} + (1/\text{Donor})$ . Far out values are represented by the white filled circle (o) which did not affect the result (see main body text) and therefore analysis including this sample is presented in **C**. \* $P \leq 0.05$ , \*\* $P \leq 0.01$ , \*\*\* $P \leq 0.001$ . Legend: Ctrl=Control; 65% MS=65% mechanical stress; T3=triiodothyronine.

Osteoarthritis  
and Cartilage

the remodelling marker *POSTN* (FC = 0.67, 95%CI [0.42; 0.92], *P*-value = 0.011). Together these results show that addition of warfarin to aged osteochondral explants resulted in a significant upregulation of hypertrophic signalling among articular chondrocytes and reduced bone formation and altered remodelling signalling.

## Discussion

In the current paper we explored (dynamic) changes of *MGP* expression in relation to the OA risk allele rs1800801-T, in preserved and lesioned OA cartilage, as well as in a human *ex vivo* explant model subjected to OA related stimuli, such as inflammation, hypertrophy and mechanical stress. Furthermore, we studied the direct effect of the frequently used vitamin K antagonist, warfarin, on articular chondrocyte and subchondral bone

signalling. In doing so, we confirm that *MGP* expression, as inhibitor of calcification via high affinity of calcium to its Gla-residues, should be considered a beneficial marker of articular cartilage. Consequently, the significantly upregulated *MGP* expression with ongoing OA pathophysiology is likely an attempt of chondrocytes to halt the OA associated osteo-induction. Noteworthy is our observation that the OA risk allele may also hamper adequate dynamic change in expression of *MGP* in response to OA and relevant cues like mechanical stress (65%MS) and this effect was most pronounced in cartilage. Finally, warfarin treatment to the aged human cartilage explants resulted in a significant upregulation of hypertrophic signalling among articular chondrocytes and reduced bone formation while altering remodelling.

Similar to previous reports<sup>7,9</sup> we here confirmed in a large RNA-sequencing dataset, that the OA risk allele rs1800801-T is associated with lower (overall) expression of *MGP* in articular cartilage




Fig. 5

**MGP expression as function of the transcript and OA risk SNP rs1800801.** MGP expression ( $-\Delta CT$ ) in an *ex vivo* osteochondral explant model stratified by rs1800801 genotype in articular cartilage (A, B) and subchondral bone (C, D). MGP gene expression in articular cartilage upon perturbation with [A] T3 (CC:  $n_{\text{Control}} = 10$  vs  $n_{\text{Treated}} = 8$ ; CT:  $n_{\text{Control}} = 11$  vs  $n_{\text{Treated}} = 13$ ) and [B] posttraumatic OA upon 65% mechanical stress (CC:  $n_{\text{Control}} = 11$  vs  $n_{\text{Treated}} = 7$ ; CT + TT:  $n_{\text{Control}} = 19$  vs  $n_{\text{Treated}} = 16$ ). MGP gene expression in subchondral bone upon perturbation with [C] T3 (CC:  $n_{\text{Control}} = 8$  vs  $n_{\text{Treated}} = 9$ ; CT:  $n_{\text{Control}} = 8$  vs  $n_{\text{Treated}} = 7$ ) and [D] posttraumatic OA upon 65% mechanical stress (CC:  $n_{\text{Control}} = 10$  vs  $n_{\text{Treated}} = 7$ ; CT + TT:  $n_{\text{Control}} = 13$  vs  $n_{\text{Treated}} = 12$ ). The boxplots represent the median and 25th, 50th and 75th percentiles, and whiskers reflects the total range. Independent samples are depicted by black dots in each graph. Numeric values associated with this Figure are shown in Supplementary Table 4. To adjust for donor variation *P*-values were determined by performing linear generalized estimation equations, with MGP levels as dependent variable and treatment as factor:  $MGP \text{ level} \sim \text{Treatment} + (1|Donor)$ . Far out values are represent by the white filled circle (o) which did not affect the result (see main body text) and therefore analysis including this sample is presented in B. \* $P \leq 0.05$ . Legend: 65% MS=Mechanical stress; T3=triiodothyronine.

Osteoarthritis  
and Cartilage

(Supplementary Fig. 3(A)). In addition, we confirmed that MGP gene expression is significantly upregulated in both articular cartilage and subchondral bone in OA pathophysiology. Although, our results showed that this effect was mainly driven by carriers of the rs1800801-T OA risk allele in both tissues, the expression of MGP does not reach the level of that in carriers of the reference allele rs1800801-C [Fig. 2(A) and (B)]. We advocate that MGP upregulation with OA pathophysiology in cartilage is an attempt of chondrocytes to compensate for the osteo-inductive effect of low MGP levels and that this is not sufficient among the MGP OA risk allele carriers. On the other hand, the upregulation of MGP in bone may be a marker of active bone resorption, as it was previously found that MGP inhibits mineralization by osteoblasts while increased MGP expression in osteoclasts mark increased osteoclastic commitment<sup>25</sup>. Together our data highlights that, similar to vascular calcification and bone loss<sup>26</sup>, also articular cartilage calcification and bone loss in OA could share a common pathogenetic mechanism involving MGP.

We also explored the dynamic response of MGP in a human *ex vivo* explant model while applying OA relevant perturbing cues

such as inflammation, hypertrophy, and mechanical stress. The strength of our explant model is that it represents physiological relevant aged human articular cartilage prone to OA pathophysiology, hence suitable to study the initial process of OA related cartilage destruction. Moreover, and despite the inherent heterogeneity between donors, we found in cartilage a consistent downregulation of MGP, associated with matrix mineralization, as general response to OA related perturbations (Fig. 4). Additionally, we showed that the rs1800801-T OA risk allele may hamper such innate dynamic change in MGP expression upon stress. A possible mechanism by which the genetic risk variant modifies response to stress lies in the fact that rs1800801 is localized in the transcription factor binding site (POLR2A, CTCF, p300) of the MGP promoter (Supplementary Fig. 4). In addition, the OA risk allele rs1800801-T was shown to reduce expression between 34% and 47% in a luciferase reporter assay and *in silico* prediction suggested this to be due to a loss of binding site for the transcription factor c-Ets<sup>12</sup>. In the subchondral bone compartment of the human *ex vivo* explants, the MGP response to the OA like perturbing cues were smaller and less consistent although a similar MGP response appeared for T3






Fig. 6

**Gene expression after 10 days of warfarin treatment.** mRNA expression of genes depicted as a fold change (FC) following 50  $\mu\text{M}$  warfarin treatment relative to controls in **[A]** cartilage ( $n_{\text{Control}} = 15$  vs  $n_{\text{Treated}} = 15$ ) and **[B]** subchondral bone ( $n_{\text{Control}} = 13$  vs  $n_{\text{Treated}} = 13$ ). Controls are depicted by the dotted line, while each gray dot represents a warfarin treated sample. The number and percentage of missing data points per gene are summarized in [Supplementary Tables 2 and 3](#). The light gray bars represent the mean  $\pm$  standard error of the mean (SEM) of the Fold change (FC) in which  $\text{FC} > 1$  represent upregulation and  $\text{FC} < 1$  represents downregulation of warfarin treated samples relative to its paired control. The x-axis is given in a  $\log_2$  scale to depict the up and down regulation in the same scale. Differences in gene levels between warfarin exposure and controls were calculated by means of a paired *t*-test. \* $P \leq 0.05$ ; \*\*\* $P \leq 0.001$ .

Osteoarthritis  
and Cartilage

exposure. This is likely the result of (slightly) lower sample sizes but, more importantly, a more complex innate regulation and signalling of *MGP* in bone as multicellular tissue type. As such, the observed variation in the *MGP* response in bone remains inconclusive and needs to be repeated in larger sample sizes.

Upon identifying *MGP*, encoding an inhibitor of ectopic calcifications, as strong OA risk gene, it was hypothesized that the OA risk was conferred via calcification of cartilage tissue<sup>10,11</sup>. Moreover, as *MGP* protein is activated by vitamin K dependent carboxylation (c-*MGP*) this finding underscored the relevance of previously found associations between OA and low vitamin K status<sup>14,15</sup>. Here, we showed that exposure of the vitamin K inhibitor warfarin to intact human articular cartilage explants provoked unbeneficial functional chondrocyte signalling towards hypertrophy, as reflected by upregulation of *COL10A1* and almost significant upregulation of *RUNX2* and *ALPL*. Moreover, we showed a modest but significant downregulation of *SOX9*, a transcription factor marking healthy articular cartilage. These observed effects of warfarin on chondrocyte signalling were similar to those previously found during *in vitro* knockdown of *MGP* in chondrocyte monolayer cultures<sup>9</sup>. With regard to the seemingly increased *MMP13* and reduced *MMP3* gene expression, it has been suggested that *MMP3* plays a role mainly in healthy cartilage remodeling, while *MMP13* more so in pathological processes. This was confirmed by performing a lookup in our RNA-sequencing data set<sup>18</sup> where *MMP3* showed a marked downregulation in lesioned compared to preserved OA cartilage. Exposure of warfarin to subchondral bone of osteochondral explants also provoked unbeneficial functional signalling towards reduced bone formation, as reflected by downregulation of *COL1A1* and the suggestive downregulation of *BGLAP*, whereas the upregulation of the osteoclastogenesis inhibitor *TNFRSF11B* (although not significant) and downregulation of the vitamin K dependent protein *POSTN* suggests altered bone remodeling likely resulting in bone loss<sup>25,27</sup>. Due to the low numbers and heterogeneity of patients, future studies are necessary to investigate if rs1800801 genotype influences response of cells to warfarin. In light of our result we advocate that the frequent prescription of warfarin as vitamin K

dependent blood anticoagulant<sup>16</sup> may have clinical consequences in evoking OA comorbidity. As such, the risk of OA comorbidity may be considerably reduced by preferred prescription of non-vitamin K antagonist as anticoagulants<sup>28</sup>. In addition, vitamin K supplementation should be considered a potential novel OA-modifying treatment option. In this respect, there has been one underpowered clinical trial studying the effect of vitamin K supplementation on OA progression. This ancillary study, originally designed to study vascular calcification, reported no overall beneficial effects of vitamin K supplementation. However, in individuals with insufficient vitamin K levels at baseline a beneficial effect was observed<sup>29</sup>.

Although the human aged macroscopically normal osteochondral explants used in our study may represent physiological relevant human articular cartilage and subchondral bone model, prone to OA pathophysiology, hence suitable to study the initial process of OA related destruction, the model is inherently subject to heterogeneity. Moreover, it does not provide insight into the *MGP* effect of such environmental perturbations to healthy cartilage and bone. Another limiting factor was the low sample size of T3, IL-1 $\beta$  and warfarin treated explants upon stratifying for rs1800801 genotype, resulting in no or less robust results than upon investigating the response in the larger mechanically stressed group. It should also be noted that the modifying effect of the *MGP* OA risk allele rs1800801-T as function of the OA status in articular cartilage and subchondral bone was only measured as a static effect i.e., differential expression of *MGP* between paired preserved and lesioned OA cartilage samples. Finally, the focus of our paper was on exploring gene expression changes of *MGP* only. Although studying protein levels of *MGP* as function of the OA risk SNP and the OA relevant cues in joint tissue would be an interesting addition and a preferred next step, such analyses should involve the detection of activated (hence carboxylated) *MGP* protein.

Together our data highlight that, similar to the bi-directional interplay of vascular calcification and bone loss in osteoporosis and atherosclerosis<sup>26</sup>, also articular cartilage calcification and bone loss in OA might share a common pathogenetic mechanism likely involving *MGP*. Moreover, warfarin on human osteochondral

explant cultures functionally underscores the previously found association between vitamin K deficiency and OA.

### Contributions

All authors have made contributions to the completion of this study. Study concept and design: EH, RCA, JM, YFM, IM. Acquisition of material and data: EH, MT, HED, DB, RGHHN. Data analysis: EH, RCA, MT, IM. Preparation of the manuscript: EH, IM. Critical reviewing and approval of the manuscript: All authors.

### Conflicts of interest

None declared.

### Funding source

The study was funded by the Dutch Scientific Research council NWO/ZonMW VICI scheme (nr 91816631/528), Dutch Arthritis Society/ReumaNederland (DAF-15-4-401 and DAA\_10\_1-402) and European Commission Seventh Framework program (TreatOA, 200800).

### Patient consent for publication

Not required.

### Data availability statement

Data are available on reasonable request.

### Acknowledgements

We thank all the participants of the RAAK study. The LUMC was and is supporting the RAAK study. We also thank Enrike van der Linden, Robert van der Wal, Peter van Schie, Anika Rabelink-Hoogenstraaten, Shaho Hasan, Maartje Meijer, Daisy Latijnhouwers and Geert Spierenburg for collecting surgical waste material. Data is generated within the scope of the Medical Delta Programs Regenerative Medicine 4D: Generating complex tissues with stem cells and printing technology and Improving Mobility with Technology. We thank the Sequence Analysis Support Core (SASC) of the Leiden University Medical Center for their support.

### Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.joca.2021.05.001>.

### References

1. Spector TD, Cicuttini F, Baker J, Loughlin J, Hart D. Genetic influences on osteoarthritis in women: a twin study. *BMJ* 1996;312(7036):940–3.
2. Felson DT. Risk factors for osteoarthritis: understanding joint vulnerability. *Clin Orthop Relat Res* 2004;427(Suppl 1):S16–21.
3. Bos SD, Bovee JV, Duijnisveld BJ, Raine EV, van Dalen WJ, Ramos YF, et al. Increased type II deiodinase protein in OA-affected cartilage and allelic imbalance of OA risk polymorphism rs225014 at DIO2 in human OA joint tissues. *Ann Rheum Dis* 2012;71(7):1254–8.
4. Reynard LN, Bui C, Canty-Laird EG, Young DA, Loughlin J. Expression of the osteoarthritis-associated gene GDF5 is modulated epigenetically by DNA methylation. *Hum Mol Genet* 2011;20(17):3450–60.
5. Loughlin J. Genetic contribution to osteoarthritis development: current state of evidence. *Curr Opin Rheumatol* 2015;27(3):284–8.
6. Shepherd C, Zhu D, Skelton AJ, Combe J, Threadgold H, Zhu L, et al. Functional characterisation of the osteoarthritis genetic risk residing at ALDH1A2 identifies rs12915901 as a key target variant. *Arthritis Rheum* 2018;70(10):1577–87.
7. den Hollander W, Boer CG, Hart DJ, Yau MS, Ramos YFM, Metrustry S, et al. Genome-wide association and functional studies identify a role for matrix Gla protein in osteoarthritis of the hand. *Ann Rheum Dis* 2017;76(12):2046–53.
8. Misra D, Booth SL, Crosier MD, Ordovas JM, Felson DT, Neogi T. Matrix Gla protein polymorphism, but not concentrations, is associated with radiographic hand osteoarthritis. *J Rheumatol* 2011;38(9):1960–5.
9. Shepherd C, Reese AE, Reynard LN, Loughlin J. Expression analysis of the osteoarthritis genetic susceptibility mapping to the matrix Gla protein gene MGP. *Arthritis Res Ther* 2019;21(1):149.
10. Luo G, Ducey P, McKee MD, Pinero GJ, Loyer E, Behringer RR, et al. Spontaneous calcification of arteries and cartilage in mice lacking matrix GLA protein. *Nature* 1997;386(6620):78–81.
11. Yagami K, Suh JY, Enomoto-Iwamoto M, Koyama E, Abrams WR, Shapiro IM, et al. Matrix GLA protein is a developmental regulator of chondrocyte mineralization and, when constitutively expressed, blocks endochondral and intramembranous ossification in the limb. *J Cell Biol* 1999;147(5):1097–108.
12. Tunon-Le Poultel D, Cannata-Andia JB, Roman-Garcia P, Diaz-Lopez JB, Coto E, Gomez C, et al. Association of matrix Gla protein gene functional polymorphisms with loss of bone mineral density and progression of aortic calcification. *Osteoporos Int* 2014;25(4):1237–46.
13. Sheng K, Zhang P, Lin W, Cheng J, Li J, Chen J. Association of Matrix Gla protein gene (rs1800801, rs1800802, rs4236) polymorphism with vascular calcification and atherosclerotic disease: a meta-analysis. *Sci Rep* 2017;7(1):8713.
14. Neogi T, Booth SL, Zhang YQ, Jacques PF, Terkeltaub R, Aliabadi P, et al. Low vitamin K status is associated with osteoarthritis in the hand and knee. *Arthritis Rheum* 2006;54(4):1255–61.
15. Shea MK, Kritchevsky SB, Hsu FC, Nevitt M, Booth SL, Kwok CK, et al. The association between vitamin K status and knee osteoarthritis features in older adults: the Health, Aging and Body Composition Study. *Osteoarthritis Cartilage* 2015;23(3):370–8.
16. Holbrook A, Schulman S, Witt DM, Vandvik PO, Fish J, Kovacs MJ, et al. Evidence-based management of anticoagulant therapy: antithrombotic therapy and prevention of thrombosis, 9th ed: American college of chest physicians evidence-based clinical practice guidelines. *Chest* 2012;141(2 Suppl 1):e152S–84S.
17. Chin KY. The relationship between vitamin K and osteoarthritis: a review of current evidence. *Nutrients* 2020;12(5).
18. Coutinho de Almeida R, Ramos YFM, Mahfouz A, den Hollander W, Lakenberg N, Houtman E, et al. RNA sequencing data integration reveals a miRNA interactome of osteoarthritis cartilage. *Ann Rheum Dis* 2019;78(2):270–7.
19. Tuerlings M, van Hoolwerff M, Houtman E, Suchiman E, Lakenberg N, Mei H, et al. RNA sequencing reveals interacting key determinants of osteoarthritis acting in subchondral bone and articular cartilage. *Arthritis Rheum* 2021;73(5):789–99.
20. Houtman E, van Hoolwerff M, Lakenberg N, Suchiman EHD, van der Linden-van der Zwaag E, Nelissen R, et al. Human osteochondral explants: reliable biomimetic models to investigate disease mechanisms and develop personalized treatments for osteoarthritis. *Rheumatol Ther* 2021;20(8):499–515.

21. Ramos YF, den Hollander W, Bovee JV, Bomer N, van der Breggen R, Lakenberg N, *et al.* Genes involved in the osteoarthritis process identified through genome wide expression analysis in articular cartilage; the RAAK study. *PLoS One* 2014;9(7), e103056.
22. den Hollander W, Pulyakhina I, Boer C, Bomer N, van der Breggen R, Arindrarto W, *et al.* Annotating transcriptional effects of genetic variants in disease-relevant tissue: transcriptome-wide allelic imbalance in osteoarthritic cartilage. *Arthritis Rheum* 2019;71(4):561–70.
23. Zeger SL, Liang KY. Longitudinal data analysis for discrete and continuous outcomes. *Biometrics* 1986;42(1):121–30.
24. Diggle P, Liang K-Y, Zeger SL. In: *Analysis of Longitudinal Data*, xi. Oxford New York: Clarendon Press; Oxford University Press; 1994:253.
25. Zhang Y, Zhao L, Wang N, Li J, He F, Li X, *et al.* Unexpected role of matrix Gla protein in osteoclasts: inhibiting osteoclast differentiation and bone resorption. *Mol Cell Biol* 2019;39(12):e00012–9.
26. Vassalle C, Mazzone A. Bone loss and vascular calcification: a bi-directional interplay? *Vasc Pharmacol* 2016;86:77–86.
27. Bonnet N, Gineyts E, Ammann P, Conway SJ, Garnero P, Ferrari S. Periostin deficiency increases bone damage and impairs injury response to fatigue loading in adult mice. *PLoS One* 2013;8(10), e78347.
28. Zhu J, Alexander GC, Nazarian S, Segal JB, Wu AW. Trends and variation in oral anticoagulant choice in patients with atrial fibrillation, 2010–2017. *Pharmacotherapy* 2018;38(9):907–20.
29. Neogi T, Felson DT, Sarno R, Booth SL. Vitamin K in hand osteoarthritis: results from a randomised clinical trial. *Ann Rheum Dis* 2008;67(11):1570–3.