



Universiteit
Leiden
The Netherlands

Within-person variation in serum thyrotropin concentration: main sources, potential underlying biological mechanisms, and clinical implications

Spoel, E. van der; Roelfsema, F.; Heemst, D. van

Citation

Spoel, E. van der, Roelfsema, F., & Heemst, D. van. (2021). Within-person variation in serum thyrotropin concentration: main sources, potential underlying biological mechanisms, and clinical implications. *Frontiers In Endocrinology*, 12.
doi:10.3389/fendo.2021.619568

Version: Publisher's Version
License: [Creative Commons CC BY 4.0 license](#)
Downloaded from: <https://hdl.handle.net/1887/3235983>

Note: To cite this publication please use the final published version (if applicable).



Within-Person Variation in Serum Thyrotropin Concentrations: Main Sources, Potential Underlying Biological Mechanisms, and Clinical Implications

Evie van der Spoel^{1*}, Ferdinand Roelfsema² and Diana van Heemst¹

¹ Section Gerontology and Geriatrics, Department of Internal Medicine, Leiden University Medical Center, Leiden, Netherlands, ² Section Endocrinology, Department of Internal Medicine, Leiden University Medical Center, Leiden, Netherlands

OPEN ACCESS

Edited by:

Douglas Forrest,
National Institutes of Health (NIH),
United States

Reviewed by:

Masami Murakami,
Gunma University, Japan
H. I. Marris,
Marrif Biotech, Canada
Michihiko Aramaki,
National Institutes of Health (NIH),
United States

*Correspondence:

Evie van der Spoel
e.van_der_spoel@lumc.nl

Specialty section:

This article was submitted to
Thyroid Endocrinology,
a section of the journal
Frontiers in Endocrinology

Received: 20 October 2020

Accepted: 08 January 2021

Published: 24 February 2021

Citation:

van der Spoel E, Roelfsema F and van Heemst D (2021) Within-Person Variation in Serum Thyrotropin Concentrations: Main Sources, Potential Underlying Biological Mechanisms, and Clinical Implications. *Front. Endocrinol.* 12:619568. doi: 10.3389/fendo.2021.619568

Background: Individuals exhibit fluctuations in the concentration of serum thyroid-stimulating hormone (TSH) over time. The scale of these variations ranges from minutes to hours, and from months to years. The main factors contributing to the observed within-person fluctuations in serum TSH comprise pulsatile secretion, circadian rhythm, seasonality, and ageing. In clinical practice and clinical research however, such within-person biological variation in serum TSH concentrations is often not considered. The aim of this review is to present an overview of the main sources of within-person variation in TSH levels, as well as the potential underlying biological mechanisms, and the clinical implications.

Summary: In euthyroid individuals, the circadian rhythm, with a nocturnal surge around 02:00–04:00 h and a nadir during daytime has the greatest impact on variations in serum TSH concentrations. Another source of within-person variation in TSH levels is seasonality, with generally higher levels during the cold winter months. Since TSH is secreted in a pulsatile manner, TSH levels also fluctuate over minutes. Furthermore, elevated TSH levels have been observed with ageing. Other factors that affect TSH levels include thyroid peroxidase (TPO)-antibody positivity, BMI, obesity, smoking, critical illness, and many xenobiotics, including environmental pollutants and drugs. Potential underlying biological mechanisms of within-person variation in TSH levels can be safely concluded from the ability of TSH to respond quickly to changes in cues from the internal or external environment in order to maintain homeostasis. Such cues include the biological clock, environmental temperature, and length of day. The observed increase in TSH level with ageing can be explained at a population level and at an organism level. In clinical practice, the season for thyroid testing can influence a patient's test result and it occurs frequently that subclinical hypothyroid patients normalize to euthyroid levels over time without intervention.

Conclusions: Serum TSH concentrations vary over time within an individual, which is caused by multiple different internal and external factors. It is important to take the within-person variations in serum TSH concentrations into account when testing a patient in clinical practice, but also in performing clinical research.

Keywords: thyrotropin, thyroid-stimulating hormone, biological variation, circadian rhythm, ageing, seasonality, within-person variation

INTRODUCTION

Hormones of the hypothalamus-pituitary-thyroid (HPT) axis are part of a feedforward and feedback system. Thyroid stimulating hormone (TSH) is a key player in this system, which is designed to respond quickly to changes in the environment aiming to maintain homeostasis in the human body. By adapting the level of TSH, circulating levels of thyroxine (T₄) and partly triiodothyronine (T₃) are controlled and kept within a normal range (1, 2). Indeed, it is known that TSH levels can vary over time within a person.

Among other, TSH levels are influenced by drugs, acute and chronic illness, undernutrition, the biological clock, seasonality, pregnancy, and by other hormones, e.g., cortisol. However, the magnitude and the importance of variation in TSH levels within a person over time are not completely clear yet. Consequently, the influence of variation in TSH levels on diagnosis and study outcomes might be underestimated. In this review, we discuss the magnitude of the variation in TSH levels and the potential sources (internal and external) that may contribute.

Variation in TSH levels can be caused by biological and non-biological variation, the latter includes pre-analytical and analytical variations, and has been reviewed by others (3). In the current review we focus on biological variation, in particular on within-person biological variation. Within-person variation in TSH levels is caused by rhythms ranging from minutes to years, including pulsatile secretion, circadian rhythm, monthly changes, and seasonality (3, 4). Furthermore, TSH levels change with age, with in general higher levels with increasing age (5). Independent of time, within-person variation in TSH is also caused by effects of among others medication, illness, TPO antibody positivity, and iodine intake.

Medical practice and guidelines have been traditionally based on the “average” patient and how the “average” patient responds to various treatments, as informed by randomized clinical trials. Especially vulnerable patients, which include older patients with comorbidities and polypharmacy, are underrepresented in clinical trials. Consequently, evidence on how such vulnerable patients respond to treatment is lacking (6). As it is becoming increasingly clear that patients are a heterogeneous group, in current practice and guidelines, there is increasing recognition of the need to individualize management of a patient’s disease. As a promising strategy toward personalized medicine, N-of-1 studies have been proposed as a design to determine the best treatment for individual patients (7). In a N-of-1 trial, each individual receives several treatments in a multiple crossover design, allowing determination of the best intervention for an

individual patient. However, a drawback of this type of study is that different points in time are compared with one another, and thus results may be influenced by fluctuations that occur within an individual over time (8).

In this context, a lack of awareness among clinicians of variation in TSH levels within individual patients over time could lead to under- or overdiagnosis of (subclinical) thyroid diseases, or to inadequate prescription of thyroid medication to patients with thyroid disease. Furthermore, if the variation in TSH levels is not considered, this could lead to a distorted interpretation of outcomes and results in scientific research. Therefore, in this review we give an overview of the main sources of within-person variation in euthyroid individuals to provide insight into TSH variation within the normal range. Additionally, available research on TSH variation in patients with thyroid disease and/or using thyroid medication will be discussed. Moreover, the potential underlying biological mechanisms and the clinical implications of within-person variation in TSH levels are reviewed. Much is known about the short-term within-person variation in TSH levels. However, long-term variations in TSH levels are rarely investigated in longitudinal studies and more often in cross-sectional studies. In this review, evidence from both cross-sectional and longitudinal studies is described. Literature research was mainly performed using PubMed in the first 8 months of 2020.

SOURCES OF WITHIN-PERSON VARIATION IN TSH LEVELS

Pulsatile Secretion

TSH is secreted in a basal (non-pulsatile) and in a pulsatile manner by the anterior pituitary gland (9, 10). The feedback control by mainly T₃ and T₄ from the thyroid gland, together with the feedforward control by mainly thyrotropin-releasing hormone (TRH) from the hypothalamus, tightly regulate TSH secretion. It is also important to note that other hormones, including leptin, somatostatin, and dopamine, are known to influence the secretion of TSH, albeit to a lesser extent (11–15). The frequency, size, and duration of a secretory burst (i.e., pulse) of TSH is determined by the interplay between these feedback and feedforward signals.

Studies on Pulsatile Secretion

In 1978, Weeke et al. found short-term variations with a mean cycle-length of 31 min in serum TSH concentrations measured

in blood withdrawn every 5 min starting in the evening for 6 to 7 h in five healthy men (4). Using spectral analysis together with periodic regression analysis in a study with frequent blood sampling, pulsatile secretion of TSH was detected with periodicities of 60 or 85–100 min in six out of the ten healthy volunteers (10). The mean pulse frequency of TSH levels was found to be 13 pulses (range 10–18) per 24 h in blood withdrawn every 10 min in six healthy subjects (9).

Deconvolution Analysis to Detect Pulsatile Secretion

Several techniques can be used to determine pulsatility in TSH serum concentrations, but deconvolution analysis is most commonly used in 24-h time series data. Keenan et al. developed a deconvolution analysis algorithm by which a 24-h hormone concentration profile is decomposed into underlying secretory bursts, basal secretion, elimination of previously secreted hormone, and random experimental variability (16–18). Therefore, deconvolution analysis can be used to assess the frequency, size, and duration of a hormone pulse. The algorithm in the software program MATLAB (the MathWorks, Inc., Natick, MA) first detrends the data and normalizes concentrations by converting them to values between 0 and 1 (18). This step is performed to normalize the span of diverse values in separate datasets so that all have a comparable chance of detecting pulses, despite variable baselines. Second, successive potential pulse-time sets, each containing one fewer burst, are created by a smoothing process. Finally, a maximum-likelihood expectation deconvolution method estimated all secretion and elimination rates simultaneously for each candidate pulse-time set. Deconvolution analysis determines secretory bursts and thereby calculates which part of TSH was secreted in a pulsatile manner within a 24 h time period. The “remaining part” of the 24-h TSH concentration that cannot be explained by secretory bursts, is called non-pulsatile or basal. It is therefore assumed in this model that the basal secretion is constant during the day. However, it is not known whether there is a constant tonic release of TSH from the pituitary or that it might be variable over time like insulin, which was studied by using portal vein sampling (19).

Example of Concentration Profile and Secretion Rate

Figure 1 presents an example of a fictional 24-h TSH concentration profile and a corresponding fictional secretion rate diagram. In the top figure, a pulse, and the mode, which is the time it takes before the pulse reaches its maximum, are indicated. In the bottom figure, the secretion rate is depicted, with an indication of the pulsatile and the non-pulsatile part of the TSH secretion.

Characteristics of Pulsatile Secretion

In a study with 38 healthy individuals, including 20 women and 18 men, with a mean age of 41 years (range, 25–64 years), the mean [standard error of the mean (SEM)], number of pulses (secretory-burst frequency) was determined as 16.7 (0.091) per 24 h with a mean mass per pulse of 0.90 (0.06) mU/L (20). Furthermore, the mean (SEM) time before the pulse reaches its




FIGURE 1 | Schematic representation of a fictional 24-h thyroid-stimulating hormone (TSH) concentration profile (top) and secretion rate (bottom). The top figure represents an example of a 24-h TSH concentration profile starting at 09:00 h. A pulse and the mode of one pulse, which is the time it takes before the pulse reaches its maximum, are indicated. The bottom figure is a schematic representation of an example of the secretion rate, which can be determined by deconvolution analysis. The basal tonic (or non-pulsatile) secretion and the pulsatile secretion together contribute to the total TSH secretion.

maximum (mode) was calculated as 20.0 (2.0) min. The fast component of the half-life of TSH, which is effected by the advection and diffusion of TSH, is generally set to be estimated

by deconvolution analysis between 17 to 26 min (21). The slow half-life, which is affected by the irreversible metabolism and elimination of TSH, is set between 66 to 93 min. The mean (SEM) fast half-life was calculated as 22.6 (1.0) min and the slow-phase half-life 104 (2.9) min in these healthy individuals (20). No differences were found between men and women (22, 23). Similar results were obtained in a group of 38 healthy older individuals, including 20 men and 18 women, with an average age of 65 (range 52–76) years, although the slow half-life was lower in this group (on average 78.1 min) (24).

Circadian Rhythm

The greatest impact on variations in serum TSH concentrations has been observed for the sampling time (25, 26), which is caused by the circadian rhythm of TSH. This daily cycle of 24 h is driven by the biological clock of the suprachiasmatic nucleus (SCN). In general, it has been observed that TSH levels reach a maximum between 02:00 and 04:00 h in healthy individuals (5, 20, 27–30). The minimum (i.e., the nadir) of the circadian rhythm of TSH occurs during daytime (4, 5, 27, 28, 31).

Cosinor Analysis to Detect Circadian Rhythmicity

To illustrate the circadian rhythm of TSH, we included an example (see **Figure 2**) of a fictional 24-h TSH concentration profile and drawn a cosinor model and its parameters. Cosinor analysis can be used to determine whether a 24-h concentration profile displays a




FIGURE 2 | Schematic representation of a fictional 24-h thyroid-stimulating hormone (TSH) concentration profile with cosinor model parameters. A cosinor model with its parameters is drawn on the same fictional 24-h TSH concentration profile of **Figure 1** to illustrate the circadian rhythm of TSH. Cosinor analysis calculates the midline estimating statistic of rhythm (MESOR), which is a circadian rhythm-adjusted mean, the maximum, the nadir (minimum), the amplitude, and the acrophase, which is the time at which the cosinor model reaches its maximal value.

sinusoidal circadian rhythm. Cosinor analysis is a model-dependent method which fits a cosinor function to the raw data (32). First, the rhythm detection test, also called the zero-amplitude test, is performed to test the overall significance of the cosinor model. If this model fits significantly, it determines among others the midline estimating statistic of rhythm (MESOR), which is a circadian rhythm-adjusted mean based on the parameters of a cosine function fitted to the raw data. In addition, the amplitude is provided, which is the difference between the maximum and MESOR of the fitted curve. The acrophase represents the time of the maximal value assumed by the curve. Furthermore, the nadir, the lowest point of the cosinor model, is calculated based on the MESOR and the amplitude. Other methods that can be used to examine the circadian rhythmicity of a 24-h TSH concentration profile are for example a modified cosine function (21, 33), a locally weighted polynomial regression analysis (24, 34), or a periodic spline analysis (5).

Circadian Rhythm in Thyroid Disease

TSH secretion patterns have also been studied in treated and untreated primary hypothyroidism patients, which has been reviewed previously (29). In seven patients with primary hypothyroidism on T4 therapy and eight patients with primary hypothyroidism on combined therapy (T3/T4) for more than 3 months, a circadian rhythm with a nocturnal rise persisted (35). Also another study in eleven primary hypothyroid patients on stable T4 replacement therapy showed retention of the circadian rhythm in TSH secretion (36). However, in untreated primary hypothyroid patients, 24-h TSH secretion was increased and the nocturnal surge was often decreased or absent while the pulse frequency was similar compared to euthyroid controls (37–41).

Circadian Rhythm and Ageing

In a population of euthyroid individuals, mean levels of TSH differ with age, but some studies showed that the circadian rhythm does not change with age (5, 27). In contrast, others reported a diminished nocturnal TSH surge in older subjects compared to young controls (42, 43) and an advanced (earlier) shift of the onset of the nocturnal surge of TSH with increasing age (23).

Seasonality

Multiple studies showed that TSH levels are subject to change of season. Evidence for such circannual variation in serum TSH comes from cross-sectional studies, but also from longitudinal studies. In general, TSH levels were highest in winter season.

Cross-Sectional Studies on Seasonality

Several cross-sectional studies found higher TSH levels in euthyroid individuals without thyroid disease whose blood withdrawal for TSH measurements was performed in autumn or winter seasons compared to individuals with blood withdrawals for TSH measurements in spring or summer seasons. An impressively large American study using 465,593 TSH measurements of 324,750 individuals without thyroid disease aged 1–104 years showed that the upper TSH limit of the 95% reference interval

was significantly higher in December (6.06 mU/L) than in August (4.31 mU/L) for both sexes and all age groups (5). The lower limit of the 95% reference interval was 0.5 throughout the year. Barchetta et al. found in 294 Italian euthyroid subjects with a mean age (SD) of 48.5 (12.4) years higher mean TSH levels in individuals evaluated in autumn or winter seasons (2.3 ± 1.3 μ U/ml) compared with individuals who had their blood samples taken in spring or summer seasons (1.8 ± 1.1 μ U/ml, $p = 0.03$) (44). Also in a large group ($N=206,486$) of patients (health check-ups, inpatient and outpatient visits) in Peking, it was observed that median TSH concentrations were highest in winter (1.96 ± 0.128 mU/L) and lowest in summer (1.86 ± 0.111 mU/L) (45). Likewise, mean monthly TSH levels were highest in December in groups of middle-aged and older Italian euthyroid individuals using 8,310 laboratory TSH measurements (46).

Longitudinal Studies on Seasonality

Similar results were obtained when the seasonality of TSH levels was investigated in longitudinal studies, with repeated measurements in the same individuals. Maes et al. showed that 08:00 h fasting TSH levels were highest in December and lowest in June when performing monthly blood sampling for 1 year in 26 Belgian healthy subjects with a mean age (SD) of 38.7 (13.4) years (47). When clustered per season, highest mean (SD) TSH levels (1.85 (1.02) mU/L) were observed in autumn (21 September–20 December) and lowest (1.48 (0.72) mU/L) in spring (21 March–20 June) (47). Furthermore, TSH levels were higher in the winter and spring seasons compared to summer and fall seasons (maximal difference of 0.30 mU/L) in a large group of Korean euthyroid subjects ($N=28,096$) who had on average 3.1 tests per person during a median (interquartile range (IQR)) follow-up of 36 (22–53) months (48).

Studies With No Statistically Significant Seasonality

A few studies found higher TSH levels in winter/autumn compared to summer/spring, but were unable to demonstrate statistically significant seasonality in TSH. For example, no significant seasonal variation in TSH levels was found in a cross-sectional Italian study of a large cohort of 11,806 euthyroid subjects with a median (IQR) age of 49 (37–61) years (49). Nevertheless, median (IQR) TSH values were slightly higher in December [1.46 (0.90–2.20) mU/L] and January [1.46 (0.90–2.16) mU/L] compared to all other months, with lowest median (IQR) TSH levels of 1.39 (0.90–2.10) in April (49). Likewise, no significant seasonal difference was present in a small subset of 159 subjects of this study in which TSH was measured twice in the same year (49). In 152 Iraqi euthyroid subjects with a mean (SD) age of 32.24 (12.05) years who had their TSH levels measured in both summer and winter seasons, no significant difference was found between summer [mean (SD) of 2.34 (1.25) mU/L] and winter [2.25 (1.25) mU/L] measurements (50). Andersen et al. found 8% higher serum TSH concentrations in autumn and winter than in spring and summer using monthly blood samples in 15 healthy Danish men with an age range of 24–53 years, but this difference was not statistically significant (51).

Seasonality in Thyroid Disease

In addition, comparable annual changes were observed in Japanese treated and untreated patients with thyroid disease aged > 20 years (using 1,637,721 TSH measurements) with median TSH levels of 1.46 (range 0.01–769.1) μ U/ml in winter and 1.31 (0.01–594.8) μ U/ml in summer (52), and in Korean patients (mean (SD) age of 48.6 (11.6) years) with subclinical hypothyroidism without medication ($N=1,751$) with a maximal TSH difference of 0.69 mU/L (48). Median TSH levels were highest in January compared to other months in 3,934 L-T4-treated athyreotic patients with a median (IQR) age of 50 (41–59) years of an Italian cross-sectional study (49). Additionally, in a subset of 119 patients [mean (IQR) age of 46 (39–59) years] with one blood withdrawal taken in winter and one in summer, median (IQR) TSH levels were higher in winter season [0.80 (0.22–1.44) mU/L] compared to summer season [0.20 (0.06–0.70) mU/L] (49). Furthermore, although the 10 (aged 32 to 66 years) (53) and 7 (aged 27 to 66 years) (54) primary hypothyroid patients in two longitudinal Japanese studies received constant doses of levothyroxine (L-T4), their unstimulated TSH levels and TSH levels after TRH stimulation were higher in winter compared to summer. In contrast, no significant difference was found in mean (SD) TSH levels measured in 25 subclinical hypothyroid patients with a mean (SD) age of 36.44 (14.82) years in winter [6.04 (1.17) mU/L] compared to summer [6.64 (1.60) mU/L] in Iraq (50).

Seasonality and Ageing

Whether the circannual rhythm of TSH changes with ageing is not clear, one study showed that the seasonal changes were independent from age (55), but another study showed that the circannual rhythm of TSH was stronger in middle-aged and older subjects compared to young subjects (46). In summary, studies showed that TSH levels were higher during winter than during other seasons in both euthyroid individuals as well as in patients with thyroid disease, but the influence of age on the seasonality of TSH concentrations is unsettled.

Ageing

Numerous studies, both cross-sectional and longitudinal, have shown that TSH levels differ with age, with in general higher mean levels with increasing age. A longitudinal study ($N=908$) with a follow-up of 13 years found that mean (SD) serum TSH concentrations increased from 1.49 (0.79) to 1.81 (0.96) mU/L with the largest difference in subjects aged 60+ years (56). Another longitudinal study in 533 healthy older individuals (65+ years) found a mean (SD) increase of 0.28 (1.4) mU/L in 13 years of follow-up (57). In a cross-sectional retrospective analysis of 465,593 TSH measurements of 324,750 individual without thyroid disease, the 97.5% reference interval increased from 6.25 mU/L in subjects aged 21–40 years, to 6.55 (41–60 years), to 6.80 (61–80 years), to 7.55 mU/L in subjects aged 80+ years (5). A similar increase was observed in another cross-sectional retrospective study in 1,388 healthy subjects with a mean (95% reference interval) of 1.30 (0.39–4.29) mU/L in the group of 20–29 years and 1.96 (0.63–6.15) mU/L in the 70+ group (58). Boucai et al. observed an increase in serum TSH concentrations over different age groups with median (2.5–97.5%

interval) levels of 1.30 (0.40–3.98) in the 20-to-29-years-old-group and 1.99 (0.44–6.92) mU/L in over-80-years-old-group (59).

Clinical Trials in Older Persons With Subclinical Hypothyroidism

One potential school of thought states that age-related changes in hormone levels, including TSH, contribute to health problems in older individuals. It is therefore an ongoing debate whether TSH levels should be “normalized” in older individuals to prevent age-related loss in functioning or whether the upper reference limit for TSH should be adapted according to age (60). In line with the increase in mean TSH levels with age, the prevalence of subclinical hypothyroidism is also increasing, which is defined as an elevated TSH level that occurs in conjunction with a serum fT4 concentration within the normal range (61, 62). Therefore, recently, two relatively large ($N=737$ and $N=251$) randomized placebo-controlled clinical trials (RCTs) have been performed in which older persons with subclinical hypothyroidism received levothyroxine (L-T4) or placebo supplementation (63, 64). The two primary study outcomes were the change in the hypothyroid symptoms score and tiredness score from the thyroid-related quality of life patient-reported outcome (ThyPRO) questionnaire from baseline to 1-year follow-up. Although TSH levels in participants receiving L-T4 treatment declined, there was no beneficial effect on their thyroid-related symptoms compared to the placebo group (63, 64). Along the same line, a systematic review and meta-analysis provided evidence that the use of thyroid hormone therapy was not associated with improvements in general quality of life or thyroid-related symptoms in RCTs with a follow-up of 3–15 months (65).

TSH and Human Longevity

In fact, higher TSH levels might even be beneficial for healthy ageing and longevity. Older individuals from the Leiden 85-Plus Study with elevated serum TSH levels without thyroid disease were not at risk of increased morbidity and may have a prolonged life span (66). We have previously reported that the offspring of long-lived families, who have the propensity to reach an advanced age with limited clinical issues, seem to have higher levels (within normal range) of serum TSH and total 24-h secretion of TSH (67). Other studies have also shown that centenarians as well as their offspring had a higher level of TSH when compared to age matching controls (68, 69). Thus, these studies support the hypothesis that elevated TSH levels with ageing are not necessarily unfavorable.

Other Sources of Within-Person Variation in TSH Levels

TPO-Ab Positivity

Evidence for the influence of several other factors on the within-person variation in TSH levels can be found in literature. For example, a positive association was found between thyroid peroxidase (TPO)-antibody positivity and TSH levels during early pregnancy in a large ($N=11,212$) individual participant-based meta-analysis (70). Similarly, a positive correlation

between TPO-Ab positivity and TSH levels was observed, cross-sectionally as well as over months, within 21 individuals with subclinical hypothyroidism (60, 71).

BMI and Obesity

Furthermore, a positive correlation between TSH levels and body mass index (BMI) was found in a population-based ($N=8,727$) cross-sectional study (72). Similarly, median (IQR) TSH levels in obese ($BMI \geq 30$ kg/m²) euthyroid subjects were somewhat higher [1.53 (1.07–2.23) vs. 1.47 (1.04–2.12) mU/L] than in the reference group (18.5–24.9 kg/m²) in a cross-sectional study with 11,224 participants (73). Moreover, TSH levels decreased significantly 12 months after weight loss due to bariatric surgery in 949 euthyroid patients with morbid obesity, which was more pronounced in the high-normal TSH group (74). BMI was positively associated with basal TSH secretion in healthy individuals, but not with pulsatile or total TSH secretion, and a higher BMI was associated with a delay in the nocturnal onset of the TSH surge in 117 healthy subjects with a mean age of 43 (range 22–77) years and a mean BMI of 26.8 (range 18.3–39.4) kg/m² (23). In contrast, TSH levels were not different between a small group of overweight women ($N=22$) and women with normal weight ($N=30$) (75).

Selenium, Iodine, and Smoking

Although selenium deficiency is associated with impaired thyroid function, there is no evidence for a direct effect of selenium intake on TSH levels in individuals without severe selenium deficiency (76). Also, iodine intake can influence TSH levels; a positive correlation was found between urinary iodine concentrations and TSH levels in a large ($N=6,565$) Korean dataset (77). However, more research on the direct relationship between iodine intake and TSH levels is necessary. In the same Korean study, it was observed that mean serum TSH levels were lower in current smokers than in non-smokers, which was more apparent in iodine-deficient subjects (78). Other studies also showed that smoking is associated with lower TSH levels with modestly higher fT4 levels (79–81), which was reviewed before (82, 83).

Severe Illness

Critically ill patients exhibit a decrease in serum concentrations of thyroid hormones but with TSH levels in the normal range or slightly decreased, which is known as non-thyroidal illness syndrome (NTIS) or euthyroid sick syndrome (ESS). These effects of severe illness on thyroid function have been reviewed by others (84, 85). Another study found that pulsatile TSH secretion is diminished in 26 critically ill patients with a mean (SEM) age of 63 (2) years (86, 87).

Drugs and Xenobiotics

Many drugs can also affect thyroid function, including glucocorticoids, lithium, amiodarone, and antiepileptic drugs. A comprehensive review on the effects of drugs on TSH levels can be found elsewhere (88). Other xenobiotics, such as environmental endocrine disruptors, also have a large influence on thyroid parameters and metabolism (89).

POTENTIAL UNDERLYING BIOLOGICAL MECHANISMS OF WITHIN-PERSON VARIATION IN TSH LEVELS

Potential Underlying Mechanisms of Pulsatile Secretion

The main sources of within-person variation in TSH levels have been discussed, but what are the biological mechanisms that underlie the sources of within-person variation in TSH levels? One of the main roles of the HPT axis is to respond to environmental changes in order to maintain homeostasis. TSH is a key player in this system, which is responsible for maintaining constant circulating levels of thyroid hormones within the normal ranges over time. TSH is highly responsive to different stressors, including inflammation (90) and environmental endocrine disruptors (89). To be able to respond quickly to changes in the environment, TSH is secreted in a pulsatile manner and the half-life of TSH is relatively short (17 to 93 min). It is not known whether T3 and T4 are secreted in a pulsatile manner from the thyroid gland. The half-lives of T3 and T4 are much longer (0.75 and 6.7 days respectively) than that of TSH (27). Free T3 exhibits a minor circadian rhythm and fT4 does not exhibit a circadian rhythm (27). However, it is not completely clear why TSH fluctuates in periods of minutes, hours, and months while fT4 levels stay relatively constant over time. Other pituitary hormones also have a short half-life and fluctuate over time—they are all designed to adapt in response to changes in the environment—while levels of peripheral hormones are more stable over time. For example, within the somatotrophic axis, growth hormone (GH) is secreted in a pulsatile manner and fluctuates strongly over 24 h, while insulin-like growth factor 1 (IGF-1) is relatively stable over 24 h. It is known that manipulations of GH and IGF-1 have shared, but also distinct effects (91). Therefore, a hypothesis could be that TSH affects, besides similar, also different mechanisms and processes in the human body than thyroid hormones, which may explain the distinct fluctuations in levels over time between TSH and thyroid hormones. In line with this hypothesis it was observed, among others, that TSH receptors were located on T cells in the thymus and that TSH directly—independent of thyroid hormones—has an influence on the maturation of these T cells (92).

Potential Underlying Mechanisms of Circadian Variation

The biological mechanisms that underlie the daily fluctuations in TSH levels are most likely related to the biological clock. The biological clock of the SCN drives the circadian rhythm of several processes, including hormones. The biological clock directly influences the circadian rhythm of TSH, but it was also found that the circadian rhythm of TSH is partly determined by the circadian rhythm of cortisol, which is completely dependent on the SCN (93). The circadian rhythms of TSH and cortisol are out of phase; TSH has a nocturnal surge in the early night while cortisol has its nadir around that time and its peak in the early morning. The circadian rhythms of cortisol and TSH being out-

of-phase can be explained by the inhibitory influence of cortisol on serum TSH concentrations (94–98). Additionally, a strong negative cross-correlation between cortisol and TSH concentrations, with TSH following cortisol concentrations in the opposite direction after a time delay of 170 min, was observed in 38 healthy older individuals (99). Also, the biological clock influences processes of the sympathetic and parasympathetic nervous system. Differences in sympathetic/parasympathetic tone could possibly lead to alterations in end organ sensitivity, such as the thyroid gland. Although it is speculative, it could be suggested that the thyroid gland is less sensitive to TSH during the night and during winter seasons, and therefore higher TSH levels are required to keep fT4 levels constant. It is also known that other physiological processes exhibit a circadian rhythm, including metabolic processes. TSH plays an important role in metabolism and might therefore exhibit parallel fluctuations. TSH plays an important role in energy balance and distribution. Energy requirements also fluctuate over time, during the day as well as during seasons. Furthermore, it has been hypothesized that TSH plays a role in maintenance and repair mechanisms, including bone turnover (100). These processes fluctuate over time, with a bone resorption marker, C-terminal cross-linked telopeptide of type 1 collagen (CTX-I), exhibiting a strong circadian rhythm (101).

Potential Underlying Mechanisms of Seasonality: Environmental Temperature

A number of potential mechanisms may explain the circannual changes in TSH levels, with environmental temperature being the most apparent explanation which has been reviewed before (102). Two studies in healthy individuals and one study in patients with primary hypothyroidism under L-T4 treatment indeed observed a significant negative correlation between TSH levels and the environmental temperature (45, 48, 53). It is important to note that these studies were performed in Korea, China, and Japan where the environmental temperature is very low in winter. In contrast, two Italian studies did not find a significant correlation between TSH levels and environmental temperature (46, 55). It becomes less cold in winter in Italy than in Asia, which could explain the discrepancy in results. The seasonal fluctuation might therefore be more pronounced in particularly cold climates. Most studies on circannual changes in TSH levels were performed in the United States, Asia, and Europe. However, some studies were performed in Antarctica and at the South Pole, where the effect of long-term cold exposure could be studied in personnel of the McMurdo and South Pole research stations. Mean TSH levels increased by approximately 30% during a 9-month Antarctic residence in nine healthy young subjects (103) and the TSH response to TRH increased by approximately 50% in 17 healthy young men (104). Similarly, mean TSH levels increased in ten healthy men after a 54-weeks Antarctic expedition (105) and comparable results were obtained in a larger study of 187 workers at the McMurdo and South Pole research stations (106). These changes in thyroid function are known as the polar T3 syndrome, which is not only characterized by increased TSH levels, but also by alterations in T3 kinetic parameters such as an

increase in serum clearance of T₃, possibly explained by increased tissue uptake (107). This thyroidal response to cold might be required to keep the core body temperature constant. Further support for the effect of environmental temperature on TSH levels are observations in bears. In a study with six American black bears it was found that the TSH response to TRH administration was prolonged, delayed, and exaggerated during hibernation in winter while (f)T₃ and (f)T₄ levels were decreased (108). In addition, T₃ levels increased stronger after TRH administration during hibernation than after hibernation.

Potential Underlying Mechanisms of Seasonality: Length of Day

Studies in animals indicated that length of day, and possibly melatonin (109), plays a role in the seasonal variation in TSH levels, which has been reviewed by others (110). Animals that breed seasonally exhibit seasonal variations in numerous processes and functions, including reproduction, fattening, hibernation, and migration, which are regulated *via* photoperiod-regulated systems. It was for example found that the simulation of a long-day in the quail induced the secretion of TSH from the pars tuberalis (PT) of the anterior pituitary gland (111). Moreover, light induced the expression of the gene encoding type 2-iodothyronine deiodinase (Dio2), which converts T₄ to T₃, in quails (112). The underlying mechanism is possibly that TSH and/or thyroid hormones stimulate the secretion of gonadotropin-releasing hormone and gonadotropin leading to gonadal growth, which is important for reproduction (111, 113). Humans however are not seasonal breeders, so these mechanisms could be of greater importance in animals than in humans. Nevertheless, understanding the seasonal mechanisms in animals might lead to better understanding of seasonality in humans. The thyroid also plays a role in other transitions to different physiological states that exist in nature, for example in the metamorphosis of the tadpole to a frog (114–116).

Potential Underlying Mechanisms of Age-Related Changes

Several hypotheses exist why there is an increase in population TSH levels with increasing age (117, 118). At a population level, the higher median TSH levels in older individuals compared to younger individuals might be explained by 1) age-dependent selective survival of individuals with elevated TSH levels or by 2) a birth cohort effect since different birth cohorts have been exposed to different environmental factors which could have influenced their TSH levels. At the organism level, age-related increase in TSH levels might be caused by 1) decreased TSH bioactivity, 2) diminished response of the thyroid gland to TSH stimulation, or 3) diminished sensitivity of the pituitary and/or hypothalamus to the negative feedback of thyroid hormones. The last hypothesis however would be associated with an increase in thyroid hormone levels, which is not supported by literature. The lack of evidence for increased thyroid hormone levels in response to diminished sensitivity of the pituitary and/or hypothalamus could be explained by enhanced thyroid hormone turnover in peripheral tissues or greater clearance of thyroid hormones from the circulation (67).

CLINICAL IMPLICATIONS OF WITHIN-PERSON VARIATION IN TSH LEVELS

TSH levels fluctuate over time within a person, both in euthyroid individuals and in most patients with thyroid disorders. If clinicians are not aware of fluctuations in TSH levels within an individual patient, this could lead to under- or overdiagnosis, and to inadequate prescription of thyroid medication to patients with thyroid disease. Since the mean TSH levels and the prevalence of subclinical hypothyroidism is increasing with age, this is especially important for the diagnosis and follow-up of older patients. Also, for clinical scientific research it is important to consider within-person variations in TSH levels over time, since this could cause more heterogeneity in research outcomes.

Season of Testing and the Transition of Subclinical Hypothyroidism to Euthyroid

Whether the sampling time has an influence on the assignment of an individual to a clinical category of thyroid status is not known. In contrast, it was shown that the season of thyroid testing played a role in the transition of patients between subclinical hypothyroidism and euthyroid status (48). In a large ($N=29,847$) retrospective longitudinal study, 57.9% of subclinical hypothyroid individuals ($N=1,751$) reverted to euthyroid TSH levels during a median follow-up of 36 months. Normalization of subclinical hypothyroidism to euthyroidism was increased during summer–fall follow-up. The opposite was true as well; the transition from euthyroidism to subclinical hypothyroidism was more likely in individuals who had their follow-up TSH measurement in winter or spring (48). Also, other studies showed that TSH levels of patients with subclinical hypothyroidism can normalize to euthyroid levels without intervention. In total, 40% of the participants ($N=2,936$) aged above 65 years from a longitudinal follow-up study, the Birmingham Elderly Thyroid Study, who classified as being subclinical hypothyroid at baseline normalized to euthyroid status without intervention over a period of 5 years (119). However, the authors also concluded that TSH concentrations were quite stable over time; 61% of the participants had a repeated TSH concentration within 0.5 mU/L of their original result. A small study ($N=107$) found that 37.4% of older (above 55 years) patients with subclinical hypothyroidism showed normalization of their TSH levels during a mean (range) follow-up of 31.7 (6–72) months (120, 121). Whether the normalization was related to the season in which follow-up measurements were taken was not reported in these two studies.

Clinical Reference Range

Studying the within- and between-person variations in TSH levels revealed that the within-person variation is smaller than the between-person variation (3, 60, 122). A seemingly small change in TSH level within the population reference interval might therefore be large—and potentially even pathological—for an individual. Indeed, a meta-analysis showed that variations in

TSH levels within the population reference range are associated with adverse health outcomes, including cardiovascular risk factors, metabolic parameters, and increased risk of osteoporosis and fracture (123). It has therefore been recommended to determine individual trends and variations in experimental clinical research, and thereby even establish one's individual TSH reference range, by performing repeated measurements (124). Although this strategy is not feasible in routine clinical practice, it might be worthwhile to explore the possibility to monitor patients who are diagnosed with

subclinical hypothyroidism more regularly since their TSH level is often close to the upper reference limit. Procedures for regular monitoring of a subclinical hypothyroid patient's TSH levels have been recommended by others (60). Moreover, Hoermann et al. stated that there should be more attention in clinical practice for the interrelationships between TSH and thyroid hormones (125). Mathematical models have shown that the relationship between TSH and fT4 is not log-linear as previously thought, but much more complex, dynamic, and individual (125).




FIGURE 3 | Graphical summary of 1) the potential underlying biological mechanisms of within-person variation in thyroid-stimulating hormone (TSH) levels over time, 2) the main sources, and 3) the clinical implications. The main topics discussed in this review are illustrated. The main sources of within-person variation in TSH levels over time are pulsatile secretion, circadian rhythm, seasonality, and ageing, as illustrated in 3.2. The main hypotheses for the underlying mechanisms of within-person variations are illustrated in 3.1. The pituitary needs to respond quickly to changes in the environment to maintain homeostasis, which is one of the potential underlying biological mechanisms of pulsatile secretion of TSH. The biological clock regulates the circadian rhythm of many physiological processes including pituitary hormones and energy metabolism. The seasonality in TSH levels are possibly caused by changes in environmental temperature, with studies indicating generally highest levels during winter season. Multiple studies showed that on a population level, TSH levels generally increase with age. However, the size and rate of the increase within an individual is not completely clear. Finally, the implications of within-person variation in TSH levels over time for clinical practice and scientific research are illustrated in 3.3. Within-person variation in TSH levels is not only present in healthy euthyroid individuals, but also seen in patients with (un)treated thyroid disease. It occurs frequently that subclinical hypothyroid patients normalize to euthyroid levels over time without intervention and the season of thyroid testing plays a role in the transition of patients between subclinical hypothyroidism and the euthyroid status. We recommend regular monitoring of an individual's TSH levels by obtaining repeated measurements so different sources of within-person variations can be considered.

CONCLUSIONS

A graphical summary of this review is provided in **Figure 3**. TSH levels fluctuate over time within a person, in euthyroid individuals as well as in most (un)treated patients with thyroid disorders, ranging from minutes, hours, months, to years. These within-person variations in TSH levels are mainly caused by pulsatile secretion, circadian rhythm, seasons, and ageing. If TSH is fluctuating around the upper reference range limit, these marginal changes could lead to diagnosis of subclinical hypothyroidism, especially in older adults. However, elevated TSH levels can return to euthyroid levels in a short period of time without intervention, which can be explained by the different sources of within-person variation. Therefore, we recommend considering the fluctuations in TSH levels and their potential underlying biological mechanisms in clinical practice and when performing clinical research. For example, by adaptation of the TSH reference limits, specifically the upper TSH reference limit, for the seasonal variations, circadian rhythm, and age. Future studies should aim to find approaches to provide a more personalized reference range for thyroid disease diagnosis and clinical treatment which considers the different sources of within-person variation in TSH levels. Moreover, we propose that there should be more attention in clinical practice for the

relationship between TSH and fT4 levels, for example using a TSH-fT4 nomogram. Ideally, an algorithm will be developed which includes the individual TSH-fT4 nomogram together with the time of day and season of the blood sampling, the age of the patient, and possibly other factors that influence TSH levels, such as pre-analytical and analytical variation.

AUTHOR CONTRIBUTIONS

All discussed the content, main message, and structure of the review. EV conducted literature search and wrote the first draft. FR and DV reviewed the text and edited some parts. All discussed the content of the figures. FR and EV made the figures. All authors contributed to the article and approved the submitted version.

FUNDING

This work was supported by the European Commission project THYRAGE (Horizon 2020 research and innovation programme, 666869).

REFERENCES

- Citterio CE, Targovnik HM, Arvan P. The role of thyroglobulin in thyroid hormonogenesis. *Nat Rev Endocrinol* (2019) 15(6):323–38. doi: 10.1038/s41574-019-0184-8
- Hoermann R, Pekker MJ, Midgley JEM, Larisch R, Dietrich JW. Triiodothyronine secretion in early thyroid failure: The adaptive response of central feedforward control. *Eur J Clin Invest* (2020) 50(2):e13192. doi: 10.1111/eci.13192
- Andersen S, Bruun NH, Pedersen KM, Laurberg P. Biologic variation is important for interpretation of thyroid function tests. *Thyroid* (2003) 13(11):1069–78. doi: 10.1089/105072503770867237
- Weeke J, Gundersen HJ. Circadian and 30 minutes variations in serum TSH and thyroid hormones in normal subjects. *Acta Endocrinol* (1978) 89(4):659–72. doi: 10.1530/acta.0.0890659
- Ehrenkranz J, Bach PR, Snow GL, Schneider A, Lee JL, Ilstrup S, et al. Circadian and Circannual Rhythms in Thyroid Hormones: Determining the TSH and Free T4 Reference Intervals Based Upon Time of Day, Age, and Sex. *Thyroid* (2015) 25(8):954–61. doi: 10.1089/thy.2014.0589
- van Deudekom FJ, Postmus I, van der Ham DJ, Pothof AB, Broekhuizen K, Blauw GJ, et al. External validity of randomized controlled trials in older adults, a systematic review. *PLoS One* (2017) 12(3):e0174053. doi: 10.1371/journal.pone.0174053
- Porcino AJ, Shamseer L, Chan AW, Kravitz RL, Orkin A, Punja S, et al. SPIRIT extension and elaboration for n-of-1 trials: SPENT 2019 checklist. *BMJ* (2020) 368:m122. doi: 10.1136/bmj.m122
- Vandenbroucke JP. The N-of-1 trial: the ideal study design that is underused. *Ned Tijdschr Tandheelkd* (2007) 114(6):260–2.
- Greenspan SL, Klibanski A, Schoenfeld D, Ridgway EC. Pulsatile secretion of thyrotropin in man. *J Clin Endocrinol Metab* (1986) 63(3):661–8. doi: 10.1210/jcem-63-3-661
- Clark PM, Clark JD, Holder R, Raggatt PR, Wheatley T. Pulsatile secretion of TSH in healthy subjects. *Ann Clin Biochem* (1987) 24(Pt 5):470–6. doi: 10.1177/000456328702400508
- Casanueva FF, Dieguez C. Neuroendocrine regulation and actions of leptin. *Front Neuroendocrinol* (1999) 20(4):317–63. doi: 10.1006/frne.1999.0187
- Mullur R, Liu YY, Brent GA. Thyroid hormone regulation of metabolism. *Physiol Rev* (2014) 94(2):355–82. doi: 10.1152/physrev.00030.2013
- Krulich L. Neurotransmitter control of thyrotropin secretion. *Neuroendocrinology* (1982) 35(2):139–47. doi: 10.1159/000123369
- Reichlin S. Somatostatin. *N Engl J Med* (1983) 309(24):1495–501. doi: 10.1056/nejm198312153092406
- Scanlon MF, Weightman DR, Shale DJ, Mora B, Heath M, Snow MH, et al. Dopamine is a physiological regulator of thyrotrophin (TSH) secretion in normal man. *Clin Endocrinol* (1979) 10(1):7–15. doi: 10.1111/j.1365-2265.1979.tb03028.x
- Keenan DM, Licinio J, Veldhuis JD. A feedback-controlled ensemble model of the stress-responsive hypothalamo-pituitary-adrenal axis. *Proc Natl Acad Sci U S A* (2001) 98(7):4028–33. doi: 10.1073/pnas.051624198
- Keenan DM, Roelfsema F, Biersema N, Veldhuis JD. Physiological control of pituitary hormone secretory-burst mass, frequency, and waveform: a statistical formulation and analysis. *Am J Physiol Regul Integr Comp Physiol* (2003) 285(3):R664–73. doi: 10.1152/ajpregu.00195.2003
- Liu PY, Keenan DM, Kok P, Padmanabhan V, O'Byrne KT, Veldhuis JD. Sensitivity and specificity of pulse detection using a new deconvolution method. *Am J Physiol Endocrinol Metab* (2009) 297(2):E538–44. doi: 10.1152/ajpendo.00071.2009
- Pørksen N, Grøfte T, Greisen J, Mengel A, Juhl C, Veldhuis JD, et al. Human insulin release processes measured by intraportal sampling. *Am J Physiol Endocrinol Metab* (2002) 282(3):E695–702. doi: 10.1152/ajpendo.00516.2000
- Roelfsema F, Pereira AM, Adriaanse R, Endert E, Fliers E, Romijn JA, et al. Thyrotropin secretion in mild and severe primary hypothyroidism is distinguished by amplified burst mass and Basal secretion with increased spikiness and approximate entropy. *J Clin Endocrinol Metab* (2010) 95(2):928–34. doi: 10.1210/jc.2009-1959
- Joustra SD, Roelfsema F, Endert E, Ballieux BE, van Trotsenburg AS, Fliers E, et al. Pituitary Hormone Secretion Profiles in IGSF1 Deficiency Syndrome. *Neuroendocrinology* (2016) 103(3-4):408–16. doi: 10.1159/000439433
- Roelfsema F, Pereira AM, Veldhuis JD, Adriaanse R, Endert E, Fliers E, et al. Thyrotropin secretion profiles are not different in men and women. *J Clin Endocrinol Metab* (2009) 94(10):3964–7. doi: 10.1210/jc.2009-1155

23. Roelfsema F, Pijl H, Kok P, Endert E, Fliers E, Biermasz NR, et al. Thyrotropin secretion in healthy subjects is robust and independent of age and gender, and only weakly dependent on body mass index. *J Clin Endocrinol Metab* (2014) 99(2):570–8. doi: 10.1210/jc.2013-2858
24. Jansen SW, Roelfsema F, van der Spoel E, Akintola AA, Postmus I, Ballieux BE, et al. Familial Longevity Is Associated With Higher TSH Secretion and Strong TSH-fT3 Relationship. *J Clin Endocrinol Metab* (2015) 100(10):3806–13. doi: 10.1210/jc.2015-2624
25. Wang D, Li D, Guo X, Yu S, Qiu L, Cheng X, et al. Effects of sex, age, sampling time, and season on thyroid-stimulating hormone concentrations: A retrospective study. *Biochem Biophys Res Commun* (2018) 506(3):450–4. doi: 10.1016/j.bbrc.2018.10.099
26. Mirjanic-Azaric B, Stojakovic-Jelisavac T, Vukovic B, Stojanovic D, Vujnic M, Uletilovic S. The impact of time of sample collection on the measurement of thyroid stimulating hormone values in the serum. *Clin Biochem* (2015) 48(18):1347–9. doi: 10.1016/j.clinbiochem.2015.08.020
27. Russell W, Harrison RF, Smith N, Darzy K, Shalet S, Weetman AP, et al. Free triiodothyronine has a distinct circadian rhythm that is delayed but parallels thyrotropin levels. *J Clin Endocrinol Metab* (2008) 93(6):2300–6. doi: 10.1210/jc.2007-2674
28. Patel YC, Alford FP, Burger HG. The 24-hour plasma thyrotrophin profile. *Clin Sci* (1972) 43(1):71–7. doi: 10.1042/cs0430071
29. Roelfsema F, Veldhuis JD. Thyrotropin secretion patterns in health and disease. *Endocr Rev* (2013) 34(5):619–57. doi: 10.1210/er.2012-1076
30. Barreca T, Franceschini R, Messina V, Bottaro L, Rolandi E. 24-hour thyroid-stimulating hormone secretory pattern in elderly men. *Gerontology* (1985) 31(2):119–23. doi: 10.1159/000212690
31. Lucke C, Hehrmann R, von Mayersbach K, von zur Muhlen A. Studies on circadian variations of plasma TSH, thyroxine and triiodothyronine in man. *Acta Endocrinol* (1977) 86(1):81–8. doi: 10.1530/acta.0.0860081
32. Refinetti R, Lissen GC, Halberg F. Procedures for numerical analysis of circadian rhythms. *Biol Rhythm Res* (2007) 38(4):275–325. doi: 10.1080/09291010600903692
33. Van Someren EJ, Nagtegaal E. Improving melatonin circadian phase estimates. *Sleep Med* (2007) 8(6):590–601. doi: 10.1016/j.sleep.2007.03.012
34. Cleveland WS, Devlin SJ. Locally Weighted Regression - an Approach to Regression-Analysis by Local Fitting. *J Am Stat Assoc* (1988) 83(403):596–610. doi: 10.2307/2289282
35. Saravanan P, Siddique H, Simmons DJ, Greenwood R, Dayan CM. Twenty-four hour hormone profiles of TSH, Free T3 and free T4 in hypothyroid patients on combined T3/T4 therapy. *Exp Clin Endocrinol Diabetes* (2007) 115(4):261–7. doi: 10.1055/s-2007-973071
36. Sturgess I, Thomas SH, Pennell DJ, Mitchell D, Croft DN. Diurnal variation in TSH and free thyroid hormones in patients on thyroxine replacement. *Acta Endocrinol* (1989) 121(5):674–6. doi: 10.1530/acta.0.1210674
37. Samuels MH, Veldhuis JD, Henry P, Ridgway EC. Pathophysiology of pulsatile and copulsatile release of thyroid-stimulating hormone, luteinizing hormone, follicle-stimulating hormone, and alpha-subunit. *J Clin Endocrinol Metab* (1990) 71(2):425–32. doi: 10.1210/jcem-71-2-425
38. Weeke J, Laurberg P. Diurnal TSH variations in hypothyroidism. *J Clin Endocrinol Metab* (1976) 43(1):32–7. doi: 10.1210/jcem-43-1-32
39. Adriaanse R, Brabant G, Prank K, Endert E, Wiersinga WM. Circadian changes in pulsatile TSH release in primary hypothyroidism. *Clin Endocrinol* (1992) 37(6):504–10. doi: 10.1111/j.1365-2265.1992.tb01481.x
40. Adriaanse R, Romijn JA, Endert E, Wiersinga WM. The nocturnal thyroid-stimulating hormone surge is absent in overt, present in mild primary and equivocal in central hypothyroidism. *Acta Endocrinol* (1992) 126(3):206–12. doi: 10.1530/acta.0.1260206
41. Adriaanse R, Brabant G, Endert E, Wiersinga WM. Pulsatile thyrotropin release in patients with untreated pituitary disease. *J Clin Endocrinol Metab* (1993) 77(1):205–9. doi: 10.1210/jcem.77.1.8325943
42. van Coevorden A, Laurent E, Decoster C, Kerkhofs M, Neve P, van Cauter E, et al. Decreased basal and stimulated thyrotropin secretion in healthy elderly men. *J Clin Endocrinol Metab* (1989) 69(1):177–85. doi: 10.1210/jcem-69-1-177
43. Greenspan SL, Klibanski A, Rowe JW, Elahi D. Age-related alterations in pulsatile secretion of TSH: role of dopaminergic regulation. *Am J Physiol* (1991) 260(3 Pt 1):E486–91. doi: 10.1152/ajpendo.1991.260.3.E486
44. Barchetta I, Baroni MG, Leonetti F, De Bernardinis M, Bertocchini L, Fontana M, et al. TSH levels are associated with vitamin D status and seasonality in an adult population of euthyroid adults. *Clin Exp Med* (2015) 15(3):389–96. doi: 10.1007/s10238-014-0290-9
45. Wang D, Cheng X, Yu S, Qiu L, Lian X, Guo X, et al. Data mining: Seasonal and temperature fluctuations in thyroid-stimulating hormone. *Clin Biochem* (2018) 60:59–63. doi: 10.1016/j.clinbiochem.2018.08.008
46. Simoni M, Velardo A, Montanini V, Faustini Fustini M, Seghedoni S, Marrama P. Circannual rhythm of plasma thyrotropin in middle-aged and old euthyroid subjects. *Horm Res* (1990) 33(5):184–9. doi: 10.1159/000181506
47. Maes M, Mommen K, Hendrickx D, Peeters D, D'Hondt P, Ranjan R, et al. Components of biological variation, including seasonality, in blood concentrations of TSH, TT3, FT4, PRL, cortisol and testosterone in healthy volunteers. *Clin Endocrinol* (1997) 46(5):587–98. doi: 10.1046/j.1365-2265.1997.1881002.x
48. Kim TH, Kim KW, Ahn HY, Choi HS, Won H, Choi Y, et al. Effect of seasonal changes on the transition between subclinical hypothyroid and euthyroid status. *J Clin Endocrinol Metab* (2013) 98(8):3420–9. doi: 10.1210/jc.2013-1607
49. Gullo D, Latina A, Frasca F, Squatrito S, Belfiore A, Vigneri R. Seasonal variations in TSH serum levels in athyreotic patients under L-thyroxine replacement monotherapy. *Clin Endocrinol* (2017) 87(2):207–15. doi: 10.1111/cen.13351
50. Mahwi TO, Abdulateef DS. Relation of Different Components of Climate with Human Pituitary-Thyroid Axis and FT3/FT4 Ratio: A Study on Euthyroid and SCH Subjects in Two Different Seasons. *Int J Endocrinol* (2019) 2019:2762978. doi: 10.1155/2019/2762978
51. Andersen S, Pedersen KM, Pedersen IB, Laurberg P. Variations in urinary iodine excretion and thyroid function. A 1-year study in healthy men. *Eur J Endocrinol* (2001) 144(5):461–5. doi: 10.1530/eje.0.1440461
52. Yoshihara A, Noh JY, Watanabe N, Iwaku K, Kunii Y, Ohye H, et al. Seasonal Changes in Serum Thyrotropin Concentrations Observed from Big Data Obtained During Six Consecutive Years from 2010 to 2015 at a Single Hospital in Japan. *Thyroid* (2018) 28(4):429–36. doi: 10.1089/thy.2017.0600
53. Konno N, Morikawa K. Seasonal variation of serum thyrotropin concentration and thyrotropin response to thyrotropin-releasing hormone in patients with primary hypothyroidism on constant replacement dosage of thyroxine. *J Clin Endocrinol Metab* (1982) 54(6):1118–24. doi: 10.1210/jcem-54-6-1118
54. Hamada N, Ohno M, Morii H, Jaedok N, Yamakawa J, Inaba M, et al. Is it necessary to adjust the replacement dose of thyroid hormone to the season in patients with hypothyroidism? *Metabolism* (1984) 33(3):215–8. doi: 10.1016/0026-0495(84)90039-8
55. Santi D, Spaggiari G, Brigante G, Setti M, Tagliavini S, Trenti T, et al. Semi-annual seasonal pattern of serum thyrotropin in adults. *Sci Rep* (2019) 9(1):10786. doi: 10.1038/s41598-019-47349-4
56. Bremner AP, Feddema P, Leedman PJ, Brown SJ, Beilby JP, Lim EM, et al. Age-related changes in thyroid function: a longitudinal study of a community-based cohort. *J Clin Endocrinol Metab* (2012) 97(5):1554–62. doi: 10.1210/jc.2011-3020
57. Waring AC, Arnold AM, Newman AB, Buzkova P, Hirsch C, Cappola AR. Longitudinal changes in thyroid function in the oldest old and survival: the cardiovascular health study all-stars study. *J Clin Endocrinol Metab* (2012) 97(11):3944–50. doi: 10.1210/jc.2012-2481
58. Yoshihara A, Noh JY, Ohye H, Sato S, Sekiya K, Kosuga Y, et al. Reference limits for serum thyrotropin in a Japanese population. *Endocr J* (2011) 58(7):585–8. doi: 10.1507/endocrj.k11e-082
59. Boucai L, Hollowell JG, Surks MI. An approach for development of age-, gender-, and ethnicity-specific thyrotropin reference limits. *Thyroid* (2011) 21(1):5–11. doi: 10.1089/thy.2010.0092
60. Karmisholt J, Andersen S, Laurberg P. Variation in thyroid function in subclinical hypothyroidism: importance of clinical follow-up and therapy. *Eur J Endocrinol* (2011) 164(3):317–23. doi: 10.1530/eje-10-1021
61. Calsolaro V, Nicolai F, Pasqualetti G, Calabrese AM, Polini A, Okoye C, et al. Overt and Subclinical Hypothyroidism in the Elderly: When to Treat? *Front Endocrinol* (2019) 10:177. doi: 10.3389/fendo.2019.00177

62. Biondi B, Cappola AR, Cooper DS. Subclinical Hypothyroidism: A Review. *JAMA* (2019) 322(2):153–60. doi: 10.1001/jama.2019.9052
63. Stott DJ, Rodondi N, Bauer DC. Thyroid Hormone Therapy for Older Adults with Subclinical Hypothyroidism. *N Engl J Med* (2017) 377(14):e20. doi: 10.1056/NEJMc1709989
64. Mooijaart SP, Du Puy RS, Stott DJ, Kearney PM, Rodondi N, Westendorp RGJ, et al. Association Between Levothyroxine Treatment and Thyroid-Related Symptoms Among Adults Aged 80 Years and Older With Subclinical Hypothyroidism. *JAMA* (2019) 322(20):1–11. doi: 10.1001/jama.2019.17274
65. Feller M, Snel M, Moutzouri E, Bauer DC, de Montmollin M, Aujesky D, et al. Association of Thyroid Hormone Therapy With Quality of Life and Thyroid-Related Symptoms in Patients With Subclinical Hypothyroidism: A Systematic Review and Meta-analysis. *JAMA* (2018) 320(13):1349–59. doi: 10.1001/jama.2018.13770
66. Gussekloo J, van Exel E, de Craen AJ, Meinders AE, Frölich M, Westendorp RG. Thyroid status, disability and cognitive function, and survival in old age. *JAMA* (2004) 292(21):2591–9. doi: 10.1001/jama.292.21.2591
67. Jansen SW, Akintola AA, Roelfsema F, van der Spoel E, Cobbaert CM, Ballieux BE, et al. Human longevity is characterised by high thyroid stimulating hormone secretion without altered energy metabolism. *Sci Rep* (2015) 5:11525. doi: 10.1038/srep11525
68. Atzmon G, Barzilai N, Hollowell JG, Surks MI, Gabriely I. Extreme longevity is associated with increased serum thyrotropin. *J Clin Endocrinol Metab* (2009) 94(4):1251–4. doi: 10.1210/jc.2008-2325
69. Atzmon G, Barzilai N, Surks MI, Gabriely I. Genetic predisposition to elevated serum thyrotropin is associated with exceptional longevity. *J Clin Endocrinol Metab* (2009) 94(12):4768–75. doi: 10.1210/jc.2009-0808
70. Korevaar TIM, Pop VJ, Chaker L, Goddijn M, de Rijke YB, Bisschop PH, et al. Dose Dependency and a Functional Cutoff for TPO-Antibody Positivity During Pregnancy. *J Clin Endocrinol Metab* (2018) 103(2):778–89. doi: 10.1210/jc.2017-01560
71. Karmisholt J, Laurberg P. Serum TSH and serum thyroid peroxidase antibody fluctuate in parallel and high urinary iodine excretion predicts subsequent thyroid failure in a 1-year study of patients with untreated subclinical hypothyroidism. *Eur J Endocrinol* (2008) 158(2):209–15. doi: 10.1530/EJE-07-0407
72. Chen Y, Chen Y, Wang N, Chen C, Nie X, Li Q, et al. Thyroid Stimulating Hormone Within the Reference Range is Associated with Visceral Adiposity Index and Lipid Accumulation Product: A Population-Based Study of SPECT-China. *Horm Metab Res* (2018) 50(1):29–36. doi: 10.1055/s-0043-122235
73. Diniz M, Beilegoli AMR, Bensenor IM, Lotufo PA, Goulart AC, Barreto SM. Association between TSH levels within the reference range and adiposity markers at the baseline of the ELSA-Brasil study. *PLoS One* (2020) 15(2):e0228801. doi: 10.1371/journal.pone.0228801
74. Neves JS, Castro Oliveira S, Souteiro P, Pedro J, Magalhaes D, Guerreiro V, et al. Effect of Weight Loss after Bariatric Surgery on Thyroid-Stimulating Hormone Levels in Patients with Morbid Obesity and Normal Thyroid Function. *Obes Surg* (2018) 28(1):97–103. doi: 10.1007/s11695-017-2792-5
75. Zantut-Wittmann DE, Quintino-Moro A, Dos Santos PNS, Melhado-Kimura V, Bahamondes L, Fernandes A. Lack of Influence of Thyroid Hormone on Bone Mineral Density and Body Composition in Healthy Euthyroid Women. *Front Endocrinol* (2019) 10:890. doi: 10.3389/fendo.2019.00890
76. Köhrle J. Selenium and the thyroid. *Curr Opin Endocrinol Diabetes Obes* (2015) 22(5):392–401. doi: 10.1097/med.0000000000000190
77. Park SY, Kim HI, Oh HK, Kim TH, Jang HW, Chung JH, et al. Age- and gender-specific reference intervals of TSH and free T4 in an iodine-replete area: Data from Korean National Health and Nutrition Examination Survey IV (2013–2015). *PLoS One* (2018) 13(2):e0190738. doi: 10.1371/journal.pone.0190738
78. Park S, Kim WG, Jeon MJ, Kim M, Oh HS, Han M, et al. Serum thyroid-stimulating hormone levels and smoking status: Data from the Korean National Health and Nutrition Examination Survey VI. *Clin Endocrinol* (2018) 88(6):969–76. doi: 10.1111/cen.13606
79. Nakajima Y, Yamada S, Nishikido A, Katano-Toki A, Ishida E, Akuzawa M, et al. Influence of Smoking on Thyroid Function in Japanese Subjects: Longitudinal Study for One Year of On-Off Smoking. *J Endocr Soc* (2019) 3(12):2385–96. doi: 10.1210/je.2019-00155
80. Zhang Y, Shi L, Zhang Q, Peng N, Chen L, Lian X, et al. The association between cigarette smoking and serum thyroid stimulating hormone, thyroid peroxidase antibodies and thyroglobulin antibodies levels in Chinese residents: A cross-sectional study in 10 cities. *PLoS One* (2019) 14(11):e0225435. doi: 10.1371/journal.pone.0225435
81. Gruppen EG, Kootstra-Ros J, Kobold AM, Connelly MA, Touw D, Bos JHJ, et al. Cigarette smoking is associated with higher thyroid hormone and lower TSH levels: the PREVENT study. *Endocrine* (2020) 67(3):613–22. doi: 10.1007/s12020-019-02125-2
82. Bertelsen JB, Hegedus L. Cigarette smoking and the thyroid. *Thyroid* (1994) 4(3):327–31. doi: 10.1089/thy.1994.4.327
83. Wiersinga WM. Smoking and thyroid. *Clin Endocrinol* (2013) 79(2):145–51. doi: 10.1111/cen.12222
84. Fliers E, Bianco AC, Langouche L, Boelen A. Thyroid function in critically ill patients. *Lancet Diabetes Endocrinol* (2015) 3(10):816–25. doi: 10.1016/s2213-8587(15)00225-9
85. McIver B, Gorman CA. Euthyroid sick syndrome: an overview. *Thyroid* (1997) 7(1):125–32. doi: 10.1089/thy.1997.7.125
86. Van den Berghe G, de Zegher F, Veldhuis JD, Wouters P, Gouwy S, Stockman W, et al. Thyrotrophin and prolactin release in prolonged critical illness: dynamics of spontaneous secretion and effects of growth hormone-secretagogues. *Clin Endocrinol* (1997) 47(5):599–612. doi: 10.1046/j.1365-2265.1997.3371118.x
87. Van den Berghe G, de Zegher F, Bouillon R. Clinical review 95: Acute and prolonged critical illness as different neuroendocrine paradigms. *J Clin Endocrinol Metab* (1998) 83(6):1827–34. doi: 10.1210/jcem.83.6.4763
88. Haugen BR. Drugs that suppress TSH or cause central hypothyroidism. *Best Pract Res Clin Endocrinol Metab* (2009) 23(6):793–800. doi: 10.1016/j.beem.2009.08.003
89. Mughal BB, Fini JB, Demeneix BA. Thyroid-disrupting chemicals and brain development: an update. *Endocr Connect* (2018) 7(4):R160–r86. doi: 10.1530/ec-18-0029
90. Straub RH. Interaction of the endocrine system with inflammation: a function of energy and volume regulation. *Arthritis Res Ther* (2014) 16(1):203. doi: 10.1186/ar4484
91. Bartke A, Sun LY, Longo V. Somatotrophic signaling: trade-offs between growth, reproductive development, and longevity. *Physiol Rev* (2013) 93(2):571–98. doi: 10.1152/physrev.00006.2012
92. van der Weerd K, van Hagen PM, Schrijver B, Heuvelmans SJ, Hofland LJ, Swagemakers SM, et al. Thyrotropin acts as a T-cell developmental factor in mice and humans. *Thyroid* (2014) 24(6):1051–61. doi: 10.1089/thy.2013.0396
93. Van Cauter E. Diurnal and ultradian rhythms in human endocrine function: a minireview. *Horm Res* (1990) 34(2):45–53. doi: 10.1159/000181794
94. Samuels MH. Effects of variations in physiological cortisol levels on thyrotropin secretion in subjects with adrenal insufficiency: a clinical research center study. *J Clin Endocrinol Metab* (2000) 85(4):1388–93. doi: 10.1210/jcem.85.4.6540
95. Re RN, Kourides IA, Ridgway EC, Weintraub BD, Maloof F. The effect of glucocorticoid administration on human pituitary secretion of thyrotropin and prolactin. *J Clin Endocrinol Metab* (1976) 43(2):338–46. doi: 10.1210/jcem-43-2-338
96. Brabant G, Brabant A, Ranft U, Ocran K, Köhrle J, Hesch RD, et al. Circadian and pulsatile thyrotropin secretion in euthyroid man under the influence of thyroid hormone and glucocorticoid administration. *J Clin Endocrinol Metab* (1987) 65(1):83–8. doi: 10.1210/jcem-65-1-83
97. Samuels MH, Luther M, Henry P, Ridgway EC. Effects of hydrocortisone on pulsatile pituitary glycoprotein secretion. *J Clin Endocrinol Metab* (1994) 78(1):211–5. doi: 10.1210/jcem.78.1.8288706
98. Samuels MH. Effects of metyrapone administration on thyrotropin secretion in healthy subjects—a clinical research center study. *J Clin Endocrinol Metab* (2000) 85(9):3049–52. doi: 10.1210/jcem.85.9.6788
99. van der Spoel E, Roelfsema F, Akintola AA, Jansen SW, Slagboom PE, Westendorp RGJ, et al. Interrelationships Between Pituitary Hormones as Assessed From 24-hour Serum Concentrations in Healthy Older Subjects.

- J Clin Endocrinol Metab* (2020) 105(4):e1201–14. doi: 10.1210/clinem/dgz253
100. Bassett JH, Williams GR. Role of Thyroid Hormones in Skeletal Development and Bone Maintenance. *Endocr Rev* (2016) 37(2):135–87. doi: 10.1210/er.2015-1106
 101. van der Spoel E, Oei N, Cachucho R, Roelfsema F, Berbee JFP, Blauw GJ, et al. The 24-hour serum profiles of bone markers in healthy older men and women. *Bone* (2019) 120:61–9. doi: 10.1016/j.bone.2018.10.002
 102. Reed HL. Circannual changes in thyroid hormone physiology: the role of cold environmental temperatures. *Arctic Med Res* (1995) 54 Suppl 2:9–15.
 103. Harford RR, Reed HL, Morris MT, Sapien IE, Warden R, D'Alesandro MM. Relationship between changes in serum thyrotropin and total and lipoprotein cholesterol with prolonged Antarctic residence. *Metabolism* (1993) 42(9):1159–63. doi: 10.1016/0026-0495(93)90274-r
 104. Reed HL, Burman KD, Shakir KM, O'Brian JT. Alterations in the hypothalamic-pituitary-thyroid axis after prolonged residence in Antarctica. *Clin Endocrinol* (1986) 25(1):55–65. doi: 10.1111/j.1365-2265.1986.tb03595.x
 105. Xu C, Zhu G, Xue Q, Zhang S, Du G, Xi Y, et al. Effect of the Antarctic environment on hormone levels and mood of Chinese expeditioners. *Int J Circumpolar Health* (2003) 62(3):255–67. doi: 10.3402/ijch.v62i3.17562
 106. Palinkas LA, Reedy KR, Shepanek M, Smith M, Anghel M, Steel GD, et al. Environmental influences on hypothalamic-pituitary-thyroid function and behavior in Antarctica. *Physiol Behav* (2007) 92(5):790–9. doi: 10.1016/j.physbeh.2007.06.008
 107. Reed HL, Silverman ED, Shakir KM, Dons R, Burman KD, O'Brian JT. Changes in serum triiodothyronine (T3) kinetics after prolonged Antarctic residence: the polar T3 syndrome. *J Clin Endocrinol Metab* (1990) 70(4):965–74. doi: 10.1210/jcem-70-4-965
 108. Azizi F, Mannix JE, Howard D, Nelson RA. Effect of winter sleep on pituitary-thyroid axis in American black bear. *Am J Physiol* (1979) 237(3):E227–30. doi: 10.1152/ajpendo.1979.237.3.E227
 109. Ono H, Hoshino Y, Yasuo S, Watanabe M, Nakane Y, Murai A, et al. Involvement of thyrotropin in photoperiodic signal transduction in mice. *Proc Natl Acad Sci U S A* (2008) 105(47):18238–42. doi: 10.1073/pnas.0808952105
 110. Dardente H, Hazlerigg DG, Ebling FJ. Thyroid hormone and seasonal rhythmicity. *Front Endocrinol* (2014) 5:19. doi: 10.3389/fendo.2014.00019
 111. Nakayama T, Yoshimura T. Seasonal Rhythms: The Role of Thyrotropin and Thyroid Hormones. *Thyroid* (2018) 28(1):4–10. doi: 10.1089/thy.2017.0186
 112. Yoshimura T, Yasuo S, Watanabe M, Iigo M, Yamamura T, Hirunagi K, et al. Light-induced hormone conversion of T4 to T3 regulates photoperiodic response of gonads in birds. *Nature* (2003) 426(6963):178–81. doi: 10.1038/nature02117
 113. Nakao N, Ono H, Yamamura T, Anraku T, Takagi T, Higashi K, et al. Thyrotrophin in the pars tuberalis triggers photoperiodic response. *Nature* (2008) 452(7185):317–22. doi: 10.1038/nature06738
 114. Ikegami K, Yoshimura T. The hypothalamic-pituitary-thyroid axis and biological rhythms: The discovery of TSH's unexpected role using animal models. *Best Pract Res Clin Endocrinol Metab* (2017) 31(5):475–85. doi: 10.1016/j.beem.2017.09.002
 115. Furlow JD, Neff ES. A developmental switch induced by thyroid hormone: *Xenopus laevis* metamorphosis. *Trends Endocrinol Metab* (2006) 17(2):40–7. doi: 10.1016/j.tem.2006.01.007
 116. Gudernatsch JF. Feeding experiments on tadpoles. *Arch Entwickl Mech Org* (1912) 35:457–83. doi: 10.1007/BF02277051
 117. Tabatabaie V, Surks MI. The aging thyroid. *Curr Opin Endocrinol Diabetes Obes* (2013) 20(5):455–9. doi: 10.1097/01.med.0000433055.99570.52
 118. Barbesino G. Thyroid Function Changes in the Elderly and Their Relationship to Cardiovascular Health: A Mini-Review. *Gerontology* (2019) 65(1):1–8. doi: 10.1159/000490911
 119. Roberts L, McCahon D, Johnson O, Haque MS, Parle J, Hobbs FR. Stability of thyroid function in older adults: the Birmingham Elderly Thyroid Study. *Br J Gen Pract* (2018) 68(675):e718–e26. doi: 10.3399/bjgp18X698861
 120. Diez JJ, Iglesias P. Spontaneous subclinical hypothyroidism in patients older than 55 years: an analysis of natural course and risk factors for the development of overt thyroid failure. *J Clin Endocrinol Metab* (2004) 89(10):4890–7. doi: 10.1210/jc.2003-032061
 121. Diez JJ, Iglesias P, Burman KD. Spontaneous normalization of thyrotropin concentrations in patients with subclinical hypothyroidism. *J Clin Endocrinol Metab* (2005) 90(7):4124–7. doi: 10.1210/jc.2005-0375
 122. Andersen S, Pedersen KM, Bruun NH, Laurberg P. Narrow individual variations in serum T(4) and T(3) in normal subjects: a clue to the understanding of subclinical thyroid disease. *J Clin Endocrinol Metab* (2002) 87(3):1068–72. doi: 10.1210/jcem.87.3.8165
 123. Taylor PN, Razvi S, Pearce SH, Dayan CM. Clinical review: A review of the clinical consequences of variation in thyroid function within the reference range. *J Clin Endocrinol Metab* (2013) 98(9):3562–71. doi: 10.1210/jc.2013-1315
 124. Waise A, Price HC. The upper limit of the reference range for thyroid-stimulating hormone should not be confused with a cut-off to define subclinical hypothyroidism. *Ann Clin Biochem* (2009) 46(Pt 2):93–8. doi: 10.1258/acb.2008.008113
 125. Hoermann R, Midgley JE, Larisch R, Dietrich JW. Homeostatic Control of the Thyroid-Pituitary Axis: Perspectives for Diagnosis and Treatment. *Front Endocrinol* (2015) 6:177. doi: 10.3389/fendo.2015.00177
- Conflict of Interest:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2021 van der Spoel, Roelfsema and van Heemst. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.