



Universiteit
Leiden
The Netherlands

SATB1, genomic instability and Gleason grading constitute a novel risk score for prostate cancer

Dumke, C.; Gemoll, T.; Oberlander, M.; Freitag-Wolf, S.; Thorns, C.; Glaessgen, A.; ... ; Habermann, J.K.

Citation

Dumke, C., Gemoll, T., Oberlander, M., Freitag-Wolf, S., Thorns, C., Glaessgen, A., ... Habermann, J. K. (2021). SATB1, genomic instability and Gleason grading constitute a novel risk score for prostate cancer. *Scientific Reports*, 11(1).
doi:10.1038/s41598-021-03702-0

Version: Publisher's Version
License: [Creative Commons CC BY 4.0 license](#)
Downloaded from: <https://hdl.handle.net/1887/3256608>

Note: To cite this publication please use the final published version (if applicable).



OPEN

SATB1, genomic instability and Gleason grading constitute a novel risk score for prostate cancer

Christoph Dumke^{1,10}, Timo Gemoll^{1,10}, Martina Oberländer^{1,2}, Sandra Freitag-Wolf³, Christoph Thorns⁴, Axel Glaessgen⁵, Rinse Klooster⁶, Silvere M. van der Maarel⁶, Jerker Widengren⁷, Christian Doehn⁸, Gert Auer⁹ & Jens K. Habermann^{1,2,9}✉

Current prostate cancer risk classifications rely on clinicopathological parameters resulting in uncertainties for prognostication. To improve individual risk stratification, we examined the predictive value of selected proteins with respect to tumor heterogeneity and genomic instability. We assessed the degree of genomic instability in 50 radical prostatectomy specimens by DNA-Image-Cytometry and evaluated protein expression in related 199 tissue-microarray (TMA) cores. Immunohistochemical data of SATB1, SPIN1, TPM4, VIME and TBB5 were correlated with the degree of genomic instability, established clinical risk factors and overall survival. Genomic instability was associated with a GS ≥ 7 ($p = 0.001$) and worse overall survival ($p = 0.008$). A positive SATB1 expression was associated with a GS ≤ 6 ($p = 0.040$), genomic stability ($p = 0.027$), and was a predictor for increased overall survival ($p = 0.023$). High expression of SPIN1 was also associated with longer overall survival ($p = 0.048$) and lower preoperative PSA-values ($p = 0.047$). The combination of SATB1 expression, genomic instability, and GS lead to a novel Prostate Cancer Prediction Score (PCP-Score) which outperforms the current D'Amico et al. stratification for predicting overall survival. Low SATB1 expression, genomic instability and GS ≥ 7 were identified as markers for poor prognosis. Their combination overcomes current clinical risk stratification regimes.

Abbreviations

AEC	Aminoethylcarbazole
CI	Confidence Interval
DAB	Diaminobenzidine
EMT	Epithelial-Mesenchymal-Transition
GS	Gleason Score
HE	Hematoxylin and Eosin
IRS	Immunoreactive Score
OS	Overall Survival
PCP-Score	Prostate Cancer Prediction Score
PSA	Prostate-Specific-Antigen
SATB1	Special-AT-rich-binding-Protein-1

¹Section for Translational Surgical Oncology and Biobanking, Department of Surgery, University of Lübeck and University Hospital Schleswig-Holstein, Campus Lübeck, Ratzeburger Allee 160, 23538 Lübeck, Germany. ²Interdisciplinary Center for Biobanking-Lübeck (ICB-L), University of Lübeck, Lübeck, Germany. ³Institute of Medical Informatics and Statistics, University Hospital Schleswig-Holstein, Campus Kiel, Kiel, Germany. ⁴Institute of Pathology, University of Lübeck and University Hospital Schleswig-Holstein, Campus Lübeck, Lübeck, Germany. ⁵Department of Clinical Pathology and Cytology, Unilabs AB, Stockholm, Sweden. ⁶Department of Human Genetics, Leiden University Medical Center, Leiden, The Netherlands. ⁷Department of Applied Physics, Royal Institute of Technology (KTH), Stockholm, Sweden. ⁸Urologikum Lübeck, Lübeck, Germany. ⁹Department of Oncology-Pathology, Karolinska Institutet, Stockholm, Sweden. ¹⁰These authors contributed equally: Christoph Dumke and Timo Gemoll. ✉email: Jens.Habermann@uni-luebeck.de

SPIN1	Spindlin-1
TBB5	β -Tubulin
TPM4	Tropomyosin-4
TMA	Tissue-Microarray
VIME	Vimentin

With 1,276,106 new cases in 2018, prostate cancer is the second most frequent malignant disease among males worldwide¹. Further, it remains the sixth leading cause of malignancy associated death¹. In clinical practice, the classification according to D'Amico containing preoperative PSA serum level, biopsy-based Gleason Score (GS) and clinical tumor stage is routinely used to distinguish between high, intermediate and low risk for PSA recurrence after radical prostatectomy (RPE), external radiotherapy or brachytherapy². More contemporary validation studies of the D'Amico classification confirmed its ability to predict patients' risk of PSA recurrence after RPE and death from prostate cancer^{3,4}. However, preoperatively classified low risk patients showed heterogeneous histology and were frequently upgraded or upstaged at final examination after RPE⁵. Furthermore, intermediate and high risk patients still remain heterogeneous groups with different prognosis depending on whether clinical tumor stage or biopsy-based GS are considered⁶. Thus, this classification can fail resulting in possible over- or undertreatment^{6,7}. Improved prognostication for individualized therapy can be achieved by assessment of intra-tumor heterogeneity on the molecular level which is underrepresented in D'Amico's classification but yet an important trait of solid tumors and associated with genomic instability^{8,9}. Consequently, there is an unmet opportunity to account and address tumor biology more precisely. For this purpose many genes and their products have been evaluated as potential biomarkers for prostate cancer¹⁰. Our literature review revealed special-AT-rich-protein-1 (SATB1), Spindlin-1 (SPIN1), Vimentin (VIME), β -Tubulin (TBB5), and Tropomyosin-4 (TPM4) to be the most promising biomarkers^{11–15}.

SATB1 is a nuclear chromatin organizer regulating the expression of genes involved in cell differentiation and apoptosis¹⁶. SATB1 seems to be upregulated in malignant tumors of the breast, colon, and bladder cancer causing aggressive tumor growth and limited prognosis^{17–19}. SATB1 levels also appear to be increased in prostate cancer with implications for tumor progression and metastasis through Epithelial-Mesenchymal-Transition (EMT)^{12,20}.

SPIN1 represents a nuclear factor that associates with the spindle apparatus during meiosis and is highly expressed in early embryonic tissues²¹. Moreover, its levels were found to be increased in ovarian cancer and liposarcoma^{13,22}.

TPM4, VIME, and TBB5 are dynamic parts of the cytoskeleton and play an important role in cell motility and metastasis^{11,23,24}. Despite their tumor biology functions, neither SATB1, SPIN1, VIME, TBB5 nor TPM4 are yet being used in routine risk assessment for prostate cancer. Therefore, we examined their protein expression at multiple tumor sites to address potential intra-tumor heterogeneity^{8,25} and correlated their protein expression to the degree of genomic instability, established clinical risk factors, as well as overall survival.

Material and methods

Patient collective. Formalin fixed and paraffin embedded prostate cancers were obtained from 50 patients that had undergone radical prostatectomy at the University Hospital Schleswig–Holstein, Campus Lübeck between 1989 and 1992. After histological confirmation, all operations were carried out using the open-surgery technique with diagnostic lymphadenectomy. In case of PSA values ≥ 10 ng/ml a preoperative bone scintigraphy was performed in order to exclude bone metastases. Neoadjuvant chemo- and/or antihormonal therapy were not administered. Postoperative treatment was performed in an outpatient setting and is—retrospectively—not retrievable. However, it can be envisioned that patients at that time (1989–1992) were treated comparably. The median follow-up was 15 years (range 1–22 years). Survival data was obtained from the residences' registration offices. All clinical and pathological data are summarized in Table 1. Risk groups were assessed as follows: low risk (T1–T2a, Gleason Score (GS) ≤ 6 , and PSA < 10 ng/ml), intermediate risk (T2b–T2c or GS 7 or PSA 10–20 ng/ml), and high risk ($\geq T3$ or GS 8–10 or PSA > 20 ng/ml). However, despite D'Amico's original classification being based on preoperative biopsy-based GS and clinical stage, only pathological GS and pathological stage were available for this study. The GS was retrospectively re-evaluated by one senior pathologist (C.T.) according to the recommendations of the International Society of Urological Pathology 2005 using 4 μ m Hematoxylin and Eosin (HE) stained tissue sections²⁶. The study was conducted according to the guidelines of the Declaration of Helsinki. All prostate cancer samples were collected within clinical routine diagnostics at the University Hospital Schleswig–Holstein in Lübeck, Germany, between 1989 and 1992. At that time, no informed consent for research purposes was obtained. The study was approved by the local Ethics Committee of the University of Lübeck which also approved the use of anonymized samples and anonymized data for this retrospective study without contacting the patients (#08–011 and #20–508).

Tissue microarray (TMA). The tissue microarray was constructed as described²⁷. Tissue areas containing cancer and benign cells were marked by one senior pathologist (C.T.). Subsequently, 1.5 mm cores of both, cancer and benign tissue from each patient, were received by a TMArrayer™ (Pathology Devices, Inc. Westminster, USA) and built into a new custom-made TMA. Based on previous reports on optimal tissue representativity for heterogeneous tumors²⁵, we decided to obtain four tissue cores from different sites of the marked tumor areas and one core from benign tissue for each patient. Thus, 199 tumor and 48 benign tissue cores were distributed onto two TMA blocks. Benign tissue was not available for two patients. Tissue integrity and histology were examined after HE-staining.

Parameter	n	%
Age (median)		
≤ 63	27	54
> 63	23	46
Tumor stage		
≤ pT2b	13	26
pT3–pT4	35	70
Not available	2	4
Lymph node stage		
pN0	47	94
pN1	1	2
Not available	2	4
Metastasis		
cM0	41	82
cM1	0	0
Not available	9	18
Gleason Score		
4–6	24	48
7	16	32
8–10	10	20
Preoperative PSA [ng/ml]		
< 10	12	24
10–20	16	32
> 20	11	22
Not available	11	22
Postoperative survival after 22 years		
Alive	15	30
Dead	33	66
Not available	2	4
Genomic stability		
Stable	32	64
Unstable	16	32
Not available	2	4

Table 1. Patients data.

Assessing the degree of genomic instability. Genomic instability was assessed by DNA image cytometry using 8 µm Feulgen stained whole tissue sections of each tumor. Two patients lacked enough material for DNA image cytometry. Cell selection criteria, quantitative measurement of nuclear DNA content, and internal standardization were based on methods described²⁸. Detection of representative tumor cell nuclei on Feulgen stained tissue sections was performed using marked tumor areas on corresponding HE slides using an ICM imaging system (Ahrens ICM Cytometry System, Hamburg, Germany). The DNA content of at least 100 nuclei per specimen was quantitatively measured and expressed in relation to the DNA content of lymphocytes as reference which was given the value 2c reflecting a physiological diploid DNA content. The resulting DNA histograms were classified according to Auer et al.^{28,29} (Supplemental Data 1, Figure S1).

Immunohistochemistry. After immunohistochemical staining (Supplemental Data 2), scoring was performed semiquantitatively by one independent senior pathologist (A.G.) according to the immunoreactive Score (IRS)³⁰ (Figure S2–S5). For VIME, an alternative score regarding the percentage of positive tumor cells was used: Score 0—negative staining of all tumor cells; Score 1—up to 20% stained tumor cells; Score 2—21 to 50% stained tumor cells; Score 3—more than 50% stained tumor cells (Figure S6). A high expression was defined as IRS 6–12 for SATB1 and TPM4, and as IRS 9–12 for SPIN1 and TBB5. For VIME a tissue core was assessed for high expression when showing a Score of 1–3. A tumor was designated as positive when at least one of the four obtained tissue cores showed a high expression for the evaluated protein, whereas a low or negative expression in all four tissue cores defined a tumor as negative. To determine intra-tumor heterogeneity, we additionally evaluated all TMA tissue cores per patient. Homogenous expression was declared if all cores per patient showed either only high or only low protein expression. In contrast, in-homogenous expression was assessed with two independent approaches: we (i) accounted for the number of tissue cores with a high protein expression and (ii) considered the expression of only one randomly chosen tissue core (random).

No. of patients (%)	SATB1 N = 50	SPIN1 N = 50	TBB5 N = 50	VIME N = 50	TPM4 N = 50
Homogeneous expression 4 of 4 tissue cores high or low	26 (52%)	39 (78%)	26 (52%)	43 (86%)	43 (46%)
Heterogeneous expression in total	24 (48%)	11 (22%)	24 (48%)	7 (14%)	7 (14%)
3 of 4 tissue cores high	7	0	5	0	1
2 of 4 tissue cores high	8	2	5	0	0
1 of 4 tissue cores high	9	9	14	7	6

Table 2. Tumor heterogeneity in protein expression.

Statistics. Statistical analysis was performed using SPSS version 21 (IBM Corporation, Somers, NY, USA). Continuous variables were dichotomized or categorized based on clinical and pathological parameters. Associations between categorical variables were assessed by Fisher's exact test, and two-sided P -values < 0.05 were considered to be statistically significant.

For the analysis of the survival times, Kaplan Meier estimators, log-rank tests and age-adjusted cox regression models were used to identify potential risk factors. Their impact upon patients' mean overall survival (OS) was expressed as hazard ratio (HR) provided with 95% confidence intervals (CI). For multivariable analysis, we used the Cox proportional hazard model and included the established prognostic factors tumor stage, preoperative PSA, Gleason Score, as well as prognostic factors and their combinations that were significant in univariable Cox regression (SATB1 positive, SATB1 ≥ 2 cores, GS ≥ 7 or genomic instability, GS ≥ 7 or SATB1 negative, GS ≥ 7 or SATB1 ≤ 1 core, genomic instability or SATB1 negative, genomic instability or SATB1 ≤ 1 core, GS/genomic instability/SATB1 ≤ 1 core; Supplemental Table S1) as well as the new PCP-Score. All possible predictors were selected on the likelihood ratio criteria in the following backward selection. Furthermore, the prediction of the new Score (PCP) was analyzed in the strata with a low and a high number of SATB1 expressing tissue cores (≤ 1 vs. ≥ 2) in order to investigate possible interactions. Beyond this, a model with and without genomic instability was fitted for a comparison of the prediction performance. For all models the proportional hazard assumption was checked using weighted residuals and none of the prognostic factors were found to violate this assumption.

Ethics approval and consent to participate. Approval from the local ethics committee of the University of Lübeck (#08-011 and #20-508).

Results

Associations of clinical parameters with survival. The mean overall survival (OS) of the entire patient collective was 14.3 years (CI, 12.3–16.2).

A low Gleason Score (GS) of 4–6 was found in 24 cases (48%), whereas 26 (52%) showed a GS of 7 or higher. Patients with a GS of 4–6 had a longer OS of 17.9 years (CI, 15.9–20) compared to 11 years (CI, 8.5–13.6) for patients with a GS of 7–10 ($p = 0.002$, Table S1). Furthermore, the GS was a significant predictor of OS in univariable cox regression analysis (age-adjusted HR 3.394, CI 1.615–7.132, $p = 0.001$, Table S1). Neither pathological stage nor preoperative PSA value showed a significant association with OS (Table S1).

Detecting the degree of genomic instability. Overall, 16 of 48 (33.3%) tumors showed nuclear DNA aneuploidy denoting genomic instability. Patients with euploid, genomically stable tumor cell populations had a longer OS with 16.5 years (CI, 14.3–18.8) compared to aneuploid, genomically unstable tumors with 10.8 years (CI, 7.6–14) ($p = 0.006$, Table S1, Figure S7). Furthermore, we could show that genomic instability was significantly associated with a higher tumor grade: 14 tumors (56%) with GS 7–10 showed genomic instability in contrast to only 2 (8.7%) of GS 4–6 carcinomas ($p = 0.001$, Table S2). Patients with genomically unstable tumor cell populations showed a 2.7-fold increased mortality risk in univariable analysis (age-adjusted HR 2.672, CI 1.294–5.515, $p = 0.008$, Table S1).

Determining intra-tumor heterogeneity by protein expression. Extensive heterogeneity of up to 48% was observed for the expression of SATB1 and TBB5 (Table 2). For SATB1, nuclear and cytoplasmic immunoreactivity occurred in epithelial tumor cells or normal epithelium of the prostate but not in stromal tissue cells (Figure S2). While 39 (78%) carcinomas were SATB1 positive, only 11 (22.9%) benign tissue samples were strongly immunoreactive for SATB1 ($p < 0.001$, Table 3). SATB1 positive patients had a longer OS of 15.5 years (CI, 13.3–17.6) compared to 10.3 years (CI, 6.5–14) for SATB1 negative ones ($p = 0.020$, Table S1, Fig. 1A). In line, a positive SATB1 expression was associated with a GS of 4–6 ($p = 0.040$, Table 3) and genomic stability ($p = 0.027$, Table 3).

Further, patients with ≥ 2 tissue cores with high SATB1 expression had an OS of 16.3 years (CI, 14.1–18.5) compared to 11.1 years for patients with only ≤ 1 high SATB1 tissue cores (CI, 8.0–14.2) ($p = 0.018$, Table S1, Fig. 1B). Interestingly, age adjusted cox regression analysis showed that SATB1 positive patients (at least one tissue core with a high SATB1 expression) and patients with ≥ 2 tissue cores with high SATB1 expression had a significantly lower risk of dying during the observation period (age-adjusted HR 0.413, CI 0.192–0.886, $p = 0.023$ & HR 0.479, CI 0.238–0.965, $p = 0.039$, Table S1). No significant results were obtained when SATB1 expression was evaluated in only one tissue core selected at random from each tumor ($p = 0.068$, Fig. 1C,D).

	Gleason Score [N = 50]			Tumor stage [N = 48]			PSA (ng/ml) [N = 39]			Ploidy [N = 48]			Histopathology [N = 98]**		
Parameter	4–6	7–10	P*	≤ T2b	≥ T3	P*	≤ 10	> 10	P*	Genomically stable	Genomically instable	P*	Benign	Cancer	P*
SATB1															
Positive	22 (91.7%)	17 (65.4%)	0.040	10 (76.9%)	27 (77.1%)	1.000	12 (85.7%)	17 (68%)	0.279	28 (87.5%)	9 (56.2%)	0.027	11 (22.9%)	39 (78%)	< 0.001
Negative	2 (8.3%)	9 (34.6%)		3 (23.1%)	8 (22.9%)		2 (14.3%)	8 (32%)		4 (12.5%)	7 (43.8%)		37 (77.1%)	11 (22%)	
SPIN1															
Positive	7 (29.2%)	4 (15.4%)	0.314	3 (23.1%)	8 (22.9%)	1.000	6 (42.9%)	3 (12%)	0.047	8 (25%)	3 (18.7%)	0.729	3 (6.4%)	11 (22%)	0.042
Negative	17 (70.8%)	22 (84.6%)		10 (76.9%)	27 (77.1%)		8 (57.1%)	22 (88%)		24 (75%)	13 (81.3%)		44 (93.6%)	39 (78%)	
TPM4															
Positive	7 (29.2%)	2 (7.7%)	0.069	3 (23.1%)	6 (17.1%)	0.687	3 (21.4%)	4 (16%)	0.686	7 (21.9%)	2 (12.5%)	0.697	1 (2.1%)	9 (18%)	0.016
Negative	17 (70.8%)	24 (92.3%)		10 (76.9%)	29 (82.9%)		11 (78.6%)	21 (84%)		25 (78.1%)	14 (87.5%)		47 (97.9%)	41 (82%)	
VIME															
Positive	3 (12.5%)	7 (26.9%)	0.294	2 (15.4%)	7 (20%)	1.000	5 (35.7%)	2 (8%)	0.075	5 (15.6%)	4 (25%)	0.457	20 (42.5%)	10 (20%)	0.027
Negative	21 (87.5%)	19 (73.1%)		11 (84.6%)	28 (80%)		9 (64.3%)	23 (92%)		27 (84.4%)	12 (75%)		27 (57.5%)	40 (80%)	
TBB5															
Positive	16 (66.7%)	11 (42.3%)	0.098	7 (53.8%)	19 (54.3%)	1.000	5 (35.7%)	15 (60%)	0.191	22 (68.7%)	4 (25%)	0.006	10 (20.8%)	27 (54%)	0.001
Negative	8 (33.3%)	15 (57.7%)		6 (56.2%)	16 (45.7%)		9 (64.3%)	10 (40%)		10 (31.3%)	12 (75%)		38 (79.2%)	23 (46%)	

Table 3. Significant associations between proteins and clinicopathological parameters. *P-Value for Fisher's Exact Test. **Because of tissue loss during staining N = 97 for SPIN1 and VIME. Significant *p*-values are highlighted in bold.




Figure 1. Overall Survival stratified by (A) at least one tissue core with a high SATB1 expression per tumor (SATB1 positive), *p* = 0.02; (B) the number of tissue cores with a high SATB1 expression per tumor, *p* = 0.018; (C) and the SATB1 expression in one randomly chosen tissue core per tumor, *p* = 0.068; (D) TMA slide with SATB1 stained tissue cores (one TMA tissue core enlarged). *P*-values for the log-rank test.




Figure 2. Overall Survival stratified according to (A) D'Amico classification: low and intermediate risk (GS ≤ 7 and PSA ≤ 20 ng/ml and $\leq$ T2b) versus high risk (GS > 7 or PSA > 20 ng/ml or $\geq$ T3), $p = 0.285$ and (B) new PCP-Score: low risk (GS ≤ 6 and genomic stability and high SATB1 expression in at least one tissue core) versus high risk (GS ≥ 7 or genomic instability or low/negative SATB1 expression in all four obtained cores), $p = 0.001$; P -values for the log-rank test.

For SPIN1, TBB5, VIME and TPM4, significance in their cytoplasmic staining could be detected between cancer and normal epithelium cells: while VIME had higher protein levels in benign prostate tissue ($p = 0.027$, Table 3), SPIN1, TBB5 and TPM4 showed higher expression in cancer tissue ($p = 0.042$, $p = 0.001$ and $p = 0.016$, respectively, Table 3). Furthermore, patients with genomically stable tumor cell populations had a higher TBB5 expression ($p = 0.006$, Table 3). However, only SPIN1 positivity was associated with a low preoperative PSA value (Table 3) and a longer OS (Figure S7), but none of these proteins were significant predictors of OS in univariable cox regression analysis.


Developing a new prediction score. Grouping our patients according to D'Amico et al., six patients were at low or intermediate risk, whereas 39 were at high risk. Interestingly, this classification failed in predicting postoperative OS ($p = 0.285$, Fig. 2A).

Our findings prompted us to combine all risk factors with a strong impact on OS, namely SATB1 expression, degree of genomic instability and GS for a new Prostate Cancer Prediction Score (PCP-Score). While low risk tumors were defined by SATB1 positivity (high SATB1 expression in at least one tissue core), genomic stability and a GS ≤ 6 , high risk patients were either SATB1 negative or showed genomic instability or had a GS ≥ 7 . From the present collective 47 patients could be stratified using this new PCP-Score, of whom 30 were assigned as high risk and 17 as low risk tumors. Patients with high risk tumors showed a significant shorter OS of 11.7 years (CI, 9.4–14) compared to 19 years (CI, 16.9–21.1) for low risk patients ($p = 0.001$, Fig. 2B) and a fourfold increased risk of dying in age adjusted univariable Cox regression analysis (age-adjusted HR 4, CI 1.701–9.407, $p = 0.001$, Table S1). Hazard ratios for the risk factors and their combinations are shown in Fig. 3. Indeed, this new score was the strongest independent predictor for OS in a multivariable model including the established prognostic factors tumor stage, preoperative PSA, Gleason Score and combinations of prognostic factors that were significant in univariable Cox regression (HR 3.277, CI 1.182–9.087, $p = 0.023$). In order to ascertain the contribution of genomic instability to the predictive value of the new score, we fitted a model with and without genomic instability yielding a noticeable gain in the score when including genomic instability (Table S3). The statistical performance decreased when immunopositivity was defined by ≥ 2 tissue cores with high SATB1 expression ($p = 0.010$, Table S1, Fig. 3).

When considering only patients with ≤ 1 tissue core with high SATB1 expression ($n = 19$), the age adjusted HR for the PCP-Score was 5.346 ($p = 0.036$, CI 1.112–25.695) whereas it was 3.045 ($p = 0.042$, CI 1.042–8.902) for patients with ≥ 2 tissue cores with high SATB1 expression. Notably, grouping patients according to GS, genomic instability and SATB1 expression in one randomly chosen tissue core was not predictive for OS (Fig. 3).

Discussion

An accurate estimation of possible tumor progression and survival is of unmet clinical need in order to offer individualized therapy. In this context, differences in protein expression (SATB1, SPIN1, TBB5, VIME, TPM4) between benign prostate tissue and prostate cancer with respect to clinicopathological risk factors, tumor heterogeneity and genomic instability were assessed. In publicly available databases (proteinatlas.org) the expression of SATB1, TBB5 and VIME is stated as being moderate to weak in prostate cancer and for SPIN1 and TPM4 as strong to weak depending on the antibody used³¹. These results are generally comparable to ours, however, due to missing information on tumor grade and clinical parameters in the human protein atlas a validation of our results is not fully possible.



* PCP-Score: Independent predictor in multivariable cox regression analysis including established risk factors as well as significant parameters in univariable analysis.

Figure 3. Age-adjusted hazard ratios of univariable cox regression analysis for all investigated parameters including combined parameters and the new Prostate Cancer Prediction Score (PCP). Positive = at least one tissue core with a high protein expression. SATB1 ≥ 2 cores = ≥ 2 tissue cores with a high protein expression. SATB1 ≤ 1 core = ≤ 1 tissue core with a high SATB1 expression. SATB1 random = high protein expression in only one randomly chosen tissue core. Red line: Hazard Ratio of new PCP-Score.

Evaluating aneuploidy has emerged as a means to predict aggressiveness and the clinical course of (prostate) cancer^{32–34}. We could show that tumor aneuploidy reflecting genomic instability was strongly associated with a GS ≥ 7 indicating a higher frequency in dedifferentiated prostate cancer tissue with worse prognosis. Furthermore, we found a significant association of genomic instability and a decreased OS. This is in line with findings that aneuploid radical prostatectomy specimens show a strong correlation with higher tumor grade and shorter recurrence free survival^{35,36}. Nevertheless, detection of aneuploidy in one randomly chosen tissue section of a tumor is not predictive for disease recurrence whereas multiple sampling denoted aneuploidy as a prognostic factor⁸. In this context, it is important that underestimation of the tumor's fate due to tumor heterogeneity is not only minimized by taking multiple samples but also by combining the assessment of genomic instability and GS⁸. This is consistent with our results showing that not a single parameter alone but the combination of GS, SATB1 expression, and genomic instability has the strongest predictive power.

While TBB5, TPM4 and VIME only showed differential expression between normal and cancerous tissue, high expression of SATB1 and SPIN1 was also significantly associated with good prognosis.

SATB1 is a global chromatin organizer that regulates gene expression over long distances by chromatin looping and recruitment of chromatin remodeling enzymes^{16,37}. We found it to be elevated in prostate cancer tissue compared to benign tissue and associated with longer OS for patients with high expression. These results were in accordance with the human protein atlas, where immunohistochemical SATB1 expression is reported as negative to weak in normal prostate tissue and as weak to moderate in prostate cancer, though higher than in normal tissue³¹. We could therefore achieve comparable results as this publicly available database which reflects the representativity of our study cohort and methods. Additionally, the proteindatlas.org indicates that the loss of SATB1 or weak expression in tumor tissue was associated with worse overall survival for renal cell carcinoma and pancreatic cancer, while a higher expression is associated with longer overall survival³¹. These results also coincide with our findings concerning prostate cancer. However, our results were in contrast to other studies in which SATB1 overexpression correlated to aggressive tumor biology and metastasis in prostate cancer^{12,20,38}. SATB1 has been suggested to promote tumor growth and metastasis by upregulation of MMP2 and Vimentin with simultaneous downregulation of E-Cadherin as hallmarks for EMT^{39,40}. However, and in accordance with our results, increased SATB1 protein levels also correlated with a better prognosis in lung and colorectal cancer^{41,42}. Furthermore, there is evidence that SATB1 overexpression did not promote breast cancer progression and was associated with a benefit in disease-free survival in estrogen receptor positive breast cancer patients^{43,44}.

SATB1 is a highly phosphorylated and acetylated protein⁴⁵ so that controversial expression differences may be explained by posttranslational modifications⁴⁶. Additionally, SATB1 molecules showed a wide spatial variation in their distribution in thymocyte cell nuclei, indicating that the majority of SATB1 molecules are highly dynamic with varying ability to access different regions of the nucleus⁴⁷. Super-resolution stimulated emission depletion (STED) imaging might identify distinct spatial distribution patterns of SATB1 in tumor cell nuclei

which could further explain its function and prognostic value⁴⁸. However, the present study suggests a positive SATB1 expression when this expression was detected in the cytoplasm and/or the cell nucleus of glandular tumor cells which was in accordance with the human protein atlas³¹.

Additionally, well differentiated proliferating cancer cells might express SATB1 in contrast to benign tissue or poorly differentiated, aneuploid cancer cells. Especially the loss of SATB1 could explain a deregulated chromatin organization and thus favor aneuploid, genomically unstable cell populations.

Furthermore, prostate cancer is a highly heterogenic disease characterized by different molecular traits across different cell populations within one tumor⁴⁹. These facts coincide with the frequent incidence of genomic instability and the high degree of variability in the protein expression of SATB1 found in this study. Only one single biopsy might therefore neither be representative nor sufficient for diagnostics⁸. This prompted us to evaluate the prognostic value of SATB1 by choosing either one randomly chosen tissue core, counting if at least one of four tissue cores shows high SATB1 expression, or using the number of tissue cores with high SATB1 expression. Particularly, we could show that the expression of SATB1 in only one randomly chosen tissue core cannot predict OS, whereas counting the number of tissue cores with a high SATB1 expression or defining a tumor as SATB1 positive when at least one of four tissue cores per patient showed a high SATB1 expression could reveal SATB1 as an predictor for longer OS. Therefore, our results support a multiple sampling strategy to address the diagnostic challenge of intra-tumor heterogeneity.

Most notably, the parameters SATB1 (when assessed by multiple samples from different tumor sites), pathological GS and genomic instability could be combined as a new Prostate Cancer Prediction Score (PCP-Score) which is a significant predictor for OS. By differentiating high risk patients from low risk patients with the novel PCP-Score, long term survivors with a mean survival of 19 years could be separated from patients with shorter postoperative survival of 11.7 years after radical prostatectomy. Interestingly, grouping our patients based on D'Amico's classification for risk of PSA recurrence showed no significant association with overall survival.

However, limitations of our study are the small sample size and that we used stored cancer tissue of a retrospective cohort after radical prostatectomy without information on disease free survival. It has also to be noted though that the patients evaluated here had a mean age of 63 years which is comparably younger compared to the average age of 72 years for the occurrence of prostate cancer⁵⁰. While we are not aware of any potential bias, we cannot exclude that our results might be specifically benefitting a younger age group.

Although this PCP-Score was built on postoperative data after radical prostatectomy, it will likely also predict disease progression in preoperative settings: For SATB1 the pretreatment situation was simulated by taking four TMA cores of representative tumor areas comparable to needle biopsies at initial diagnosis. Moreover, a good correlation between the ploidy status of preoperative needle biopsy and postoperative radical prostatectomy specimen was shown⁵¹. We assume that especially in preoperative biopsy specimens with an underestimated GS⁷, also assessing SATB1 expression and the degree of genomic instability could be of high value in predicting clinical outcome. However, for implementation of the new Score in preoperative routine diagnostics, large validation studies using external datasets and patient material from preoperative biopsies would be necessary.

In conclusion, we demonstrated that our Score containing SATB1 expression, examined by immunohistochemical analysis in a multiple sample approach, combined with genomic instability and GS is a predictor for prognosis in prostate cancer potentially outperforming risk stratifications in current use. Further multi-center validation experiments including the evaluation of disease-free survival are warranted.

Data availability

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Received: 25 May 2021; Accepted: 23 November 2021

Published online: 27 December 2021

References

- Bray, F. *et al.* Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J. Clin.* **68**(6), 394–424. <https://doi.org/10.3322/caac.21492> (2018).
- D'Amico, A. V. *et al.* Biochemical outcome after radical prostatectomy, external beam radiation therapy, or interstitial radiation therapy for clinically localized prostate cancer. *JAMA* **280**(11), 969–974. <https://doi.org/10.1001/jama.280.11.969> (1998).
- Hernandez, D. J., Nielsen, M. E., Han, M. & Partin, A. W. Contemporary evaluation of the D'Amico risk classification of prostate cancer. *Urology* **70**(5), 931–935. <https://doi.org/10.1016/j.urol.2007.08.055> (2007).
- Boorjian, S. A., Karnes, R. J., Rangel, L. J., Bergstralh, E. J. & Blute, M. L. Mayo clinic validation of the D'Amico risk group classification for predicting survival following radical prostatectomy. *J. Urol.* **179**(4), 1354–1361. <https://doi.org/10.1016/j.juro.2007.11.061> (2008).
- Schiffmann, J. *et al.* Heterogeneity in D'Amico classification-based low-risk prostate cancer: Differences in upgrading and upstaging according to active surveillance eligibility. *Urol. Oncol. Semin. Orig. Investig.* **33**(7), 329.e13. <https://doi.org/10.1016/j.urolonc.2015.04.004> (2015).
- Reese, A. C., Pierorazio, P. M., Han, M. & Partin, A. W. Contemporary evaluation of the national comprehensive cancer network prostate cancer risk classification system. *Urology* **80**(5), 1075–1079. <https://doi.org/10.1016/j.urol.2012.07.040> (2012).
- Heidegger, I. *et al.* High Risk of under-grading and -staging in prostate cancer patients eligible for active surveillance. *PLoS ONE* **10**(2), e0115537. <https://doi.org/10.1371/journal.pone.0115537> (2015).
- Cyll, K. *et al.* Tumour heterogeneity poses a significant challenge to cancer biomarker research. *Br. J. Cancer* <https://doi.org/10.1038/bjc.2017.171> (2017) (Published online June 15, 2017).
- Sansregret, L. & Swanton, C. The role of aneuploidy in cancer evolution. *Cold Spring Harb. Perspect. Med.* **7**(1), a028373. <https://doi.org/10.1101/cshperspect.a028373> (2017).
- Esfahani, M., Ataei, N. & Panjehpour, M. Biomarkers for evaluation of prostate cancer prognosis. *Asian Pac. J. Cancer Prev.* **16**(7), 2601–2611. <https://doi.org/10.7314/APJCP.2015.16.7.2601> (2015).

11. Lang, S. H. *et al.* Enhanced expression of vimentin in motile prostate cell lines and in poorly differentiated and metastatic prostate carcinoma. *Prostate* **52**(4), 253–263. <https://doi.org/10.1002/pros.10088> (2002).
12. Shukla, S. *et al.* Upregulation of SATB1 is associated with prostate cancer aggressiveness and disease progression. *PLoS ONE* **8**(1), e53527. <https://doi.org/10.1371/journal.pone.0053527> (2013).
13. Yue, W., Sun, L.-Y., Li, C.-H., Zhang, L.-X. & Pei, X.-T. Screening and identification of ovarian carcinomas related genes. *Ai Zheng Aizheng Chin. J. Cancer* **23**(2), 141–145 (2004).
14. Li, D.-Q. *et al.* Identification of breast cancer metastasis-associated proteins in an isogenic tumor metastasis model using two-dimensional gel electrophoresis and liquid chromatography-ion trap-mass spectrometry. *Proteomics* **6**(11), 3352–3368. <https://doi.org/10.1002/pmic.200500617> (2006).
15. Ranganathan, S., Salazar, H., Benetatos, C. A. & Hudes, G. R. Immunohistochemical analysis of β -tubulin isoforms in human prostate carcinoma and benign prostatic hypertrophy. *Prostate* **30**(4), 263–268. [https://doi.org/10.1002/\(SICI\)1097-0045\(19970301\)30:4<3c263::AID-PROS6%3e3.0.CO;2-M](https://doi.org/10.1002/(SICI)1097-0045(19970301)30:4<3c263::AID-PROS6%3e3.0.CO;2-M) (1997).
16. Gong, F. *et al.* The BCL2 gene is regulated by a special AT-rich sequence binding protein 1-mediated long range chromosomal interaction between the promoter and the distal element located within the 3'-UTR. *Nucleic Acids Res.* **39**(11), 4640–4652. <https://doi.org/10.1093/nar/gkr023> (2011).
17. Han, H.-J., Russo, J., Kohwi, Y. & Kohwi-Shigematsu, T. SATB1 reprogrammes gene expression to promote breast tumour growth and metastasis. *Nature* **452**, 187–193 (2008).
18. Zhang, Y. *et al.* Expression of SATB1 promotes the growth and metastasis of colorectal cancer. *PLoS ONE* **9**(6), e100413. <https://doi.org/10.1371/journal.pone.0100413> (2014).
19. Han, B., Luan, L., Xu, Z. & Wu, B. Expression and biological roles of SATB1 in human bladder cancer. *Tumor Biol.* **34**(5), 2943–2949. <https://doi.org/10.1007/s13277-013-0857-1> (2013).
20. Mao, L. *et al.* SATB1 is overexpressed in metastatic prostate cancer and promotes prostate cancer cell growth and invasion. *J. Transl. Med.* **11**, 111 (2013).
21. Oh, B., Hwang, S.-Y., Solter, D. & Knowles, B. B. Spindlin, a major maternal transcript expressed in the mouse during the transition from oocyte to embryo. *Development* **124**(2), 493–503 (1997).
22. Franz, H. *et al.* The histone code reader SPIN1 controls RET signaling in liposarcoma. *Oncotarget* **6**(7), 4773 (2015).
23. Gunning, P., O'Neill, G. & Hardeman, E. Tropomyosin-based regulation of the actin cytoskeleton in time and space. *Physiol. Rev.* **88**(1), 1–35. <https://doi.org/10.1152/physrev.00001.2007> (2008).
24. Portyanko, A., Akalovich, S. & Doroshenko, T. Beta-III tubulin as a potential target for blocking invasive growth of malignant epithelial tumors. *Eksp Klin Farmakol.* **78**(8), 25–28 (2015).
25. Rubin, M. A., Dunn, R., Strawderman, M. & Pienta, K. J. Tissue microarray sampling strategy for prostate cancer biomarker analysis. *Am. J. Surg. Pathol.* **26**(3), 312–319 (2002).
26. Epstein, J. I., Allsbrook, W. C. J., Amin, M. B., Egevad, L. L., the ISUP Grading Committee. The 2005 International Society of Urological Pathology (ISUP) consensus conference on gleason grading of prostatic carcinoma. *Am. J. Surg. Pathol.* **29**(9), 1228–1242 (2005).
27. Oberländer, M. *et al.* A 'Waterfall' transfer-based workflow for improved quality of tissue microarray construction and processing in breast cancer research. *Pathol. Oncol. Res.* **20**(3), 719–726. <https://doi.org/10.1007/s12253-014-9752-3> (2014).
28. Auer, G. U., Caspersson, T. O. & Wallgren, A. S. DNA content an survival in mammary carcinoma. *Anal. Quant. Cytol. J.* **2**(2), 161–165 (1980).
29. Auer, G., Askensten, U. & Ahrens, O. Cytophotometry. *Hum. Pathol.* **20**(6), 518–527. [https://doi.org/10.1016/0046-8177\(89\)90243-8](https://doi.org/10.1016/0046-8177(89)90243-8) (1989).
30. Remmele, W. & Stegner, H. Recommendation for uniform definition of an immunoreactive score (IRS) for immunohistochemical estrogen receptor detection (ER-ICA) in breast cancer tissue. *Pathologie* **8**(3), 138–140 (1987).
31. www.proteinatlas.org. Date of access: 09 September 2021.
32. Böcking, A., Tils, M., Schramm, M., Dietz, J. & Biesterfeld, S. DNA-cytometric grading of prostate cancer systematic review with descriptive data analysis. *Pathol. Discov.* **2**(1), 7. <https://doi.org/10.7243/2052-7896-2-7> (2014).
33. Yildirim-Assaf, S. *et al.* The prognostic significance of determining DNA content in breast cancer by DNA image cytometry: The role of high grade aneuploidy in node negative breast cancer. *J. Clin. Pathol.* **60**(6), 649–655. <https://doi.org/10.1136/jcp.2005.035550> (2007).
34. Araujo, S. E. A., Bernardo, W. M., Habr-Gama, A., Kiss, D. R. & Ceccanello, I. DNA Ploidy status and prognosis in colorectal cancer: A meta-analysis of published data. *Dis. Colon Rectum.* **50**(11), 1800–1810. <https://doi.org/10.1007/s10350-007-9013-6> (2007).
35. Bantis, A. *et al.* Telomerase RNA expression and DNA ploidy as prognostic markers of prostate carcinomas. *Tumori* **95**(6), 744–752 (2009).
36. Bantis, A. *et al.* Prognostic value of DNA analysis of prostate adenocarcinoma: Correlation to clinicopathologic predictors. *J. Exp. Clin. Cancer Res.* **24**(2), 273–278 (2005).
37. Cai, S., Lee, C. C. & Kohwi-Shigematsu, T. SATB1 packages densely looped, transcriptionally active chromatin for coordinated expression of cytokine genes. *Nat. Genet.* **38**(11), 1278–1288. <https://doi.org/10.1038/ng1913> (2006).
38. Mao, L. *et al.* Oncolytic virus carrying shRNA targeting SATB1 inhibits prostate cancer growth and metastasis. *Tumor Biol.* **36**(11), 9073–9081. <https://doi.org/10.1007/s13277-015-3658-x> (2015).
39. Mao, L. *et al.* SATB1 promotes prostate cancer metastasis by the regulation of epithelial–mesenchymal transition. *Biomed. Pharmacother.* **79**, 1–8. <https://doi.org/10.1016/j.biopha.2016.01.038> (2016).
40. Qi, H. *et al.* SATB1 promotes epithelial–mesenchymal transition and metastasis in prostate cancer. *Oncol. Lett.* <https://doi.org/10.3892/ol.2017.5765> (2017).
41. Selinger, C. I. *et al.* Loss of special AT-rich binding protein 1 expression is a marker of poor survival in lung cancer. *J. Thorac. Oncol.* **6**(7), 1179–1189. <https://doi.org/10.1097/JTO.0b013e31821b4ce0> (2011).
42. Al-Sohaily, S. *et al.* Loss of special AT-rich sequence-binding protein 1 (SATB1) predicts poor survival in patients with colorectal cancer. *Histopathology* **65**(2), 155–163. <https://doi.org/10.1111/his.12295> (2014).
43. Iorns, E. *et al.* The role of SATB1 in breast cancer pathogenesis. *JNCI J. Natl. Cancer Inst.* **102**(16), 1284–1296. <https://doi.org/10.1093/jnci/djq243> (2010).
44. Hanker, L. C. *et al.* SATB1 gene expression and breast cancer prognosis. *The Breast* **20**(4), 309–313. <https://doi.org/10.1016/j.breast.2010.10.002> (2011).
45. Pavan Kumar, P. *et al.* Phosphorylation of SATB1, a global gene regulator, acts as a molecular switch regulating its transcriptional activity in vivo. *Mol. Cell.* **22**(2), 231–243. <https://doi.org/10.1016/j.molcel.2006.03.010> (2006).
46. Han, S., Xia, J., Qin, X., Han, S. & Wu, A. Phosphorylated SATB1 is associated with the progression and prognosis of glioma. *Cell Death Dis.* **4**(10), e901. <https://doi.org/10.1038/cddis.2013.433> (2013).
47. Ghosh, R. P. *et al.* Satb1 integrates DNA binding site geometry and torsional stress to differentially target nucleosome-dense regions. *Nat. Commun.* **10**(1), 1–16. <https://doi.org/10.1038/s41467-019-11118-8> (2019).
48. Bergstrand, J. *et al.* Super-resolution microscopy can identify specific protein distribution patterns in platelets incubated with cancer cells. *Nanoscale* **11**(20), 10023–10033. <https://doi.org/10.1039/C9NR01967G> (2019).
49. Shoaib, J. & Barbieri, C. E. Clinical variability and molecular heterogeneity in prostate cancer. *Asian J. Androl.* **18**(4), 543 (2016).

50. Shao, Y.-H. *et al.* Contemporary risk profile of prostate cancer in the United States. *JNCI J. Natl. Cancer Inst.* **101**(18), 1280–1283. <https://doi.org/10.1093/jnci/djp262> (2009).
51. Ross, J. S. *et al.* Prediction of pathologic stage and postprostatectomy disease recurrence by DNA ploidy analysis of initial needle biopsy specimens of prostate cancer. *Cancer* **74**(10), 2811–2818. [https://doi.org/10.1002/1097-0142\(19941115\)74:10%3c2811::AID-CNCR2820741012%3e3.0.CO;2-B](https://doi.org/10.1002/1097-0142(19941115)74:10%3c2811::AID-CNCR2820741012%3e3.0.CO;2-B) (1994).

Acknowledgements

Grants from the Ad Infinitum Foundation, Werner & Clara Kreitz Foundation and the European Union 7th Framework Programme (FLUODIAMON #201837) are gratefully acknowledged. This study was performed in collaboration with the *Surgical Center for Translational Oncology-Lübeck (SCTO-L)* and the *Interdisciplinary Center for Biobanking-Lübeck (ICB-L)*. The authors would like to thank Katja Klempt-Giessing and Gisela Grosser-Pape for excellent technical assistance.

Author contributions

C.D. and C.D. were responsible for acquisition of the patient collective (tissue and corresponding patient data). C.T. performed the histological examination and Gleason Grading. C.D. and M.O. were responsible for the laboratory work (HE staining, TMA-construction, immunohistochemistry and DNA-image-cytometry). C.D. and A.G. was responsible for the evaluation of the immunohistochemical staining. C.D. and J.H. analyzed and interpreted the DNA cytometry data. C.D., S.F.W. and T.G. were responsible for analyzing the patient's datasets and statistics. C.D., T.G. and J.H. wrote the final manuscript. All authors read and approved the final manuscript.

Funding

Open Access funding enabled and organized by Projekt DEAL. Christoph Dumke received a scholarship from the Ad Infinitum Foundation.

Competing interests

The authors declare no competing interests.

Additional information

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1038/s41598-021-03702-0>.

Correspondence and requests for materials should be addressed to J.K.H.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2021