



Universiteit
Leiden
The Netherlands

Diastolic dyssynchrony by SPECT: a novel parameter to predict post-infarct adverse remodeling

Bijl, P. van der; Delgado, V.; Bax, J.J.

Citation

Bijl, P. van der, Delgado, V., & Bax, J. J. (2021). Diastolic dyssynchrony by SPECT: a novel parameter to predict post-infarct adverse remodeling. *Journal Of Nuclear Cardiology*. doi:10.1007/s12350-020-02484-8

Version: Publisher's Version

License: [Licensed under Article 25fa Copyright Act/Law \(Amendment Taverne\)](#)

Downloaded from: <https://hdl.handle.net/1887/3279692>

Note: To cite this publication please use the final published version (if applicable).



Diastolic dyssynchrony by SPECT: A novel parameter to predict post-infarct adverse remodeling

Pieter van der Bijl, MD, PhD,^a Victoria Delgado, MD, PhD,^a and Jeroen J. Bax, MD, PhD^a

^a Department of Cardiology, Heart Lung Center, Leiden University Medical Center, Leiden, The Netherlands

Received Dec 1, 2020; accepted Dec 2, 2020
doi:10.1007/s12350-020-02484-8

See related article, <https://doi.org/10.1007/s12350-020-02483-9>.

Left ventricular (LV) post-infarct remodeling is caused by an inflammatory response, which leads to the degradation of myocardial matrix, slippage of muscle bundles, wall thinning, infarct expansion, and eventually LV adverse remodeling.¹ LV post-infarct remodeling carries prognostic implications, with a higher rate of heart failure hospitalization and possibly increased mortality.^{1,2} Identification of post-infarct survivors who will experience adverse LV remodeling might, therefore, provide an opportunity to focus preventative strategies on such patients. Many different predictors of LV post-infarct remodeling have been proposed, including clinical features (e.g., ST-segment resolution on the surface ECG³), biochemical markers (e.g., baseline N-terminal pro-brain natriuretic peptide (NT-proBNP)⁴) and imaging biomarkers (e.g., baseline LV conic index⁵).

In the current issue of the journal, Zhang⁶ now adds a novel imaging biomarker (diastolic dyssynchrony [DD]) assessed with phase analysis of ^{99m}technetium-sestamibi-gated single-photon emission computed tomography (SPECT)) to the armamentarium of LV post-infarct remodeling predictors. LV infarcts were created in a swine model by percutaneous coronary balloon occlusion of the left anterior descending

coronary artery. DD was assessed with phase contrast histogram bandwidth (PBW) and phase standard deviation (PSD) before the infarct, and subsequently at 1 day, 1 week, and 4 weeks post-infarct. The progression of LV remodeling was defined as the percentage change of LV end-systolic volume on echocardiography at 4 weeks post-infarct, compared to 1 day after the infarct. Both PBW ($\beta = 0.004$, 95% confidence interval [CI] 0.001 to 0.007, $P = .024$) and PSD ($\beta = 0.008$, 95% CI 0.000 to 0.017, $P = .049$) at day 1 post-infarct were independently associated with progressive LV post-infarct remodeling at 4 weeks.

SPECT is an accepted technique to quantify DD.⁷ The development of DD is well documented in the presence of coronary artery disease (with or without a previous infarct) as well as in hypertensive heart disease and heart failure patients.^{8–10} DD was defined as (i) the standard deviation (SD) of the time from the start of the QRS complex to the peak myocardial early diastolic velocity on tissue Doppler imaging and (ii) the maximum difference between the start of the QRS complex to the peak myocardial early diastolic velocity on tissue Doppler imaging). DD was not found to correlate with systolic dyssynchrony ($r = 0.03$, $P = .68$ and $r = 0.03$, $P = .74$, respectively, when using the abovementioned definitions), when analyzing 110 asymptomatic hypertensive individuals, thereby suggesting that these phenomena (systolic and DD) may have different underlying mechanisms.⁹ Similarly, DD (defined identically to Chang et al.⁹) was not strongly linked to systolic dyssynchrony in a cohort of heart failure patients of mixed ischemic and non-ischemic etiology.¹⁰ Since diastolic dysfunction occurs earlier during the ischemic cascade than systolic dysfunction and revascularization of acute coronary artery lesions improves DD (while revascularization of chronic lesions does

Reprint requests: Pieter van der Bijl, MD, PhD, Department of Cardiology, Heart Lung Center, Leiden University Medical Center, Albinusdreef 2, 2300 RC Leiden, The Netherlands; pieter.vanderbijl@gmail.com

J Nucl Cardiol
1071-3581/\$34.00

Copyright © 2021 American Society of Nuclear Cardiology.

not), DD may reflect stunned (but viable) myocardium.¹¹

DD (defined as the visually apprehended segmental early relaxation phenomenon on echocardiography) was not found to be predictive of angina, myocardial infarction, heart failure, stroke, or death in a group of 244 individuals referred for stress echocardiography.¹² In contrast, DD was associated with survival in patients with coronary disease and heart failure, when assessed by SPECT PBW and PSD.¹³ Systolic dyssynchrony (defined as the maximum difference of the time from the start of the QRS complex to peak radial strain by speckle tracking echocardiography) was linked to LV post-infarct remodeling (adjusted odds ratio [OR] 1.03 ms, 95% CI 1.02 to 1.05, $P < .001$) in 178 patients presenting with acute myocardial infarction.¹⁴ Similarly, when systolic dyssynchrony was defined as the SD of the time from the start of the QRS complex to the peak myocardial systolic velocity on tissue Doppler imaging, it was also an independent predictor (OR 1.19, 95% CI 1.07 to 1.32, $P = .001$) of post-infarct LV remodeling.¹⁵ In contrast, DD (defined as the SD of the time from the start of the QRS complex to the peak myocardial early diastolic velocity on tissue Doppler imaging) was not independently predictive of adverse LV remodeling in post-infarct patients ($P = .6$).¹⁵ In a group of 48 ST-segment myocardial infarction patients, DD (defined as the maximum difference of the time from the start of the QRS complex to the peak myocardial early diastolic velocity on tissue Doppler imaging) was also not independently associated with adverse LV remodeling.¹¹

Early identification of post-infarct patients who will develop adverse LV remodeling has potential clinical utility. No comprehensive, head-to-head comparison of the multitude of predictors of LV post-infarct remodeling has ever been performed. In the current issue of the journal, Zhang et al⁶ introduce DD, which was assessed with phase analysis of ^{99m}technetium-sestamibi-gated SPECT, as a tool for the early identification of progressive adverse LV remodeling post-infarct. The choice of progressive LV remodeling, rather than remodeling per se, is interesting since most studies have focused on LV remodeling which was defined at a specific time point post-infarct. Few data exist on predictors of progressive LV remodeling, and as such the current analysis adds valuable insight to post-infarct remodeling. Although the authors could show SPECT-derived DD to be associated with progression of adverse LV remodeling, there are some issues which will be of interest for future studies. While Zhang et al⁶ compared DD (assessed with SPECT) to echocardiographic measures of DD (the maximum difference of the time from the start of the QRS complex to the peak myocardial early diastolic velocity on tissue Doppler imaging and the SD

of the time from the start of the QRS complex to the peak myocardial early diastolic velocity on tissue Doppler imaging), there is no evidence presented that DD is superior to any other marker of future adverse LV remodeling. Two prior echocardiographic studies could not confirm DD to be a predictor of LV post-infarct remodeling, which is in contradiction to the results presented and which raises the question if SPECT is a more sensitive technique as compared to echocardiography.^{11,15} The study by Zhang et al⁶ is preclinical and still has to be applied in the clinical setting. The novelty of the current study lies in the focus on DD—a phenomenon which has not been extensively investigated in coronary artery disease, and which may not only have a unique underlying mechanism, but also different clinical implications as compared to systolic dyssynchrony. Direct comparison of post-infarct biomarkers is required to determine which will most accurately predict adverse LV remodeling. Finally, the clinical value of targeting the population, which will undergo such remodeling in terms of effective prevention, will have to be prospectively studied before application to routine practice can become a reality.

Disclosures

The Department of Cardiology of Leiden University Medical Center received research grants from Abbott Vascular, Bayer, Biotronik, Bioventrix, Boston Scientific, Edwards Lifesciences, GE Healthcare, and Medtronic. Jeroen Bax received speaker fees from Abbott Vascular. Victoria Delgado received speaker fees from Abbott Vascular, Edwards Lifesciences, GE Healthcare, Medtronic and MSD. The remaining author has nothing to disclose.

Ethical approval

This article does not contain any studies with human participants performed by any of the authors.

References

1. Van der Bijl P, Abou R, Goedemans L, Gersh BJ, Holmes DR Jr, Ajmone Marsan N, et al. Left ventricular post-infarct remodeling: Implications for systolic function improvement and outcomes in the modern era. *JACC Heart Fail* 2020;8:131-40. <https://doi.org/10.1016/j.jchf.2019.08.014>.
2. Bolognese L, Neskovic AN, Parodi G, Cerisano G, Buonamici P, Santoro GM, et al. Left ventricular remodeling after primary coronary angioplasty: Patterns of left ventricular dilation and long-term prognostic implications. *Circulation* 2002;106:2351-7. <https://doi.org/10.1161/01.cir.0000036014.90197.f0>.
3. Kim JS, Ko YG, Yoon SJ, Moon JY, Kim YJ, Choi BW, et al. Correlation of serial cardiac magnetic resonance imaging parameters with early resolution of ST-segment elevation after primary percutaneous coronary intervention. *Circ J* 2008;72:1621-6. <https://doi.org/10.1253/circj.cj-08-0232>.
4. Nilsson JC, Groenning BA, Nielsen G, Fritz-Hansen T, Trawinski J, Hildebrandt PR, et al. Left ventricular remodeling in the first

- year after acute myocardial infarction and the predictive value of N-terminal pro brain natriuretic peptide. *Am Heart J* 2002;143:696-702. <https://doi.org/10.1067/mhj.2002.120293>.
5. Li F, Chen YG, Yao GH, Li L, Ge ZM, Zhang M, et al. Usefulness of left ventricular conic index measured by real-time three-dimensional echocardiography to predict left ventricular remodeling after acute myocardial infarction. *Am J Cardiol* 2008;102:1433-7. <https://doi.org/10.1016/j.amjcard.2008.07.034>.
 6. Zhang F WJ, Shao X, Chen Y, Fan S, Shi Y, Liu B, Yu W, Li X, Xu M, Yang M, Xi X, Wu Z, Li S, Wang Y. Longitudinal evaluation of diastolic dyssynchrony by SPECT gated myocardial perfusion imaging early after acute myocardial infarction and the relationship with progression of left ventricular remodeling in swine model. *J Nucl Cardiol* 2020.
 7. Henneman MM, Chen J, Dibbets-Schneider P, Stokkel MP, Bleeker GB, Ypenburg C, et al. Can LV dyssynchrony as assessed with phase analysis on gated myocardial perfusion SPECT predict response to CRT? *J Nucl Med* 2007;48:1104-11. <https://doi.org/10.2967/jnumed.107.039925>.
 8. Lee PW, Zhang Q, Yip GW, Wu L, Lam YY, Wu EB, et al. Left ventricular systolic and diastolic dyssynchrony in coronary artery disease with preserved ejection fraction. *Clin Sci (Lond)* 2009;116:521-9. <https://doi.org/10.1042/CS20080100>.
 9. Chang SA, Kim HK, Kim DH, Kim YJ, Sohn DW, Oh BH, et al. Left ventricular systolic and diastolic dyssynchrony in asymptomatic hypertensive patients. *J Am Soc Echocardiogr* 2009;22:337-42. <https://doi.org/10.1016/j.echo.2009.01.024>.
 10. Yu CM, Zhang Q, Yip GW, Lee PW, Kum LC, Lam YY, et al. Diastolic and systolic asynchrony in patients with diastolic heart failure: A common but ignored condition. *J Am Coll Cardiol* 2007;49:97-105. <https://doi.org/10.1016/j.jacc.2006.10.022>.
 11. Turan B, Dasli T, Erkol A, Erden I, Basaran Y. Diastolic dyssynchrony in acute ST segment elevation myocardial infarction and relationship with functional recovery of left ventricle. *J Cardiovasc Ultrasound* 2016;24:208-14. <https://doi.org/10.4250/jcu.2016.24.3.208>.
 12. Obeidat O, Arida M, Al-Mallah M, Alam M, Ananthasubramanian K. Segmental early relaxation phenomenon: Incidence, clinical characteristics, and significance in stress echocardiography. *Chest* 2004;125:1218-23. <https://doi.org/10.1378/chest.125.4.1218>.
 13. Fudim M, Fathallah M, Shaw LK, James O, Samad Z, Piccini JP, et al. The prognostic value of diastolic and systolic mechanical left ventricular dyssynchrony among patients with coronary artery disease and heart failure. *J Nucl Cardiol*. 2019. <https://doi.org/10.1007/s12350-019-01843-4>.
 14. Mollema SA, Liem SS, Suffoletto MS, Bleeker GB, van der Hoeven BL, van de Veire NR, et al. Left ventricular dyssynchrony acutely after myocardial infarction predicts left ventricular remodeling. *J Am Coll Cardiol* 2007;50:1532-40. <https://doi.org/10.1016/j.jacc.2007.07.025>.
 15. Zhang Y, Yip GW, Chan AK, Wang M, Lam WW, Fung JW, et al. Left ventricular systolic dyssynchrony is a predictor of cardiac remodeling after myocardial infarction. *Am Heart J* 2008;156:1124-32. <https://doi.org/10.1016/j.ahj.2008.07.019>.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.