



Universiteit  
Leiden  
The Netherlands

## **Renal function decline in older men and women with advanced chronic kidney disease-results from the EQUAL study**

Chesnaye, N.C.; Dekker, F.W.; Evans, M.; Caskey, F.J.; Torino, C.; Postorino, M.; ... ; EQUAL Study Investigators

### **Citation**

Chesnaye, N. C., Dekker, F. W., Evans, M., Caskey, F. J., Torino, C., Postorino, M., ... Jager, K. J. (2021). Renal function decline in older men and women with advanced chronic kidney disease-results from the EQUAL study. *Nephrology Dialysis Transplantation*, 36(9), 1656-1663. doi:10.1093/ndt/gfaa095

Version: Publisher's Version

License: [Creative Commons CC BY-NC 4.0 license](#)

Downloaded from: <https://hdl.handle.net/1887/3276458>

**Note:** To cite this publication please use the final published version (if applicable).

3. Brown MJ, McInnes GT, Papst CC *et al.* Aliskiren and the calcium channel blocker amlodipine combination as an initial treatment strategy for hypertension control (ACCELERATE): a randomised, parallel-group trial. *Lancet* 2011; 377: 312–320
4. Parving HH, Persson F, Lewis JB *et al.* Aliskiren combined with losartan in type 2 diabetes and nephropathy. *N Engl J Med* 2008; 358: 2433–2446
5. Persson F, Lewis JB, Lewis EJ *et al.*; for the AVOID Study Investigators. Impact of baseline renal function on the efficacy and safety of aliskiren added to losartan in patients with type 2 diabetes and nephropathy. *Diabetes Care* 2010; 33: 2304–2309
6. Tang SC, Lin M, Tam S *et al.* Aliskiren combined with losartan in immunoglobulin A nephropathy: an open-labeled pilot study. *Nephrol Dial Transplant* 2012; 27: 613–618
7. Parving HH, Brenner BM, McMurray JJ *et al.* Aliskiren trial in type 2 diabetes using cardio-renal endpoints (ALTITUDE): rationale and study design. *Nephrol Dial Transplant* 2009; 24: 1663–1671
8. Parving HH, Brenner BM, McMurray JJ *et al.* Cardiorenal end points in a trial of aliskiren for type 2 diabetes. *N Engl J Med* 2012; 367: 2204–2213
9. McMurray JJ, Abraham WT, Dickstein K *et al.* Aliskiren, ALTITUDE, and the implications for ATMOSPHERE. *Eur J Heart Failure* 2012; 14: 341–343
10. McMurray JJ, Krum H, Abraham WT *et al.* Aliskiren, enalapril, or aliskiren and enalapril in heart failure. *N Engl J Med* 2016; 374: 1521–1532
11. Levey AS, Bosch JP, Lewis JB *et al.* A more accurate method to estimate glomerular filtration rate from serum creatinine: a new prediction equation. Modification of Diet in Renal Disease Study Group. *Ann Intern Med* 1999; 130: 461–470
12. Levey AS, Inker LA, Matsushita K *et al.* GFR decline as an end point for clinical trials in CKD: a scientific workshop sponsored by the National Kidney Foundation and the US Food and Drug Administration. *Am J Kidney Dis* 2014; 64: 821–835
13. Coresh J, Turin TC, Matsushita K *et al.* Decline in estimated glomerular filtration rate and subsequent risk of end-stage renal disease and mortality. *JAMA* 2014; 311: 2518–2531
14. Randomised placebo-controlled trial of effect of ramipril on decline in glomerular filtration rate and risk of terminal renal failure in proteinuric, non-diabetic nephropathy. The GISEN Group (Gruppo Italiano di Studi Epidemiologici in Nefrologia). *Lancet* 1997; 349: 1857–1863
15. Imrey PB. Compromising outcomes. *J Am Soc Nephrol* 2019; 30: 1147–1150
16. Cravedi P, Ruggenenti P, Remuzzi G. Proteinuria should be used as a surrogate in CKD. *Nat Rev Nephrol* 2012; 8: 301–306
17. Villamil A, Chrysant SG, Calhoun D *et al.* Renin inhibition with aliskiren provides additive antihypertensive efficacy when used in combination with hydrochlorothiazide. *J Hypertens* 2007; 25: 217–226
18. Bekassy ZD, Kristoffersson AC, Rebetz J *et al.* Aliskiren inhibits renin-mediated complement activation. *Kidney Int* 2018; 94: 689–700
19. Chung S, Kim S, Kim M *et al.* Treatment combining aliskiren with paricalcitol is effective against progressive renal tubulointerstitial fibrosis via dual blockade of intrarenal renin. *PLoS One* 2017; 12: e0181757
20. Hollenberg NK. Renin inhibition: what are the clinical perspectives? *Semin Nephrol* 2007; 27: 511–518
21. Krop M, Garrelts IM, de Bruin RJ *et al.* Aliskiren accumulates in renin secretory granules and binds plasma prorenin. *Hypertension* 2008; 52: 1076–1083

Received: 17.1.2020; Editorial decision: 26.3.2020

*Nephrol Dial Transplant* (2021) 36: 1656–1663

doi: 10.1093/ndt/gfaa095

Advance Access publication 27 June 2020

## Renal function decline in older men and women with advanced chronic kidney disease—results from the EQUAL study

Nicholas C. Chesnaye<sup>1</sup>, Friedo W. Dekker<sup>2</sup>, Marie Evans <sup>3</sup>, Fergus J. Caskey<sup>4</sup>, Claudia Torino<sup>5</sup>, Maurizio Postorino<sup>5</sup>, Maciej Szymczak<sup>6</sup>, Chava L. Ramspek <sup>2</sup>, Christiane Drechsler<sup>7</sup>, Christoph Wanner<sup>7</sup> and Kitty J. Jager<sup>1</sup>; the EQUAL study investigators

<sup>1</sup>ERA-EDTA Registry, Department of Medical Informatics, Academic Medical Center, University of Amsterdam, Amsterdam Public Health Research Institute, Amsterdam, The Netherlands, <sup>2</sup>Department of Clinical Epidemiology, Leiden University Medical Center, Leiden, The Netherlands, <sup>3</sup>Renal Unit, Department of Clinical Intervention and Technology, Karolinska Institutet and Karolinska University Hospital, Stockholm, Sweden, <sup>4</sup>Population Health Sciences, Bristol Medical School, University of Bristol, Bristol, UK, <sup>5</sup>IFC-CNR, Clinical Epidemiology and Pathophysiology of Renal Diseases and Hypertension and G.O.M., Bianchi Melacrino Morelli, Reggio Calabria, Italy, <sup>6</sup>Department of Nephrology and Transplantation Medicine, Wroclaw Medical University, Wroclaw, Poland and <sup>7</sup>Division of Nephrology, University Hospital of Wurzburg, Wurzburg, Germany

Correspondence to: Nicholas C. Chesnaye; E-mail: n.c.chesnaye@amsterdamumc.nl;

Twitter handles: @jager\_kitty, @ChristophWanne4

### ABSTRACT

**Introduction.** Understanding the mechanisms underlying the differences in renal decline between men and women may improve sex-specific clinical monitoring and management. To this end, we aimed to compare the slope of renal function decline in older men and women in chronic kidney

disease (CKD) Stages 4 and 5, taking into account informative censoring related to the sex-specific risks of mortality and dialysis initiation.

**Methods.** The European QUALity Study on treatment in advanced CKD (EQUAL) study is an observational prospective cohort study in Stages 4 and 5 CKD patients  $\geq 65$  years not on

dialysis. Data on clinical and demographic patient characteristics were collected between April 2012 and December 2018. Estimated glomerular filtration rate (eGFR) was calculated using the CKD Epidemiology Collaboration equation. eGFR trajectory by sex was modelled using linear mixed models, and joint models were applied to deal with informative censoring.

**Results.** We included 7801 eGFR measurements in 1682 patients over a total of 2911 years of follow-up. Renal function declined by 14.0% [95% confidence interval (CI) 12.9–15.1%] on average each year. Renal function declined faster in men (16.2%/year, 95% CI 15.9–17.1%) compared with women (9.6%/year, 95% CI 6.3–12.1%), which remained largely unchanged after accounting for various mediators and for informative censoring due to mortality and dialysis initiation. Diabetes was identified as an important determinant of renal decline specifically in women.

**Conclusion.** In conclusion, renal function declines faster in men compared with women, which remained similar after adjustment for mediators and despite a higher risk of informative censoring in men. We demonstrate a disproportional negative impact of diabetes specifically in women.

**Keywords:** EQUAL, renal function decline, sex disparities

## INTRODUCTION

The epidemiology of chronic kidney disease (CKD) differs by sex. Population-based studies across the globe consistently show a higher prevalence of CKD in women compared with men [1–7], yet ~60% of those starting renal replacement therapy for end-stage kidney disease (ESKD) are men [8, 9]. This paradox has several potential explanations [10]. First, the longer life expectancy in women along with the natural decline of glomerular filtration rate (GFR) with age may partly explain the higher prevalence of CKD in women. Secondly, several (population-based) studies [11–14] as well as a large meta-analysis of studies in non-diabetic CKD patients [15], point towards a faster decline of renal function in men. In contrast, others have demonstrated a more rapid progression in women in various (sub-)populations [16–18], whereas some found no difference between the sexes at all [19, 20]. A meta-analysis of randomized controlled trials (RCTs) found that women progress at an equal speed as men, with adjusted analyses even suggesting a faster progression in women [21]. Given these inconclusive results, it is clear that the estimated sex-specific decline in renal function depends on the population studied; CKD stage, the presence of diabetes mellitus (postmenopausal), age, and population-based cohorts versus referred patients, are all factors that likely contribute to the variation in current evidence.

Studying renal function decline by sex is complicated by a sex-specific selection process caused by a higher mortality risk in men across all ranges of pre-ESKD estimated GFR (eGFR) [22, 23]. The effect of eGFR decline and albuminuria on mortality risk seems stronger in women, adding complexity to the selection process [22]. Furthermore, as men and women start dialysis at different levels of eGFR [24], censoring at dialysis initiation may be deemed informative when studying CKD progression. Consequently, it is important

## KEY LEARNING POINTS

### What is already known about this subject?

- It is known that the epidemiology of chronic kidney disease (CKD) differs by sex; however, the current evidence on sex-specific slopes of renal decline in advanced CKD remains inconclusive.
- Studying renal function decline by sex is complicated by informative censoring caused by sex-specific risks of mortality and dialysis initiation.

### What this study adds?

- Men progress faster than women, even after adjustment for important mediators, and despite having a higher risk of censoring.
- Diabetes is an important determinant of renal decline, with a disproportional negative impact specifically in women.

### What impact this may have on practice or policy?

- Our results improve understanding of the mechanisms underlying the differences in renal function decline between the sexes and help achieve individualized and sex-specific management and treatment in advanced CKD.

when investigating this topic to take into account informative censoring caused by mortality and dialysis initiation, as the estimated slopes of renal function decline by sex may otherwise be biased.

Understanding the mechanisms underlying the differences in renal function decline between the sexes may aid sex-specific clinical monitoring and management. To date, very few studies have investigated renal function decline by sex specifically during pre-dialysis Stages 4 and 5 in referred CKD patients of older age, and none has taken into account the potential bias caused by the sex-specific risk of mortality and dialysis initiation [25]. Consequently, here we aim to compare the slope of renal function decline in older men and women with advanced CKD, taking into account informative censoring due to mortality and dialysis. As a secondary aim, we will explore sex-specific determinants of renal function decline in this population.

## MATERIALS AND METHODS

### Study design and population

The European QUALity Study on treatment in advanced CKD (EQUAL) is an ongoing observational cohort study including Stages 4 and 5 CKD patients not on dialysis receiving routine medical care in Germany, Italy, the Netherlands, Poland, Sweden and the UK. Patients of  $\geq 65$  years of age were included with an incident eGFR  $< 20$  mL/min/1.73 m<sup>2</sup> calculated by the Modification of Diet in Renal Disease equation.

Patients were excluded if the drop in eGFR resulted from an acute event or if they had previously received dialysis or a kidney transplant. Approval was obtained from the medical ethical committees in each country. Informed consent was obtained from all patients. A full description of the study has been published elsewhere [26].

### Data collection

Clinical data were collected between April 2012 and December 2018 on patient demographics, primary renal disease, laboratory data and cardiovascular risk factors [smoking status, body mass index (BMI), haemoglobin, blood pressure, cholesterol and diabetes mellitus]. Data on the following pre-existing cardiovascular co-morbid conditions were also collected (definitions provided in the [Supplementary data](#)): cerebrovascular disease, peripheral vascular disease, myocardial infarction, angina pectoris, congestive heart failure, left ventricular hypertrophy, hypertension and cardiac arrhythmias. Study visits were scheduled at 6-month intervals, and patients were followed until dialysis initiation, kidney transplantation, death, refusal for further participation, loss to follow-up or end of follow-up. The eGFR was calculated from serum creatinine level standardized to isotope dilution mass spectrometry using the CKD Epidemiology Collaboration (CKD-EPI) equation [27]. In addition, GFR was estimated during follow-up from routine 24-h urine collection by taking the average of creatinine clearance and urea clearance, normalized to body surface area following the Dubois and Dubois formula. Albumin-creatinine ratio (ACR) was also determined following routine 24-h urine collection, or a single sample if 24-h urinary collection, was unavailable. Primary kidney disease was classified using the codes of the ERA-EDTA and grouped as glomerulonephritis, diabetes mellitus, tubulo-interstitial disease, hypertension and miscellaneous kidney diseases.

### Statistical analysis

Patient characteristics were reported by sex as mean values with standard deviations (SDs) for normally distributed continuous variables, as medians with interquartile ranges (IQRs) for skewed continuous variables and as proportions for categorical variables. Linear mixed models (LMMs) were used to model the eGFR trajectory. A random intercept was included to capture the variation in eGFR baseline value between patients and a random slope for time to capture variability in the patient's eGFR trajectory. Due to nonlinear patient trajectories of eGFR, the latter was included as a cubic B-spline with two equally spaced knots positioned between the minimum and the maximum of follow-up. The unadjusted model includes time, sex and their interaction, and describes the sex-specific trajectory of eGFR over time. In subsequent models, we investigate to which extent the effect of sex on the eGFR trajectory is mediated by various groups of *a priori* defined covariates (i.e. mediators). All models were adjusted for baseline eGFR and age at inclusion.

We followed patients until death or dialysis initiation. Missing eGFR values may be introduced when patients drop out of the study due to mortality or are censored due to dialysis initiation. As the level of renal function is related to these

events, drop out is deemed informative [28–30]. We applied joint models (JMs) for longitudinal and time-to-event data to avoid biased estimates of eGFR decline [31]. The JM links the LMM described above to a Cox survival model, which captures the risk of either mortality or dialysis. In this manner, the JM informs the longitudinal eGFR trajectory on missingness caused by either of these events. To determine whether the difference in eGFR slope between men and women had changed after taking into account informative censoring due to mortality or dialysis, we tested for equality between the time-sex interaction coefficients in the LMM and JM using a Z-score test [32].

Sex-specific determinants of eGFR decline were studied through effect modification using interaction analyses, specifically through three-way interactions between sex, time and the characteristics of interest. Q-Q plots were used to check whether the residuals were normally distributed, and eGFR was log-transformed to fulfil this assumption. Consequently, regression coefficients were exponentiated and interpreted as the mean percent change in eGFR per year. Only complete cases were analysed, and missing values are reported in the [Supplementary data](#). All analyses were performed with SAS version 9.4 and R version 3.4.1.

### Sensitivity analyses

We performed a number of sensitivity analyses. First, in addition to the CKD-EPI equation, we also repeated the analyses using the full age spectrum equation and the revised Lund-Malmö equation to estimate GFR [33, 34]. Secondly, we studied the association between sex and GFR decline estimated from 24-h urine collection. Thirdly, as age is an important variable in all estimating GFRs, we also considered the relationship between sex and 1/creatinine over time. Lastly, due to the wide range in individual follow-up time, we also repeated the analyses in patients with at least 1 year of follow-up.

## RESULTS

### Patient characteristics

**Table 1** describes the baseline characteristics of 1682 patients by sex. On average, patients were 76 years old at inclusion (IQR 71–81), two-thirds were men and the eGFR at baseline was 17.0 mL/min/1.73 m<sup>2</sup> (IQR 14.5–20.4). Women were older, had a slightly higher BMI, higher values of serum calcium, cholesterol and potassium, but lower levels of haemoglobin. Diabetes and glomerular disease accounted for a lower proportion of primary renal disease in women compared with men, whereas tubulo-interstitial disease and hypertension were more common in women. Women had higher baseline renal function and lower ACR. Regarding co-morbidity, more men had diabetes, peripheral vascular disease, myocardial infarction and angina pectoris.


### The effect of sex on the eGFR trajectory

We included 7801 eGFR measurements over a total of 2911 years of follow-up, with a median (IQR) of 4 (2–7)

**Table 1. Baseline patient characteristics by sex**

Baseline patient characteristics	Overall ( <i>n</i> = 1682)	Men ( <i>n</i> = 1099)	Women ( <i>n</i> = 583)	P-value
<b>Demographics</b>				
Age, mean (SD)	76.30 (6.76)	75.97 (6.45)	76.94 (7.28)	0.006
<b>Primary renal disease, <i>n</i> (%)</b>				
Diabetes	341 (20.5)	238 (21.9)	103 (17.9)	0.006
Glomerular disease	152 (9.1)	111 (10.2)	41 (7.1)	
Tubulo-interstitial disease	138 (8.3)	76 (7.0)	62 (10.8)	
Hypertension	596 (35.8)	378 (34.7)	218 (37.9)	
Miscellaneous renal disorders	436 (26.2)	285 (26.2)	151 (26.3)	
Weight, mean (SD), kg	79.70 (17.16)	83.44 (16.10)	72.51 (16.88)	<0.001
Height, mean (SD), cm	167.57 (9.94)	172.17 (7.86)	158.81 (7.25)	<0.001
BMI, mean (SD), kg/m <sup>2</sup>	28.42 (5.34)	28.20 (4.81)	28.86 (6.23)	0.023
<b>Blood chemistry</b>				
Albumin, mean (SD), g/dL	37.70 (5.91)	37.66 (5.89)	37.78 (5.97)	0.708
Calcium, mean (SD), mmol/L	2.24 (0.32)	2.23 (0.32)	2.27 (0.33)	0.013
Cholesterol, mean (SD), mmol/L	4.53 (1.28)	4.34 (1.17)	4.89 (1.41)	<0.001
PO <sub>4</sub> , mean (SD), mmol/L	1.30 (0.32)	1.30 (0.33)	1.31 (0.30)	0.303
Potassium, mean (SD), mmol/L	4.64 (0.61)	4.67 (0.62)	4.60 (0.60)	0.037
<b>Cardiovascular</b>				
Systolic blood pressure, mean (SD), mmHg	142.85 (21.96)	143.31 (21.61)	141.99 (22.61)	0.245
Diastolic blood pressure, mean (SD)	73.83 (11.26)	74.00 (11.35)	73.49 (11.10)	0.379
Hb, mean (SD), g/dL	0.72 (0.09)	0.73 (0.10)	0.71 (0.09)	<0.001
Current smoker, <i>n</i> (%)	119 (9.3)	82 (9.7)	37 (8.4)	0.538
Ex-smoker, <i>n</i> (%)	752 (63.1)	587 (74.5)	165 (40.8)	<0.001
<b>Renal function</b>				
CKD-EPI, median (IQR), mL/min/1.73 m <sup>2</sup>	17.01 (13.79–20.11)	16.69 (13.67–19.63)	17.63 (14.40–21.08)	<0.001
MDRD, median (IQR), mL/min/1.73 m <sup>2</sup>	18.57 (15.27–21.92)	18.45 (15.05–21.62)	18.99 (15.54–22.62)	0.036
ACR, median (IQR)	33.67 (4.90–154.67)	41.36 (7.47–161.10)	19.66 (2.99–119.00)	0.002
<b>Comorbidities, <i>n</i> (%)</b>				
Diabetes	693 (42)	480 (44.5)	213 (37.4)	0.006
Chronic heart failure	290 (18.1)	195 (18.7)	95 (17.0)	0.443
Cerebrovascular disease	257 (15.7)	171 (15.9)	86 (15.3)	0.781
Peripheral vascular disease	279 (17.2)	203 (19.2)	76 (13.5)	0.005
Myocardial infarction	287 (17.4)	222 (20.6)	65 (11.4)	<0.001
Angina pectoris	239 (14.7)	178 (16.8)	61 (10.9)	0.002
Left ventricular hypertrophy	349 (23.7)	244 (25.3)	105 (20.8)	0.062
Atrial fibrillation	297 (18.2)	190 (17.9)	107 (18.9)	0.644
Hypertension	1432 (89.1)	935 (89.0)	497 (89.4)	0.860

MDRD, Modification of Diet in Renal Disease.



**FIGURE 1: (a)** The average eGFR trajectory by sex with 95% CIs. **(b)** The average eGFR trajectory by sex (LMM) adjusted for censoring due to death (JM: Death) and dialysis (JM: Dialysis). The adjusted trajectories represent the average eGFR trajectory in the hypothetical situation that all patients had remained alive/had not started dialysis. The top group of lines corresponds to the eGFR trajectory in women and the bottom lines to that in men.


**Table 2. The average annual percent decline in eGFR by sex, adjusted for various groups of mediators**

Model	Crude		Adjusted	
	Estimates	P-value	Estimates	P-value
Unadjusted model ( <i>N</i> = 1682, <i>n</i> = 7801)				
Slope men	16.2 (14.9–17.4%)	<0.001	–	–
Slope women	9.6 (5.8–13.1%)	<0.001	–	–
Difference	<b>6.6 (4.3–9.1%)</b>	<b>&lt;0.001</b>	–	–
Adjusted for age ( <i>N</i> = 1647, <i>n</i> = 7691)				
Slope men	16.2 (15.0–17.5%)	<0.001	16.3 (15.0–17.6%)	<0.001
Slope women	9.6 (5.8–13.2%)	<0.001	9.6 (5.9–13.3%)	<0.001
Difference	<b>6.7 (4.3–9.1%)</b>	<b>&lt;0.001</b>	<b>6.7 (4.3–6.7%)</b>	<b>&lt;0.001</b>
Adjusted for ACR ( <i>N</i> = 1006, <i>n</i> = 2956)				
Slope men	14.3 (12.7–16.0%)	<0.001	14.2 (12.5–15.8%)	<0.001
Slope women	8.2 (3.3–12.9%)	<0.001	8.0 (3.1–12.6%)	<0.001
Difference	<b>6.1 (3.1–9.4%)</b>	<b>&lt;0.001</b>	<b>6.2 (3.2–9.4%)</b>	<b>&lt;0.001</b>
Adjusted for comorbidities <sup>a</sup> ( <i>N</i> = 1386, <i>n</i> = 6547)				
Slope men	17.1 (15.6–18.5%)	<0.001	17.0 (15.6–18.4%)	<0.001
Slope women	10.1 (6.0–14.1%)	<0.001	10.1 (5.9–14.1%)	<0.001
Difference	<b>6.9 (4.3–9.7%)</b>	<b>&lt;0.001</b>	<b>6.9 (4.3–9.7%)</b>	<b>&lt;0.001</b>
Adjusted for CVD risk factors <sup>b</sup> ( <i>N</i> = 1158, <i>n</i> = 5444)				
Slope men	16.8 (15.3–18.3)	<0.001	16.0 (14.5–17.5%)	<0.001
Slope women	9.9 (5.3–14.2%)	<0.001	9.6 (5.2–13.8%)	<0.001
Difference	<b>6.9 (4.1–10.0%)</b>	<b>&lt;0.001</b>	<b>6.4 (3.7–9.4%)</b>	<b>&lt;0.001</b>
Adjusted for blood chemistry <sup>c</sup> ( <i>N</i> = 937, <i>n</i> = 2633)				
Slope men	14.9 (13.1–16.6%)	<0.001	12.3 (10.8–13.8%)	<0.001
Slope women	9.0 (3.7–14%)	<0.001	7.4 (2.8–11.4%)	<0.001
Difference	<b>5.8 (2.6–9.4%)</b>	<b>&lt;0.001</b>	<b>5.0 (2.2–7.9%)</b>	<b>&lt;0.001</b>
Adjusted for PO <sub>4</sub> ( <i>N</i> = 1622, <i>n</i> = 6986)				
Slope men	16.3 (15.0–17.6%)	<0.001	13.5 (12.3–14.6%)	<0.001
Slope women	9.6 (5.7–13.3%)	<0.001	7.9 (4.5–7.9%)	<0.001
Difference	<b>6.7 (4.3–9.3%)</b>	<b>&lt;0.001</b>	<b>5.5 (3.4–7.8%)</b>	<b>&lt;0.001</b>
Adjusted for diabetes ( <i>N</i> = 1645, <i>n</i> = 7687)				
Slope men	16.3 (15.0–17.6%)	<0.001	16.3 (15.0–17.6%)	<0.001
Slope women	9.6 (5.8–13.2%)	<0.001	9.6 (5.9–13.2%)	<0.001
Difference	<b>6.7 (4.3–9.2%)</b>	<b>&lt;0.001</b>	<b>6.7 (4.3–9.2%)</b>	<b>&lt;0.001</b>

Difference in annual percent decline between sex in bold. <sup>a</sup>Diabetes, chronic heart failure, cerebrovascular disease, peripheral vascular disease, myocardial infarction, angina pectoris, left ventricular hypertrophy, atrial fibrillation and hypertension.

<sup>b</sup>Systolic and diastolic blood pressure, smoking status and history, and haemoglobin.

<sup>c</sup>Serum albumin, cholesterol, calcium, phosphate and potassium.



**FIGURE 2: (a) Effect modification by age on renal function decline by sex. (b) Effect modification by diabetes on renal function decline by sex.**

measurements per patient, and a median follow-up time of 18.6 months (6.7–32.6). Renal function declined 14.0% [95% confidence intervals (CI) 12.9–15.1%] on average each year. Figure 1a shows a faster unadjusted annual decline in renal

function in men (16.2%/year, 95% CI 15.9–17.1%) compared with women (9.6%/year, 95% CI 6.3–12.1%), with a difference of 6.6% (95% CI 4.3–9.1%). These estimates remained largely unchanged after accounting for various groups of mediators

(Table 2). For the purpose of comparison with existing literature, we determined the linear sex-specific slopes of renal function decline, without log-transformation, as  $-1.82$  (95% CI  $-1.63$ – $2.01$ ) mL/min/1.73 m<sup>2</sup> per year for men and  $-0.91$  (95% CI  $-0.40$ – $1.43$ ) mL/min/1.73 m<sup>2</sup> per year for women. Sensitivity analyses using GFR estimated following routine 24-h urine collection, 1/creatinine and eGFR calculated using the full age spectrum and the revised Lund–Malmö equations provided similar results. Estimated renal function decline in a subgroup of patients with at least 1 year of follow-up was also similar to the main results (Supplementary data).

### The effect of sex on the eGFR trajectory adjusted for informative censoring

Figure 1b shows the eGFR trajectory in men and women after accounting for informative censoring due to death or dialysis initiation. The adjusted trajectories represent the average eGFR trajectory in the hypothetical situation that all patients had remained alive/had not started dialysis. After accounting for death (5-year cumulative incidence of 21.4% in men and 19.6% in women,  $P = 0.32$ ), the difference in renal decline between men (16.1%/year, 95% CI 15.0–17.1%) and women (9.5%/year, 95% CI 6.3–12.6%) remained 6.6% ( $P$ -value for change in coefficient = 0.97). Accounting for drop out due to dialysis initiation (5-year cumulative incidence of 31.9% in men and 21.4% in women,  $P$ -value for difference  $< 0.0001$ ) also had little effect on the difference in renal function decline between men (17.2%/year, 95% CI 16.1–18.5%) and women (10.4%/year, 95% CI 6.9–14.2%), increasing the difference in slope between men and women marginally from 6.6% to 6.8% ( $P$ -value for change in coefficient = 0.81).

### The sex-specific determinants of the eGFR trajectory

We identified effect modification by age, diabetes and myocardial infarction at inclusion on the slope of renal function decline by sex. We found that women of older age had slower declines in renal function compared with younger women (Figure 2a), whereas age had little effect on renal decline in men ( $P$ -value for interaction = 0.03). In addition, women with diabetes had significantly faster declines in renal function compared with non-diabetics, whereas this was not the case in men (Figure 2b,  $P$ -value for interaction = 0.05). The differential effect of diabetes seemed more pronounced in women under the age of 82 years ( $P$ -value for interaction = 0.02, Supplementary data, Figure S1). Other baseline characteristics did not differentially affect the slope of renal decline in men and women.

## DISCUSSION

In our population of elderly CKD Stages 4 and 5 patients not on dialysis, we demonstrate a faster decline of renal function in men compared with women, which persisted after taking into account important mediators. By applying JMs to account for the sex-specific risks of informative censoring due to death and dialysis, we demonstrate that men progress faster than women despite having a higher risk of drop out. Furthermore, we identified diabetes as an important determinant of renal decline specifically in women, demonstrating that renal function in female

diabetics deteriorated at a similar pace as in men. Interestingly, older women had slower declines of renal function, indicative of a certain degree of selection bias in our cohort.

To our knowledge, this is the first study to explore renal decline by sex in a referred cohort of incident CKD patients with an eGFR of  $< 20$  mL/min/1.73 m<sup>2</sup>. We found that renal function in men declined approximately twice as fast as in women ( $-1.82$  mL/min/1.73 m<sup>2</sup> per year and  $-0.89$  mL/min/1.73 m<sup>2</sup> per year, respectively), which remained similar after adjustment for various mediators and informative censoring. Comparable studies on sex-specific renal decline during the transition period from Stages 4 and 5 CKD to dialysis are scarce [11, 25]. Nonetheless, our results are in line with studies published in cohorts consisting of patients in earlier stages of CKD [15]; a Swedish population-based study of CKD Stage 3 patients estimated similar differences in renal declines between men ( $-1.26$  mL/min/1.73 m<sup>2</sup> per year for a 70 year old) and women ( $-0.76$  mL/min/1.73 m<sup>2</sup> per year) [13]. More recently, in a referred cohort of CKD Stages 2 and 3 patients, the CRIC study also found faster declines in men ( $-1.43$  mL/min/1.73 m<sup>2</sup> per year) compared with women ( $-1.09$  mL/min/1.73 m<sup>2</sup> per year), although this difference was somewhat smaller compared with our estimates [12]. Even in the ‘healthy’ general population (cohort baseline eGFR of 80.7 mL/min/1.73 m<sup>2</sup>), the PREVENT study found an eGFR slope of  $-0.55$  mL/min/year/1.73 m<sup>2</sup> in men and  $-0.33$  mL/min/year/1.73 m<sup>2</sup> in women [14]. Altogether, most available evidence points towards a faster decline of renal function in men, seemingly regardless of CKD stage. Nonetheless, a handful of studies exist that have found either a faster progression in women [16] or no difference at all between the sexes [19, 20]. One of these studies, a large meta-analysis of RCTs, found that women progress at an equal speed as men, with adjusted analyses (baseline creatinine, blood pressure, urinary protein, age and treatment assignment), suggesting a faster progression in women [21], although this discrepancy may be attributed to stringent patient selection common to RCTs and erroneous adjustment within the causal pathway.

This sex difference in renal decline has several potential explanations related to biological and/or sociocultural aspects [10]. Risk factors related to lifestyle, such as a poor diet and smoking, may partly be responsible for faster decline as seen in men [18, 35]. Although more men had a history of smoking and a higher burden of cardiovascular co-morbidities in our cohort, adjustment for these factors had little effect on the sex difference in renal decline. Others have demonstrated differential effects of albuminuria, cholesterol, blood pressure and glycaemic control, on renal function decline in men and women [14, 16, 35], although most of these studies applied methodology corresponding to prognostic research, thus not contributing to mechanistic evidence. Lastly, sex hormones also likely play a role, as animal studies have demonstrated renoprotective effects of oestrogens and damaging effects of testosterone [25, 36–38].

We demonstrate that diabetes has a stronger effect on renal decline in women compared with men, to the extent that renal decline was similar between the sexes in those with diabetes. The literature surrounding this topic is inconsistent, with some reporting faster declines in diabetic men [39, 40], and others

finding no differences between the sexes [41]. In line with our findings, a Japanese cohort of Type 2 diabetics described faster declines in women (−3.5 per year) compared with men (−2.0 per year), attributing this finding to a poorer metabolic control in women [42]. Similarly, a UK RCT in Type 2 diabetics found that women had an 88% increased risk over men of declining to <60 mL/min/1.73 m<sup>2</sup> [43]. Moreover, excess mortality risk in diabetic women has been described in the dialysis population [44] as well as in non-renal cohorts [45, 46], confirming a disproportional negative impact of diabetes in women. Diminished protection of oestrogens in the hyperglycaemic state may explain this disparate effect, even though the women in our population were likely post-menopausal [47].

Missing eGFR values are introduced over time as patients are censored due to dialysis initiation or death. As the level of renal function is related to these events, censoring is deemed informative [28]. More importantly, as the risks of dialysis initiation or death are specific to men and women, informative censoring may affect the estimated slopes for men and women differentially, potentially introducing bias. We are unaware of any previous studies that have taken into account the sex-specific risks of drop out when studying renal decline by sex. Here, we were able to account for this issue by modelling both eGFR decline and the risk of drop out simultaneously, providing eGFR slopes corrected for both censoring due to death and dialysis. As the risk of death did not differ substantially between men and women in our cohort, adjustment had little effect on the difference in slopes between men and women. However, as Nitsch *et al.* demonstrated in their meta-analysis, the mortality risk difference between men and women is far larger in earlier stages of CKD [22]. In such populations, accounting for mortality would have likely had a larger effect on the difference in renal slopes between men and women, accounting for more of the difference in renal decline compared with our cohort. Conversely, accounting for censoring caused by dialysis initiation led to marginally steeper adjusted slopes, reflecting the faster renal decline in patients that were censored due to dialysis initiation. As the risk of dialysis was higher in men, the unbiased difference in renal function decline between the sexes was amplified slightly after accounting for this event, although this change in effect was not statistically significant.

Studying renal decline by sex is complicated by a sex-specific selection process throughout the pre-dialysis period. Contrary to our expectations, we found slower renal declines in older women. The literature on the effect of age on renal decline is inconsistent, with some reporting faster renal declines with increasing age [17, 48], and others reporting the opposite [49, 50]. Potential explanations for our findings may be a differential mortality rate in men and women (prior to inclusion), which may be inclined to select the healthier surviving women with slowly progressing CKD. One may also hypothesize that this finding may be caused by a sex-dependent decrease in muscle mass with age, biasing our eGFRs. Lastly, considering all patients in the EQUAL cohort are referred, there may be selection mechanisms at play in the referral patterns.

The main strength of our study is that we apply JMs to deal with informative censoring caused by mortality and dialysis initiation, providing unbiased estimates of renal decline.

Furthermore, patients in our cohort were prospectively included when their eGFR dropped below the pre-defined level of 20 mL/min/1.73 m<sup>2</sup>, thus minimizing the risk of survivor bias. Our study is also subject to several limitations. Preferably, we would have used measured GFR by a reference method to estimate the slope of renal decline; however, measuring GFR with a tracer technology was unfortunately not feasible in a cohort study of the size of EQUAL. The use of eGFR in the main analysis may partly reflect muscle mass, which may disproportionately bias eGFR estimates in women [51]. Nonetheless, others have shown mGFR to perform similarly to eGFR. Lastly, due to the observational nature of our study, residual confounding may play a role, and therefore the results should be interpreted accordingly.

In older patients with advanced CKD, we demonstrate faster declines in renal function in men compared with women, even after adjustment for multiple groups of mediators. Importantly, informative events such as death and dialysis initiation explained little of the difference in renal decline between the sexes in our advanced CKD cohort. In diabetics, however, both men and women declined at a similar rate, demonstrating a disproportional negative impact of diabetes in women. Our results improve understanding of the mechanisms underlying the differences in renal function decline between the sexes and warrant further research to develop the sex-specific interventions needed to achieve individualized management and treatment.

## SUPPLEMENTARY DATA

Supplementary data are available at [ndt](https://academic.oup.com/ndt) online.

## ACKNOWLEDGEMENTS

We would like to thank all the patients and health professionals participating in the EQUAL study.

## FUNDING

Funding was received from the European Renal Association–European Dialysis and Transplant Association (ERA-EDTA), the Swedish Medical Association (SLS), the Stockholm County Council ALF, Njurfonden (Sweden), the Italian Society of Nephrology (SIN-Reni), the Dutch Kidney Foundation (SB 142), the Young Investigators grant in Germany and the National Institute for Health Research (NIHR) in the UK.

## CONFLICT OF INTEREST STATEMENT

M.E. received payment for lectures and participated in advisory board meetings for Astellas, received lecture honoraria from Vifor, participated in the advisory board and received study grants for AstraZeneca.

## REFERENCES

1. Roth M, Roderick P, Mindell J. Kidney disease and renal function. In: Craig R, Mindell J, eds. *Health survey for England 2010*. Leeds: NHS Information Centre, 2011. Chapter 8, 1–27
2. Murphy D, McCulloch CE, Lin F et al.; for the Centers for Disease Control and Prevention Chronic Kidney Disease Surveillance Team. Trends in prevalence of chronic kidney disease in the United States. *Ann Intern Med* 2016; 165: 473–481



3. Ong-Ajyooth L, Vareesangthip K, Khonputsa P *et al*. Prevalence of chronic kidney disease in Thai adults: a national health survey. *BMC Nephrol* 2009; 10: 35
4. Gasparini A, Evans M, Coresh J *et al*. Prevalence and recognition of chronic kidney disease in Stockholm healthcare. *Nephrol Dial Transplant* 2016; 31: 2086–2094
5. Tanamas SK, Magliano DJ, Lynch B *et al*. AUSDIAB 2012: the Australian diabetes, obesity and lifestyle study. Melbourne: Baker IDI Heart and Diabetes Institute, 2013: 1–92
6. Foley RN, Chen S-C, Solid CA *et al*. Early mortality in patients starting dialysis appears to go unregistered. *Kidney Int* 2014; 86: 1–7
7. United States Renal Data System. *2018 USRDS Annual Data Report: Epidemiology of Kidney Disease in the United States*. National Institutes of Health, National Institute of Diabetes and Digestive and Kidney Diseases, Bethesda, MD, 2018
8. ERA-EDTA Registry. *ERA-EDTA Registry Annual Report 2016*. Amsterdam UMC, location AMC, Department of Medical Informatics, Amsterdam, the Netherlands, 2018
9. Sparke C, Moon L, Green F *et al*. Estimating the total incidence of kidney failure in Australia including individuals who are not treated by dialysis or transplantation. *Am J Kidney Dis* 2013; 61: 413–419
10. Carrero JJ, Hecking M, Chesnaye NC *et al*. Sex and gender disparities in the epidemiology and outcomes of chronic kidney disease. *Nat Rev Nephrol* 2018; 14: 151–164
11. Levin A, Djurdjev O, Beaulieu M *et al*. Variability and risk factors for kidney disease progression and death following attainment of stage 4 CKD in a referred cohort. *Am J Kidney Dis* 2008; 52: 661–671
12. Ricardo AC, Yang W, Sha D *et al*; on behalf of the CRIC Investigators. Sex-related disparities in CKD progression. *J Am Soc Nephrol* 2019; 30: 137–146
13. Eriksen BO, Ingebreten OC. The progression of chronic kidney disease: a 10-year population-based study of the effects of gender and age. *Kidney Int* 2006; 69: 375–382
14. Halbesma N, Brantsma AH, Bakker SJL *et al*. Gender differences in predictors of the decline of renal function in the general population. *Kidney Int* 2008; 74: 505–512
15. Neugarten J, Acharya A, Silbiger SR. Effect of gender on the progression of nondiabetic renal disease: a meta-analysis. *J Am Soc Nephrol* 2000; 11: 319–329
16. Chang PY, Chien LN, Lin YF *et al*. Risk factors of gender for renal progression in patients with early chronic kidney disease. *Medicine (Baltimore)* 2016; 95: e4203
17. Imai E, Horio M, Yamagata K *et al*. Slower decline of glomerular filtration rate in the Japanese general population: a longitudinal 10-year follow-up study. *Hypertens Res* 2008; 31: 433–441
18. Coggins CH, Lewis JB, Caggiula AW *et al*. Nephrology dialysis transplantation differences between women and men with chronic renal disease. *Nephrol Dial Transplant* 1998; 13: 1430–1437
19. Norris KC, Greene T, Kopple J *et al*. Baseline predictors of renal disease progression in the African American study of hypertension and kidney disease. *J Am Soc Nephrol* 2006; 17: 2928–2936
20. John R, Webb M, Young A *et al*. Unreferred chronic kidney disease: a longitudinal study. *Am J Kidney Dis* 2004; 43: 825–835
21. Jafar TH, Schmid CH, Stark PC *et al*. The rate of progression of renal disease may not be slower in women compared with men: a patient-level meta-analysis. *Nephrol Dial Transplant* 2003; 18: 2047–2053
22. Nitsch D, Grams M, Sang Y *et al*; for the Chronic Kidney Disease Prognosis Consortium. Associations of estimated glomerular filtration rate and albuminuria with mortality and renal failure by sex: a meta-analysis. *BMJ* 2013; 346: f324
23. Ravani P, Fiocco M, Liu P *et al*. Influence of mortality on estimating the risk of kidney failure in people with Stage 4 chronic kidney disease *J Am Soc Nephrol* 2019; 30: 2219–2227
24. Wright S, Klausner D, Baird B *et al*. Timing of dialysis initiation and survival in ESRD. *Clin J Am Soc Nephrol* 2010; 5: 1828–1835
25. Neugarten J, Golestaneh L. Influence of sex on the progression of chronic kidney disease. *Mayo Clin Proc* 2019; 94: 1339–1356
26. Jager KJ, Ocak G, Drechsler C *et al*. The EQUAL study: a European study in chronic kidney disease stage 4 patients. *Nephrol Dial Transplant* 2012; 27 (Suppl 3): 27–31
27. Levey AS, Stevens LA, Schmid CH *et al*; for the CKD-EPI (Chronic Kidney Disease Epidemiology Collaboration). A new equation to estimate glomerular filtration rate. *Ann Intern Med* 2009; 150: 604–612
28. Henderson R, Diggle P, Dobson A. Joint modelling of longitudinal measurements and event time data. *Biostatistics* 2000; 1: 465–480
29. Hu C, Sale ME. A joint model for nonlinear longitudinal data with informative dropout. *J Pharmacokinet Pharmacodyn* 2003; 30: 83–103
30. Wu MC, Carroll RJ. Estimation and comparison of changes in the presence of informative right censoring by modeling the censoring process. *Biometrics* 1988; 44: 175
31. Rizopoulos D. The R Package JMbayes for fitting joint models for longitudinal and time-to-event data using MCMC. *J Stat Softw* 2016; 72: 1–46; doi: 10.18637/jss.v072.i07
32. Paternoster R, Brame R, Mazerolle P *et al*. Using the correct statistical test for the equality of regression coefficients. *Criminology* 1998; 36: 859–866
33. Pottel H, Hoste L, Dubourg L *et al*. An estimated glomerular filtration rate equation for the full age spectrum. *Nephrol Dial Transplant* 2016; 31: 798–806
34. Nyman U, Grubb A, Larsson A *et al*. The revised Lund-Malmö GFR estimating equation outperforms MDRD and CKD-EPI across GFR, age and BMI intervals in a large Swedish population. *Clin Chem Lab Med* 2014; 52: 815–824
35. Briganti EM, Branley P, Chadban SJ *et al*. Smoking is associated with renal impairment and proteinuria in the normal population: the AusDiab kidney study. *Am J Kidney Dis* 2002; 40: 704–712
36. Rk D, Ek J. Estrogen-induced cardiorenal protection: potential cellular, biochemical, and molecular mechanisms. *Am J Physiol Ren Physiol* 2001; 280: 365–388
37. Carrero JJ. Gender differences in chronic kidney disease: underpinnings and therapeutic implications. *Kidney Blood Press Res* 2010; 33: 383–392
38. Silbiger S, Neugarten J. Gender and human chronic renal disease. *Gen Med* 2008; 5 (Suppl 1): 3–10
39. Hemmelgarn BR, Zhang J, Manns BJ *et al*. Progression of kidney dysfunction in the community-dwelling elderly. *Kidney Int* 2006; 69: 2155–2161
40. de Hauteclouque A, Ragot S, Slaoui Y *et al*; The SURDIAGENE Study group. The influence of sex on renal function decline in people with Type 2 diabetes. *Diabet Med* 2014; 31: 1121–1128
41. Krolewski AS, Skupien J, Rossing P *et al*. Fast renal decline to end-stage renal disease: an unrecognized feature of nephropathy in diabetes. *Kidney Int* 2017; 91: 1300–1311
42. Kajiwara A, Kita A, Saruwatari J *et al*. Sex differences in the renal function decline of patients with type 2 diabetes. *J Diabetes Res* 2016; 2016: 1–8
43. Margaret K, Yu Lyles CR, Bent-Shaw LA *et al*. Risk factor, age and sex differences in chronic kidney disease prevalence in a diabetic cohort: the Pathways Study. *Am J Nephrol* 2012; 36: 245–251
44. Carrero JJ, de Mutsert R, Axelsson J *et al*; for the NECOSAD Study Group. Sex differences in the impact of diabetes on mortality in chronic dialysis patients. *Nephrol Dial Transplant* 2011; 26: 270–276
45. Huxley RR, Peters SAE, Mishra GD *et al*. Risk of all-cause mortality and vascular events in women versus men with type 1 diabetes: a systematic review and meta-analysis. *Lancet Diabetes Endocrinol* 2015; 3: 198–206
46. Xu G, You D, Wong L *et al*. Risk of all-cause mortality in women versus men with type 2 diabetes: a systematic review and meta-analysis. *Eur J Endocrinol* 2019; 180: 243–255
47. Maric C, Sullivan S. Estrogens and the diabetic kidney. *Gen Med* 2008; 5: S103–S113
48. Cohen E, Nardi Y, Krause I *et al*. A longitudinal assessment of the natural rate of decline in renal function with age. *J Nephrol* 2014; 27: 635–641
49. O'Hare AM, Choi AI, Bertenthal D *et al*. Age affects outcomes in chronic kidney disease. *J Am Soc Nephrol* 2007; 18: 2758–2765
50. Evans M, Fryzek JP, Elinder CG *et al*. The natural history of chronic renal failure: results from an unselected, population-based, inception cohort in Sweden. *Am J Kidney Dis* 2005; 46: 863–870
51. Zaman T, Filipowicz R, Beddhu S. Implications and importance of skeletal muscle mass in estimating glomerular filtration rate at dialysis initiation. *J Ren Nutr* 2013; 23: 233–236

Received: 21.1.2020; Editorial decision: 6.4.2020