



Universiteit
Leiden
The Netherlands

Near-infrared fluorescence imaging with indocyanine green in vascular surgery

Hoven, P. van den

Citation

Hoven, P. van den. (2022, June 9). *Near-infrared fluorescence imaging with indocyanine green in vascular surgery*. Retrieved from <https://hdl.handle.net/1887/3309684>

Version: Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/3309684>

Note: To cite this publication please use the final published version (if applicable).

Chapter 4

Perfusion patterns in patients with chronic limb-threatening ischemia versus control patients using near-infrared fluorescence imaging with indocyanine green

P. van den Hoven ¹, L.N. Goncalves ¹, P.H.A. Quax ¹, C.S.P. van Rijswijk ¹, J. van Schaik ¹, A. Schepers ¹, A.L. Vahrmeijer ¹, J.F. Hamming ¹, J.R. van der Vorst ¹

1. Leiden University Medical Center, Leiden, The Netherlands

Published in Biomedicines, October 2021.

Abstract

Introduction

In assessing the severity of lower extremity arterial disease (LEAD), physicians rely on clinical judgement supported by conventional measurements of macrovascular blood flow. However, current diagnostic techniques provide no information about regional tissue perfusion and are of limited value in patients with chronic limb-threatening ischemia (CLTI). Near-infrared (NIR) fluorescence imaging using indocyanine green (ICG) has been used extensively in perfusion studies and is a possible modality for tissue perfusion measurement in patients with CLTI.

Methods

In this prospective cohort study ICG NIR fluorescence imaging was performed in patients with CLTI and control patients using the Quest Spectrum Platform[®]. The time-intensity curves were analyzed using the Quest Research Framework[®]. Fourteen parameters were extracted.

Results

Successful ICG NIR fluorescence imaging was performed in 19 patients with CLTI and 16 control patients. The time to maximum intensity was lower for CLTI patients (90.5 vs 143.3 seconds, $p=0.002$). For inflow parameters, the maximum slope, the normalized maximum slope and the ingress rate were all significantly higher in the CLTI group.

Conclusion

Inflow parameters observed in patients with CLTI were superior to the control group. Possible explanations for the increased inflow include damage to the regulatory mechanisms of the microcirculation, arterial stiffness and transcapillary leakage.

Introduction

Lower-extremity arterial disease (LEAD) is most often caused by atherosclerosis (1, 2). Subsequent hemodynamic alterations leading to hypoxia can trigger a cascade of events leading to macro- and microvascular changes in the affected limb (3). In the most advanced stage, chronic limb-threatening ischemia (CLTI), blood supply to the lower extremity is insufficient to meet metabolic needs (2, 4). For these patients, a common finding in physical examination of the lower extremities is the appearance of “dependent rubor” or “blanching”, which is presumably caused by dysfunction of the venoarteriolar reflex (5). In assessing the severity of LEAD, physicians often rely on their clinical judgement of the extremities. The diagnosis is confirmed using conventional measurements of macrovascular blood flow including the ankle-brachial index (ABI), toe pressure measurement, computed tomography (CT) angiography, magnetic resonance angiography and digital subtraction angiography. However, these techniques provide no information about regional tissue perfusion and have shown to be of limited value in patients with CLTI (6). New emerging methods for the assessment of regional tissue perfusion include dynamic volume perfusion CT, laser speckle imaging (LSI) and near-infrared (NIR) fluorescence imaging using indocyanine green (ICG) (7-9). ICG NIR fluorescence imaging has been used in various medical fields for assessment of tissue perfusion, including cardiac and reconstructive surgery (10, 11). This imaging technique measures fluorescence in the NIR light spectrum (700-1000nm), which is characterized by low tissue autofluorescence and deep tissue penetration (12). Upon intravenous administration of ICG, which has a peak emission of 814nm, the camera measures the NIR fluorescence intensity over time. The feasibility of ICG as a fluorophore in perfusion assessment is explained by its confinement to the intravascular compartment due to the binding with plasma proteins (13). For assessment of skin perfusion, ICG NIR fluorescence imaging is currently used intraoperatively in reconstructive surgery to predict flap viability (14). For patients with LEAD, similar results were seen in predicting skin necrosis following amputation surgery (15). However, these findings rely on qualitative analyses, meaning the observer subjectively grades the visualized NIR fluorescence intensity. To quantify and grade regional tissue perfusion, better understanding of the different perfusion patterns as observed with ICG NIR fluorescence imaging is needed. Several studies have been performed to quantify ICG NIR fluorescence imaging between patients with different stages of LEAD (16-18). However, inconsistency is seen between stages and it is unclear whether advanced stages of LEAD alter the in- and outflow of ICG (16). Furthermore, there is limited information about the perfusion patterns of ICG NIR fluorescence imaging in control patients. Therefore, as a first step in the quantification of tissue perfusion using ICG NIR fluorescence imaging, the aim of this study was to analyze the perfusion patterns seen in patients with CLTI and compare these to non-LEAD control patients.

Methods

This prospective cohort study was approved by the Medical Research and Ethics Committee of the Leiden University Medical Center and registered in the Dutch Trial Register with number NL7531. Patients with CLTI as classified according to the Global Vascular Guidelines on the management of CLTI, were included (4). These were patients diagnosed with either Fontaine stage 3 or stage 4 LEAD. The control group consisted of patients who underwent intravenous ICG administration prior to liver metastasectomy. Patients were included from December 2018 until April 2021 in a single academic hospital in the Netherlands. Exclusion criteria were allergy or hypersensitivity to sodium iodide, iodide, or ICG; known hyperthyroidism or autonomous thyroid adenoma, pregnancy, kidney failure (eGFR <45) and/or severe liver failure. Informed consent was obtained in all patients. ABI and toe pressure measurements were performed in all patients. As additional measurement for patients with CLTI, duplex ultrasound measurements of the feet were performed and the highest acceleration in either the dorsalis pedis artery or posterior tibial artery was reported. These acceleration measurements are described in detail in an earlier study by Brouwers et al. and are performed to assess the severity of arterial stenosis (19). performed as routine measurements for The Quest Spectrum Platform® (Quest Medical Imaging, Middenmeer, The Netherlands) was used to perform ICG NIR fluorescence imaging (Figure 1).




Figure 1. ICG NIR fluorescence imaging camera setup.

This imaging system is capable of measuring both visible light as well as the NIR signal of ICG. Patients with CLTI were administered an intravenous bolus injection of 0.1 mg/kg ICG (VERDYE 25 mg, Diagnostic Green GmbH, Aschheim-Dornach, Germany) using a peripheral venous line in the cubital fossa or on the dorsum of the hand. Patients in the control group were administered a bolus injection of 10mg ICG according to local hospital guidelines. Following administration of ICG, the NIR fluorescence intensity in both feet was recorded for 10 minutes (Figure 2).




Figure 2. ICG NIR fluorescence imaging in a control patient showing the visual image (left), merged image (middle) and NIR fluorescence intensity at the dorsum of both feet.

Measurements were performed on patients in a supine position following a rest period of at least 10 minutes in a room cleared of ambient light. The camera was placed perpendicular to the dorsum of both feet at a distance of 50cm. The NIR fluorescence videos were analyzed using the Quest Research Framework® (Quest Medical Imaging, Middenmeer, the Netherlands). The whole foot was selected as region of interest (ROI). Upon selection of the ROI, the software creates a time-intensity curve of the measured intensity in arbitrary units (a.u.). A tracker was used to ensure the ROI was synchronized with leg movement. Fourteen parameters were extracted from these curves, of which an explanation is given in Figure 3. The ingress rate was defined as the intensity increase per second from baseline to maximum intensity. The Tmax was measured starting at the point of 10% intensity increase upon baseline. The time-intensity curves were also analyzed after normalization for maximum intensity. The curves extracted from these curves were, in percentage per second, the maximum slope ingress and the maximum slope egress. Starting time was defined as an increase of 1 arbitrary unit for the intensity curves and 1% for the normalized curves. Statistical analyses were performed using IBM SPSS Statistics 25 (IBM Corp. Released 2017. IBM SPSS Statistics for Windows, Version 25.0. Armonk, NY, USA: IBM Corp.). Parameters were compared using the Mann-Whitney U test.




Figure 3. Time-intensity curve with extracted parameters.

Results

Successful ICG NIR fluorescence imaging measurements were performed in 35 patients. Nineteen patients presented with LEAD of whom 28 limbs were classified as CLTI. The control group consisted of 16 patients with a total of 32 limbs. The characteristics for each group are displayed in Table 1. For the CLTI group, 10 limbs were classified as Fontaine stage 4. Compared to the control group, patients in the CLTI group were more likely to present with diabetes, hypertension and smoking. The mean ABI in the CLTI group was 0.77 versus 1.11 in the control group. The ABI in the CLTI group was not measurable in 9 out of 28 limbs. The acceleration measured on duplex ultrasonography was measured in 22 CLTI limbs with a mean of 0.93 m/s².

ICG NIR fluorescence imaging parameters

The results on ICG NIR fluorescence imaging for the 14 extracted parameters are displayed in Table 2. The mean maximum intensity was significantly lower in the control group (37.9 vs 25.8 a.u., $p < 0.001$). Furthermore, the time to maximum

intensity (i.e. Tmax) was reached earlier in the CLTI group (90.5 vs 143.3 seconds, $p=0.002$). When taking a closer look at the inflow parameters, the maximum slope, the normalized maximum slope and the ingress rate were all significantly higher in the CLTI group (2.0 vs 0.6 a.u./s, $p<0.001$; 4.2 vs 2.4 %/s, $p<0.001$; 1.0 vs 0.2 a.u./s, $p<0.001$). For outflow parameters, a significant difference was seen for the maximum slope egress, which was higher in the control group (0.5 vs 0.2 a.u./s, $p=0.005$). No significant difference was observed for the normalized maximum slope egress (1.0 vs 0.8 %/s, $p=0.733$). Comparison of the AUC for different intervals following the Tmax displayed no significant difference between the CLTI and control group.

Table 1. Patient characteristics.

	CLTI	Control group
N (limbs)	19 (28)	16 (32)
Age (SD)	70.4 (7.5)	66.6 (12.3)
Diabetes Mellitus (%)	9 (47.4)	3 (18.8)
Hypertension (%)	15 (78.9)	7 (43.8)
Active smoking (%)	5 (26.3)	1 (6.3)
Fontaine stage limbs, n(%)		
3	18 (64.3)	-
4	10 (35.7)	-
Mean ABI (SD)	0.77 (0.34)	1.11 (0.10)
Mean TP (SD)	44 (25)	106 (22)
Acceleration (SD)	0.93 (1.23)	-

Abbreviations: SD, standard deviation; CLTI, chronic limb-threatening ischemia; ABI, ankle-brachial index; TP, Toe Pressure).

Time-intensity curves

The time-intensity curves for the control group and CLTI group are displayed in Figure 4. Results are displayed for the intensity-related curves (above) and the normalized curves (below) for both groups. Time-intensity curves displaying the absolute intensity change over time show an overall higher absolute intensity for the CLTI group. Following a steep incline in intensity increase for the CLTI group, the outflow seems comparable with control patients. The intensity-related curves show a widespread distribution, especially in the CLTI group. In this group, the maximum slope ingress (2.0%/s) has a standard deviation of 2.5 (Table 2). For the AUC egress parameters, standard deviations between 10.0% and 13.5% are observed. When normalizing these time-intensity curves for maximum intensity, both groups display a more narrow distribution in all parameters. For the normalized maximum slope in the CLTI group (4.2%/s), a standard deviation of 3.1% was observed. When looking at AUC egress parameters, the standard deviations have a distribution of 1.8 to 6.1%.

Table 2. ICG NIR fluorescence imaging parameters.

Parameter	CLTI	Controls	p-value
Maximum intensity (SD)	37.9 (14.4)	25.8 (10.8)	<0.001
Maximum slope ingress (SD)	2.0 (2.5)	0.6 (0.4)	<0.001
Normalized maximum slope (SD)	4.2 (3.1)	2.4 (1.2)	<0.001
Ingress rate (SD)	1.0 (1.7)	0.2 (0.2)	<0.001
AUC ingress 10 (SD)	47.4 (2.2)	48.8 (3.3)	0.073
AUC ingress (SD)	71.4 (6.3)	70.6 (3.8)	0.213
Tmax (SD)	90.5 (53.4)	143.3 (64.5)	0.002
Maximum slope egress (SD)	0.5 (0.7)	0.2 (0.1)	0.005
Normalized maximum slope egress (SD)	1.0 (0.9)	0.8 (0.3)	0.733
AUC egress 60 (SD)	92.8 (10.0)	96.7 (1.8)	0.113
AUC egress 120 (SD)	87.9 (11.9)	92.8 (2.3)	0.127
AUC egress 180 (SD)	82.9 (12.9)	88.3 (4.3)	0.164
AUC egress 240 (SD)	78.2 (13.3)	83.8 (5.4)	0.168
AUC egress 300 (SD)	73.7 (13.5)	73.3 (6.0)	0.271

Abbreviations: SD, standard deviation; CLTI, chronic limb-threatening ischemia; AUC, area under the curve.




Figure 4. Absolute intensity - and normalized time-intensity curves for the CLTI group and control group: (a) Absolute time-intensity curve CLTI group, (b) Absolute time-intensity curve control group, (c) Normalized time-intensity curve CLTI group, (d) Normalized time-intensity curve control group.

Discussion

This study demonstrates the different perfusion patterns as seen on ICG NIR fluorescence imaging between patients with CLTI and control patients. Interestingly, most inflow parameters observed in patients with CLTI were higher compared to the control group. Concerning the outflow of ICG, however, no significant differences were observed. Furthermore, there is a widespread distribution of measured intensity over time in both groups. There are several earlier studies reporting the use of ICG NIR fluorescence imaging for perfusion assessment in patients with LEAD as well as control patients (7, 16, 18, 20-25). In these studies, an abundance of parameters have been examined, which have been compared to varying diagnosis measurements, including the ABI, TP and transcutaneous oxygen pressure measurements. Patterns of foot perfusion in non-LEAD control patients were analyzed in one study (18). Regarding inflow parameters, Igari et al. found a prolonged time to maximum intensity for patients with LEAD compared to control patients (18). No statistical differences were seen for the maximum intensity and T1/2 between the two groups. The differences in perfusion patterns amongst various stages of LEAD were analyzed in several studies. When comparing inflow parameters between different stages of LEAD, Terasaki et al. observed a prolonged T1/2 for Fontaine stage 3 compared to stage 2, however, this was not observed for stage 4. Regarding outflow, their study concluded that a percentage decrease of 90% in maximum measured intensity was the most accurate parameter in diagnosing LEAD. For patients with CLTI, Venermo et al. found an increase in inflow, the PDE10, to be strongly correlated to the transcutaneous oxygen pressure in patients with diabetes mellitus (23). The same parameter was moderately correlated in patients without diabetes mellitus, suggesting a difference in perfusion patterns between these groups. According to the findings in these earlier studies and the results found in this study, the hypothesis that LEAD progression leads to diminished in- and outflow of ICG is debatable. Several mechanisms might contribute to the increased inflow of ICG seen in patients with CLTI in this study. First, ICG NIR fluorescence imaging is able to penetrate tissue to a depth of several millimeters (26). Therefore, this imaging technique mainly visualizes the skin with superficial vessels and the superior part of the subcutaneous tissue, i.e. the microcirculation. Nutritional capillaries of this microcirculation in the foot account for approximately 15% of total foot blood flow, which is regulated by various mechanisms including arteriovenous (AV) shunts (27). For patients with LEAD and CLTI in particular, the diminished blood flow has led to hypoxia altering microcirculatory function and damaging these regulatory mechanisms (3, 5). The dysfunction of AV shunts might lead to a relative increase of blood flow to the skin in patients with CLTI, which also explains the “dependent rubor” seen in this group. Secondly, atherosclerosis leads to stiffness of the arterial wall which is a common finding in patients with CLTI and leading to an increased pulse wave velocity (28). In a healthy arterial system, the blood flow is gradually transmitted

to the peripheral tissue due to compliance of the vessel wall (29). This might explain the more gradual perfusion pattern seen in the control group. Furthermore, damage to the microcirculation in CLTI leads to transcapillary leakage, which might further enhance the measured NIR fluorescence intensity. Although a higher dosage of ICG was administered in the majority of patients in the control group, it is unlikely that this would have influenced the perfusion pattern. Besides, an overall lower absolute intensity was seen in this group. To confirm the findings on increased inflow, a larger cohort of patients with CLTI is needed. Therefore, due to the small sample size, the conclusions in this study must be perceived as a proof of concept. Besides the small cohort size of patients with CLTI, this study is limited by the heterogenous aspect of the CLTI population. In particular for patients with diabetes mellitus, skin perfusion follows a different pattern than LEAD Fontaine stage 4 patients without diabetes mellitus. Therefore, future studies should distinguish between CLTI patients with and without diabetes mellitus. Furthermore, the control group used in the present study were patients scheduled for liver metastasectomy and therefore might not resemble healthy volunteers in terms of comorbidities. Although LEAD was excluded based on medical history and ABI measurements, there could be differences in perfusion patterns with healthy volunteers. Therefore, in future patient selection and further understanding perfusion patterns, healthy volunteers should be taken into account as well. With regards to the NIR fluorescence intensity analysis, the use of normalized time-intensity curves seems rational, since intensity-related parameters are prone to multiple influencing factors, including camera distance and ICG dosage (30, 31). This normalization minimizes the effect of these influencing factors on the measured intensity and contributes to a more narrow distribution, as can be seen in the time-intensity curves in this study. The use of this normalization might be of use in future research on quantification of tissue perfusion with ICG NIR fluorescence imaging.

Conclusion

An increase in inflow parameters was observed with ICG NIR fluorescence imaging in patients with CLTI compared to control patients. This is possibly explained by damage to the regulatory mechanisms of the microcirculation and arterial stiffness. In the search for providing cut-off values for adequate perfusion, more research in larger cohorts is needed on in- and outflow patterns for control patients and various stages of LEAD.

Reference list

1. Weitz JI, Byrne J, Clagett GP, Farkouh ME, Porter JM, Sackett DL, et al. Diagnosis and treatment of chronic arterial insufficiency of the lower extremities: a critical review. *Circulation*. 1996;94(11):3026-49.
2. Aboyans V, Ricco JB, Bartelink MEL, Bjorck M, Brodmann M, Cohnert T, et al. 2017 ESC Guidelines on the Diagnosis and Treatment of Peripheral Arterial Diseases, in collaboration with the European Society for Vascular Surgery (ESVS). *Rev Esp Cardiol (Engl Ed)*. 2018;71(2):111.
3. Krishna SM, Moxon JV, Golledge J. A review of the pathophysiology and potential biomarkers for peripheral artery disease. *Int J Mol Sci*. 2015;16(5):11294-322.
4. Conte MS, Bradbury AW, Kolh P, White JV, Dick F, Fitridge R, et al. Global vascular guidelines on the management of chronic limb-threatening ischemia. *J Vasc Surg*. 2019;69(6S):3S-125S e40.
5. Abularrage CJ, Sidawy AN, Aidinian G, Singh N, Weiswasser JM, Arora S. Evaluation of the microcirculation in vascular disease. *J Vasc Surg*. 2005;42(3):574-81.
6. Misra S, Shishehbor MH, Takahashi EA, Aronow HD, Brewster LP, Bunte MC, et al. Perfusion Assessment in Critical Limb Ischemia: Principles for Understanding and the Development of Evidence and Evaluation of Devices: A Scientific Statement From the American Heart Association. *Circulation*. 2019;140(12):e657-e72.
7. van den Hoven P, Ooms S, van Manen L, van der Bogt KEA, van Schaik J, Hamming JF, et al. A systematic review of the use of near-infrared fluorescence imaging in patients with peripheral artery disease. *J Vasc Surg*. 2019;70(1):286-97 e1.
8. Kikuchi S, Miyake K, Tada Y, Uchida D, Koya A, Saito Y, et al. Laser speckle flowgraphy can also be used to show dynamic changes in the blood flow of the skin of the foot after surgical revascularization. *Vascular*. 2019;27(3):242-51.
9. Cindil E, Erbas G, Akkan K, Cerit MN, Sendur HN, Zor MH, et al. Dynamic Volume Perfusion CT of the Foot in Critical Limb Ischemia: Response to Percutaneous Revascularization. *AJR Am J Roentgenol*. 2020;214(6):1398-408.
10. Driessen C, Arnardottir TH, Lorenzo AR, Mani MR. How should indocyanine green dye angiography be assessed to best predict mastectomy skin flap necrosis? A systematic review. *J Plast Reconstr Aesthet Surg*. 2020;73(6):1031-42.
11. Dupree A, Riess H, Detter C, Debus ES, Wipper SH. Utilization of indocyanine green fluorescent imaging (ICG-FI) for the assessment of microperfusion in vascular medicine. *Innov Surg Sci*. 2018;3(3):193-201.
12. Vahrmeijer AL, Hutteman M, van der Vorst JR, van de Velde CJ, Frangioni JV. Image-guided cancer surgery using near-infrared fluorescence. *Nat Rev Clin Oncol*. 2013;10(9):507-18.
13. Schaafsma BE, Mieog JS, Hutteman M, van der Vorst JR, Kuppen PJ, Lowik CW, et al. The clinical use of indocyanine green as a near-infrared fluorescent contrast agent for image-guided oncologic surgery. *J Surg Oncol*. 2011;104(3):323-32.
14. Parmeshwar N, Sultan SM, Kim EA, Piper ML. A Systematic Review of the Utility of Indocyanine Angiography in Autologous Breast Reconstruction. *Ann Plast Surg*. 2020; Publish Ahead of Print.

15. Zimmermann A, Roenneberg C, Wendorff H, Holzbach T, Giunta RE, Eckstein HH. Early postoperative detection of tissue necrosis in amputation stumps with indocyanine green fluorescence angiography. *Vasc Endovascular Surg.* 2010;44(4):269-73.
16. Terasaki H, Inoue Y, Sugano N, Jibiki M, Kudo T, Lepantalo M, et al. A quantitative method for evaluating local perfusion using indocyanine green fluorescence imaging. *Ann Vasc Surg.* 2013;27(8):1154-61.
17. Kang Y, Lee J, Kwon K, Choi C. Application of novel dynamic optical imaging for evaluation of peripheral tissue perfusion. *Int J Cardiol.* 2010;145(3):e99-101.
18. Igari K, Kudo T, Uchiyama H, Toyofuku T, Inoue Y. Indocyanine green angiography for the diagnosis of peripheral arterial disease with isolated infrapopliteal lesions. *Ann Vasc Surg.* 2014;28(6):1479-84.
19. Brouwers J, van Doorn LP, van Wissen RC, Putter H, Hamming JF. Using maximal systolic acceleration to diagnose and assess the severity of peripheral artery disease in a flow model study. *J Vasc Surg.* 2020;71(1):242-9.
20. Kang Y, Lee J, Kwon K, Choi C. Dynamic fluorescence imaging of indocyanine green for reliable and sensitive diagnosis of peripheral vascular insufficiency. *Microvasc Res.* 2010;80(3):552-5.
21. Zimmermann A, Roenneberg C, Reeps C, Wendorff H, Holzbach T, Eckstein HH. The determination of tissue perfusion and collateralization in peripheral arterial disease with indocyanine green fluorescence angiography. *Clin Hemorheol Microcirc.* 2012;50(3):157-66.
22. Igari K, Kudo T, Uchiyama H, Toyofuku T, Inoue Y. Quantitative evaluation of microvascular dysfunction in peripheral neuropathy with diabetes by indocyanine green angiography. *Diabetes Res Clin Pract.* 2014;104(1):121-5.
23. Venermo M, Settembre N, Alback A, Vikatmaa P, Aho PS, Lepantalo M, et al. Pilot Assessment of the Repeatability of Indocyanine Green Fluorescence Imaging and Correlation with Traditional Foot Perfusion Assessments. *Eur J Vasc Endovasc Surg.* 2016;52(4):527-33.
24. Nishizawa M, Igari K, Kudo T, Toyofuku T, Inoue Y, Uetake H. A Comparison of the Regional Circulation in the Feet between Dialysis and Non-Dialysis Patients using Indocyanine Green Angiography. *Scand J Surg.* 2016.
25. Goncalves LN, van den Hoven P, van Schaik J, Leeuwenburgh L, Hendricks CHF, Verduijn PS, et al. Perfusion Parameters in Near-Infrared Fluorescence Imaging with Indocyanine Green: A Systematic Review of the Literature. *Life (Basel).* 2021;11(5).
26. Frangioni JV. In vivo near-infrared fluorescence imaging. *Curr Opin Chem Biol.* 2003;7(5):626-34.
27. Rossi M, Carpi A. Skin microcirculation in peripheral arterial obliterative disease. *Biomed Pharmacother.* 2004;58(8):427-31.
28. Mendes-Pinto D, Rodrigues-Machado MDG, Navarro TP, Dardik A. Association Between Critical Limb Ischemia, the Society for Vascular Surgery Wound, Ischemia and Foot Infection (WIfI) Classification System and Arterial Stiffness. *Ann Vasc Surg.* 2020;63:250-8 e2.
29. Yu S, McEniery CM. Central Versus Peripheral Artery Stiffening and Cardiovascular Risk. *Arterioscler Thromb Vasc Biol.* 2020;40(5):1028-33.

30. Pruijboom T, van Kuijk SMJ, Qiu SS, van den Bos J, Wieringa FP, van der Hulst R, et al. Optimizing Indocyanine Green Fluorescence Angiography in Reconstructive Flap Surgery: A Systematic Review and Ex Vivo Experiments. *Surg Innov.* 2020;27(1):103-19.
31. Lutken CD, Achiam MP, Svendsen MB, Boni L, Nerup N. Optimizing quantitative fluorescence angiography for visceral perfusion assessment. *Surg Endosc.* 2020;34(12):5223-33.

