



**Universiteit
Leiden**
The Netherlands

Association of female genital schistosomiasis with the cervicovaginal microbiota and sexually transmitted infections in Zambian women

Sturt, A.S.; Webb, E.L.; Himschoot, L.; Phiri, C.R.; Mapani, J.; Mudenda, M.; ... ; Bustinduy, A.L.

Citation

Sturt, A. S., Webb, E. L., Himschoot, L., Phiri, C. R., Mapani, J., Mudenda, M., ... Bustinduy, A. L. (2021). Association of female genital schistosomiasis with the cervicovaginal microbiota and sexually transmitted infections in Zambian women. *Open Forum Infectious Diseases*, 8(9). doi:10.1093/ofid/ofab438

Version: Publisher's Version
License: [Creative Commons CC BY 4.0 license](#)
Downloaded from: <https://hdl.handle.net/1887/3249790>

Note: To cite this publication please use the final published version (if applicable).

Association of Female Genital Schistosomiasis With the Cervicovaginal Microbiota and Sexually Transmitted Infections in Zambian Women

Amy S. Sturt,¹ Emily L. Webb,² Lisa Himschoot,³ Comfort R. Phiri,⁴ Joyce Mapani,⁵ Maina Mudenda,⁵ Eyrun F. Kjetland,^{6,7} Tobias Mweene,⁴ Bruno Levecke,⁸ Govert J. van Dam,⁹ Paul L. A. M. Corstjens,¹⁰ Helen Ayles,^{1,4} Richard J. Hayes,² Lisette van Lieshout,⁹ Isaiah Hansingo,⁵ Suzanna C. Francis,² Piet Cools,^{3,8} and Amaya L. Bustinduy¹

¹Department of Clinical Research, London School of Hygiene and Tropical Medicine, London, UK, ²MRC International Statistics and Epidemiology Group, London School of Hygiene and Tropical Medicine, London, UK, ³Department of Diagnostic Sciences, Faculty of Medicine and Health Sciences, Ghent University, Ghent, Belgium, ⁴Zambart, Lusaka, Zambia, ⁵Department of Obstetrics and Gynecology, Livingstone Central Hospital, Livingstone, Zambia, ⁶Department of Infectious Diseases, Oslo University Hospital, Oslo, Norway, ⁷University of KwaZulu-Natal, Durban, South Africa, ⁸Department of Virology, Parasitology, and Immunology, Ghent University, Merelbeke, Belgium, ⁹Department of Parasitology, Leiden University Medical Center, Leiden, the Netherlands, and ¹⁰Department of Cell and Chemical Biology, Leiden University Medical Center, Leiden, the Netherlands

Background. The cervicovaginal microbiota, including sexually transmitted infections (STIs), have not been well described in female genital schistosomiasis (FGS).

Methods. Women (aged 18–31, sexually active, nonpregnant) were invited to participate at the final follow-up of the HPTN 071 (PopART) Population Cohort in January–August 2018. We measured key species of the cervicovaginal microbiota (*Lactobacillus crispatus*, *L. iners*, *Gardnerella vaginalis*, *Atopobium vaginae*, and *Candida*) and STIs (*Chlamydia trachomatis*, *Neisseria gonorrhoeae*, *Trichomonas vaginalis*, and *Mycoplasma genitalium*) using quantitative PCR (qPCR). We evaluated associations of the microbiota and STI presence and concentration with FGS (qPCR-detected *Schistosoma* DNA in any of 3 genital specimens).

Results. The presence and concentration of key cervicovaginal species did not differ between participants with (n = 30) or without FGS (n = 158). A higher proportion of participants with FGS had *T. vaginalis* compared with FGS-negative women ($P = .08$), with further analysis showing that *T. vaginalis* was more prevalent among women with ≥ 2 *Schistosoma* qPCR-positive genital specimens (50.0%, 8/16) than among FGS-negative women (21.5%, 34/158; $P = .01$).

Conclusions. We found weak evidence of an association between the presence of *T. vaginalis* and FGS, with a stronger association in women with a higher-burden FGS infection. Additional research is needed on potential between-parasite interactions, especially regarding HIV-1 vulnerability.

Keywords. cervicovaginal microbiota; female genital schistosomiasis; *Schistosoma haematobium*; sexually transmitted infection.

Female genital schistosomiasis (FGS) is caused when eggs from the waterborne parasite *S. haematobium* are entrapped in genital tissue [1]. FGS is prevalent in Sub-Saharan Africa and is associated with adverse reproductive outcomes, including ectopic pregnancy, infertility, and prevalent HIV-1 [1, 2]. The cervicovaginal environment has been described as “optimal” when it is dominated by lactic acid-producing lactobacilli, commensal microorganisms that adhere to an intact vaginal squamous epithelium [3, 4], protecting against pathogens by acidifying the vagina and producing antimicrobial substances such as bacteriocins [3]. Bacterial vaginosis (BV) and

vulvovaginal candidiasis are examples of “nonoptimal” microbiota. BV is prevalent in women in Sub-Saharan Africa [5, 6] and is characterized by a shift from lactobacilli dominance to an increase in anaerobic species or yeast [4]. BV has important sexual and reproductive health consequences, including increased risk of pelvic inflammatory disease [5]. Additionally, BV has been associated with adverse pregnancy outcomes such as preterm delivery [5, 7], a leading cause of under-5 mortality in Sub-Saharan Africa [8]. BV is also strongly associated with HIV-1 acquisition and other sexually transmitted pathogens infecting the genital tract [5, 9, 10]. Studies employing 16S rRNA sequencing to evaluate the cervicovaginal microbiota suggest that high-intensity urinary *S. haematobium* infection, in the absence of investigation for genital involvement, may alter cervicovaginal microbiota diversity [11]. However, the relationship between cervicovaginal microbiota and FGS is not well characterized.

Prevalent sexually transmitted infections (STIs) have been reported in women with [12] or in populations endemic for [13,

Received 9 April 2021; editorial decision 17 August 2021; accepted 20 August 2021.

Correspondence: Amy Sturt, MD, PhD, London School of Hygiene and Tropical Medicine, Department of Clinical Research, Keppel Street, Room 358, London, WC1E 7HT, UK (amy.sturt@lshtm.ac.uk).

Open Forum Infectious Diseases® 2021

© The Author(s) 2021. Published by Oxford University Press on behalf of Infectious Diseases Society of America. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited. <https://doi.org/10.1093/ofid/ofab438>

[14] urinary schistosomiasis. However, evaluation of genital involvement (FGS) in studies of urinary *S. haematobium* and STI co-infection is not universally performed or reported [12–14]. The biopsy prevalence of FGS in participants with urinary *S. haematobium* infection ranges from 23.0% to 75.0% ([15, 16]), making the assumption of genital involvement (FGS) in urinary *S. haematobium* infection tenuous. A reference standard diagnostic does not exist for FGS. Thus, an FGS evaluation is often based on a combination of diagnostic tests including circulating anodic antigen (CAA), urine microscopy, colposcopy, tissue-based diagnostics (biopsy, Papanicolaou smear, wet prep), and polymerase chain reaction (PCR) [17–19]. Where an FGS evaluation has been performed, studies reporting PCR-defined FGS have either not investigated or reported STI prevalence [20–22] or STI prevalence has been correlated with visual FGS findings [23]. In this cross-sectional study, we utilized PCR to detect *Schistosoma* DNA in the female genital tract and evaluated the association of PCR-defined FGS with the concentration and presence of key markers of the cervicovaginal microbiota, including STI.

METHODS

Study Setting and Participants

The cross-sectional bilharzia and HIV (BILHIV) study [18] was nested in HPTN 071 (PopART), a cluster randomized trial to measure the impact of an HIV-1 combination prevention package [24]. In HPTN 071 (PopART), HIV-1 incidence was measured in a population cohort at baseline and 12, 24, and 36 months [24]. Between January and August 2018, after the 36-month HPTN 071 (PopART) visit, community workers made home visits to women expressing interest in the BILHIV study [18]. Eligible women were aged 18–31 years, not pregnant, sexually active, and resident in 1 of 2 urban communities that participated in HPTN 071 (PopART) in Livingstone, Zambia.

Home and Clinic-Based Sample Collection

The home visit included written informed consent, a questionnaire, genital self-sampling (cervical and vaginal), and urine specimen collection, as previously described [18]. Enrolled women not currently menstruating were invited to attend Livingstone Central Hospital cervical cancer clinic, where midwives performed cervicovaginal lavage (CVL). After speculum insertion, a bulb syringe was used to flush normal saline (10 mL) across the cervix and vaginal walls for 1 minute. CVL fluid was collected from the posterior fornices (Supplementary Data). CVL and vaginal and cervical swab specimens were used for quantitative PCR (qPCR) detection of *Schistosoma*; cervical swabs were used for characterization of the microbiota and STI by qPCR; urine was used for detection of CAA and *S. haematobium* eggs by microscopy.

Cervicovaginal images were captured with a portable colposcope (MobileODT, Tel Aviv, Israel) and were evaluated (by E.F.K.) for any of the 4 recognized FGS cervicovaginal manifestations: grainy sandy patches, homogenous yellow sandy patches, rubbery papules, and abnormal blood vessels [25]. Women with at least 1 of these manifestations [25] or with any positive urine or genital *Schistosoma* diagnostic were treated free of charge with 40 mg/kg of praziquantel. Testing for STI was not performed at the point of care, and participants with suspected STI were offered syndromic management, as per local guidelines [26].

HIV-1

Laboratory-based fourth-generation HIV-1 testing (Abbott Architect HIV Ag/Ab Combo Assay) was performed for HPTN 071 (PopART) Population Cohort participants at each study visit [24].

Urine Microscopy and Circulating Anodic Antigen

Urine was centrifuged and examined by microscopy for *S. haematobium* eggs. The participant was considered positive if a pellet contained at least 1 *S. haematobium* egg [18]. A lateral flow assay utilizing up-converting reporter particles for the quantification of CAA was performed on urine samples, as previously described [18, 27]. Analyzing the equivalent of 417 μ L of urine (wet reagent, UCAAhT417), a test result indicating a CAA value >0.6 pg/mL was considered positive [28].

qPCR for Detection of *Schistosoma* DNA

Detection of the *Schistosoma*-specific internal-transcribed-spacer-2 (ITS2) target by qPCR was performed at LUMC, as previously described (Supplementary Data) [18, 29]. DNA extraction of 200 μ L of CVL or cervical or vaginal swab fluid was done with QIAamp spin columns (QIAGEN Benelux, Venlo, the Netherlands) according to manufacturer's guidelines. The qPCR output was reported in cycle threshold values (Ct values), and parasite DNA loads were categorized by the following prespecified values: high (Ct < 30), moderate ($30 \leq$ Ct < 35), low ($35 \leq$ Ct < 50), and negative (no amplification) [30].

Cervicovaginal Microbiota Characterization and STI Detection

We quantified *Lactobacillus crispatus* as a key marker of vaginal health. Additionally, we characterized markers of a “nonoptimal” cervicovaginal microbiota (*Gardnerella vaginalis* and *Atopobium vaginae*), as well as *L. iners* (an enigmatic and highly prevalent lactobacillus), *Candida spp.* and STI (*Chlamydia trachomatis*, *Neisseria gonorrhoeae*, *Mycoplasma genitalium*, and *Trichomonas vaginalis*). STI were quantified by qPCR using the S-DiaCTNG (for *C. trachomatis* and *N. gonorrhoea*) and S-DiaMGTV kits (for *M. genitalium* and *T. vaginalis*; Diagenode Diagnostics, Seraing, Belgium) on DNA from cervical swabs at Ghent

University (Ghent, Belgium) according to the manufacturer's instructions. Quantification of *A. vaginae*, *G. vaginalis*, *L. crispatus*, *L. iners*, and *Candida* species was performed in the Laboratory Bacteriology Research, Ghent, using LightCycler480, version 1.5 (Roche, Basel, Switzerland) (Supplementary Data). The concentration of each species was expressed as genome-equivalents per mL (ge/mL) [31].

FGS Definitions

In this study, we employed various diagnostic tests to evaluate urinary schistosome infection (CAA and urine microscopy) and FGS (portable colposcopy and *Schistosoma* DNA on CVL and genital swabs). As previously described [32], participants were grouped by diagnostic test results into 3 mutually exclusive categories: FGS, at least 1 positive *Schistosoma* qPCR on a genital specimen (cervical swab, vaginal swab, and/or CVL); FGS negative, negative results on all diagnostic methods; probable FGS, genital *Schistosoma* qPCRs negative but urinary schistosomiasis positive (as defined above), in combination with 1 of 4 clinical findings suggestive of FGS on any colposcope-obtained photograph [25].

Patient Consent

The study was approved by the University of Zambia Biomedical Research Ethics Committee (011-08-17), the Zambia National Health Research Authority, and the London School of Hygiene and Tropical Medicine Ethics Committee (14506). Permission to conduct the study was given by Livingstone District Health Office and the Livingstone Central Hospital superintendent. Participants provided written informed consent.

Statistical Methods

All participants with FGS ($n = 30$) and all participants with probable FGS ($n = 25$) were selected for characterization of the cervicovaginal microbiota and STI by qPCR on cervical swabs. Three FGS-negative participants were selected for every FGS and probable FGS participant using a random number generator. The FGS-negative participants were frequency-matched by age to participants with FGS (age groups 18–19, 20–21, 22–23, 24–25, 26–27, 28–29, 30–31).

Participant characteristics were summarized by median and interquartile range (IQR) for continuous variables and by frequency and percentage for categorical variables. Differences in characteristics between FGS categories were evaluated using the Fisher exact or chi-square test. For cervicovaginal microbiota and STI species with at least 20% of sample results detectable by qPCR (ie, $\geq 20\%$ prevalence), P values for comparison of presence, median (IQR), and \log_{10} concentration mean between FGS and FGS-negative groups were calculated using the chi-square test, rank-sum test, and t test, respectively. For species with $< 20\%$ prevalence, species presence was compared between FGS

groups using the Fisher exact test. To enable investigation of potential confounding, concentrations of cervicovaginal microbiota with $\geq 20\%$ prevalence were log-transformed to normalize their distribution, and linear regression was used to evaluate the association between FGS and mean log-concentration of each organism (ge/mL) in univariable and multivariable analysis. We developed a causal framework (Supplementary Figure 1) to inform our minimal adjustment set and adjusted for age, education, and community of residence. For all species, logistic regression was used to calculate the crude and adjusted odds ratio (OR) for presence vs absence by FGS group; due to the relatively low number of participants with detectable concentrations, logistic regression analyses only adjusted for age. Given the exploratory nature of this work, we did not correct for multiple comparisons.

Our primary analysis focused on the detection of *Schistosoma* DNA in the genital tract (FGS vs FGS negative). A secondary analysis compared the FGS and probable FGS groups with the FGS-negative participants. To evaluate the possible association between FGS burden and changes in presence and concentration of cervicovaginal key species, 2 ad hoc analyses were performed: (1) participants with ≥ 2 genital samples with detectable *Schistosoma* DNA were compared with the FGS-negative group, (2) participants with a moderate/high genital *Schistosoma* DNA concentration (Ct < 35 in at least 1 of 3 samples) were compared with the FGS-negative group. Data were analyzed using STATA 15.1 (Stata Corporation, College Station, TX, USA).

RESULTS

A total of 603 eligible women were enrolled in the BILHIV study, and 213 (35.3%) were included in the present study (Supplementary Figure 1). Of those included, 14.1% (30/213) had FGS, defined by a positive genital *Schistosoma* qPCR from any of the following sites: 9.4% (20/213) cervical swab, 7.0% (15/213) vaginal swab, and 6.6% (14/211) CVL. In participants with FGS, portable colposcopy revealed sandy patches in 20% (6/30), abnormal blood vessels in 20% (6/30), and no clinical signs of FGS in 50% (15/30). Two women with FGS did not present to the clinic for portable colposcopy, and 1 had uninterpretable images. Of participants with FGS, 53.3% (16/30) had *Schistosoma* qPCR detected from multiple sites, and 53.3% (16/30) had moderate/high genital *Schistosoma* DNA loads; these groups overlapped by 12 participants. Twenty-five women had probable FGS, and 74.2% (158/262) of the women who were negative on all diagnostic tests were randomly selected for inclusion in this study. Consistent with the FGS definitions, portable colposcopy in 100.0% (25/25) of the participants with probable FGS showed FGS-associated cervicovaginal manifestations. In the majority of participants with probable FGS, portable colposcopy showed sandy patches (76.0%, 19/25), with 24% (6/25) of participants having abnormal blood vessels.

Baseline Characteristics

The majority of the participants were married/cohabitating, had received some secondary education, and were using hormonal contraception (Table 1). At the conclusion of HPTN 071 (PopART), HIV-1 prevalence was 17.4% (37/213) among the women included in this study, and one-third had at least 1 STI (Table 1). Urinary schistosome infection, defined as either a positive urine microscopy (11.7%, 25/213) or positive CAA (20.7%, 44/213), was reported in 21.6% (46/213) of participants. There were differences between the 3 categories of FGS status for age ($P = .002$), marital status ($P = .04$), education ($P = .06$), and employment ($P = .04$), with participants in the probable FGS group being more likely to be older, employed, and married than FGS and FGS-negative participants. There was strong evidence of a difference in community of residence by FGS status ($P < .001$), with participants with FGS and probable FGS more likely to live in Community A than participants in the FGS-negative group (Table 1). Other characteristics were similar by FGS status.

Primary Comparison: FGS vs FGS Negative

Concentrations of evaluated species are shown in Figure 1. Compared with FGS-negative women, there was no evidence of a difference in the presence or concentration of cervicovaginal *L. crispatus*, *L. iners*, *A. vaginae*, *G. vaginalis*, or *C. albicans* in participants with FGS (Table 2). A higher proportion of participants with FGS had *T. vaginalis* present, although there was not strong evidence of an association (OR, 2.11; 95% CI, 0.92–4.86; $P = .08$) (Supplementary Table 1). This result was similar after adjusting for age (Supplementary Table 1). Otherwise, compared with FGS-negative women, the presence and concentration of other STIs were similar in women with FGS (Table 2).

Combining the FGS and probable FGS groups, participants with FGS/probable FGS similarly had a higher prevalence of *T. vaginalis* compared with FGS-negative participants ($P = .05$) (Supplementary Table 2). Otherwise, compared with FGS-negative women, the presence and concentrations of cervicovaginal microbiota and the presence of STI were similar in women with FGS/probable FGS compared with FGS-negative participants (Supplementary Table 2).

Ad Hoc Analysis—Schistosoma DNA Concentration

In participants ($n = 16$) with FGS and a moderate/high genital *Schistosoma* DNA concentration (*Schistosoma* qPCR Ct < 35), the presence of *T. vaginalis* was higher than among the FGS-negative participants ($P = .01$) (Table 3). Women with FGS and a moderate/high *Schistosoma* DNA concentration (Ct < 35) had a higher mean concentration of *G. vaginalis* compared with FGS-negative women ($P = .03$) (Table 3).

Ad Hoc Analysis—Clinical Disease Burden

In an ad hoc analysis, participants ($n = 16$) with a higher FGS burden, defined as ≥ 2 *Schistosoma* qPCR-positive genital

specimens, had higher prevalence of *T. vaginalis* compared with FGS-negative participants ($P = .01$) (Table 4). There was also evidence of a difference in the median concentration and the mean log-concentration of *L. iners* (both $P = .03$) compared with FGS-negative women, with lower levels among the higher-FGS burden group (Table 4).

DISCUSSION

In this study, we describe the association of FGS with the cervicovaginal microbiota, including lactobacilli, *Candida* spp., markers of a “nonoptimal” cervicovaginal environment, and STI. We did not find evidence that the presence or concentration of key cervicovaginal species was associated with FGS. While FGS was not associated with *C. trachomatis*, *M. genitalium*, or *N. gonorrhoeae*, there was weak evidence of an association of presence of *T. vaginalis* with FGS, which remained after adjusting for age. This association was also present when participants with FGS and probable FGS were combined and was strengthened in the ad hoc analyses of participants with higher-burden FGS.

We performed 2 ad hoc analyses. High-intensity *S. haematobium* infection, in the absence of evaluation for FGS, has been associated with higher cervicovaginal alpha diversity [11]. Thus, first we investigated whether *Schistosoma* DNA concentrations might be associated with the cervicovaginal microbiota in 16 participants with a higher FGS burden, indicated by moderate/high genital *Schistosoma* DNA concentrations. In this ad hoc analysis, we found that the *G. vaginalis* log-concentration mean was higher in women with a higher FGS burden. Participants underwent CVL when they were not menstruating, and we have previously described that 66.2% (139/210) of women in this cohort had detectable CVL hemoglobin. Iron sources, like hemoglobin, are often required for bacterial growth [33]. *G. vaginalis* is well adapted to harvest iron from the environment [34], and higher concentrations of *G. vaginalis* coincide with menses [33]. Cervical tissue in women with FGS is more vascularized than non-egg-containing tissue, and thus the abnormal cervical vessels and contact bleeding seen in clinical FGS provide a plausible link to increased concentrations of *G. vaginalis* in high-burden FGS [35].

In a second ad hoc analysis, we examined participants with multiple qPCR-positive genital specimens as a potential proxy marker of high-burden FGS and found that reduced a *L. iners* concentration (median and log mean) was associated with high-burden FGS. The cervicovaginal microbiota was characterized with 16S rRNA sequencing in Tanzanian women with *S. haematobium* infection ($n = 16$). Although power was limited, women with high-intensity *S. haematobium* infection had reduced abundance of *L. iners* compared with women with low-intensity infection, albeit without evidence of a difference ($P = .39$) [11]. Further research is needed to evaluate the

Table 1. Baseline Characteristics of the Study Population (n = 213) by FGS Status

Participant Characteristics		FGS Negative, No. (%) (n = 158)	Probable FGS, No. (%) (n = 25)	FGS, No. (%) (n = 30)	P Value ^a
Age, median (IQR), y		23 (22–24)	27 (23–31)	22 (21–24)	.002 ^b
Marital status	Single	72 (45.6)	4 (16.0)	13 (43.3)	.04
	Married or cohabitating	81 (51.3)	20 (80.0)	17 (56.7)	
	Divorced or separated	5 (3.2)	1 (4.0)	0 (0.0)	
Education (highest level)	None or any primary school	35 (22.2)	12 (48.0)	10 (33.3)	.06
	Any secondary school	111 (70.3)	13 (52.0)	19 (63.3)	
	Trade, degree or higher	12 (7.6)	0 (0.0)	1 (3.3)	
District	Community A	66 (41.8)	20 (80.0)	22 (73.3)	<.01
	Community B	92 (58.2)	5 (20.0)	8 (26.7)	
Household members	1–3	51 (32.3)	4 (16.0)	12 (40.0)	.3
	4–5	61 (38.6)	13 (52.0)	8 (26.7)	
	6+	46 (29.1)	8 (32.0)	10 (33.3)	
Employment status	Not working	117 (74.1)	14 (56.0)	26 (86.7)	.04
	Working	41 (25.9)	11 (44.0)	4 (13.3)	
Sexual behavior characteristics					
Age at sexual debut, y	8–16	64 (40.5)	11 (44.0)	17 (56.7)	.4
	17–19	74 (46.8)	13 (52.0)	10 (33.3)	
	20–24	20 (12.7)	1 (4.0)	3 (10.0)	
Lifetime sexual partners	1	57 (36.1)	8 (32.0)	5 (16.7)	.3
	2	36 (22.8)	6 (24.0)	9 (30.0)	
	3	23 (14.6)	6 (24.0)	8 (26.7)	
	4+	42 (26.6)	5 (20.0)	8 (26.7)	
Currently sexually active ^{c,d}	No	26 (16.6)	2 (8.0)	3 (10.0)	.5
	Yes	131 (83.4)	23 (92.0)	27 (90.0)	
Condom use with last sex ^e	No	120 (76.9)	16 (66.7)	22 (73.3)	.6 ^f
	Yes	36 (23.1)	8 (33.3)	8 (26.7)	
HIV-1 status	Not detected	132 (83.5)	20 (80.0)	24 (80.0)	.8
	Detected	26 (16.5)	5 (20.0)	6 (20.0)	
Any STI ^g	Not detected	106 (67.1)	14 (56.0)	18 (60.0)	.5
	Detected	52 (32.9)	11 (44.0)	12 (40.0)	
Contraceptive use					
Condoms	No	132 (83.5)	20 (80.0)	26 (86.7)	.8
	Yes	26 (16.5)	5 (20.0)	4 (13.3)	
OCP	No	148 (93.7)	22 (88.0)	29 (96.7)	.5
	Yes	10 (6.3)	3 (12.0)	1 (3.3)	
Injectable	No	82 (51.9)	13 (52.0)	15 (50.0)	1.0 ^f
	Yes	76 (48.1)	12 (48.0)	15 (50.0)	
Implant	No	144 (91.1)	23 (92.0)	28 (93.3)	1.0
	Yes	14 (8.9)	2 (8.0)	2 (6.7)	
Any hormonal contraception ^h	No	58 (36.7)	8 (32.0)	10 (35.7)	.9 ^f
	Yes	100 (63.3)	17 (68.0)	18 (64.3)	
Schistosomiasis-related factors					
Urine microscopy	Not detected	158 (100.0)	19 (76.0)	11 (36.7)	<.001 ⁱ
	Detected	0 (0.0)	6 (24.0)	19 (63.3)	
Urine CAA	Negative	158 (100.0)	0 (0.0)	11 (36.7)	<.001 ⁱ
	Positive	0 (0.0)	25 (100.0)	19 (63.3)	

Abbreviations: CAA, circulating anodic antigen; FGS, female genital schistosomiasis; IQR, interquartile range; OCP, oral contraceptive pill; STI, sexually transmitted infection.

^aFisher's exact *P* value unless otherwise indicated.

^bKruskal-Wallis *P* value.

^cAny sexual activity in the last 6 months.

^dParticipants who responded with "no answer" (n = 1) are not shown in the table.

^eParticipants who responded with "no answer" (n = 3) are not shown in the table.

^fChi-square *P* value.

^gAny STI defined as the presence of at least 1 of *N. gonorrhoeae*, *C. trachomatis*, *M. genitalium*, or *T. vaginalis*.

^hAny hormonal contraception is defined as use of injectable agents, implants, or oral contraceptive pills.

ⁱPart of the definition for FGS categories.




Figure 1. Stacked bar chart of concentrations of cervicovaginal microbiota and taxa causing sexually transmitted infection by female genital schistosomiasis status. Abbreviation: FGS, female genital schistosomiasis.

relationship between the presence and concentration of *L. iners* in FGS.

In both ad hoc analyses, we found that presence of *T. vaginalis* was higher among the participants with a higher FGS burden. Our finding supports results from a small South African study (n = 45) that reported an association between FGS (diagnosed by identification of sandy patches on colposcopy) and presence of *T. vaginalis* in young women (ages 15–23) [36]. Acquisition of *T. vaginalis* and *S. haematobium* may share common risk factors, like age and socioeconomic

status [18]. We have previously described higher FGS prevalence among younger age groups [18]. Epidemiologic data from Zambian adolescents, sex workers, and pregnant women (aged 13–45 years) describe a *T. vaginalis* prevalence between 24.6% and 33.2% [37], consistent with the prevalence in our population (24.9%). While *S. haematobium* is geographically restricted to Africa and the Middle East, it is associated with poverty [38] and is acquired through contact with cercariae-infested fresh water [1]. *T. vaginalis* is primarily sexually transmitted, is prevalent worldwide, with the highest prevalence in

Table 2. Presence and Concentration of Vaginal Lactobacilli, Other Key Microbiota, and Sexually Transmitted Infection, Overall and by FGS Status

Organism	No. (%) and Concentration ^a	All Participants (n = 213)	FGS Negative (n = 158)	FGS (n = 30)	P Value ^a
<i>L. crispatus</i>	Presence	69 (32.4)	49 (31.0)	13 (43.3)	.19
	Median (IQR)	8.7×10 ⁶ (2.6×10 ⁵ –4.2×10 ⁶)	1.1×10 ⁷ (2.6×10 ⁵ –7.0×10 ⁶)	7.5×10 ⁵ (1.9×10 ⁵ –2.7×10 ⁷)	.20
	Log-concentration mean	16.01	16.38	14.82	.19
<i>L. iners</i>	Presence	156 (73.2)	113 (71.5)	24 (80.0)	.34
	Median (IQR)	2.7×10 ⁸ (3.8×10 ⁷ –1.5×10 ⁹)	2.8×10 ⁸ (3.9×10 ⁷ –1.6×10 ⁹)	1.7×10 ⁸ (1.6×10 ⁷ –1.0×10 ⁹)	.39
	Log-concentration mean	18.93	18.98	18.36	.33
<i>G. vaginalis</i>	Presence	156 (73.2)	115 (72.8)	23 (76.7)	.66
	Median (IQR)	7.7×10 ⁶ (8.3×10 ⁵ –5.2×10 ⁷)	8.1×10 ⁶ (7.7×10 ⁵ –4.8×10 ⁷)	3.8×10 ⁶ (6.8×10 ⁵ –6.4×10 ⁷)	.93
	Log-concentration mean	15.85	15.74	15.91	.79
<i>A. vaginae</i>	Presence	152 (71.4)	112 (70.9)	19 (63.3)	.41
	Median (IQR)	5.8×10 ⁷ (8.2×10 ⁶ –2.1×10 ⁸)	5.8×10 ⁷ (8.7×10 ⁶ –2.0×10 ⁸)	3.0×10 ⁷ (1.06×10 ⁶ –2.8×10 ⁸)	.83
	Log-concentration mean	17.31	17.33	17.06	.66
<i>T. vaginalis</i>	Presence	53 (24.9)	34 (21.5)	11 (36.7)	.08
	Median (IQR)	4.2×10 ⁴ (173.0–2.3×10 ⁶)	1.7×10 ⁵ (56.9–6.3×10 ⁶)	1.0×10 ⁴ (5030.0–4.5×10 ⁵)	.53
	Log-concentration mean	10.39	10.88	10.19	.69
Vaginal microbiota with prevalence <20%					
<i>N. gonorrhoeae</i>	Presence	13 (6.1)	12 (7.6)	0.0 (0)	.22
<i>C. trachomatis</i>	Presence	17 (8.0)	13 (8.2)	3 (10.0)	.72
<i>M. genitalium</i>	Presence	8 (3.8)	7 (4.4)	0 (0.0)	.60
<i>C. albicans</i>	Presence	12 (5.63)	8 (5.1)	2 (6.7)	.66

Abbreviations: FGS, female genital schistosomiasis; IQR, interquartile range.

^aConcentrations are expressed in genome equivalents/mL.

^bFor species with >20% prevalence, P values for presence, median (IQR), and log-concentration mean are from the chi-square test, rank-sum test, and t test, respectively. For species with <20% prevalence, P values for presence are from the Fisher exact test.

Table 3. Ad Hoc Analysis of the Presence and Concentration of Vaginal *Lactobacilli*, Other Cervicovaginal Microbiota, and Sexually Transmitted Infection in Participants With a Moderate to High Concentration of *Schistosoma* DNA (Ct < 35) Compared With FGS-Negative Participants

Organism	No. (%) and Concentration ^a	FGS Negative (n = 158)	FGS With Ct < 35 (n = 16)	P Value ^b
<i>L. crispatus</i>	Presence	49 (31.0)	6 (37.5)	.60
	Median (IQR)	1.1×10 ⁷ (2.6×10 ⁵ –7.0×10 ⁶)	1.2×10 ⁶ (3.0×10 ⁵ –2.0×10 ⁷)	.28
	Log-concentration mean	16.38	14.27	.19
<i>L. iners</i>	Presence	113 (71.5)	11 (68.75)	.82
	Median (IQR)	2.8×10 ⁸ (3.9×10 ⁷ –1.6×10 ⁹)	7.5×10 ⁷ (3.6×10 ⁶ –7.5×10 ⁸)	.30
	Log-concentration mean	18.98	17.86	.21
<i>G. vaginalis</i>	Presence	115 (72.8)	13 (81.3)	.46
	Median (IQR)	8.1×10 ⁶ (7.7×10 ⁵ –4.8×10 ⁷)	1.5×10 ⁷ (3.8×10 ⁶ –3.1×10 ⁸)	.11
	Log-concentration mean	15.74	17.45	.03
<i>A. vaginae</i>	Presence	112 (70.9)	13 (81.3)	.38
	Median (IQR)	5.8×10 ⁷ (8.7×10 ⁶ –2.0×10 ⁸)	1.6×10 ⁸ (1.6×10 ⁷ –2.8×10 ⁸)	.70
	Log-concentration mean	17.33	17.38	.95
<i>T. vaginalis</i>	Presence	34 (21.5)	8 (50.0)	.01
	Median (IQR)	1.7×10 ⁵ (56.9–6.3×10 ⁶)	8085.0 (5210.0–1.0×10 ⁶)	.56
	Log-concentration mean	10.88	10.09	.70
Vaginal microbiota with prevalence <20%				
<i>N. gonorrhoeae</i>	Presence	12 (7.6)	0 (0.0)	.61
<i>C. trachomatis</i>	Presence	13 (8.2)	2 (12.5)	.63
<i>M. genitalium</i>	Presence	7 (4.4)	0 (0.0)	1.0
<i>C. albicans</i>	Presence	8 (5.1)	2 (12.5)	.23

Abbreviations: Ct, cycle threshold; FGS, female genital schistosomiasis; IQR, interquartile range.

^aConcentrations are expressed in genome equivalents/mL.

^bFor species with >20% prevalence, P values for presence, median (IQR), and log-concentration mean are from the chi-square test, rank-sum test, and t test, respectively. For species with <20% prevalence, P values for presence are from the Fisher exact test.

Table 4. Ad Hoc Analysis of the Presence and Concentration of Vaginal *Lactobacilli*, Cervicovaginal Microbiota, and Sexually Transmitted Infection in Participants With *Schistosoma* DNA Detected in ≥2 Genital Specimens Compared With FGS-Negative Participants

Organism	No. (%) and Concentration ^a	FGS Negative (n = 158)	≥2 PCR FGS (n = 16)	P Value ^b
<i>L. crispatus</i>	Presence	49 (31.0)	7 (43.8)	.30
	Median (IQR)	1.1×10 ⁷ (2.6×10 ⁵ –7.0×10 ⁶)	1.9×10 ⁶ (7.7×10 ⁴ –2.7×10 ⁷)	.32
	Log-concentration mean	16.38	14.90	.34
<i>L. iners</i>	Presence	113 (71.5)	10 (62.5)	.45
	Median (IQR)	2.8×10 ⁸ (3.9×10 ⁷ –1.6×10 ⁹)	5.1×10 ⁷ (9.0×10 ⁵ –1.1×10 ⁸)	.03
	Log-concentration mean	18.98	17.00	.03
<i>G. vaginalis</i>	Presence	115 (72.8)	14 (87.5)	.20
	Median (IQR)	8.1×10 ⁶ (7.7×10 ⁵ –4.8×10 ⁷)	1.6×10 ⁷ (1.1×10 ⁶ –3.6×10 ⁸)	.29
	Log-concentration mean	15.74	16.85	.16
<i>A. vaginae</i>	Presence	112 (70.9)	11 (68.8)	.86
	Median (IQR)	5.8×10 ⁷ (8.7×10 ⁶ –2.0×10 ⁸)	1.6×10 ⁸ (2.7×10 ⁷ –3.2×10 ⁸)	.33
	Log-concentration mean	17.33	18.04	.36
<i>T. vaginalis</i>	Presence	34 (21.5)	8 (50.0)	.01
	Median (IQR)	1.7×10 ⁵ (56.9–6.3×10 ⁶)	5680.0 (2967.5–1.0×10 ⁶)	.50
	Log-concentration mean	10.88	9.71	.56
Vaginal microbiota with prevalence <20%				
<i>N. gonorrhoeae</i>	Presence	12 (7.6)	0 (0.0)	.61
<i>C. trachomatis</i>	Presence	13 (8.2)	2 (12.5)	.63
<i>M. genitalium</i>	Presence	7 (4.4)	0 (0.0)	1.0
<i>C. albicans</i>	Presence	8 (5.1)	1 (6.3)	.59

Abbreviations: FGS, female genital schistosomiasis; IQR, interquartile range; PCR, polymerase chain reaction.

^aConcentrations are expressed in genome equivalents/mL.

^bFor species with >20% prevalence, P values for presence, median (IQR), and log-concentration mean are from the chi-square test, rank-sum test and t test, respectively. For species with <20% prevalence, P values for presence are from the Fisher exact test.

women from low-income countries [39]. Both *S. haematobium* [40] and *T. vaginalis* infection [41, 42] have been reported as risk factors for HIV-1 acquisition, although these associations are not universally reported [43]. Additionally, *T. vaginalis* may influence HIV-1 transmission. Studies in women not uniformly receiving antiretroviral therapy (ART) with *T. vaginalis* show a decline in genital HIV-1 shedding after metronidazole therapy [44–46]. However, this decline in cervical HIV-1 shedding was not seen in a study of Kenyan women receiving ART [47]. The highest global *T. vaginalis* incidence rates have been reported in Africa [39]. Overlapping synergies between *T. vaginalis* and *S. haematobium* cervicovaginal pathogenesis may begin with, but are not limited to, disruption of the cervicovaginal epithelium [48]. Both parasites have been associated with characteristic cervicovaginal manifestations [17, 49], the “strawberry cervix” in *T. vaginalis* [49] and sandy patches, rubbery papules, and abnormal blood vessels in FGS [17]. Breaches in the cervicovaginal epithelium in *T. vaginalis* infection potentially represent one mechanism of HIV-1 vulnerability in what is likely a multifactorial cascade, including modulating the cervicovaginal immune environment [50–52], disabling innate immunity [51], and disrupting the local microbiota [52]. To disentangle the association between *S. haematobium*, *T. vaginalis*, and HIV-1, future studies using sensitive molecular methods for both parasites are needed. Additionally, macrophage polarization can be influenced by the local immune environment, schistosomes, and *T. vaginalis* [53, 54]. We have previously shown that, compared with FGS-negative women, high-burden FGS is associated with higher concentrations of cervicovaginal Th2 cytokines [32]. A mouse model of urogenital *S. haematobium* infection suggested that the Th2 immune environment may be associated with delayed pathogen clearance [55]. Thus, further research is needed regarding the interaction between the immune environment and macrophage phenotypes in FGS and their role in potentially influencing *T. vaginalis* persistence.

Our study is the first to evaluate the cervicovaginal microbiota and STI in FGS defined by qPCR. This is particularly relevant in a population of sexually active, nonpregnant women in the context of high HIV-1 prevalence. However, there are also some relevant limitations. FGS does not have an accepted diagnostic reference standard, and the sensitivity of cervicovaginal PCR for the detection of *Schistosoma* DNA is imperfect. Thus, studies using *Schistosoma* DNA to detect FGS may be subject to misclassification bias. To maximize sensitivity, we used the results of 3 genital specimens. This decision was supported by previous work showing a sensitivity of 80.0% (61.4%–92.3%) with a specificity of 100.0% (99.3%–100.0%) when vaginal and cervical swab PCR were compared with any positive genital PCR (CVL, cervical swab, or vaginal swab) as the reference standard [18]. The study was conducted in an urban location with relatively low *S. haematobium* prevalence; thus the numbers of FGS

cases in the primary and ad hoc analyses were small, limiting precision in the effect sizes. This analysis also included multiple statistical comparisons; thus, we focused on the species that showed a consistent pattern of association across primary and ad hoc analyses, rather than overinterpreting significance testing for any 1 species in isolation. Evidence for these associations in the ad hoc analyses should be viewed as hypothesis-generating. Additionally, the cross-sectional study design limited our ability to assess FGS duration and the long-term impact on the prevalence and concentrations of key species or STIs. There were a number of behavioral and biological factors that were not measured in our study including tobacco use [56], viral STIs (human papillomavirus and herpes simplex virus–2) [56], and intravaginal cleansing practices [57]. As these factors may be associated with the cervicovaginal microbiota, we cannot exclude residual or unmeasured confounding. Genital swabs were self-collected by participants, raising the potential for false-negative genital swabs. In future work, β -globin PCR could be implemented as a positive control to confirm the presence of human DNA [18]. Lastly, we defined FGS by *Schistosoma* DNA detection on qPCR; however, we cannot exclude that cervicovaginal qPCR detected *S. haematobium* eggs from a sexual partner’s semen [18].

In conclusion, we report weak evidence of an association between presence of *T. vaginalis* and FGS, with a consistently stronger association in women with a higher burden FGS infection. Additional research is needed to understand the interactions between *S. haematobium* and *T. vaginalis*, especially regarding HIV-1 vulnerability.

Acknowledgments

We wish to acknowledge the participants for devoting their time and trust to this study. We would like to recognize the work performed by the BILHIV project supervisor Namakau Chola and the BILHIV community workers Ethel Mwansa, Mwiingana Lukonga, Ruth Mwanza, Mervis Kantukaleza, and Judith Lungu. Our work in Livingstone would not have been possible without support from Clement Mwakamui (Zambart). We gratefully acknowledge Eric A. T. Brienen (LUMC) for performing the genital *Schistosoma* PCR analysis and Claudia J. de Dood (LUMC) and Pytsje T. Hoekstra (LUMC) for performing the CAA analysis.

Financial support. Dr. A. Bustinduy received funding from the Wellcome Trust (Award 205954/Z/17/Z). Dr. P. Cools received funding from the Research Foundation–Flanders (BOFSTG2019010101) and was financially supported by a Bill and Melinda Gates Foundation Project (OPP1120972). Dr. E. Webb and Professor R. Hayes received funding from MRC Grant Reference MR/K012126/1, and Dr. S. C. Francis received salary from MRC Grant Reference MR/N023692/1. These awards are jointly funded by the UK Medical Research Council (MRC) and the UK Department for International Development (DFID) under the MRC/DFID Concordat agreement and are also part of the EDCTP2 program supported by the European Union. HPTN 071 (PopART) was supported by the National Institute of Allergy and Infectious Diseases (NIAID) under Cooperative Agreements UM1-AI068619, UM1-AI068617, and UM1-AI068613, with funding from the US President’s Emergency Plan for AIDS Relief (PEPFAR); the International Initiative for Impact Evaluation with support from the Bill and Melinda Gates Foundation; the NIAID, the National Institute on Drug Abuse, and the National Institute of Mental Health, all part of the National Institutes of Health. Professor Eyrun

Kjetland was supported by South-Eastern Regional Health Authority, Norway, project #2016055. Bruno Levecke received funding from the Fund for Scientific Research–Flanders (Grant No. 1285316N).

Potential conflicts of interest. The authors report no conflicts of interest. All authors have submitted the ICMJE Form for Disclosure of Potential Conflicts of Interest. Conflicts that the editors consider relevant to the content of the manuscript have been disclosed.

Author contributions. Amy S. Sturt—conceptualization, data curation, formal analysis, BILHIV project administration, visualization, original manuscript preparation, manuscript editing and revision. Emily L. Webb—conceptualization, data curation, formal analysis, supervision, visualization, original manuscript preparation, manuscript editing and revision. Lisa Himschoot—investigation, manuscript editing and revision. Comfort R. Phiri—BILHIV project administration, writing—review and editing. Maina Mudenda—investigation, writing—review and editing. Joyce Mapani—investigation, writing—review and editing. Eyrun F. Kjetland—investigation, writing—review and editing. Tobias Mweene—investigation, writing—review and editing. Bruno Levecke—investigation, resources, writing—review and editing. Govert J. van Dam—investigation, writing—review and editing. Paul L. A. M. Corstjens—investigation, writing—review and editing. Helen Ayles—resources, writing—review and editing. Richard J. Hayes—supervision, resources, writing—review and editing. Lisette van Lieshout—investigation, writing—review and editing. Isaiah Hansingo—resources, supervision, writing—review and editing. Suzanna C. Francis—conceptualization, supervision, original manuscript preparation, manuscript editing and revision. Piet Cools—conceptualization, resources, investigation, original manuscript preparation, manuscript editing and revision. Amaya L. Bustinduy—conceptualization, funding acquisition, supervision, original manuscript preparation, manuscript editing and revision.

References

- Sturt AS, Webb EL, Francis SC, et al. Beyond the barrier: female genital schistosomiasis as a potential risk factor for HIV-1 acquisition. *Acta Trop* **2020**; 209:105524.
- Kjetland EF, Ndhlovu PD, Gomo E, et al. Association between genital schistosomiasis and HIV in rural Zimbabwean women. *AIDS* **2006**; 20:593–600.
- Boris S, Barbés C. Role played by lactobacilli in controlling the population of vaginal pathogens. *Microbes Infect* **2000**; 2:543–6.
- McKinnon LR, Achilles SL, Bradshaw CS, et al. The evolving facets of bacterial vaginosis: implications for HIV transmission. *AIDS Res Hum Retroviruses* **2019**; 35:219–28.
- van de Wijgert J, Jaspers V. The global health impact of vaginal dysbiosis. *Res Microbiol* **2017**; 168:859–64.
- Torrone EA, Morrison CS, Chen PL, et al; STIMA Working Group. Prevalence of sexually transmitted infections and bacterial vaginosis among women in Sub-Saharan Africa: an individual participant data meta-analysis of 18 HIV prevention studies. *PLoS Med* **2018**; 15:e1002511.
- Hillier SL, Nugent RP, Eschenbach DA, et al. Association between bacterial vaginosis and preterm delivery of a low-birth-weight infant. The Vaginal Infections and Prematurity Study Group. *N Engl J Med* **1995**; 333:1737–42.
- Liu L, Oza S, Hogan D, et al. Global, regional, and national causes of child mortality in 2000–13, with projections to inform post-2015 priorities: an updated systematic analysis. *Lancet* **2015**; 385:430–40.
- Atashili J, Poole C, Ndumbe PM, et al. Bacterial vaginosis and HIV acquisition: a meta-analysis of published studies. *AIDS* **2008**; 22:1493–501.
- Jaspers V, Crucitti T, Menten J, et al; Vaginal Biomarkers Study Group. Prevalence and correlates of bacterial vaginosis in different sub-populations of women in Sub-Saharan Africa: a cross-sectional study. *PLoS One* **2014**; 9:e109670.
- Bullington BW, Lee MH, Mlingi J, et al. Cervicovaginal bacterial communities in reproductive-aged Tanzanian women with *Schistosoma mansoni*, *Schistosoma haematobium*, or without schistosome infection. *ISME J* **2021**; 15:1539–50.
- Gadoth A, Mvumbi G, Hoff NA, et al. Urogenital schistosomiasis and sexually transmitted coinfections among pregnant women in a schistosome-endemic region of the Democratic Republic of Congo. *Am J Trop Med Hyg* **2019**; 101:828–36.
- Leutscher PD, Ramarokoto CE, Hoffmann S, et al. Coexistence of urogenital schistosomiasis and sexually transmitted infection in women and men living in an area where *Schistosoma haematobium* is endemic. *Clin Infect Dis* **2008**; 47:775–82.
- Yirenya-Tawiah D, Annang TN, Apea-Kubi KA, et al. *Chlamydia trachomatis* and *Neisseria gonorrhoeae* prevalence among women of reproductive age living in urogenital schistosomiasis endemic area in Ghana. *BMC Res Notes* **2014**; 7:85–91.
- Poggensee G, Kiwelu I, Saria M, et al. Schistosomiasis of the lower reproductive tract without egg excretion in urine. *Am J Trop Med Hyg* **1998**; 59:782–3.
- Renaud G, Devidas A, Develoux M, et al. Prevalence of vaginal schistosomiasis caused by *Schistosoma haematobium* in an endemic village in Niger. *Trans R Soc Trop Med Hyg* **1989**; 83:797.
- Kjetland EF, Ndhlovu PD, Mduluzi T, et al. Simple clinical manifestations of genital *Schistosoma haematobium* infection in rural Zimbabwean women. *Am J Trop Med Hyg* **2005**; 72:311–9.
- Sturt AS, Webb EL, Phiri CR, et al. Genital self-sampling compared with cervicovaginal lavage for the diagnosis of female genital schistosomiasis in Zambian women: the BILHIV study. *PLoS Negl Trop Dis* **2020**; 14:e0008337.
- Downs JA, Mguta C, Kaatano GM, et al. Urogenital schistosomiasis in women of reproductive age in Tanzania's Lake Victoria region. *Am J Trop Med Hyg* **2011**; 84:364–9.
- Pillay P, van Lieshout L, Taylor M, et al. Cervical cytology as a diagnostic tool for female genital schistosomiasis: correlation to cervical atypia and *Schistosoma* polymerase chain reaction. *Cytojournal* **2016**; 13:1–10.
- Kjetland EF, Hove RJ, Gomo E, et al. Schistosomiasis PCR in vaginal lavage as an indicator of genital *Schistosoma haematobium* infection in rural Zimbabwean women. *Am J Trop Med Hyg* **2009**; 81:1050–5.
- Galappaththi-Arachchige HN, Holmen S, Koukounari A, et al. Evaluating diagnostic indicators of urogenital *Schistosoma haematobium* infection in young women: a cross sectional study in rural South Africa. *PLoS One* **2018**; 13:e0191459.
- Randrianasolo BS, Jourdan PM, Ravoniarimbinina P, et al. Gynecological manifestations, histopathological findings, and *Schistosoma*-specific polymerase chain reaction results among women with *Schistosoma haematobium* infection: a cross-sectional study in Madagascar. *J Infect Dis* **2015**; 212:275–84.
- Hayes RJ, Donnell D, Floyd S, et al; HPTN 071 (PopART) Study Team. Effect of universal testing and treatment on HIV incidence - HPTN 071 (PopART). *N Engl J Med* **2019**; 381:207–18.
- World Health Organization. *Female Genital Schistosomiasis: A Pocket Atlas for Clinical Health-Care Professionals*. World Health Organization; **2015**. Available at: <http://www.who.int/iris/handle/10665/180863>. Accessed April 30, 2021.
- Zambian Ministry of Health. *Guidelines for the Etiological and Clinical Management of Sexually Transmitted Infections in Zambia*. Zambian Ministry of Health; **2017**.
- Corstjens PL, De Dood CJ, Kornelis D, et al. Tools for diagnosis, monitoring and screening of *Schistosoma* infections utilizing lateral-flow based assays and upconverting phosphor labels. *Parasitology* **2014**; 141:1841–55.
- Corstjens PLAM, de Dood CJ, Knopp S, et al. Circulating anodic antigen (CAA): a highly sensitive diagnostic biomarker to detect active *Schistosoma* infections—improvement and use during SCORE. *Am J Trop Med Hyg* **2020**; 103:50–7.
- Obeng BB, Aryeetey YA, de Dood CJ, et al. Application of a circulating-cathodic-antigen (CCA) strip test and real-time PCR, in comparison with microscopy, for the detection of *Schistosoma haematobium* in urine samples from Ghana. *Ann Trop Med Parasitol* **2008**; 102:625–33.
- Pillay P, Taylor M, Zulu SG, et al. Real-time polymerase chain reaction for detection of *Schistosoma* DNA in small-volume urine samples reflects focal distribution of urogenital schistosomiasis in primary school girls in KwaZulu Natal, South Africa. *Am J Trop Med Hyg* **2014**; 90:546–52.
- Jaspers V, van de Wijgert J, Cools P, et al; Vaginal Biomarkers Study Group. The significance of *Lactobacillus crispatus* and *L. vaginalis* for vaginal health and the negative effect of recent sex: a cross-sectional descriptive study across groups of African women. *BMC Infect Dis* **2015**; 15:1–14.
- Sturt AS, Webb EL, Patterson C, et al. Cervicovaginal immune activation in Zambian women with female genital schistosomiasis. *Front Immunol* **2021**; 12:1–15.
- Srinivasan S, Liu C, Mitchell CM, et al. Temporal variability of human vaginal bacteria and relationship with bacterial vaginosis. *PLoS One* **2010**; 5:e10197.
- Jarosik GP, Land CB, Duhon P, et al. Acquisition of iron by *Gardnerella vaginalis*. *Infect Immun* **1998**; 66:5041–7.
- Jourdan PM, Roald B, Poggensee G, et al. Increased vascularity in cervicovaginal mucosa with *Schistosoma haematobium* infection. *PLoS Negl Trop Dis* **2011**; 5:e1170.
- Kleppa E, Ramsuran V, Zulu S, et al. Effect of female genital schistosomiasis and anti-schistosomal treatment on monocytes, CD4+ T-cells and CCR5 expression in the female genital tract. *PLoS One* **2014**; 9:e98593.
- Crucitti T, Jaspers V, Mulenga C, et al. *Trichomonas vaginalis* is highly prevalent in adolescent girls, pregnant women, and commercial sex workers in Ndola, Zambia. *Sex Transm Dis* **2010**; 37:223–7.
- Hotez PJ, Kamath A. Neglected tropical diseases in Sub-Saharan Africa: review of their prevalence, distribution, and disease burden. *PLoS Negl Trop Dis* **2009**; 3:e412.
- Rowley J, Vander Hoorn S, Korenromp E, et al. Chlamydia, gonorrhoea, trichomoniasis and syphilis: global prevalence and incidence estimates, 2016. *Bull World Health Organ* **2019**; 97:548–62P.

40. Wall KM, Kilembe W, Vwalika B, et al. Schistosomiasis is associated with incident HIV transmission and death in Zambia. *PLoS Negl Trop Dis* **2018**; 12:e0006902.
41. Mavedzenge SN, Pol BV, Cheng H, et al. Epidemiological synergy of *Trichomonas vaginalis* and HIV in Zimbabwean and South African women. *Sex Transm Dis* **2010**; 37:460–6.
42. Masha SC, Cools P, Sanders EJ, et al. *Trichomonas vaginalis* and HIV infection acquisition: a systematic review and meta-analysis. *Sex Transm Infect* **2019**; 95:36–42.
43. Bochner AF, Baeten JM, Secor WE, et al. Associations between schistosomiasis and HIV-1 acquisition risk in four prospective cohorts: a nested case-control analysis. *J Int AIDS Soc* **2020**; 23:e25534.
44. Wang CC, McClelland RS, Reilly M, et al. The effect of treatment of vaginal infections on shedding of human immunodeficiency virus type 1. *J Infect Dis* **2001**; 183:1017–22.
45. Kissinger P, Amedee A, Clark RA, et al. *Trichomonas vaginalis* treatment reduces vaginal HIV-1 shedding. *Sex Transm Dis* **2009**; 36:11–6.
46. Anderson BL, Firnhaber C, Liu T, et al. Effect of trichomoniasis therapy on genital HIV viral burden among African women. *Sex Transm Dis* **2012**; 39:638–42.
47. Masese LN, Graham SM, Gitau R, et al. A prospective study of vaginal trichomoniasis and HIV-1 shedding in women on antiretroviral therapy. *BMC Infect Dis* **2011**; 11:1–6.
48. Guenther PC, Secor WE, Dezzutti CS. *Trichomonas vaginalis*-induced epithelial monolayer disruption and human immunodeficiency virus type 1 (HIV-1) replication: implications for the sexual transmission of HIV-1. *Infect Immun* **2005**; 73:4155–60.
49. Kissinger P. *Trichomonas vaginalis*: a review of epidemiologic, clinical and treatment issues. *BMC Infect Dis* **2015**; 15:1–8.
50. Thurman AR, Doncel GF. Innate immunity and inflammatory response to *Trichomonas vaginalis* and bacterial vaginosis: relationship to HIV acquisition. *Am J Reprod Immunol* **2011**; 65:89–98.
51. Mercer F, Diala FG, Chen YP, et al. Leukocyte lysis and cytokine induction by the human sexually transmitted parasite *Trichomonas vaginalis*. *PLoS Negl Trop Dis* **2016**; 10:e0004913.
52. Fichorova RN, Buck OR, Yamamoto HS, et al. The villain team-up or how *Trichomonas vaginalis* and bacterial vaginosis alter innate immunity in concert. *Sex Transm Infect* **2013**; 89:460–6.
53. Cortes-Selva D, Fairfax K. Schistosome and intestinal helminth modulation of macrophage immunometabolism. *Immunology* **2021**; 162:123–34.
54. Han IH, Song HO, Ryu JS. IL-6 produced by prostate epithelial cells stimulated with *Trichomonas vaginalis* promotes proliferation of prostate cancer cells by inducing M2 polarization of THP-1-derived macrophages. *PLoS Negl Trop Dis* **2020**; 14:e0008126.
55. Hsieh YJ, Fu CL, Hsieh MH. Helminth-induced interleukin-4 abrogates invariant natural killer T cell activation-associated clearance of bacterial infection. *Infect Immun* **2014**; 82:2087–97.
56. Chernes TL, Hillier SL, Meyn LA, et al. A delicate balance: risk factors for acquisition of bacterial vaginosis include sexual activity, absence of hydrogen peroxide-producing lactobacilli, black race, and positive herpes simplex virus type 2 serology. *Sex Transm Dis* **2008**; 35:78–83.
57. Low N, Chersich MF, Schmidlin K, et al. Intravaginal practices, bacterial vaginosis, and HIV infection in women: individual participant data meta-analysis. *PLoS Med* **2011**; 8:e1000416.