



Universiteit
Leiden
The Netherlands

A multi-level assessment of the bidirectional relationship between aging and the circadian clock

Buijink, M.R.; Michel, S.

Citation

Buijink, M. R., & Michel, S. (2021). A multi-level assessment of the bidirectional relationship between aging and the circadian clock. *Journal Of Neurochemistry*, 157(1), 73-94. doi:10.1111/jnc.15286

Version: Publisher's Version

License: [Creative Commons CC BY-NC-ND 4.0 license](#)

Downloaded from: <https://hdl.handle.net/1887/3209338>

Note: To cite this publication please use the final published version (if applicable).

REVIEW

A multi-level assessment of the bidirectional relationship between aging and the circadian clock

M. Renate Buijink  | Stephan Michel 

Department of Cellular and Chemical Biology, Laboratory for Neurophysiology, Leiden University Medical Center, Leiden, the Netherlands

Correspondence

Stephan Michel, Department of Cellular and Chemical Biology, Laboratory for Neurophysiology, Leiden University Medical Center, Postzone S5-P, PO Box 9600, 2300 RC Leiden, The Netherlands.
Email: s.michel@lumc.nl

Funding information

Velux Stiftung, Grant/Award Number: 1029

This is a Review for the special issue "Circadian Rhythms in the Brain"

Abstract

The daily temporal order of physiological processes and behavior contribute to the wellbeing of many organisms including humans. The central circadian clock, which coordinates the timing within our body, is located in the suprachiasmatic nucleus (SCN) of the hypothalamus. Like in other parts of the brain, aging impairs the SCN function, which in turn promotes the development and progression of aging-related diseases. We here review the impact of aging on the different levels of the circadian clock machinery—from molecules to organs—with a focus on the role of the SCN. We find that the molecular clock is less effected by aging compared to other cellular components of the clock. Proper rhythmic regulation of intracellular signaling, ion channels and neuronal excitability of SCN neurons are greatly disturbed in aging. This suggests a disconnection between the molecular clock and the electrophysiology of these cells. The neuronal network of the SCN is able to compensate for some of these cellular deficits. However, it still results in a clear reduction in the amplitude of the SCN electrical rhythm, suggesting a weakening of the output timing signal. Consequently, other brain areas and organs not only show aging-related deficits in their own local clocks, but also receive a weaker systemic timing signal. The negative spiral completes with the weakening of positive feedback from the periphery to the SCN. Consequently, chronotherapeutic interventions should aim at strengthening overall synchrony in the circadian system using life-style and/or pharmacological approaches.

KEYWORDS

chronotherapy, neurodegeneration, neuronal network, peripheral clocks, senescence, suprachiasmatic nucleus

Abbreviations: AD, Alzheimer's disease; AVP, Arginine vasopressin; BDNF, brain-derived neurotrophic factor; Bmal1, Brain and Muscle ARNT-Like 1; cAMP, Cyclic adenosine monophosphate; Cry1, Cryptochrome 1; Cry2, Cryptochrome 2; LD, light-dark; mTOR, mammalian target of rapamycin; NAD⁺, nicotinamide adenine dinucleotide; NAD⁺/NADH, nicotinamide adenine dinucleotide, oxidized/reduced; NF-κB, Nuclear Factor kappa-light-chain-enhancer of activated B cells; Per1, Period 1; Per2, Period 2; SCN, suprachiasmatic nucleus; SIRT1, sirtuin 1 silent mating type information regulation 2 homolog 1; VIP, vasoactive intestinal protein; VIP-KO, VIP knock-out.

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2020 The Authors. *Journal of Neurochemistry* published by John Wiley & Sons Ltd on behalf of International Society for Neurochemistry.



1 | INTRODUCTION

One of the great achievements of modern medicine is the development of successful treatments of many diseases that meant a death sentence in earlier times. As a consequence, the average life span has increased steadily in the past century, creating new opportunities, but also many challenges. One of the biggest challenges is to abate age-related diseases and stay healthy, even productive, at an old age. Aging affects most physiological processes and parts of our body to some degree and we are currently unable to halt or reverse the aging process. Still, we can strive to achieve 'healthy aging' by changes in lifestyle or smart medication. One of the potential interventions to achieve healthier aging is to support the regulation of daily rhythms of processes in our body. For instance, strengthening the circadian clock function will alleviate clock-related sleep disorders that often occur during aging and worsen in neurodegenerative diseases. The negative interaction between circadian clock and age-related disease often creates a downward spiral in health. For instance, for Alzheimer's and other neurodegenerative diseases, sleep disorders seem to be already present at the preclinical state (Musiek et al., 2018) and worsen in the course of the disease (Leng et al., 2019). Importantly, the sleep disruptions also aggravate the home situation for the patient and his family, often leading to early institutionalisation and significant reduction in the patient's quality of life.

The suprachiasmatic nucleus (SCN), the central clock that organizes temporal order of many physiological processes including sleep/wake behavior is situated in the hypothalamus and—not surprisingly—is also affected by aging. Conversely, there is mounting evidence that the circadian clock in return is influencing the course of aging, the progression of age-related disease and general life expectancy (Fonseca Costa & Ripperger, 2015; Kondratova & Kondratov, 2012; Musiek & Holtzman, 2016). A functional central clock in the SCN is beneficial to the performance of other brain areas, like the hippocampus, and organs like heart and liver. Understanding the impact of aging on the central clock and how to restore its function can therefore be advantageous for developing interventions for many age-related symptoms and disorders. After a brief overview of the circadian system, this review will focus on the age-related deficits in the SCN and underline the differential effect aging has at various levels of the SCN organization; from molecules to networks. We will then discuss the effects this compromised central clock has on so-called peripheral clocks, but also consider the consequences of age-related loss of local peripheral clock function. Lastly, we will briefly recapitulate the evidence for the interaction between the circadian clock and neurodegenerative diseases and close with some considerations of potential chronotherapeutic interventions.

2 | THE ESTABLISHMENT OF TEMPORAL ORDER IN OUR BODY

While there is no evolutionary advantage to an increase in life span, selection pressure was at first very high on environmental stressors

with daily fluctuations, like ultraviolet radiation, extreme temperature difference and later the presence of predators. Consequently, molecular clock mechanisms were established in very early life forms, like cyanobacteria, and further developed into a specialized circadian system we now find in almost all organisms on this planet. In mammals, a dedicated central clock in the SCN seems to work as a conductor in an orchestra of clocks, controlling the temporal harmony in our body to ensure the optimal time of performance of all our physiological functions and behavior.

The endogenous circadian clocks run with a slightly different pace than the environmental cycle of 24 hr and therefore have to be synchronized or entrained by external zeitgebers like the daily cycles of light and darkness. The SCN receives information on the light-dark cycle from specialized retinal ganglion cells and distributes its 'corrected' information on the time of day (and day-length) to other brain areas and organs. Importantly, the SCN is not just relaying light-information, but integrates this into its neuronal network, together with feedback from the rest of the body, like sleep quality, physical activity, metabolic state and stress level. The resulting SCN timing signal therefore not only holds information on the time of day and season, but also of the state of many bodily functions. It is therefore inevitable that the effects of aging on brain and organ function will impact SCN function and vice versa, which often complicates the interpretation of the data on the aging circadian clock.

In addition to this clockshop of central and peripheral clocks, the circadian signal of the SCN is also shaped and influenced on a number of organizational levels, namely the molecular, cellular, membrane, and network level (Evans, 2016; Michel & Meijer, 2020).

As mentioned above, the basis for a circadian regulation of cellular function was established early on in evolution with a molecular mechanism creating an oscillation with a transcriptional-translational feedback loop (Buhr & Takahashi, 2013). The clock proteins BMAL1 and CLOCK enhance the transcription of *cry* and *per*, which will inhibit their own transcription with a sufficient delay to ensure the ~24 hr rhythms of the clock gene expression. This generates oscillations of proteins that either directly control other cellular functions, or control the transcription of other clock-controlled genes. An estimated 40%–50% of all protein encoding genes in mice and humans are regulated by the circadian clock in an organ specific way (Ruben et al., 2018; Zhang et al., 2014). For instance, in the SCN, rhythmic regulation of intracellular signaling pathways like cAMP/ Ca^{2+} seem to serve the stability and precision of clock function (O'Neill & Reddy, 2012), while in the liver circadian timing of metabolic pathways is optimizing hepatocyte activity to food availability and energy demand.

3 | THE AGING SCN

The SCN is a small brain area consisting of a relatively homogeneous network of neurons and glia, that is in appearance, its functioning and adaptability surprisingly diverse and remains the subject of a growing number of studies. Like all brain areas, the

SCN is affected by aging (Farajnia et al., 2014). However, because of its regulatory role in temporal cellular processes, a compromised SCN function influences the process of aging in other brain areas and organs. The importance of the SCN in aging is illustrated by the effect of transplantation of fetal SCN tissue in the third ventricle, alleviating aging deficits in circadian rhythms in rodents (Cai & Wise, 1996; Van Reeth et al., 1994) and increasing life expectancy (Hurd & Ralph, 1998). In addition, manipulation of the central clock by for example phase shifts, SCN ablation and genetic manipulation, results in health deterioration similar to that seen in aging, and often leads to a shortened lifespan (Davidson et al., 2006; Hurd & Ralph, 1998). It is important to note that, when using animal models for aging, the lifespan and aging dynamics of each species and strain has to be known and evaluated. In mice (C57bl6), life stages have been defined as matured adult (3–6 months), middle-aged (10–14 months) and old (18–24 months) (Flurkey et al., 2007). The specifics of age, animal model and study parameters for the relevant studies discussed by us can be found in table 1 (Table 1). Below we will discuss the effect of aging on different levels of the circadian system, from the molecular clock in individual SCN cells, via the SCN network, to the output signal of the SCN toward the periphery (Figure 1).

3.1 | Input to the SCN—Aging affects light transduction to the SCN

Environmental light information reaches the SCN through intrinsically photoreceptive retinal ganglion cells (ipRGCs) in the eye and the retinal hypothalamic tract (Hattar et al., 2002). Aging is accompanied by an attenuated light response in the SCN, quantified by the induction of immediate early genes, like c-Fos (Sutin et al., 1993). The decline in light sensitivity is likely a consequence of an overall loss of function of the light transduction pathway. First of all, aging is associated with a yellowing of the lens, which reduces the transmission of—mainly short wavelength—light to the retina (Kessel et al., 2010). Secondly, in the course of aging, the number of ipRGCs and their dendritic arborisation are reduced (Esquivia et al., 2017; Lax et al., 2019; Semo et al., 2003). Moreover, the loss of ipRGCs in aging is accompanied by a loss of projections to the SCN (Engelberth et al., 2017; Lupi et al., 2012). Despite the multi-level deterioration of light transduction to the SCN, there are several studies indicating that light is still capable of modulating circadian rhythms in humans (Benloucif et al., 2006; Najjar et al., 2014). Therefore, light therapy remains a suitable candidate for chronotherapeutic purposes in the elderly, although possibly with adjusted light wavelengths or intensities (Rubiño et al., 2020).

3.2 | The molecular clock of the SCN ages well

While the light-input from the eyes to the SCN is impaired in aging, it seems that the core molecular clock in SCN neurons remains largely functional. However, the results of studies on the effect of aging on the

molecular clock components are ambiguous and only rather consistent for the core clock genes Period 1 (Per1) and Brain and Muscle ARNT-Like 1 (Bmal1). While almost all studies report no change in Per1 expression rhythm or level, aging-related changes in expression of Bmal1 are consistently found (Per1: Asai et al., 2001; Bonaconsa et al., 2014; Davidson et al., 2008; Kolker et al., 2003; Weinert et al., 2001; Yamazaki et al., 2002; Bmal1: Bonaconsa et al., 2014; Chang & Guarente, 2013; Duncan et al., 2013; Kolker et al., 2003; Wyse & Coogan, 2010). There are only a few studies that have investigated the effect of aging on Cryptochrome 1/2 and Clock genes and most do not report any age-related effects for the SCN (Asai et al., 2001; Bonaconsa et al., 2014; Kolker et al., 2003; Weinert et al., 2001; Wyse & Coogan, 2010). By contrast, Per2 has been studied extensively with respect to aging. Under an equinoxial light regime (Light-Dark; LD 12:12), almost half of the studies did not find an effect of aging on Per2 expression (Asai et al., 2001; Buijink et al., 2020; Kolker et al., 2003; Leise et al., 2013; Polidarova et al., 2017), while others report a lower level (exclusively for mRNA; Bonaconsa et al., 2014; Chang & Guarente, 2013; Weinert et al., 2001), or a lower amplitude of Per2 expression (Nakamura et al., 2015; Nakamura et al., 2012). One study actually reported an increase in Per2 amplitude (Sellix et al., 2012). In a study from our laboratory we found that a moderate, and naturally occurring deviation from the LD 12:12 light regime, namely LD 16:8 and LD 8:16, resulted in a similar adaptation in Per2 expression rhythms in young and old mice, demonstrating that the aging molecular clock retains its plasticity (Buijink et al., 2020). However, two other recent studies show that more extreme lighting conditions—constant light and constant darkness—did affect Per2 rhythms in aging (Nakamura et al., 2015; Polidarova et al., 2017). This suggests that there are limits to the flexibility of the aging molecular clock. Nonetheless, the currently available literature indicates that the effect of aging on the molecular clock is relatively subtle, and it is unlikely to account for the severe deficits seen in other parts of the circadian system. Therefore, it is uncertain if strengthening the molecular clock, for example with pharmaceuticals, would benefit the elderly.

3.3 | Other clock components in the cytosol of aging SCN neurons are affected by aging

Core clock gene expression in transcriptional-translational feedback loops were long considered to be essential for generating a circadian rhythm in circadian pacemaker cells of the SCN. Despite studies revealing cytosolic oscillators like Ca²⁺ (Noguchi et al., 2017), cyclic adenosine monophosphate (cAMP; Doi et al., 2011; O'Neill et al., 2008) and nicotinamide adenine dinucleotide (NAD⁺/NADH; Huang et al., 2016; Ramsey et al., 2009) contributing to cellular clock function, the core clock genes are still the main players for initiating the circadian modulation of cellular metabolites and signaling pathways. For the SCN neuron and its network, this results in the circadian modulation of ion channels regulating membrane excitability (Harvey et al., 2020), forming the basis for the rhythm in electrical activity, which is viewed as an important timing signal issued by the SCN (see 3.4.). However,



TABLE 1 Effect of aging on the SCN

	Results (aged compared to young)	Model	Light condition	Target	Age (m) young	Age (m) old	# Time points	References	
Molecular clock									
Per1	-	No change in level and rhythm. Decreased induction of Per1 in response to light pulse in old.	Rat	DD (2d)	mRNA	2-3	22-26	6	(Asai et al., 2001)
	-	No change in level.	Mouse	DD (1d)	mRNA	4	15	2	(Weinert et al., 2001)
	-	No change in level. Shorter period in old.	Rat	12:12	mRNA	2	24-26	≥24	(Yamazaki et al., 2002)
	-	No change in level and rhythm. Decreased induction of Per1 in response to light pulse.	Hamster	DD (1-3d)	mRNA	2-3	17-20	4	(Kolker et al., 2003)
	x	Slightly reduced response to phase advance in old. No change in response to phase delay.	Rat	12:12 & shift	protein	4-8	>24	≥24.	(Davidson et al., 2008)
	-	No change in level. Possibly slightly phase advanced.	Mouse	12:12	mRNA	3	22	4	(Bonaconsa et al., 2014)
Per2	-	No change in level and rhythm.	Rat	DD (2d)	mRNA	2-3	22-26	6	(Asai et al., 2001)
	↓	Lower level at ZT7, but not at ZT 21.	Mouse	DD (1d)	mRNA	4	15	2	(Weinert et al., 2001)
	-	No change in level and rhythm. No change in response to light pulse	Hamster	DD (1-3d)	mRNA	2-3	17-20	4	(Kolker et al., 2003)
	-	No change in level. Reduced amplitude of 3rd cycle.	Mouse	12:12	protein	3-6	13-16	≥24/4	(Nakamura et al., 2011)
	↑	Higher amplitude in old. Increased response to phase advance in old.	Mouse	12:12 & shift	protein	3-6	22-28	≥24.	(Sellix et al., 2012)
	↓	Lower level at ZT4	Mouse	DD (2d)	mRNA	6	21	1	(Chang & Guarente, 2013)
	-	No change in amplitude at baseline. Changed response to phase shift in old.	Mouse	12:12 & shift	protein	3.5	19	≥24.	(Leise et al., 2013)
	↓	Possibly slightly reduced level and phase advanced.	Mouse	12:12	mRNA	3	22	4	(Bonaconsa et al., 2014)
	-	Faster decay of amplitude in old (12:12 & DD). Small phase delay and longer period in DD.	Mouse	12:12 & DD	protein	3-5	13-15	≥24.	(Nakamura et al. 2015)
	-	No change in amplitude under LD. Higher incidence of arrhythmicity in LL.	Mouse	12:12, LL & DD	protein	8-13	24-26	≥24.	(Polidarova et al., 2017)
-	No change in rhythm under LD 12:12 and long and short photoperiod (16:8/8:16)	Mouse	12:12, 16:8, 8:16	protein	4-8	22-28	≥24.	(Buijink et al., 2020)	
Cry1	↓	Lower amplitude.	Rat	DD (2d)	mRNA	2-3	22-26	6	(Asai et al., 2001)
	-	No change in level. Rhythmic in old, not in young.	Mouse	DD (1d)	mRNA	4	15	2	(Weinert et al., 2001)
	-	No change in level. Rhythmic in old, not in young	Mouse	12:12	mRNA	3	22	4	(Bonaconsa et al., 2014)
Cry2	-	No change in level. Not rhythmic in young and old.	Mouse	12:12	mRNA	3	22	4	(Bonaconsa et al., 2014)
Clock	-	No change in level. Not rhythmic in young and old.	Mouse	DD (1d)	mRNA	4	15	2	(Weinert et al., 2001)
	↓	Lower level in old. Not rhythmic in young and old.	Hamster	DD (1-3d)	mRNA	2-3	17-20	4	(Kolker et al., 2003)

(Continues)

TABLE 1 (Continued)

	Results (aged compared to young)	Model	Light condition	Target	Age (m) young	Age (m) old	# Time points	References
	- No change in level. Rhythmic in old, not in young.	Mouse	12:12	protein	4	16	6	(Wyse & Coogan, 2010)
	- No change in level. Not rhythmic in young and old.	Mouse	12:12	mRNA	3	22	4	(Bonaconsa et al., 2014)
<i>Bmal1</i>	↓ Lower level at CT2 and CT20. No changes in rhythmicity.	Hamster	DD (1-3d)	mRNA	2-3	17-20	4	(Kolker et al., 2003)
	↓ Lower level. Not rhythmic in young and old.	Mouse	12:12	protein	4	16	6	(Wyse & Coogan, 2010)
	↓ Lower level at ZT4.	Mouse	DD (2d)	protein	6	21	1	(Chang & Guarente, 2013)
	↓ Lower level. Rhythmic in young and old.	Hamster	14:10	mRNA	3-5	17-21	4	(Duncan et al., 2013)
	↓ Lower level in dark period. Rhythmic in young, not in old.	Mouse	12:12	mRNA	3	22	4	(Bonaconsa et al., 2014)
Neuropeptides								
<i>VIP</i>	↓ Reduced number of VIP neurons.	Rat	-	peptide	7-8	32-33	-	(Chee et al., 1988)
	- No change in number of VIP neurons. No effect of high/low light intensity.	Rat	12:12	peptide	3	34-37	1	(Lucassen et al., 1995)
	↓ Reduced number of VIP neurons in males, not females.	Human	-	peptide	10-40y	66-99y	-	(Zhou et al., 1995)
	↓ Lower level and loss of rhythmicity of VIP.	Rat	12:12	mRNA	3	24		(Kawakami et al., 1997)
	↓ Lower level and reduced number of VIP neurons. Loss of rhythmicity.	Rat	14:10	mRNA	2-4	18-20	7	(Krajnak et al., 1998)
	↓ Lower level in neurons. Not rhythmic in young and old.	Hamster	14:10	mRNA	3-5	19-22	5	(Duncan et al., 2001)
	↓ Lower level and loss of rhythmicity of VIP.	Rat	12:12	mRNA	2-3	19-20	8	(Kalló et al., 2004)
	- No change in level and number of VIP neurons. Shift (delay) in peak expression.	m. lemur	14:10	peptide	2y	6y	5	(Cayetanot et al., 2005)
	↓ Reduced number of VIP neurons.	Rat	12:12	peptide	6	24	1	(Pereira et al., 2005)
<i>AVP</i>	↓ Reduced number of AVP neurons. SCN area and AVP neurons larger in aging.	Rat	-	peptide	7-8	32-33	-	(Roosendaal et al., 1987)
	↓ Reduced number of AVP neurons under low light intensities, no change in high light intensities.	Rat	12:12	peptide	3	34-37	1	(Lucassen et al., 1995)
	- No change in level and rhythmicity.	Rat	14:10	mRNA	2-4	18-20	7	(Krajnak et al., 1998)
	↓ Reduced number of AVP neurons.	Vole	12:12	peptide	4-5	11-12	1	(Van der Zee et al., 1999)
	- No change in level and rhythmicity.	Hamster	14:10	mRNA	3-5	19-22	5	(Duncan et al., 2001)
	- No change in level and rhythmicity. Changed expression of V1a and V1b receptor.	Rat	12:12	mRNA	2-3	19-20	8	(Kalamatianos et al., 2004)
	↑ Higher level. No change in number of AVP neurons. Shift (delay) in peak expression.	m. lemur	14:10	peptide	2y	6y	5	(Cayetanot et al., 2005)
	↓ Reduced number of AVP neurons.	Rat	12:12	peptide	6	24	1	(Pereira et al., 2005)

(Continues)



TABLE 1 (Continued)

	Results (aged compared to young)	Model	Light condition	Target	Age (m) young	Age (m) old	# Time points	References
Electrical activity							Rec. time	
↓	Reduced averaged peak neuronal firing rates (single-unit activity)	Rat	12:12	ex vivo	5–7	23–28	30h	(Satinoff et al., 1993)
↓	Reduced averaged neuronal firing rates during day (single-unit activity)	Hamster	12:12	ex vivo	2	24	7–12h	(Watanabe et al., 1995)
↓	Reduced averaged peak neuronal firing rates (single-unit activity)	Rat	12:12	ex vivo	3	28	-	(Ruby et al., 1998)
↓	Slightly lower day-night difference of spontaneous firing rate (single-unit activity). Higher proportion of silent cells during the day.	Mouse	12:12	ex vivo	2–3	14–21	<6h	(Nygard et al., 2005)
↓	Reduced amplitude of neuronal firing (single-unit activity)	Mouse	12:12	ex vivo	4–5	15–17	48h	(Biello, 2009)
↓	Lower amplitude of multi-unit neuronal activity (multi-unit activity)	Mouse	12:12 & DD	in vivo	4	15	>7d	(Nakamura et al., 2011)
↓	Lower day-night difference of spontaneous firing rate (single-unit activity), lower amplitude of multi-unit neuronal activity (multi-unit activity)	Mouse	12:12	ex vivo	3	23	-	(Farajnia et al., 2012)
↓	Lower amplitude of firing rates during the day (single-unit activity)	Mouse	12:12	ex vivo		18–20	8–12h	(Leise et al., 2013)

Note: Comparison of outcomes and parameters of different studies on molecular, neuropeptide, and electrical rhythms.

the link between the molecular clock and channel regulation has been described for only a few of ion channels to date (Kudo et al., 2015; Meredith et al., 2006; Schmutz et al., 2014) and more work is needed to identify the direct and indirect pathways used by the clock genes to control the circadian modulation of ion channel properties.

One of the best studied intracellular signaling molecules known to influence ion channel activity is Ca^{2+} . Transmembrane Ca^{2+} influx serves to modulate spike frequency, but can also activate (clock) gene expression, for instance to mediate light-induced phase shifts of the circadian rhythm (Golombek & Rosenstein, 2010). Interestingly, the level of intracellular Ca^{2+} concentration also shows a circadian rhythm with higher levels during the day compared to the night (Aguilar-Roblero et al., 2016; Colwell, 2000; Ikeda et al., 2003; Jones et al., 2018). While the source and the function of this Ca^{2+} rhythm is still not fully understood (Enoki et al., 2017; Ikeda et al., 2003), there is even less known about the impact of aging; however, one study suggests a dysregulation of the Ca^{2+} rhythm in aging (Farajnia et al., 2015). This lack of knowledge is surprising since there is a vast amount of literature on age-related Ca^{2+} homeostasis in neurons in other brain areas, especially the hippocampus (Calvo-Rodriguez et al., 2020; Uryash et al., 2020).

Importantly, intracellular Ca^{2+} signaling—often in concert with cAMP—comprises a cytosolic oscillator that reinforces SCN clock

function (Doi et al., 2011; O'Neill et al., 2008). In addition, the paracrine feedback (e.g. of vasoactive intestinal protein; VIP) via g-coupled receptors, can amplify the cAMP/ Ca^{2+} signaling and further stabilize clock function. There is evidence that the rhythm in cAMP (Gerhold et al., 2005) as well as VIP is compromised in the aging SCN (Krajnak et al., 1998) demonstrating again the complexity of the impact of aging on clock function.

For the SCN clock, the level of cell signaling and cytosolic oscillators seems to be gravely affected by aging, which is reflected in the consequences for the single-cell pacemaker described below. Given the bridging function between the molecular clock and the electrophysiological properties of SCN neurons, this level should be one of the important research-foci for mechanisms and interventions for the aging clock.

3.4 | Electrophysiological properties of SCN neurons and network are severely disrupted with aging

The SCN neurons express a circadian rhythm in electrical activity, with most neurons producing action potentials during the day and being




FIGURE 1 Aging affect multiple levels of the SCN differentially. The onion symbolizes different layers of the central circadian clock. The effect of aging on the different layers is symbolized on the right panels. Molecular rhythms seem to be less effected while diminished amplitude or reorganization can be seen in rhythms of intracellular Ca^{2+} , membrane ion channel activity, phase distribution of single SCN units of the neuronal network and behavior (from top to bottom). Panels were modified after (Farajnia et al., 2014)

relatively silent during the night (Brown & Piggins, 2007; Gillette & Tischkau, 1999). This rhythm is maintained in the absence of inputs from the eyes (light) or other parts of the brain, as demonstrated in vivo and in brain slices (Weaver, 1998). Together with the rhythmic release of neuropeptides like arginine vasopressin (AVP), the neuronal activity rhythm of the SCN is considered to be the major output to propagate the endogenous timing signal throughout other parts of the brain and control temporal order of the brain and other organs (Colwell, 2011; Kalsbeek et al., 2011; Paul et al., 2020). Contrary to the rather mild effects of aging on the molecular clock, the amplitude of the electrical activity rhythm is severely attenuated in the aged SCN as shown by in vivo and ex vivo studies (Biello, 2009; Farajnia et al., 2012, 2015; Leise et al., 2013; Nakamura et al., 2011; Nygard et al., 2005; Satinoff et al., 1993; Watanabe et al., 1995). It seems that, while the aging molecular clock keeps ticking, processes in the membrane clocks lose track of time. The majority of SCN neurons recorded from hypothalamic slices from old mice (>24 month) showed compromised rhythms in membrane potential and membrane conductance, suggesting loss of circadian control of passive and active ion currents (Farajnia et al., 2012). In SCN neurons, the circadian modulation of ion

channel activity is necessary to control membrane excitability and generate the rhythms in electrical activity (Brown & Piggins, 2007). The electrical activity is used for synchronization within the SCN network and functions as an important output timing signal to downstream targets (Brown & Piggins, 2007; Colwell, 2011; Harvey et al., 2020; Itri et al., 2005). The activity of several voltage dependent K^+ channels express a circadian modulation and seem to be required for proper electrical rhythm, like the fast delayed rectifier current (Itri et al., 2005), the A-type current (Granados-Fuentes et al., 2012; Itri et al., 2010) and Ca^{2+} -activated K^+ channels (Kent & Meredith, 2008; Pitts et al., 2006; Whitt et al., 2016).

Aging was found to diminish the circadian modulation of a number of voltage-dependent K^+ channels, but interestingly did not alter the channels' basic conductance (Farajnia et al., 2012, 2015). The loss in circadian rhythm in channel activity of the fast delayed rectifier channel and the transient K^+ channel can contribute to a change in firing frequency modulation (Farajnia et al., 2012), while the impact on the rhythm in large-conductance Ca^{2+} -activated K^+ current led to change in the waveform of the action potential (Farajnia et al., 2015). It seems that the age-resistant molecular clock becomes uncoupled

from the downstream processes, which are needed to control membrane rhythmicity.

As we have discussed above, there are several mechanisms that are responsible for the transmission of information from the electrical activity at the membrane to the molecular clock and vice versa. It is likely that aging-associated deterioration of one or more of these coupling mechanisms is responsible for the discrepancy seen between the effect of aging on the molecular and electrical clock. Environmental light information may reach the molecular clock, whereby it retains the capability of entraining to the light-dark cycle, including adapting phase distributions to long and short photoperiods (Buijink et al., 2020). However, this information fails to reach the electrical components of the central clock, both directly via projections from the eye, as well as indirectly, through communication with the molecular clock, resulting in an uncoupling of the electrical and molecular clock components. This is partially corroborated by the finding that the aging molecular clock is unable to remain stable under constant lighting conditions (Nakamura et al. 2015; Polidarova et al., 2017). Irrespective of the underlying cause, since electrical activity is one of the two main outputs from the SCN to the rest of the brain and body, it is likely that the reduced amplitude has consequences for the signaling of the SCN to other brain areas and the periphery.

3.5 | SCN network synchronization—neurotransmitter and peptide signaling

The SCN employs not only electrical signaling, but also multiple neurotransmitters and neuropeptides for communication between neurons of the SCN, as well as towards the periphery. There are several ways these means of communication are known to be affected by aging. For instance, synaptic terminals and signaling for the neurotransmitter gamma-aminobutyric acid decline in aging, which would affect the control of synchronization between SCN neurons (Biello, 2009; Farajnia et al., 2012; Nygard et al., 2005; Palomba et al., 2008). Aging is also accompanied by a reduced expression of the NR2B subunit of the N-Methyl-d-aspartic acid receptor, affecting glutamatergic signaling, which is the main input signal for the SCN (Biello, 2009; Biello et al., 2018). Another important 'synchronizer' in the SCN, the neuropeptide VIP, displays a reduction in its mRNA rhythm and fewer cells in the old SCN express VIP (Chee et al., 1988; Duncan et al., 2001; Kalló et al., 2004; Kawakami et al., 1997; Krajnak et al., 1998; Pereira et al., 2005; Zhou et al., 1995). Similarly, a significant reduction in cells expressing AVP—another essential neuropeptide in the SCN—was described in the aging rodent SCN (Roosendaal et al., 1987; Van der Zee et al., 1999), although several other studies did not find a difference in AVP mRNA levels and its rhythmicity (Duncan et al., 2001; Kalamatianos et al., 2004; Krajnak et al., 1998). In the mouse lemur, a small primate, VIP, and AVP protein levels are not affected, but their rhythms are phase delayed (Cayetanot et al., 2005). Importantly, exposure to increased light intensities could prevent the loss of AVP, but not of VIP neurons in rats (Lucassen et al., 1995). It is remarkable that, to our knowledge,

no recent study has been conducted on neuropeptides in the aging SCN. The emergence of new techniques, enabling the cell-specific tracking of activity by targeted labelling and imaging of these neuropeptides (Jones et al., 2018; Shan et al., 2020), should facilitate new studies on the impact of aging on peptidergic signaling in the SCN and will generate more in-depth knowledge of this system.

3.6 | From cells to network—A role for astrocytes?

Although neurons are still supposed to play the main role in timekeeping, recent studies have highlighted the importance of glia cells, specifically astrocytes, in the functioning of the SCN (Brancaccio et al., 2017, 2019; Tso et al., 2017). Furthermore, the importance of astrocytes in aging, and specifically the requirement of a functional clock in astrocytes was illustrated by the finding that the selective knock-out of *Bmal1* in only astrocytes shortens the lifespan in mice (Barca-Mayo et al., 2020). The circadian clock in astrocytes is found to be capable of driving circadian rhythm in the absence of a functional molecular clock in SCN neurons (Brancaccio et al., 2019), as well as modulate circadian period in the SCN and behavior in a normal functioning SCN (Tso et al., 2017). Therefore, it is likely that under normal circumstances astrocytes play a role in fine-tuning circadian clock function in the SCN. In the aging brain, astrocytes show signs of cellular senescence and undergo transcriptional changes that indicate neuroinflammation, a process detrimental for the correct functioning of these cells (Clarke et al., 2018; Salminen et al., 2011). Unfortunately, there is a shortage of studies on astrocytes (and other glia cells) in the aging SCN, so it is unclear to what extent they are involved in its malfunctioning. One study at least indicates that the number of astrocytes is increased in the aged SCN, which warrants additional research into the functioning of astrocytes in the old SCN (Deng et al., 2010).

4 | AGING OF CLOCKS OUTSIDE THE SCN

The SCN orchestrates circadian rhythms in the rest of the brain and body through different pathways. The output pathways of the SCN are affected by aging in different ways, in turn resulting in various disturbances of circadian rhythms throughout the brain and body. In this part of the review we will discuss the effect of aging on target areas of the SCN, namely the hypothalamus and the pineal gland, as well as how dysregulated circadian rhythms in aging affects the rest of the brain and body.

4.1 | Communication with the periphery—Outputs of the SCN within the hypothalamus

Most of the direct output areas of the SCN are located in other parts of the hypothalamus, a crucial brain region for regulating a



wide range of physiological processes (Kalsbeek & Buijs, 2002). This makes the location of the SCN within the hypothalamus ideal to relay non-visual light information from the eye to the rest of the brain and body for circadian regulation of their physiological processes. The importance of circadian rhythms for functions of the hypothalamus is illustrated by the relatively strong autonomous rhythms in several hypothalamic areas outside the SCN (Guilding et al., 2009). The processes in the hypothalamus that require a circadian regulation are supported by input from the SCN. In turn, rhythmic processes in the periphery, like feeding and physical activity feedback to the SCN mainly through other areas in the hypothalamus (Buijs et al., 2017, 2019). These reciprocal processes are essential for energy preservation and for hormone homeostasis. Many of the regulatory functions of the SCN that are affected by aging, like energy homeostasis, sleep-wake rhythms and hormonal balance, are mediated through areas within the hypothalamus (Kim & Choe, 2019). Indeed, the hypothalamus is even suggested to be a key regulator in systemic aging (Zhang et al., 2013).

One of the hallmarks of aging is a deregulation of nutrient sensing, which is essential to control the organism's energy balance (López-Otín et al., 2013). A complex network of neurons in the dorsomedial, ventromedial and lateral hypothalamus, as well as the paraventricular nucleus and arcuate nucleus of the hypothalamus has developed to sense and regulate vertebrate energy expenditure and food intake. The essential circadian regulation of this network is achieved by the SCN, which has direct projections to the paraventricular nucleus and the dorsomedial hypothalamus (Kalsbeek et al., 1993). Two key proteins in nutrient sensing—the mammalian target of rapamycin (mTOR) and sirtuin 1 (SIRT1)—are critical in the aging process and are directly linked to the molecular clock.

The mTOR pathway is able to detect changes in energy status, by monitoring levels of ATP and amino acids, and modify a wide variety of essential cellular processes (Wullschlegel et al., 2006). In the hypothalamus of old mice, mTOR signaling in pro-opiomelanocortin neurons is increased, which contributes to an aging phenotype (Yang et al., 2012). On the other hand, pharmacological inhibition of mTOR activity, for example with rapamycin, promotes longevity and health span (Harrison et al., 2009; Vellai et al., 2003). Importantly, mTOR activity is also inhibited by a calorie restricted diet (Kaeberlein et al., 2005), which to date is the most efficient life extending intervention in multiple organisms, from yeast to primates (Colman et al., 2009; Lin et al., 2002). The mTOR pathway is linked to the timekeeping system in multiple ways. In the SCN, the mTOR pathway can be triggered by light (Cao et al., 2008), while the core clock gene *Bmal1* negatively regulates mTOR signaling (Khapre et al., 2014). Inhibition of mTOR signaling reduces the amplitude and lengthens the period of *Per2* expression rhythm in the SCN and liver (Ramanathan et al., 2018). Another essential regulator of cellular energy status is sirtuin 1 (SIRT1). Sirtuins are a class of proteins that detect their surrounding energy status by sensing oxidized NAD⁺, and respond with the regulation of a multitude of metabolic processes (Houtkooper et al., 2012). Like mTOR, SIRT1 is a candidate target for life extending agents, since overexpression of its monologue in

yeast, Sir2 was found to increase lifespan (Kaeberlein et al., 1999). SIRT1 is similarly thought to be involved in the effect of calorie restriction on life span, since calorie restriction enhances SIRT1 expression (Cohen et al., 2004). Interestingly, SIRT1 influences the transcription of the core clock genes *Bmal1*, *Clock*, *Per2* and *Cry*, and SIRT1 knockout or overexpression modulates circadian behavior (Chang & Guarente, 2013). Moreover, SIRT1 levels decreases in the aged SCN, which results in a longer free-running period in behavior, while overexpression of SIRT1 in the hypothalamus slows down the aging process (Chang & Guarente 2013b; Satoh et al., 2013). The function of mTOR and SIRT1 in nutrient sensing and their potential to directly modify core clock genes in both the SCN, as well as in peripheral tissue points at an important role in conveying information on energy status to the circadian system.

Disturbances of circadian rhythmicity at the level of the SCN, the SCN target brain areas, or the periphery can cause diverse health issues, mostly based on changes in energy metabolism. The close entanglement of aging with the system regulating energy homeostasis and with the circadian system, highlights the role of energy consumption in modifying circadian rhythms and influence the aging process. Healthy and moderate food intake at the correct time of day will likely support peripheral oscillator function as well as provide appropriate timing feedback to the hypothalamus, promoting energy homeostasis, overall health, and longevity.

4.2 | Circadian rhythms and memory impairment—Aging of the hippocampus

One of the most debilitating consequences of the senescent brain is the diminished memory forming and retrieval. Even healthy aging is often accompanied by memory deficits, but aging is also the prominent risk factor for developing dementia. One of the brain centres playing an important role for declarative memory is the hippocampus and aging-related memory deficits have been linked to changes in this brain area (Fan et al., 2017). The hippocampus, like many other brain regions (Paul et al., 2020), exhibits circadian rhythms in clock gene expression and physiological functions. One example hinting how clock genes in the hippocampus may contribute to neurodegeneration is demonstrated by the deletion of clock genes (e.g. *bmal1*, *clock* and *Npas2*) leading to neuronal oxidative damage, seemingly through a disturbance in the redox homeostasis in the brain (Musiek et al., 2013).

Circadian rhythms in learning and memory have been described (Chaudhury & Colwell, 2002; Snider et al., 2018), which seem to persist in VIP-KO mice with compromised SCN function (Chaudhury et al., 2008). Interestingly, the circadian modulation of long-term potentiation, which is thought to be a physiological correlate for learning, persists in hippocampal brain slices thus demonstrating that hippocampal rhythms may be independent of inputs from the central clock in the SCN (Wang et al., 2009). Together the data suggest that local oscillators can control daily rhythms of neuronal performance in the hippocampus, however, detailed studies on these independent circadian oscillators and their function are still sparse.



Similar to the SCN, aging seems to affect the clock genes in the hippocampus less severe (Duncan et al., 2013) compared to the more drastic effects on downstream rhythms in metabolites, or membrane and synaptic properties as seen in a recent proteomics study (Adler et al., 2020). However, while the impact of aging on hippocampal function is well described, the contribution of the circadian clock to aging-related dysfunctions still needs further studies. It is important to realize that this effect of the circadian system on the aging hippocampus is not necessary a direct impact from the central clock. One of the systemic timing cues, corticosterone, showed a lower amplitude of daily variation in plasma and hippocampus in aged rats (24 months), but its concentration was in both still higher during the night compared to the day. (Barrientos et al., 2015). The plasma rhythms in old rats showed a reduction of their peak values during the night, while the corticosterone rhythms in the hippocampus were blunted by an increase of the through levels during the day. It is important to note that this study used automated blood-samplers, which avoided stress-induced corticosterone release in aged animals. Interestingly, the elevation of corticosterone in the hippocampus during the inactive phase of the old animals led to a stronger activation of glucocorticoid receptors, activation of microglia, increasing neuroinflammation and potentially contributing to cognitive decline (Barrientos et al., 2015). A recent study on adult hippocampal neural stem cell pools, which represent a reserve for maintaining plasticity through neurogenesis, also shows a negative effect of the presence of glucocorticoid receptors (Schouten et al., 2020). However, using a mouse model for accelerated aging, this study also finds benefits of circadian modulation in corticosterone, which, together with ultradian oscillations, helps to sustain this reserve pool of neural stem/precursor cells. And finally, another example for the interconnectedness of the different players involved in aging is the influence of aging on microglia function in the hippocampus. These first responders of the brain immune system also have surveying function in their 'resting' state and contribute to maintenance of synaptic function in brain networks like the hippocampus (Schafer et al., 2012). In aging, microglia will change morphology and physiology and their immune response is compromised (Nissen, 2017). In addition, microglia in the aging hippocampus has been shown to have altered rhythms in period gene expression and lose their circadian rhythm in immune-challenge-induced cytokine response (Fonken et al., 2016). Again, it seems that dampened amplitude of the corticosterone rhythms contributes to the disruption of circadian rhythms in microglia and an exaggerated immune response in the aged hippocampus. It can be expected that the resulting increase in chronic inflammation will affect neuronal circuits in the hippocampus and the rest of the brain, as discussed below.

4.3 | The interaction between circadian rhythmicity and chronic (neuro)inflammation

The link between the circadian clock and the peripheral immune system has been well established (Labrecque & Cermakian, 2015;

Scheiermann et al., 2018) and the clock also plays a role in CNS immune responses (Fonken et al., 2015; McKee et al., 2020), which can be of particular interest for aging and neurodegenerative diseases (Lananna & Musiek, 2020). The peak of immune responsiveness occurs at the end of the inactive period suggesting a preparation for potential attacks on the immune system during the active period of the organism (Curtis et al., 2014). Chronic inflammation is a key hallmark of aging in periphery and the CNS, caused by malfunctioning autophagy, damaged cells and cellular debris lingering within tissue, sending out pro-inflammatory signals (Di Benedetto et al., 2017). In the brain, interactions between neurons, astrocytes and microglia regulate the immune response (Castellani & Schwartz, 2020) and this crosstalk is significantly affected by aging (Rodriguez-Arellano et al., 2016; Santoro et al., 2018). In addition, astrocytes and microglia both have circadian clocks modulating their function (Fonken et al., 2015; Prolo et al., 2005), which are altered in aging (Fonken et al., 2016). The astrocyte clock function has been best described in the SCN, where it acts in concert with neurons to ensure the stable generation of circadian rhythm in neuronal activity (Brancaccio et al., 2017; Prolo et al., 2005) even in the absence of a clock in SCN neurons (Brancaccio et al., 2019). There is some evidence that the astrocyte clock also plays a role in regulating neuroinflammation by regulating Nuclear Factor kappa-light-chain-enhancer of activated B cells (NF- κ B) proinflammatory signaling (Sugimoto et al., 2014) and brain-derived neurotrophic factor (BDNF) dependent protection of neurons from oxidative stress (Ishii et al., 2019). The circadian clock in the brain's early responders to an immune challenge, the microglia (Bilbo & Stevens, 2017), modulates gene expression of cytokines and response to an immune lipopolysaccharide challenge as found in isolated hippocampal microglia (Fonken et al., 2015). Interestingly, these rhythms of cytokine release and immune response of microglia are blunted in 24 months old rats (Fonken et al., 2016). Given that clock deficient monocytes show higher lethality after immune challenge (Nguyen et al., 2013), the reduction of clock function in aging immune cells may be contributing to the loss of function in neuronal network in the SCN and the rest of the brain.

Aging-associated inflammation is not only a problem for the central nervous system, but also a systemic problem, and there are several ways the circadian system is known to contribute to (chronic) inflammation and a subsequent diseased state in peripheral tissue. For instance, forced circadian disruption in mice causes an exaggerated inflammatory response and even death in response to an immune challenge (Castanon-Cervantes et al., 2010). In a recent study, a causal link was found between disruptions in circadian rhythms and the development of atherosclerosis through an aberrant inflammatory response (Schilperoort et al., 2020). Inflammation and aging are often linked to melatonin and glucocorticoid, both of which display a strong circadian rhythm. Not surprisingly, the link between aging, aging-related inflammation and the circadian system is often made. However, the question remains to what extend the connection is (bi) directional, and thus, if purely strengthening circadian rhythmicity could lower inflammatory burden in aging. Nevertheless, there is strong evidence that several interventions that strengthen the



circadian clock, either directly or indirectly, also decreases inflammation markers.

4.4 | Communication with the periphery—The aging pineal gland and its product, melatonin

An important relay station between the SCN and the periphery is the pineal gland. The pineal gland is innervated indirectly by the SCN, through a pathway leading from the SCN to the PVN in the hypothalamus, via ganglia in the spinal cord and to the pineal gland (Tecler-Mariam-Mesbah et al., 1999). The main function of the pineal gland is the production and release of melatonin, acting as a systemic circadian timing signal (Lewy et al., 1980; Pang & Ralph, 1975; Pelham, 1975). Melatonin production by the pineal gland normally peaks during the dark period, and induces sleepiness in humans.

The role of the pineal gland in the aging process became clear from studying the effect of pinealectomy and pineal transplantations. Pinealectomy at a young age induced an aging phenotype, and transplanting young pineal tissue to aged animals caused rejuvenation (De Butte & Pappas, 2007; Pierpaoli & Bulian, 2005; Pierpaoli & Regelson, 1994). However, it is still unknown to what extent the pineal gland itself is actually affected by aging. The pineal gland undergoes clear aging-associated calcification, but it is not certain if this affects its function (Tan et al., 2018). One indication for a functional deficit is a significant advance in the phase of *Per1* rhythms in the aging pineal gland (Yamazaki et al., 2002). Moreover there are many studies that found a dampened circadian rhythm in melatonin, because of lower melatonin levels during night-time (Iguchi et al., 1982). This could be a consequence of dysfunction of the pineal gland, or due to reduced innervations from the SCN (Jengeleski et al., 1989). The age-related decline in melatonin was questioned by a study using a rigorous method with a constant routine, and taking into account several non-prescription drugs, like ibuprofen and aspirin, which suppress melatonin and are often used by the elderly (Duffy et al., 2002; Zeitzer et al., 1999). However, the same researchers did find a reduced melatonin level when subjects were allowed to sleep, meaning that under real life conditions, melatonin rhythms do deteriorate in aging (Zeitzer et al., 2007). In addition, melatonin receptor expression in peripheral tissue is found to decrease in aging (Sánchez-Hidalgo et al., 2009).

The decline in melatonin amplitude as well as the decreased melatonin receptor expression could well be contributing to a diminished synchronization of the central and peripheral clocks (Pfeffer et al., 2018). Importantly, aging had also been associated with a decline in melatonin sensitivity of the SCN (von Gall & Weaver, 2008; Wu et al., 2007), which may partially account for the reduced rhythm strength of the SCN observed in aging (Pfeffer et al., 2017). Melatonin is linked to several aging associated disorders, like cardiovascular diseases (Dominguez-Rodríguez et al., 2010), type 2 diabetes (Karamitri & Jockers, 2019), neurodegenerative diseases (Wu & Swaab, 2005) and (neuro)inflammation (Hardeland, 2018). However, it is currently unclear if this is a direct consequence of the

attenuated melatonin signal, or indirectly acts through a weakening of the circadian system.

Melatonin, both from the pineal gland, as well as from other sources, has an important role in the immune system (Markus et al., 2018), and administration of melatonin is found to counteract age-associated inflammation (Rodríguez et al., 2007). Although most cells of the body can produce melatonin, it is the pineal gland derived melatonin that plays a role in the circadian rhythmicity of the immune response (Lopes et al., 1997). The idea of melatonin as a contributor and possible treatment for neurodegenerative disease is currently receiving much attention (Cardinali, 2019; Socaciu et al., 2020; and see below). This is on the one hand based on the finding that neurodegenerative diseases are associated with, and possibly preceded by a decline in melatonin levels (Wu & Swaab, 2005). On the other hand, melatonin has a proven neuroprotective role through its anti-oxidant and anti-inflammatory function (Alghamdi, 2018). And finally, melatonin supports the function of the SCN and may strengthen circadian rhythmicity in aging. However, the contribution of compromised systemic level or circadian rhythm amplitude of melatonin to the development of neurodegenerative diseases is still unclear (Cardinali, 2019; Zhang et al., 2016).

Thus, aging goes along with a decline in circadian rhythmicity of melatonin produced by the pineal gland. The decline in melatonin levels and rhythm amplitude is associated with several processes that are at the core of health deterioration during aging, like chronic inflammation and reduced synchronization of peripheral clocks (Pfeffer et al., 2018; Rodríguez et al., 2007). Concordantly, melatonin has many properties that can delay the aging processes, like its strong anti-oxidant function, promotion of neurogenesis and mitochondrial health (Ramírez-Rodríguez et al., 2009; Tan et al., 2016). More research is needed to confirm if treatment with melatonin leads to improve overall health in aging, and it will be very interesting to see to what extent a synchronization of central and/or peripheral clocks is involved.

4.5 | Aging clocks in peripheral tissue

The discovery of the molecular clock has had a major impact on the subsequent discovery of self-sustaining clocks in a wide variety of tissues, from the liver (Yamazaki et al., 2000) to the cochlea (Park et al., 2017) and beyond (Brown et al., 2019). It has led to the detection of differentially phased rhythms in a variety of tissues, painting the picture of a complex hierarchical system of central and peripheral pacemakers and oscillators (Brown et al., 2019). Studies using bioluminescent reporters for clock genes have shown that the molecular clock in the SCN is relatively unaffected by aging, while the rhythmicity and phase shifting capacity of the molecular clock in the periphery is considerably more affected (Davidson et al., 2008; Sellix et al., 2012). For instance, Davidson and colleagues show that the *Per2* expression rhythm in the aging liver is incapable of following a 6-hr shift in the LD cycle within the six days measured in the experiment, while the aging SCN showed an almost similar re-entrainment



as the young SCN (Davidson et al., 2008). Even under normal lighting conditions, circadian rhythms drift out of phase in aging, affecting the whole-body synchrony of temporal processes (Bonaconsa et al., 2014; Wyse & Coogan, 2010; Yamazaki et al., 2002).

From deteriorating of whole-body homeostasis to impaired molecular processes within single cells, the effects of senescence are noticeable on all levels. The affected processes often concern metabolism, from aging-associated obesity, to mitochondrial deficiency. Most, if not all, metabolic processes are highly rhythmic, for example lipid and glucose homeostasis, and these rhythmic processes are often found to be disturbed in aging. A recent study shows that aging severely affects the levels and rhythmicity of lipids in brown and white adipose tissue, indicating a disturbed lipid metabolism (Held et al. 2020). Aging is also associated with changes in rhythmic gene expression in the liver, affecting genes associated with metabolic processes (Sato et al., 2017). Interestingly, both this study, as well as several others find that the rhythmic expression of the core clock genes in peripheral tissue is relatively unaffected by aging, compared to other classes of transcripts. This indicates that the molecular clock in the periphery is more resilient to aging than their target genes, which shows parallels to the aging SCN, discussed in paragraph 3.2 and 3.3.

In line with the fact that many of the processes affected by aging concern metabolism, dietary interventions that consist of caloric restriction boost longevity, and improve overall health (see 4.1). Importantly, time restricted feeding—allowing food intake only during certain hours of the day—can have a similar beneficial health effect, which is likely a consequence of a stronger temporal regulation of metabolism. For example, a study by Hatori and colleagues showed that the detrimental effects of a high-fat diet on health, like metabolic disease, type-2 diabetes and liver damage—diseases also prevalent in aging humans—could be counteracted by only adjusting the time-window of the availability of the high-fat diet. It could even prevent weight gain, while the daily caloric intake was the same for the ad libitum and time-restricted feeding group (Hatori et al., 2012). Furthermore, time-restricted feeding is capable of restoring gene expression rhythms in the liver and WAT when circadian rhythms are disrupted (Yamamuro et al., 2020). Similarly, overall lowering of caloric intake also leads to a considerable upregulation of circadian rhythmicity in the aging liver through the important metabolic pathway NAD⁺/SIRT (Sato et al., 2017). Beneficial effects are also found with diets that mimic continuous calorie restriction with a periodic reduction of caloric intake (Brandhorst et al., 2015). Both these so-called fasting-mimicking diets as well as time-restricted feeding do not have clear negative side-effects, and are proposed as feasible interventions with health span increasing potential (Longo & Panda, 2016). It is possible that the health benefits from these diets are partially attributable to the strengthened circadian rhythmicity, since there is a direct interaction between the main proteins involved in the effect of calorie restriction and the molecular clock (see 4.1). It will be interesting to investigate what paradigms lead to the best balance between health benefits and are also feasible to implement in daily life.

5 | CIRCADIAN CLOCKS AND NEURODEGENERATIVE DISEASE

In addition to the general hallmarks of aging like mitochondrial dysfunction, dysregulated energy metabolism, oxidative damage, impaired proteostasis, inflammation, deficiencies in DNA repair and stem cell exhaustion (López-Otín et al., 2013), brain aging is marked by impaired cellular stress response, aberrant neuronal network activity and dysregulated neuronal calcium homeostasis (Mattson & Arumugam, 2018). Non-pathological brain aging leads to compromised brain function, with a reduced cognitive function as one of the most common consequences. However, aging is accompanied by an increased risk for developing neurodegenerative diseases, with growing prevalence because of the increase in life-expectancy.

The relationship between the function of the circadian system and neurodegeneration is bidirectional; neurodegeneration can impair the SCN function (Bellanti et al., 2017; Leng et al., 2019) and the functional state of the circadian clock can affect the progression of the disease (Duncan, 2020). While this constitutes a downward spiral, the timely strengthening of clock function may activate an upward spiral, slowing the process of neurodegeneration (Figure 2). Indeed, using bright light with melatonin (Riemersma-van der Lek et al., 2008) or just melatonin (Wade et al., 2014) as chronotherapeutic interventions in the elderly diagnosed with dementia, resulted in improved cognitive function. In addition, the interventions also improved sleep disorders, which are often associated with neurodegeneration. Similarly, bright light also strengthened sleep-wake rhythms in Parkinson's patients (Videnovic et al., 2017). However, given the small number of studies and inconsistencies, more studies need to be conducted to determine the value for the treatment of neurodegenerative disease.

The impact of aging on circadian clock function is not necessarily the result of neurodegeneration. Senescent-related alteration in clock function are often milder compared to deficits seen in neurodegenerative diseases like Alzheimer's disease (AD; Chauhan et al., 2017; Leng et al., 2019). At the level of the SCN, there are




FIGURE 2 Effect of aging on the organism can produce a downward spiral while chronotherapeutic interventions can counteract this trend

ambiguous reports on neuronal loss in healthy aging (Madeira et al., 1995; Roberts et al., 2012; Tsukahara et al., 2005) and the clock dysfunction is more likely caused by changes in cell physiology and network properties (see above). In contrast, volume and cell count of the central clock are reduced in AD (Swaab et al., 1985) and specifically neuropeptidergic neurons are diminished in the SCN of AD patients (Ishunina & Swaab, 2002; Zhou et al., 1995). In addition to cell loss, a reduction of melatonin receptor expression (MT-1) has also been observed in AD patients, which reduces the positive feedback of melatonin on SCN function (Wu et al., 2007). Furthermore, the number of astrocytes in relation to neurons is increased in these patients, which indicates that another culprit of AD—neuroinflammation—is also contributing to the pathogenesis of AD in the SCN (Castellani & Schwartz, 2020). One general problem here is the distinction of senescent-related effects of neuroinflammation and the additional effect of AD, since markers of neuroinflammation are also reported in the SCN of healthy aging rats (Deng et al., 2010). Studies with AD models or AD patients has yet to reveal direct evidence for the link between neuroinflammation and AD in the SCN, although the prerequisites for this—increase in astrocytes and neurofibrillary tangles have been found in the SCN (Roy et al., 2019). However, the SCN shows a somewhat untypical AD pathology with no mature plaques present, while the surrounding hypothalamus does show typical plaque development (Stopa et al., 1999).

An interesting question from both the mechanistic, as well as therapeutic view is if a dysfunctional clock promotes AD or if clock deficiencies are only caused by AD at a later stage. The current data of patients with preclinical stages of AD indicated that circadian disruptions set in very early, if not preceding the AD pathogenesis (Musiek et al., 2018). In addition, several studies confirm the correlation of sleep- or rhythms-disorders and AD pathology markers like amyloid-beta levels or neuronal atrophy (Ju et al., 2017; Van Someren et al., 2019; Sprecher et al., 2017) and even cognitive decline (Waller et al., 2016). The further investigation of this link may offer not only new mechanistic insights, but could also be useful for diagnostic purpose as well as chronotherapeutic approaches. The interventions discussed in the next paragraph could therefore not only promote healthy aging of the brain and our clock, but may also delay the onset and/or progression of neurodegenerative diseases.

6 | CHRONOTHERAPIES

Aging per se is not a disease and therefore interventions to counteract aging-related limitations in performance of the elderly can be questionable. However, aging can also present a health risk for many individuals, increasing their chances for cardiovascular and neurodegenerative diseases, among others. The degree of interventions is therefore dependent on the level of impairment caused by aging and/or aging-related diseases. The circadian clock with its unique position to regulate the temporal efficiency of many physiological functions in our body is in any case a good mediator for interventions with systemic impact.

One of the most efficient and non-invasive ways to influence the clock is light and, indeed as mentioned above, light has been successfully used to strengthen the clock in the elderly and partially restore dysregulated sleep patterns. A second important gate to strengthen the clock is lifestyle. Regular and right timing of behaviors like eating, exercise and sleeping can help in turn to synchronize central and peripheral oscillators with positive effects on physiological function especially in aging and disease (Figure 3).

Physical activity—even at a moderate level—seems to strengthen clock function in general (Hughes & Piggins, 2012; Leise et al., 2013) and in aging in particular (Dupont Rocher et al., 2016; Gu et al., 2015). Similar to light, the timing of these activities is important. For instance, eating at the wrong time of day would have adverse effects on the clock. This has been shown in a number of animal experiments using scheduled feeding (Mukherji et al., 2015), and is also evident from studies in humans (Wehrens et al., 2017). Lifestyle-adjustments like regular bedtimes are also a tool to increase temporal order in the circadian system, as misalignment can promote serious diseases like metabolic syndrome (Leproult et al., 2014). A third venue of interventions are substances that impact circadian clock function. One of the earliest agents used to influence clock function is melatonin or a synthetic agonist of it (Zisapel, 2018). Studies in both shift-workers and the elderly have shown improvement of sleep-wake rhythms after administration of melatonin (Riemersma-van der Lek et al., 2008; Skene & Arendt, 2006). More recently, high throughput drug screenings have identified so called clock enhancing molecules: synthetic small molecules with little to no side effects, which can influence phase, period or amplitude of the rhythm in central or peripheral clocks (Chen et al., 2018). Currently, some of these agents have been shown to successfully target the interface between circadian clock and serious diseases (Miller & Hirota, 2020) like diabetes (He et al., 2016), immune disorders (Solt et al., 2011) and also age-related metabolic challenges (Nohara et al., 2019) in animal models.

While controlled (day)light exposure as well as lifestyle approaches are appropriate to assist the organism in healthy aging, the role of these and other chronotherapies in neurodegenerative disease has to be thoroughly investigated. As mentioned earlier, we need to know more about the interaction between circadian system and neurodegenerative diseases. Is it a mere correlation or are they causally connected? More studies using multiple markers of the circadian clock, like body temperature and melatonin profile, in addition to behavior measurements will give better insights into the mechanisms of circadian disruption in these diseases and their potential (chrono)treatment.

7 | CONCLUSION AND FUTURE DEVELOPMENTS

The central clock has many ways to confer itemporal information to the rest of the body. It is the collective effort of molecular,




FIGURE 3 Potential intervention strategies to strengthen the circadian clock in aging. Among the lifestyle interventions regular bedtimes and mealtimes, exercise, social interaction and (day)-light exposure can synchronize the body clocks, when applied at the 'right' time. Pharmacological interventions with small molecules and melatonin may boost the amplitude of the cellular circadian rhythms, while neuroactive compounds may target network synchronisation. Inhibitors of mTOR like rapamycin acting on the cellular metabolism, may reduce the effect of aging on clock function.

cytosolic, membrane, and network oscillators that make up the precision, strength and flexibility of the SCN rhythm, but there is some level of redundancy (Figure 1). In addition, the function of the SCN is even better adapted and tuned to the challenges in environment and physiology by feedback loops from its target organs and behavior. As a result, the SCN takes the role of a conductor of a well-rehearsed orchestra of peripheral clocks that efficiently regulates the temporal order in our body. The

impact of aging on the clock function is therefore as complex as the intertwined symphony of rhythms. Therefore, strengthening of the rhythm of the SCN alone might not be sufficient to restore circadian rhythms in the rest of the body, and a multi-level intervention should be warranted. Supporting proper rhythmicity in both the SCN, the rest of the brain, as well as the periphery by means of correctly timed (and dosed) light exposure, food intake, exercise, and possibly treatment with drugs acting on



clock function, could contribute to healthy aging and slow down the progression of neurodegenerative disease (Figure 3). While the lifestyle interventions are readily applicable to humans and mostly work on a systemic level, it is still unclear which cellular or subcellular targets would be most effective for drug interventions. Aging has minor impact on the molecular clock therefore this level may be a less rewarding target for interventions. Overall, improving synchronisation within the cell, the SCN network and between central and peripheral clocks may reinstate precision and stability in the aged circadian system.

ACKNOWLEDGMENT

The authors thank L. Hausmann for her patience and support. Figures were re-drawn by Marco Bazelmans in BioRender (<https://biorender.com/>) on the basis of a draft provided by the author. This work was supported by the Velux Foundation, grant# 1029 to SM.

CONFLICT OF INTEREST

The author declare no conflict of interest.

ORCID

M. Renate Buijink  <https://orcid.org/0000-0002-0904-139X>

Stephan Michel  <https://orcid.org/0000-0001-5506-5037>

REFERENCES

- Adler, P., Chiang, C. -K., Mayne, J., Ning, Z., Zhang, X., Xu, B., Cheng, H. -Y., - M., & Figeys, D. (2020). Aging disrupts the circadian patterns of protein expression in the murine hippocampus. *Frontiers in Aging Neuroscience*, 11. <https://doi.org/10.3389/fnagi.2019.00368>.
- Aguilar-Roblero, R., Quinto, D., Báez-Ruiz, A., Chávez, J. L., Belin, A. C., Díaz-Muñoz, M., Michel, S., & Lundkvist, G. (2016). Ryanodine-sensitive intracellular Ca^{2+} channels are involved in the output from the SCN circadian clock. *European Journal of Neuroscience*, 44, 2504–2514.
- Alghamdi, B. S. (2018). The neuroprotective role of melatonin in neurological disorders. *Journal of Neuroscience Research*, 96, 1136–1149.
- Asai, M., Yoshinobu, Y., Kaneko, S., Mori, A., Nikaido, T., Moriya, T., Akiyama, M., & Shibata, S. (2001). Circadian profile of Per gene mRNA expression in the suprachiasmatic nucleus, paraventricular nucleus, and pineal body of aged rats. *Journal of Neuroscience Research*, 66, 1133–1139.
- Barca-Mayo, O., Boender, A. J., Armirotti, A., & De Pietri Tonelli, D. (2020). Deletion of astrocytic BMAL1 results in metabolic imbalance and shorter lifespan in mice. *Glia*, 68, 1131–1147. <https://doi.org/10.1002/glia.23764>
- Barrientos, R. M., Thompson, V. M., Kitt, M. M., Amat, J., Hale, M. W., Frank, M. G., Crysedale, N. Y., Stamper, C. E., Hennessey, P. A., Watkins, L. R., Spencer, R. L., Lowry, C. A., & Maier, S. F. (2015). Greater glucocorticoid receptor activation in hippocampus of aged rats sensitizes microglia. *Neurobiology of Aging*, 36, 1483–1495.
- Bellanti, F., Iannelli, G., Blonda, M., Tamborra, R., Villani, R., Romano, A., Calcagnini, S., Mazzoccoli, G., Vinciguerra, M., Gaetani, S., Giudetti, A. M., Vendemiale, G., Cassano, T., & Serviddio, G. (2017). Alterations of clock gene RNA expression in brain regions of a triple transgenic model of Alzheimer's disease. *Journal of Alzheimer's Disease*, 59, 615–631. <https://doi.org/10.3233/JAD-160942>
- Benloucif, S., Green, K., L'Hermite-Balériaux, M., Weintraub, S., Wolfe, L. F., & Zee, P. C. (2006). Responsiveness of the aging circadian clock to light. *Neurobiology of Aging*, 27, 1870–1879.
- Biello, S. M. (2009). Circadian clock resetting in the mouse changes with age. *Age (Dordr)*, 31, 293–303. <https://doi.org/10.1007/s11357-009-9102-7>
- Biello, S. M., Bonsall, D. R., Atkinson, L. A., Molyneux, P. C., Harrington, M. E., & Lall, G. S. (2018). Alterations in glutamatergic signaling contribute to the decline of circadian photoentrainment in aged mice. *Neurobiology of Aging*, 66, 75–84.
- Bilbo, S., & Stevens, B. (2017) Microglia: The brain's first responders. *Cerebrum* 2017, 1–14. <https://doi.org/10.3389/fnagi.2019.00368>.
- Bonaconsa, M., Malpeli, G., Montaruli, A., Carandente, F., Grassi-Zucconi, G., & Bentivoglio, M. (2014). Differential modulation of clock gene expression in the suprachiasmatic nucleus, liver and heart of aged mice. *Experimental Gerontology*, 55, 70–79.
- Brancaccio, M., Edwards, M. D., Patton, A. P., Smyllie, N. J., Chesham, J. E., Maywood, E. S., & Hastings, M. H. (2019). Cell-autonomous clock of astrocytes drives circadian behavior in mammals. *Science*, 363, 187–192. <https://doi.org/10.1126/science.aat4104>
- Brancaccio, M., Patton, A. P., Chesham, J. E., Maywood, E. S., & Hastings, M. H. (2017). Astrocytes control circadian timekeeping in the suprachiasmatic nucleus via glutamatergic signaling. *Neuron*, 93, 1420–1435. <https://doi.org/10.1016/j.neuron.2017.02.030>
- Brandhorst, S., Choi, I. Y., Wei, M., Cheng, C. W., Sedrakyan, S., Navarrete, G., Dubeau, L., Yap, L. P., Park, R., Vinciguerra, M., Di Biase, S., Mirzaei, H., Mirisola, M. G., Childress, P., Ji, L., Groshen, S., Penna, F., Odetti, P., Perin, L., ... Longo, V. D. (2015). A periodic diet that mimics fasting promotes multi-system regeneration, enhanced cognitive performance, and healthspan. *Cell Metabolism*, 22, 86–99. <https://doi.org/10.1016/j.cmet.2015.05.012>
- Brown, A. J., Pendergast, J. S., & Yamazaki, S. (2019). Peripheral circadian oscillators. *Yale Journal of Biology and Medicine*, 92, 327–335.
- Brown, T. M., & Piggins, H. D. (2007). Electrophysiology of the suprachiasmatic circadian clock. *Progress in Neurobiology*, 82, 229–255.
- Buhr, E. D., & Takahashi, J. S. (2013). Molecular components of the mammalian circadian clock. In A. Kramer & M. Meroz (Eds.), *Circadian Clocks* (pp. 3–27). Springer.
- Buijink, M. R., Olde Engberink, A. H. O., Wit, C. B., Almog, A., Meijer, J. H., Rohling, J. H. T., & Michel, S. (2020). Aging affects the capacity of photoperiodic adaptation downstream from the central molecular clock. *Journal of Biological Rhythms*, 35, 167–179.
- Buijs, F. N., Guzmán-Ruiz, M., León-Mercado, L., Basualdo, M. C., Escobar, C., Kalsbeek, A., & Buijs, R. M. (2017). Suprachiasmatic nucleus interaction with the arcuate nucleus; Essential for organizing physiological rhythms. *eNeuro* 4, ENEURO.0028-0017.2017. <https://doi.org/10.1523/ENEURO.0028-17.2017>
- Buijs, R. M., Guzmán Ruiz, M. A., Méndez Hernández, R., & Rodríguez Cortés, B. (2019). The suprachiasmatic nucleus; a responsive clock regulating homeostasis by daily changing the setpoints of physiological parameters. *Autonomic Neuroscience*, 218, 43–50. <https://doi.org/10.1016/j.autneu.2019.02.001>
- Cai, A., & Wise, P. M. (1996). Age-related changes in light-induced jun-b and jun-d expression: Effects of transplantation of fetal tissue containing the suprachiasmatic nucleus. *Journal of Biological Rhythms*, 11, 284–290. <https://doi.org/10.1177/074873049601100401>
- Calvo-Rodríguez, M., Hernando-Pérez, E., López-Vázquez, S., Núñez, J., Villalobos, C., & Núñez, L. (2020). Remodeling of intracellular Ca^{2+} homeostasis in rat hippocampal neurons aged in vitro. *International Journal of Molecular Sciences*, 21, 1549.
- Cao, R., Lee, B., Cho, H.-Y., Saklayen, S., & Obrietan, K. (2008). Photic regulation of the mTOR signaling pathway in the suprachiasmatic circadian clock. *Molecular and Cellular Neurosciences*, 38, 312–324.
- Cardinali, D. P. (2019). Melatonin: Clinical perspectives in neurodegeneration. *Frontiers in Endocrinology*, 10, 480. <https://doi.org/10.3389/fendo.2019.00480>
- Castanon-Cervantes, O., Wu, M., Ehlen, J. C., Paul, K., Gamble, K. L., Johnson, R. L., Besing, R. C., Menaker, M., Gewirtz, A. T., & Davidson,

- A. J. (2010). Dysregulation of inflammatory responses by chronic circadian disruption. *The Journal of Immunology*, *185*, 5796–5805. <https://doi.org/10.4049/jimmunol.1001026>
- Castellani, G., & Schwartz, M. (2020). Immunological features of non-neuronal brain cells: Implications for Alzheimer's disease immunotherapy. *Trends in Immunology*, *41*, 794–804. <https://doi.org/10.1016/j.it.2020.07.005>
- Cayetanot, F., Bentivoglio, M., & Aujard, F. (2005). Arginine-vasopressin and vasointestinal polypeptide rhythms in the suprachiasmatic nucleus of the mouse lemur reveal aging-related alterations of circadian pacemaker neurons in a non-human primate. *European Journal of Neuroscience*, *22*, 902–910.
- Chang, H.-C., & Guarente, L. (2013). SIRT1 mediates central circadian control in the SCN by a mechanism that decays with aging. *Cell*, *153*, 1448–1460. <https://doi.org/10.1016/j.cell.2013.05.027>
- Chaudhury, D., & Colwell, C. S. (2002). Circadian modulation of learning and memory in fear-conditioned mice. *Behavioral Brain Research*, *133*, 95–108.
- Chaudhury, D., Loh, D. H., Dragich, J. M., Hagopian, A., & Colwell, C. S. (2008). Select cognitive deficits in Vasoactive Intestinal Peptide deficient mice. *BMC Neuroscience*, *9*, 63. <https://doi.org/10.1186/1471-2202-9-63>
- Chauhan, R., Chen, K. F., Kent, B. A., & Crowther, D. C. (2017). Central and peripheral circadian clocks and their role in Alzheimer's disease. *Disease Models & Mechanisms*, *10*, 1187–1199.
- Chee, C. A., Roozendaal, B., Swaab, D. F., Goudsmit, E., & Mirmiran, M. (1988). Vasoactive intestinal polypeptide neuron changes in the senile rat suprachiasmatic nucleus. *Neurobiology of Aging*, *9*, 307–312.
- Chen, Z., Yoo, S. H., & Takahashi, J. S. (2018). Development and therapeutic potential of small-molecule modulators of circadian systems. *Annual Review of Pharmacology and Toxicology*, *58*, 231–252.
- Clarke, L. E., Liddelow, S. A., Chakraborty, C., Münch, A. E., Heiman, M., & Barres, B. A. (2018). Normal aging induces A1-like astrocyte reactivity. *Proceedings of the National Academy of Sciences of the United States of America*, *115*, E1896–E1905. <https://doi.org/10.1073/pnas.1800165115>
- Cohen, H. Y., Miller, C., Bitterman, K. J., Wall, N. R., Hekking, B., Kessler, B., Howitz, K. T., Gorospe, M., de Cabo, R., & Sinclair, D. A. (2004). Calorie restriction promotes mammalian cell survival by inducing the SIRT1 deacetylase. *Science*, *305*, 390–392. <https://doi.org/10.1126/science.1099196>
- Colman, R. J., Anderson, R. M., Johnson, S. C., Kastman, E. K., Kosmatka, K. J., Beasley, T. M., Allison, D. B., Cruzen, C., Simmons, H. A., Kemnitz, J. W., & Weindruch, R. (2009). Caloric restriction delays disease onset and mortality in rhesus monkeys. *Science*, *325*, 201–204. <https://doi.org/10.1126/science.1173635>
- Colwell, C. S. (2000). Circadian modulation of calcium levels in cells in the suprachiasmatic nucleus. *European Journal of Neuroscience*, *12*, 571–576.
- Colwell, C. S. (2011). Linking neural activity and molecular oscillations in the SCN. *Nature Reviews Neuroscience*, *12*, 553–569.
- Curtis, A. M., Bellet, M. M., Sassone-Corsi, P., & O'Neill, L. A. (2014). Circadian clock proteins and immunity. *Immunity*, *40*, 178–186. <https://doi.org/10.1016/j.immuni.2014.02.002>
- Davidson, A. J., Sellix, M. T., Daniel, J., Yamazaki, S., Menaker, M., & Block, G. D. (2006). Chronic jet-lag increases mortality in aged mice. *Current Biology*, *16*, R914–916. <https://doi.org/10.1016/j.cub.2006.09.058>
- Davidson, A. J., Yamazaki, S., Arble, D. M., Menaker, M., & Block, G. D. (2008). Resetting of central and peripheral circadian oscillators in aged rats. *Neurobiology of Aging*, *29*, 471–477.
- De Butte, M., & Pappas, B. A. (2007). Pinealectomy causes hippocampal CA1 and CA3 cell loss: Reversal by melatonin supplementation. *Neurobiology of Aging*, *28*, 306–313.
- Deng, X. H., Bertini, G., Palomba, M., Xu, Y. Z., Bonaconsa, M., Nygard, M., & Bentivoglio, M. (2010). Glial transcripts and immune-challenged glia in the suprachiasmatic nucleus of young and aged mice. *Chronobiology International*, *27*, 742–767.
- Di Benedetto, S., Muller, L., Wenger, E., Duzel, S., & Pawelec, G. (2017). Contribution of neuroinflammation and immunity to brain aging and the mitigating effects of physical and cognitive interventions. *Neuroscience and Biobehavioral Reviews*, *75*, 114–128.
- Doi, M., Ishida, A., Miyake, A., Sato, M., Komatsu, R., Yamazaki, F., Kimura, I., Tsuchiya, S., Kori, H., Seo, K., Yamaguchi, Y., Matsuo, M., Fustin, J.-M., Tanaka, R., Santo, Y., Yamada, H., Takahashi, Y., Araki, M., Nakao, K., ... Okamura, H. (2011). Circadian regulation of intracellular G-protein signalling mediates intercellular synchrony and rhythmicity in the suprachiasmatic nucleus. *Nature Communications*, *2*, (1). <http://dx.doi.org/10.1038/ncomms1316>
- Dominguez-Rodriguez, A., Abreu-Gonzalez, P., Sanchez-Sanchez, J. J., Kaski, J. C., & Reiter, R. J. (2010). Melatonin and circadian biology in human cardiovascular disease. *Journal of Pineal Research*, *49*, 14–22. <https://doi.org/10.1111/j.1600-079X.2010.00773.x>
- Duffy, J. F., Zeitzer, J. M., Rimmer, D. W., Klerman, E. B., Dijk, D.-J., & Czeisler, C. A. (2002). Peak of circadian melatonin rhythm occurs later within the sleep of older subjects. *American Journal of Physiology. Endocrinology and Metabolism*, *282*, E297–E303. <https://doi.org/10.1152/ajpendo.00268.2001>
- Duncan, M. J. (2020). Interacting influences of aging and Alzheimer's disease on circadian rhythms. *European Journal of Neuroscience*, *51*, 310–325. <https://doi.org/10.1111/ejn.14358>
- Duncan, M. J., Herron, J. M., & Hill, S. A. (2001). Aging selectively suppresses vasoactive intestinal peptide messenger RNA expression in the suprachiasmatic nucleus of the Syrian hamster. *Molecular Brain Research*, *87*, 196–203.
- Duncan, M. J., Prochot, J. R., Cook, D. H., Tyler Smith, J., & Franklin, K. M. (2013). Influence of aging on Bmal1 and Per2 expression in extra-SCN oscillators in hamster brain. *Brain Research*, *1491*, 44–53. <https://doi.org/10.1016/j.brainres.2012.11.008>
- Dupont Rocher, S., Bessot, N., Sesboué, B., Bulla, J., & Davenne, D. (2016). Circadian characteristics of older adults and aerobic capacity. *The Journals of Gerontology. Series A, Biological Sciences and Medical Sciences*, *71*, 817–822.
- Engelberth, R. C. G. J., Silva, K. D. D. A., Fiuza, F. P., Soares, J. G., Costa, M. S. M. O., de Melo Lima, R. R., do Nascimento, E. S., dos Santos, J. R., Cavalcanti, J. R. L. P., & Cavalcante, J. S. (2017). Retinal, NPY- and 5ht- inputs to the aged suprachiasmatic nucleus in common marmosets (*Callithrix jacchus*). *Neuroscience Research*, *121*, 54–59.
- Enoki, R., Oda, Y., Mieda, M., Ono, D., Honma, S., & Honma, K. I. (2017). Synchronous circadian voltage rhythms with asynchronous calcium rhythms in the suprachiasmatic nucleus. *Proceedings of the National Academy of Sciences of the United States of America*, *114*, E2476–E2485. <https://doi.org/10.1073/pnas.1616815114>
- Esquiva, G., Lax, P., Pérez-Santónja, J. J., García-Fernández, J. M., & Cuenca, N. (2017). Loss of melanopsin-expressing ganglion cell subtypes and dendritic degeneration in the aging human retina. *Frontiers in Aging Neuroscience*, *9*. <https://doi.org/10.3389/fnagi.2017.00079>
- Evans, J. A. (2016). Collective timekeeping among cells of the master circadian clock. *Journal of Endocrinology*, *230*, R27–49. <https://doi.org/10.1530/JOE-16-0054>
- Fan, X., Wheatley, E. G., & Villeda, S. A. (2017). Mechanisms of hippocampal aging and the potential for rejuvenation. *Annual Review of Neuroscience*, *40*, 251–272.
- Farajnia, S., Deboer, T., Rohling, J. H., Meijer, J. H., & Michel, S. (2014). Aging of the suprachiasmatic clock. *Neuroscientist*, *20*, 44–55. <https://doi.org/10.1177/1073858413498936>
- Farajnia, S., Meijer, J. H., & Michel, S. (2015). Age-related changes in large-conductance calcium-activated potassium channels in mammalian circadian clock neurons. *Neurobiology of Aging*, *36*, 2176–2183.

- Farajnia, S., Michel, S., Deboer, T., vanderLeest, H. T., Houben, T., Rohling, J. H. T., Ramkisoensing, A., Yasenkov, R., & Meijer, J. H. (2012). Evidence for neuronal desynchrony in the aged suprachiasmatic nucleus clock. *Journal of Neuroscience*, *32*, 5891–5899. <https://doi.org/10.1523/JNEUROSCI.0469-12.2012>
- Flurkey, K., Curren, M. J., & Harrison, D. E. (2007). Chapter 20 - mouse models in aging research. In J. G. Fox, M. T. Davisson, F. W. Quimby, S. W. Barthold, C. E. Newcomer, & A. L. Smith (Eds.), *The Mouse in Biomedical Research*, 2nd ed. (pp. 637–672). Academic Press.
- Fonken, L. K., Frank, M. G., Kitt, M. M., Barrientos, R. M., Watkins, L. R., & Maier, S. F. (2015). Microglia inflammatory responses are controlled by an intrinsic circadian clock. *Brain, Behavior, and Immunity*, *45*, 171–179.
- Fonken, L. K., Kitt, M. M., Gaudet, A. D., Barrientos, R. M., Watkins, L. R., & Maier, S. F. (2016). Diminished circadian rhythms in hippocampal microglia may contribute to age-related neuroinflammatory sensitization. *Neurobiology of Aging*, *47*, 102–112.
- Fonseca Costa, S. S., & Ripperger, J. A. (2015). Impact of the circadian clock on the aging process. *Frontiers in Neurology*, *6*, 43.
- Gerhold, L. M., Rosewell, K. L., & Wise, P. M. (2005). Suppression of vasoactive intestinal polypeptide in the suprachiasmatic nucleus leads to aging-like alterations in cAMP rhythms and activation of gonadotropin-releasing hormone neurons. *Journal of Neuroscience*, *25*, 62–67. <https://doi.org/10.1523/JNEUROSCI.3598-04.2005>
- Gillette, M. U., & Tischkau, S. A. (1999). Suprachiasmatic nucleus: The brain's circadian clock. *Recent Progress in Hormone Research*, *54*, 33–58; discussion 58–39.
- Golombek, D. A., & Rosenstein, R. E. (2010). Physiology of circadian entrainment. *Physiological Reviews*, *90*, 1063–1102.
- Granados-Fuentes, D., Norris, A. J., Carrasquillo, Y., Nerbonne, J. M., & Herzog, E. D. (2012). I_A channels encoded by Kv1.4 and Kv4.2 regulate neuronal firing in the suprachiasmatic nucleus and circadian rhythms in locomotor activity. *Journal of Neuroscience*, *32*, 10045–10052. <https://doi.org/10.1523/JNEUROSCI.0174-12.2012>
- Gu, C., Coomans, C. P., Hu, K., Scheer, F. A. J. L., Stanley, H. E., & Meijer, J. H. (2015). Lack of exercise leads to significant and reversible loss of scale invariance in both aged and young mice. *Proceedings of the National Academy of Sciences of the United States of America*, *112*, 2320–2324. <https://doi.org/10.1073/pnas.1424706112>
- Guilding, C., Hughes, A. T. L., Brown, T. M., Namvar, S., & Piggins, H. D. (2009). A riot of rhythms: Neuronal and glial circadian oscillators in the mediobasal hypothalamus. *Molecular Brain*, *2*, 28.
- Hardeland, R. (2018). Melatonin and inflammation—Story of a double-edged blade. *Journal of Pineal Research*, *65*, e12525. <https://doi.org/10.1111/jpi.12525>
- Harrison, D. E., Strong, R., Sharp, Z. D., Nelson, J. F., Astle, C. M., Flurkey, K., Nadon, N. L., Wilkinson, J. E., Frenkel, K., Carter, C. S., Pahor, M., Javors, M. A., Fernandez, E., & Miller, R. A. (2009). Rapamycin fed late in life extends lifespan in genetically heterogeneous mice. *Nature*, *460*, 392–395. <https://doi.org/10.1038/nature08221>
- Harvey, J. R. M., Plante, A. E., & Meredith, A. L. (2020). Ion channels controlling circadian rhythms in suprachiasmatic nucleus excitability. *Physiological Reviews*, *100*, 1415–1454.
- Hatori, M., Vollmers, C., Zarrinpar, A., DiTacchio, L., Bushong, E. A., Gill, S., Leblanc, M., Chaix, A., Joens, M., Fitzpatrick, J. A. J., Ellisman, M. H., & Panda, S. (2012). Time-restricted feeding without reducing caloric intake prevents metabolic diseases in mice fed a high-fat diet. *Cell Metabolism*, *15*, 848–860. <https://doi.org/10.1016/j.cmet.2012.04.019>
- Hattar, S., Liao, H. W., Takao, M., Berson, D. M., & Yau, K. W. (2002). Melanopsin-containing retinal ganglion cells: Architecture, projections, and intrinsic photosensitivity. *Science*, *295*, 1065. <https://doi.org/10.1126/science.1069609>
- He, B., Nohara, K., Park, N., Park, Y.-S., Guillory, B., Zhao, Z., Garcia, J. M., Koike, N., Lee, C. C., Takahashi, J. S., Yoo, S.-H., & Chen, Z. (2016). The small molecule nobletin targets the molecular oscillator to enhance circadian rhythms and protect against metabolic syndrome. *Cell Metabolism*, *23*, 610–621. <https://doi.org/10.1016/j.cmet.2016.03.007>
- Held, N. M., Buijink, M. R., Elfrink, H. L., Kooijman, S., Janssens, G. E., Vaz, F. M., Michel, S., Houtkooper, R. H., & van Weeghel, M. (2020). Aging selectively dampens oscillation of lipid abundance in white and brown adipose tissue. *bioRxiv*, 2020.2009.2004.283226. <https://doi.org/10.1101/2020.09.04.283226>
- Houtkooper, R. H., Pirinen, E., & Auwerx, J. (2012). Sirtuins as regulators of metabolism and healthspan. *Nature Reviews Molecular Cell Biology*, *13*, 225–238.
- Huang, G., Zhang, Y., Shan, Y., Yang, S., Chelliah, Y., Wang, H., & Takahashi, J. S. (2016). Circadian oscillations of nadh redox state using a heterologous metabolic sensor in mammalian Cells. *Journal of Biological Chemistry*, *291*, 23906–23914.
- Hughes, A. T., & Piggins, H. D. (2012). Feedback actions of locomotor activity to the circadian clock. *Progress in Brain Research*, *199*, 305–336.
- Hurd, M. W., & Ralph, M. R. (1998). The significance of circadian organization for longevity in the golden hamster. *Journal of Biological Rhythms*, *13*, 430–436. <https://doi.org/10.1177/074873098129000255>
- Iguchi, H., Kato, K.-I., & Ibayashi, H. (1982). Age-dependent reduction in serum melatonin concentrations in healthy human subjects. *Journal of Clinical Endocrinology and Metabolism*, *55*, 27–29.
- Ikeda, M., Sugiyama, T., Wallace, C. S., Gompf, H. S., Yoshioka, T., Miyawaki, A., & Allen, C. N. (2003). Circadian dynamics of cytosolic and nuclear Ca^{2+} in single suprachiasmatic nucleus neurons. *Neuron*, *38*, 253–263. [https://doi.org/10.1016/S0896-6273\(03\)00164-8](https://doi.org/10.1016/S0896-6273(03)00164-8)
- Ishii, T., Warabi, E., & Mann, G. E. (2019). Circadian control of BDNF-mediated Nrf2 activation in astrocytes protects dopaminergic neurons from ferroptosis. *Free Radical Biology and Medicine*, *133*, 169–178.
- Ishunina, T. A., & Swaab, D. F. (2002). Neurohypophyseal peptides in aging and Alzheimer's disease. *Ageing Research Reviews*, *1*, 537–558. [https://doi.org/10.1016/S1568-1637\(02\)00013-2](https://doi.org/10.1016/S1568-1637(02)00013-2)
- Itri, J. N., Michel, S., Vansteensel, M. J., Meijer, J. H., & Colwell, C. S. (2005). Fast delayed rectifier potassium current is required for circadian neural activity. *Nature Neuroscience*, *8*, 650–656.
- Itri, J. N., Vosko, A. M., Schroeder, A., Dragich, J. M., Michel, S., & Colwell, C. S. (2010). Circadian regulation of A-type potassium currents in the suprachiasmatic nucleus. *Journal of Neurophysiology*, *103*, 632–640. <https://doi.org/10.1152/jn.00670.2009>
- Jengeleski, C. A., Powers, R. E., O'Connor, D. T., & Price, D. L. (1989). Noradrenergic innervation of human pineal gland: Abnormalities in aging and Alzheimer's disease. *Brain Research*, *481*, 378–382. [https://doi.org/10.1016/0006-8993\(89\)90818-4](https://doi.org/10.1016/0006-8993(89)90818-4)
- Jones, J. R., Simon, T., Lones, L., & Herzog, E. D. (2018). SCN VIP neurons are essential for normal light-mediated resetting of the circadian system. *Journal of Neuroscience*, *38*, 7986–7995. <https://doi.org/10.1523/JNEUROSCI.1322-18.2018>
- Ju, Y.-E., Ooms, S. J., Sutphen, C., Macauley, S. L., Zangrilli, M. A., Jerome, G., Fagan, A. M., Mignot, E., Zempel, J. M., Claassen, J. A. H. R., & Holtzman, D. M. (2017). Slow wave sleep disruption increases cerebrospinal fluid amyloid- β levels. *Brain*, *140*, 2104–2111. <https://doi.org/10.1093/brain/awx148>
- Kaeberlein, M., McVey, M., & Guarente, L. (1999). The SIR2/3/4 complex and SIR2 alone promote longevity in *Saccharomyces cerevisiae* by two different mechanisms. *Genes & Development*, *13*, 2570–2580. <https://doi.org/10.1101/gad.13.19.2570>
- Kaeberlein, M., Powers, R. W., Steffen, K. K., Westman, E. A., Hu, D., Dang, N., Kerr, E. O., Kirkland, K. T., & Kennedy, B. K. (2005). Regulation of yeast replicative life span by TOR and Sch9 in response to nutrients. *Science*, *310*, 1193–1196. <https://doi.org/10.1126/science.1115535>

- Kalamatianos, T., Kalló, I., & Coen, C. W. (2004). Ageing and the diurnal expression of the mRNAs for vasopressin and for the V1a and V1b vasopressin receptors in the suprachiasmatic nucleus of male rats. *Journal of Neuroendocrinology*, *16*, 493–501. <https://doi.org/10.1111/j.1365-2826.2004.01196.x>
- Kalló, I., Kalamatianos, T., Piggins, H. D., & Coen, C. W. (2004). Ageing and the diurnal expression of mRNAs for vasoactive intestinal peptide and for the VPAC2 and PAC1 receptors in the suprachiasmatic nucleus of male rats. *Journal of Neuroendocrinology*, *16*, 758–766. <https://doi.org/10.1111/j.1365-2826.2004.01232.x>
- Kalsbeek, A., & Buijs, R. M. (2002). Output pathways of the mammalian suprachiasmatic nucleus: Coding circadian time by transmitter selection and specific targeting. *Cell and Tissue Research*, *309*, 109–118. <https://doi.org/10.1007/s00441-002-0577-0>
- Kalsbeek, A., Teclemariam-Mesbah, R., & Péavet, P. (1993). Efferent projections of the suprachiasmatic nucleus in the golden hamster (*Mesocricetus auratus*). *The Journal of Comparative Neurology*, *332*, 293–314.
- Kalsbeek, A., Yi, C.-X., Cailotto, C., la Fleur, S. E., Fliers, E., & Buijs, R. M. (2011). Mammalian clock output mechanisms. *Essays in Biochemistry*, *49*, 137–151. <https://doi.org/10.1042/bse0490137>
- Karamitri, A., & Jockers, R. (2019). Melatonin in type 2 diabetes mellitus and obesity. *Nature Reviews Endocrinology*, *15*, 105–125.
- Kawakami, F., Okamura, H., Tamada, Y., Maebayashi, Y., Fukui, K., & Ibata, Y. (1997). Loss of day-night differences in VIP mRNA levels in the suprachiasmatic nucleus of aged rats. *Neuroscience Letters*, *222*, 99–102.
- Kent, J., & Meredith, A. L. (2008). BK channels regulate spontaneous action potential rhythmicity in the suprachiasmatic nucleus. *PLoS One*, *3*, e3884. <https://doi.org/10.1371/journal.pone.0003884>
- Kessel, L., Lundeman, J. H., Herbst, K., Andersen, T. V., & Larsen, M. (2010). Age-related changes in the transmission properties of the human lens and their relevance to circadian entrainment. *Journal of Cataract and Refractive Surgery*, *36*, 308–312.
- Khapre, R. V., Kondratova, A. A., Patel, S., Dubrovsky, Y., Wrobel, M., Antoch, M. P., & Kondratov, R. V. (2014). BMAL1-dependent regulation of the mTOR signaling pathway delays aging. *Aging*, *6*, 48–57. <https://doi.org/10.18632/aging.100633>
- Kim, K., & Choe, H. K. (2019). Role of hypothalamus in aging and its underlying cellular mechanisms. *Mechanisms of Ageing and Development*, *177*, 74–79.
- Kolker, D. E., Fukuyama, H., Huang, D. S., Takahashi, J. S., Horton, T. H., & Turek, F. W. (2003). Aging alters circadian and light-induced expression of clock genes in golden hamsters. *Journal of Biological Rhythms*, *18*, 159–169. <https://doi.org/10.1177/0748730403251802>
- Kondratova, A. A., & Kondratov, R. V. (2012). The circadian clock and pathology of the ageing brain. *Nature Reviews Neuroscience*, *13*, 325–335.
- Krajnak, K., Kashon, M. L., Rosewell, K. L., & Wise, P. M. (1998). Aging alters the rhythmic expression of vasoactive intestinal polypeptide mRNA but not arginine vasopressin mRNA in the suprachiasmatic nuclei of female rats. *Journal of Neuroscience*, *18*, 4767–4774. <https://doi.org/10.1523/JNEUROSCI.18-12-04767.1998>
- Kudo, T., Block, G. D., & Colwell, C. S. (2015). The circadian clock gene *Period1* connects the molecular clock to neural activity in the suprachiasmatic nucleus. *ASN Neuro*, *7*, (6). <http://dx.doi.org/10.1177/1759091415610761>
- Labrecque, N., & Cermakian, N. (2015). Circadian clocks in the immune system. *Journal of Biological Rhythms*, *30*, 277–290. <https://doi.org/10.1177/0748730415577723>
- Lananna, B. V., & Musiek, E. S. (2020). The wrinkling of time: Aging, inflammation, oxidative stress, and the circadian clock in neurodegeneration. *Neurobiology of Diseases*, *139*, 104832. <https://doi.org/10.1016/j.nbd.2020.104832>
- Lax, P., Ortuño-Lizarán, I., Maneu, V., Vidal-Sanz, M., & Cuenca, N. (2019). Photosensitive melanopsin-containing retinal ganglion cells in health and disease: Implications for circadian rhythms. *International Journal of Molecular Sciences*, *20*, 3164.
- Leise, T. L., Harrington, M. E., Molyneux, P. C., Song, I., Queenan, H., Zimmerman, E., Lall, G. S., & Biello, S. M. (2013). Voluntary exercise can strengthen the circadian system in aged mice. *Age (Dordr)*, *35*, 2137–2152. <https://doi.org/10.1007/s11357-012-9502-y>
- Leng, Y., Musiek, E. S., Hu, K., Cappuccio, F. P., & Yaffe, K. (2019). Association between circadian rhythms and neurodegenerative diseases. *The Lancet Neurology*, *18*, 307–318. [https://doi.org/10.1016/S1474-4422\(18\)30461-7](https://doi.org/10.1016/S1474-4422(18)30461-7)
- Leproult, R., Holmbäck, U., & Van Cauter, E. (2014). Circadian misalignment augments markers of insulin resistance and inflammation, independently of sleep loss. *Diabetes*, *63*, 1860–1869. <https://doi.org/10.2337/db13-1546>
- Lewy, A. J., Tetsuo, M., Markey, S. P., Goodwin, F. K., & Kopin, I. J. (1980). Pinealectomy abolishes plasma melatonin in the rat. *Journal of Clinical Endocrinology and Metabolism*, *50*, 204–205.
- Lin, S.-J., Kaeberlein, M., Andalis, A. A., Sturtz, L. A., Defossez, P.-A., Culotta, V. C., Fink, G. R., & Guarente, L. (2002). Calorie restriction extends *Saccharomyces cerevisiae* lifespan by increasing respiration. *Nature*, *418*, 344–348. <https://doi.org/10.1038/nature00829>
- Longo, V. D., & Panda, S. (2016). Fasting, circadian rhythms, and time-restricted feeding in healthy lifespan. *Cell Metabolism*, *23*, 1048–1059. <https://doi.org/10.1016/j.cmet.2016.06.001>
- Lopes, C., DeLyra, J. L., Markus, R. P., & Mariano, M. (1997). Circadian rhythm in experimental granulomatous inflammation is modulated by melatonin. *Journal of Pineal Research*, *23*, 72–78. <https://doi.org/10.1111/j.1600-079X.1997.tb00338.x>
- López-Otín, C., Blasco, M. A., Partridge, L., Serrano, M., & Kroemer, G. (2013). The hallmarks of aging. *Cell*, *153*, 1194–1217. <https://doi.org/10.1016/j.cell.2013.05.039>
- Lucassen, P. J., Hofman, M. A., & Swaab, D. F. (1995). Increased light intensity prevents the age related loss of vasopressin-expressing neurons in the rat suprachiasmatic nucleus. *Brain Research*, *693*, 261–266. [https://doi.org/10.1016/0006-8993\(95\)00933-H](https://doi.org/10.1016/0006-8993(95)00933-H)
- Lupi, D., Semo, M. A., & Foster, R. G. (2012). Impact of age and retinal degeneration on the light input to circadian brain structures. *Neurobiology of Aging*, *33*, 383–392.
- Madeira, M. D., Sousa, N., Santer, R. M., Paula-Barbosa, M. M., & Gundersen, H. J. (1995). Age and sex do not affect the volume, cell numbers, or cell size of the suprachiasmatic nucleus of the rat: An unbiased stereological study. *The Journal of Comparative Neurology*, *361*, 585–601. <https://doi.org/10.1002/cne.903610404>
- Markus, R. P., Fernandes, P. A., Kinker, G. S., da Silveira Cruz-Machado, S., & Marçola, M. (2018). Immune-pineal axis – acute inflammatory responses coordinate melatonin synthesis by pinealocytes and phagocytes. *British Journal of Pharmacology*, *175*, 3239–3250.
- Mattson, M. P., & Arumugam, T. V. (2018). Hallmarks of brain aging: Adaptive and pathological modification by metabolic states. *Cell Metabolism*, *27*, 1176–1199. <https://doi.org/10.1016/j.cmet.2018.05.011>
- McKee, C. A., Lananna, B. V., & Musiek, E. S. (2020). Circadian regulation of astrocyte function: Implications for Alzheimer's disease. *Cellular and Molecular Life Sciences*, *77*, 1049–1058. <https://doi.org/10.1007/s00018-019-03314-y>
- Meredith, A. L., Wiler, S. W., Miller, B. H., Takahashi, J. S., Fodor, A. A., Ruby, N. F., & Aldrich, R. W. (2006). BK calcium-activated potassium channels regulate circadian behavioral rhythms and pacemaker output. *Nature Neuroscience*, *9*, 1041–1049.
- Michel, S., & Meijer, J. H. (2020). From clock to functional pacemaker. *European Journal of Neuroscience*, *51*, 482–493.
- Miller, S., & Hirota, T. (2020). Pharmacological interventions to circadian clocks and their molecular bases. *Journal of Molecular Biology*, *432*, 3498–3514.

- Mukherji, A., Kobiita, A., Damara, M., Misra, N., Meziane, H., Champy, M. F., & Chambon, P. (2015). Shifting eating to the circadian rest phase misaligns the peripheral clocks with the master SCN clock and leads to a metabolic syndrome. *Proceedings of the National Academy of Sciences of the United States of America*, *112*, E6691–6698. <https://doi.org/10.1073/pnas.1519807112>
- Musiek, E. S., Bhamasani, M., Zangrilli, M. A., Morris, J. C., Holtzman, D. M., & Ju, Y. S. (2018). Circadian rest-activity pattern changes in aging and preclinical alzheimer disease. *Journal of the American Medical Association Neurology*, *75*, 582–590. <https://doi.org/10.1001/jaman.euro.2017.4719>
- Musiek, E. S., & Holtzman, D. M. (2016). Mechanisms linking circadian clocks, sleep, and neurodegeneration. *Science*, *354*, 1004–1008. <https://doi.org/10.1126/science.aah4968>
- Musiek, E. S., Lim, M. M., Yang, G., Bauer, A. Q., Qi, L., Lee, Y., Roh, J. H., Ortiz-Gonzalez, X., Dearborn, J. T., Culver, J. P., Herzog, E. D., Hogenesch, J. B., Wozniak, D. F., Dikranian, K., Giasson, B. I., Weaver, D. R., Holtzman, D. M., & FitzGerald, G. A. (2013). Circadian clock proteins regulate neuronal redox homeostasis and neurodegeneration. *Journal of Clinical Investigation*, *123*, 5389–5400. <https://doi.org/10.1172/JCI70317>
- Najjar, R. P., Chiquet, C., Teikari, P., Cornut, P.-L., Claustrat, B., Denis, P., Cooper, H. M., & Gronfier, C. (2014). Aging of non-visual spectral sensitivity to light in humans: Compensatory mechanisms? *PLoS One*, *9*, e85837. <https://doi.org/10.1371/journal.pone.0085837>
- Nakamura, T. J., Michel, S., Block, G. D., & Colwell, C. S. (2012). Neural circuits underlying circadian oscillations in mammals: Clocks in a dish. In K. Ballanyi (Ed.), *Isolated Central Nervous System Circuits* (pp. 183–210). Humana Press.
- Nakamura, T. J., Nakamura, W., Tokuda, I. T., Ishikawa, T., Kudo, T., Colwell, C. S., & Block, G. D. (2015). Age-related changes in the circadian system unmasked by constant conditions. *eneuro*, *2*(4). <https://doi.org/10.1523/ENEURO.0064-15.2015>
- Nakamura, T. J., Nakamura, W., Yamazaki, S., Kudo, T., Cutler, T., Colwell, C. S., & Block, G. D. (2011). Age-related decline in circadian output. *Journal of Neuroscience*, *31*, 10201–10205. <https://doi.org/10.1523/JNEUROSCI.0451-11.2011>
- Nguyen, K. D., Fentress, S. J., Qiu, Y., Yun, K., Cox, J. S., & Chawla, A. (2013). Circadian gene Bmal1 regulates diurnal oscillations of Ly6C(hi) inflammatory monocytes. *Science*, *341*, 1483–1488. <https://doi.org/10.1126/science.1240636>
- Nissen, J. C. (2017). Microglial function across the spectrum of age and gender. *International Journal of Molecular Sciences*, *18*, 561.
- Noguchi, T., Leise, T. L., Kingsbury, N. J., Diemer, T., Wang, L. L., Henson, M. A., & Welsh, D. K. (2017). Calcium circadian rhythmicity in the suprachiasmatic nucleus: Cell autonomy and network modulation. *eneuro*, *4*(4). <https://doi.org/10.1523/ENEURO.0160-17.2017>
- Nohara, K., Mallampalli, V., Nemkov, T., Wirianto, M., Yang, J., Ye, Y., Sun, Y., Han, L., Esser, K. A., Mileyskova, E., D'Alessandro, A., Green, C. B., Takahashi, J. S., Dowhan, W., Yo, S.-H., & Chen, Z. (2019). Nobiletin fortifies mitochondrial respiration in skeletal muscle to promote healthy aging against metabolic challenge. *Nature Communications*, *10*, 3923.
- Nygaard, M., Hill, R. H., Wikstrom, M. A., & Kristensson, K. (2005). Age-related changes in electrophysiological properties of the mouse suprachiasmatic nucleus in vitro. *Brain Research Bulletin*, *65*, 149–154.
- O'Neill, J. S., Maywood, E. S., Chesham, J. E., Takahashi, J. S., & Hastings, M. H. (2008). cAMP-dependent signaling as a core component of the mammalian circadian pacemaker. *Science*, *320*, 949–953. <https://doi.org/10.1126/science.1152506>
- O'Neill, J. S., & Reddy, A. B. (2012). The essential role of cAMP/Ca²⁺ signalling in mammalian circadian timekeeping. *Biochemical Society Transactions*, *40*, 44–50.
- Palomba, M., Nygård, M., Florenzano, F., Bertini, G., Kristensson, K., & Bentivoglio, M. (2008). Decline of the presynaptic network, including gabaergic terminals, in the aging suprachiasmatic nucleus of the mouse. *Journal of Biological Rhythms*, *23*, 220–231. <https://doi.org/10.1177/0748730408316998>
- Pang, S. F., & Ralph, C. L. (1975). Pineal and serum melatonin at mid-day and midnight following pinealectomy or castration in male rats. *Journal of Experimental Zoology*, *193*, 275–280.
- Park, J. S., Cederroth, C. R., Basinou, V., Sweetapple, L., Buijink, R., Lundkvist, G. B., Michel, S., & Canlon, B. (2017). Differential phase arrangement of cellular clocks along the tonotopic axis of the mouse cochlea ex vivo. *Current Biology*, *27*(2623–2629), e2622. <https://doi.org/10.1016/j.cub.2017.07.019>
- Paul, J. R., Davis, J. A., Goode, L. K., Becker, B. K., Fusilier, A., Meador-Woodruff, A., & Gamble, K. L. (2020). Circadian regulation of membrane physiology in neural oscillators throughout the brain. *European Journal of Neuroscience*, *51*, 109–138. <https://doi.org/10.1111/ejn.14343>
- Paul, S., Hanna, L., Harding, C., Hayter, E. A., Walmsley, L., Bechtold, D. A., & Brown, T. M. (2020). Output from VIP cells of the mammalian central clock regulates daily physiological rhythms. *Nature Communications*, *11*, 1453. <https://doi.org/10.1038/s41467-020-15277-x>
- Pelham, R. W. (1975). A serum melatonin rhythm in chickens and its abolition by pinealectomy. *Endocrinology*, *96*, 543–546.
- Pereira, P. A., Cardoso, A., & Paula-Barbosa, M. M. (2005). Nerve growth factor restores the expression of vasopressin and vasoactive intestinal polypeptide in the suprachiasmatic nucleus of aged rats. *Brain Research*, *1048*, 123–130. <https://doi.org/10.1016/j.brainres.2005.04.066>
- Pfeffer, M., Korf, H. W., & Wicht, H. (2017). The role of the melatoninergic system in light-entrained behavior of mice. *International Journal of Molecular Sciences*, *18*, 530.
- Pfeffer, M., Korf, H.-W., & Wicht, H. (2018). Synchronizing effects of melatonin on diurnal and circadian rhythms. *General and Comparative Endocrinology*, *258*, 215–221.
- Pierpaoli, W., & Bulian, D. (2005). The pineal aging and death program: Life prolongation in pre-aging pinealectomized mice. *Annals of the New York Academy of Sciences*, *1057*, 133–144.
- Pierpaoli, W., & Regelson, W. (1994). Pineal control of aging: Effect of melatonin and pineal grafting on aging mice. *Proceedings of the National Academy of Sciences of the United States of America*, *91*, 787–791. <https://doi.org/10.1073/pnas.91.2.787>
- Pitts, G. R., Ohta, H., & McMahon, D. G. (2006). Daily rhythmicity of large-conductance Ca²⁺-activated K⁺ currents in suprachiasmatic nucleus neurons. *Brain Research*, *1071*, 54–62. <https://doi.org/10.1016/j.brainres.2005.11.078>
- Polidarova, L., Sladek, M., Novosadova, Z., & Sumova, A. (2017). Aging does not compromise in vitro oscillation of the suprachiasmatic nuclei but makes it more vulnerable to constant light. *Chronobiology International*, *34*, 105–117.
- Prolo, L. M., Takahashi, J. S., & Herzog, E. D. (2005). Circadian rhythm generation and entrainment in astrocytes. *Journal of Neuroscience*, *25*, 404–408. <https://doi.org/10.1523/JNEUROSCI.4133-04.2005>
- Ramanathan, C., Kathale, N. D., Liu, D., Lee, C., Freeman, D. A., Hogenesch, J. B., Cao, R., & Liu, A. C. (2018). mTOR signaling regulates central and peripheral circadian clock function. *PLoS Genetics*, *14*, e1007369. <https://doi.org/10.1371/journal.pgen.1007369>
- Ramirez-Rodríguez, G., Klempin, F., Babu, H., Benítez-King, G., & Kempermann, G. (2009). Melatonin modulates cell survival of new neurons in the hippocampus of adult mice. *Neuropsychopharmacology*, *34*, 2180–2191. <https://doi.org/10.1038/npp.2009.46>
- Ramsey, K. M., Yoshino, J., Brace, C. S., Abrassart, D., Kobayashi, Y., Marche, B., Hong, H.-K., Chong, J. L., Buhr, E. D., Lee, C., Takahashi, J. S., Imai, S.-I., & Bass, J. (2009). Circadian clock feedback cycle through NAMPT-mediated NAD⁺ biosynthesis. *Science*, *324*, 651–654. <https://doi.org/10.1126/science.1171641>



- Riemersma-van der Lek, R. F., Swaab, D. F., Twisk, J., Hol, E. M., Hoogendijk, W. J., & Van Someren, E. J. (2008). Effect of bright light and melatonin on cognitive and noncognitive function in elderly residents of group care facilities: A randomized controlled trial. *Journal of the American Medical Association*, *299*, 2642–2655. <https://doi.org/10.1001/jama.299.22.2642>
- Roberts, D. E., Killiany, R. J., & Rosene, D. L. (2012). Neuron numbers in the hypothalamus of the normal aging rhesus monkey: Stability across the adult lifespan and between the sexes. *The Journal of Comparative Neurology*, *520*, 1181–1197. <https://doi.org/10.1002/cne.22761>
- Rodríguez, M. I., Escames, G., López, L. C., López, A., García, J. A., Ortiz, F., & Acuña-Castroviejo, D. (2007). Chronic melatonin treatment reduces the age-dependent inflammatory process in senescence-accelerated mice. *Journal of Pineal Research*, *42*, 272–279. <https://doi.org/10.1111/j.1600-079X.2006.00416.x>
- Rodríguez-Arellano, J. J., Parpura, V., Zorec, R., & Verkhratsky, A. (2016). Astrocytes in physiological aging and Alzheimer's disease. *Neuroscience*, *323*, 170–182. <https://doi.org/10.1016/j.neurosci.2015.01.007>
- Roosendaal, B., van Gool, W. A., Swaab, D. F., Hoogendijk, J. E., & Mirmiran, M. (1987). Changes in vasopressin cells of the rat supra-chiasmatic nucleus with aging. *Brain Research*, *409*, 259–264. [https://doi.org/10.1016/0006-8993\(87\)90710-4](https://doi.org/10.1016/0006-8993(87)90710-4)
- Roy, U., Heredia-Muñoz, M. T., Stute, L., Höfling, C., Matysik, J., Meijer, J. H., Roßner, S., & Alia, A. (2019). Degeneration of the suprachiasmatic nucleus in an Alzheimer's disease mouse model monitored by in vivo magnetic resonance relaxation measurements and immunohistochemistry. *Journal of Alzheimer's Disease*, *69*, 363–375. <https://doi.org/10.3233/JAD-190037>
- Ruben, M. D., Wu, G., Smith, D. F., Schmidt, R. E., Francey, L. J., Lee, Y. Y., Anafi, R. C., & Hogenesch, J. B. (2018). A database of tissue-specific rhythmically expressed human genes has potential applications in circadian medicine. *Science Translational Medicine*, *10*, eaat8806. <https://doi.org/10.1126/scitranslmed.aat8806>
- Rubiño, J. A., Gamundí, A., Akaair, M., Canellas, F., Rial, R., & Nicolau, M. C. (2020). Bright light therapy and circadian cycles in institutionalized elders. *Frontiers in Neuroscience*, *14*, 359.
- Ruby, N. F., Edgar, D. M., Heller, H. C., & Miller, J. D. (1998). The aged suprachiasmatic nucleus is phase-shifted by cAMP in vitro. *Brain Research*, *779*, 338–341. [https://doi.org/10.1016/S0006-8993\(97\)01236-5](https://doi.org/10.1016/S0006-8993(97)01236-5)
- Salminen, A., Ojala, J., Kaarniranta, K., Haapasalo, A., Hiltunen, M., & Soininen, H. (2011). Astrocytes in the aging brain express characteristics of senescence-associated secretory phenotype. *European Journal of Neuroscience*, *34*, 3–11.
- Sánchez-Hidalgo, M., Guerrero Montávez, J. M., Carrascosa-Salmoral, M. D. P., Naranjo Gutierrez, M. D. C., Lardone, P. J., & de la Lastra Romero, C. A. (2009). Decreased MT1 and MT2 melatonin receptor expression in extrapineal tissues of the rat during physiological aging. *Journal of Pineal Research*, *46*, 29–35. <https://doi.org/10.1111/j.1600-079X.2008.00604.x>
- Santoro, A., Spinelli, C. C., Martucciello, S., Nori, S. L., Capunzo, M., Puca, A. A., & Ciaglia, E. (2018). Innate immunity and cellular senescence: The good and the bad in the developmental and aged brain. *J. Leukocyte Biol.*, *103*, 509–524.
- Satinoff, E., Li, H., Tchong, T. K., Liu, C., McArthur, A. J., Medanic, M., & Gillette, M. U. (1993). Do the suprachiasmatic nuclei oscillate in old rats as they do in young ones? *American Journal of Physiology*, *265*, R1216–1222.
- Sato, S., Solanas, G., Peixoto, F. O., Bee, L., Symeonidi, A., Schmidt, M. S., Brenner, C., Masri, S., Benitah, S. A., & Sassone-Corsi, P. (2017). Circadian reprogramming in the liver identifies metabolic pathways of aging. *Cell*, *170*(664–677), e611. <https://doi.org/10.1016/j.cell.2017.07.042>
- Satoh, A., Brace, C. S., Rensing, N., Cliften, P., Wozniak, D. F., Herzog, E. D., Yamada, K. A., & Imai, S.-I. (2013). Sirt1 extends life span and delays aging in mice through the regulation of Nk2 homeobox 1 in the DMH and LH. *Cell Metabolism*, *18*, 416–430. <https://doi.org/10.1016/j.cmet.2013.07.013>
- Schafer, D. P., Lehrman, E. K., Kautzman, A. G., Koyama, R., Mardinly, A. R., Yamasaki, R., Ransohoff, R. M., Greenberg, M. E., Barres, B. A., & Stevens, B. (2012). Microglia sculpt postnatal neural circuits in an activity and complement-dependent manner. *Neuron*, *74*, 691–705. <https://doi.org/10.1016/j.neuron.2012.03.026>
- Scheiermann, C., Gibbs, J., Ince, L., & Loudon, A. (2018). Clocking in to immunity. *Nature Reviews Immunology*, *18*, 423–437.
- Schilperoot, M., van den Berg, R., Bosmans, L. A., van Os, B. W., Dollé, M. E. T., Smits, N. A. M., Guichelaar, T., van Baarle, D., Koemans, L., Berbée, J. F. P., Deboer, T., Meijer, J. H., de Vries, M. R., Vreeken, D., van Gils, J. M., Willems van Dijk, K., van Kerkhof, L. W. M., Lutgens, E., Biermasz, N. R. ... Kooijmans, S. (2020). Disruption of circadian rhythm by alternating light-dark cycles aggravates atherosclerosis development in APOE*3-Leiden.CETP mice. *Journal of Pineal Research*, *68*, e12614.
- Schmutz, I., Chavan, R., Ripperger, J., Maywood, E. S., Langwieser, N., Jurik, A., Stauffer, A., Delorme, J. E., Moosmang, S., Hastings, M. H., Hofmann, F., & Albrecht, U. (2014). A specific role for the REV-ERB α -controlled L-type voltage-gated calcium channel CaV1.2 in resetting the circadian clock in the late night. *Journal of Biological Rhythms*, *29*(4), 288–298. <http://dx.doi.org/10.1177/0748730414540453>
- Schouten, M., Bielefeld, P., Garcia-Corzo, L., Passchier, E. M. J., Gradari, S., Jungenitz, T., Pons-Espinal, M., Gebara, E., Martín-Suárez, S., Lucassen, P. J., De Vries, H. E., Trejo, J. L., Schwarzacher, S. W., De Pietri Tonelli, D., Toni, N., Mira, H., Encinas, J. M., & Fitzsimons, C. P. (2020). Circadian glucocorticoid oscillations preserve a population of adult hippocampal neural stem cells in the aging brain. *Molecular Psychiatry*, *25*, 1382–1405.
- Sellix, M. T., Evans, J. A., Leise, T. L., Castanon-Cervantes, O., Hill, D. D., DeLisser, P., Block, G. D., Menaker, M., & Davidson, A. J. (2012). Aging differentially affects the re-entrainment response of central and peripheral circadian oscillators. *Journal of Neuroscience*, *32*, 16193–16202. <https://doi.org/10.1523/JNEUROSCI.3559-12.2012>
- Semo, M., Peirson, S., Lupi, D., Lucas, R. J., Jeffery, G., & Foster, R. G. (2003). Melanopsin retinal ganglion cells and the maintenance of circadian and pupillary responses to light in aged rodless/coneless (rd/rd cl) mice. *European Journal of Neuroscience*, *17*, 1793–1801.
- Shan, Y., Abel, J. H., Li, Y., Izumo, M., Cox, K. H., Jeong, B., Yoo, S.-H., Olson, D. P., Doyle, F. J., & Takahashi, J. S. (2020). Dual-Color Single-cell imaging of the suprachiasmatic nucleus reveals a circadian role in network synchrony. *Neuron*, *108*, 164–179. <https://doi.org/10.1016/j.neuron.2020.07.012>
- Skene, D. J., & Arendt, J. (2006). Human circadian rhythms: Physiological and therapeutic relevance of light and melatonin. *Annals of Clinical Biochemistry*, *43*, 344–353. <https://doi.org/10.1258/00045630678520142>
- Snider, K. H., Sullivan, K. A., & Obrietan, K. (2018). Circadian regulation of hippocampal-dependent memory: Circuits, synapses, and molecular mechanisms. *Neural Plasticity*, *2018*, 1–13. <https://doi.org/10.1155/2018/7292540>
- Socaciu, A. I., Ionuț, R., Socaciu, M. A., Ungur, A. P., Bârsan, M., Chiorean, A., Socaciu, C., & Răjnoveanu, A. G. (2020). Melatonin, an ubiquitous metabolic regulator: Functions, mechanisms and effects on circadian disruption and degenerative diseases. *Reviews in Endocrine & Metabolic Disorders*, *21*, 465–478. <https://doi.org/10.1007/s11154-020-09570-9>
- Solt, L. A., Kumar, N., Nuhant, P., Wang, Y., Lauer, J. L., Liu, J., Istrate, M. A., Kamenecka, T. M., Roush, W. R., Vidović, D., Schürer, S. C.,



- Xu, J., Wagoner, G., Drew, P. D., Griffin, P. R., & Burrell, T. P. (2011). Suppression of TH17 differentiation and autoimmunity by a synthetic ROR ligand. *Nature*, *472*, 491–494. <https://doi.org/10.1038/nature10075>
- Sprecher, K. E., Kosciak, R. L., Carlsson, C. M., Zetterberg, H., Blennow, K., Okonkwo, O. C., Sager, M. A., Asthana, S., Johnson, S. C., Benca, R. M., & Bendlin, B. B. (2017). Poor sleep is associated with CSF biomarkers of amyloid pathology in cognitively normal adults. *Neurology*, *89*, 445–453. <https://doi.org/10.1212/WNL.0000000000004171>
- Stopa, E. G., Volicer, L., Kuo-Leblanc, V., Harper, D., Lathi, D., Tate, B., & Satlin, A. (1999). Pathologic evaluation of the human suprachiasmatic nucleus in severe dementia. *Journal of Neuropathology and Experimental Neurology*, *58*, 29–39.
- Sugimoto, T., Morioka, N., Zhang, F. F., Sato, K., Abe, H., Hisaoka-Nakashima, K., & Nakata, Y. (2014). Clock gene *Per1* regulates the production of CCL2 and interleukin-6 through p38, JNK1 and NF-kappaB activation in spinal astrocytes. *Molecular and Cellular Neurosciences*, *59*, 37–46.
- Sutin, E. L., Dement, W. C., Heller, H. C., & Kilduff, T. S. (1993). Light-induced gene expression in the suprachiasmatic nucleus of young and aging rats. *Neurobiology of Aging*, *14*, 441–446.
- Swaab, D. F., Fliers, E., & Partiman, T. S. (1985). The suprachiasmatic nucleus of the human brain in relation to sex, age and senile dementia. *Brain Research*, *342*, 37–44. [https://doi.org/10.1016/0006-8993\(85\)91350-2](https://doi.org/10.1016/0006-8993(85)91350-2)
- Tan, D. X., Manchester, L. C., Qin, L., & Reiter, R. J. (2016). Melatonin: A mitochondrial targeting molecule involving mitochondrial protection and dynamics. *International Journal of Molecular Sciences*, *17*(12), 2124
- Tan, D. X., Xu, B., Zhou, X., & Reiter, R. J. (2018). Pineal calcification, melatonin production, aging, associated health consequences and rejuvenation of the pineal gland. *Molecules*, *23*, 301. <https://doi.org/10.3390/molecules23020301>
- Teclemariam-Mesbah, R., Ter Horst, G. J., Postema, F., Wortel, J., & Buijs, R. M. (1999). Anatomical demonstration of the suprachiasmatic nucleus–pineal pathway. *The Journal of Comparative Neurology*, *406*, 171–182.
- Tso, C. F., Simon, T., Greenlaw, A. C., Puri, T., Mieda, M., & Herzog, E. D. (2017). Astrocytes regulate daily rhythms in the suprachiasmatic nucleus and behavior. *Current Biology*, *27*, 1055–1061.
- Tsukahara, S., Tanaka, S., Ishida, K., Hoshi, N., & Kitagawa, H. (2005). Age-related change and its sex differences in histoarchitecture of the hypothalamic suprachiasmatic nucleus of F344/N rats. *Experimental Gerontology*, *40*, 147–155.
- Uryash, A., Flores, V., Adams, J. A., Allen, P. D., & Lopez, J. R. (2020). Memory and learning deficits are associated with Ca²⁺ dyshomeostasis in normal aging. *Frontiers in Aging Neuroscience*, *12*, 224. <https://doi.org/10.3389/fnagi.2020.00224>
- Van der Zee, E. A., Jansen, K., & Gerkema, M. P. (1999). Severe loss of vasopressin-immunoreactive cells in the suprachiasmatic nucleus of aging voles coincides with reduced circadian organization of running wheel activity. *Brain Research*, *816*, 572–579. [https://doi.org/10.1016/S0006-8993\(98\)01239-6](https://doi.org/10.1016/S0006-8993(98)01239-6)
- Van Reeth, O., Zhang, Y., Zee, P. C., & Turek, F. W. (1994). Grafting fetal suprachiasmatic nuclei in the hypothalamus of old hamsters restores responsiveness of the circadian clock to a phase shifting stimulus. *Brain Research*, *643*, 338–342. [https://doi.org/10.1016/0006-8993\(94\)90044-2](https://doi.org/10.1016/0006-8993(94)90044-2)
- Van Someren, E. J. W., Oosterman, J. M., Van Harten, B., Vogels, R. L., Gouw, A. A., Weinstein, H. C., Poggesi, A., Scheltens, P., & Scherder, E. J. A. (2019). Medial temporal lobe atrophy relates more strongly to sleep-wake rhythm fragmentation than to age or any other known risk. *Neurobiology of Learning and Memory*, *160*, 132–138. <https://doi.org/10.1016/j.nlm.2018.05.017>
- Vellai, T., Takacs-Vellai, K., Zhang, Y., Kovacs, A. L., Orosz, L., & Müller, F. (2003). Influence of TOR kinase on lifespan in *C. elegans*. *Nature*, *426*, 620. <https://doi.org/10.1038/426620a>
- Videnovic, A., Klerman, E. B., Wang, W., Marconi, A., Kuhta, T., & Zee, P. C. (2017). Timed light therapy for sleep and daytime sleepiness associated with parkinson disease: A randomized clinical trial. *JAMA Neurology*, *74*, 411–418. <https://doi.org/10.1001/jaman.2016.5192>
- von Gall, C., & Weaver, D. R. (2008). Loss of responsiveness to melatonin in the aging mouse suprachiasmatic nucleus. *Neurobiology of Aging*, *29*, 464–470.
- Wade, A. G., Farmer, M., Harari, G., Fund, N., Laudon, M., Nir, T., Frydman-Marom, A., & Zisapel, N. (2014). Add-on prolonged-release melatonin for cognitive function and sleep in mild to moderate Alzheimer's disease: A 6-month, randomized, placebo-controlled, multicenter trial. *Clinical Interventions in Aging*, *9*, 947–961.
- Waller, K. L., Mortensen, E. L., Avlund, K., Fagerlund, B., Lauritzen, M., Gammeltoft, S., & Jennum, P. (2016). Melatonin and cortisol profiles in late midlife and their association with age-related changes in cognition. *Nature and Science of Sleep*, *8*, 47–53.
- Wang, L. M. C., Dragich, J. M., Kudo, T., Odom, I. H., Welsh, D. K., O'Dell, T. J., & Colwell, C. S. (2009). Expression of the circadian clock gene *Period2* in the hippocampus: Possible implications for synaptic plasticity and learned behaviour. *ASN Neuro*, *1*, AN20090020.
- Watanabe, A., Shibata, S., & Watanabe, S. (1995). Circadian rhythm of spontaneous neuronal activity in the suprachiasmatic nucleus of old hamster in vitro. *Brain Research*, *695*, 237–239. [https://doi.org/10.1016/0006-8993\(95\)00713-Z](https://doi.org/10.1016/0006-8993(95)00713-Z)
- Weaver, D. R. (1998). The suprachiasmatic nucleus: A 25-year retrospective. *Journal of Biological Rhythms*, *13*, 100–112. <https://doi.org/10.1177/074873098128999952>
- Wehrens, S. M. T., Christou, S., Isherwood, C., Middleton, B., Gibbs, M. A., Archer, S. N., Skene, D. J., & Johnston, J. D. (2017). Meal timing regulates the human circadian system. *Current Biology*, *27*(1768–1775), e1763. <https://doi.org/10.1016/j.cub.2017.04.059>
- Weinert, H., Weinert, D., Schurov, I., Maywood, E. S., & Hastings, M. H. (2001). Impaired expression of the *mPer2* circadian clock gene in the suprachiasmatic nuclei of aging mice. *Chronobiology International*, *18*, 559–565.
- Whitt, J. P., Montgomery, J. R., & Meredith, A. L. (2016). BK channel inactivation gates daytime excitability in the circadian clock. *Nature Communications*, *7*, 10837.
- Wu, Y.-H., & Swaab, D. F. (2005). The human pineal gland and melatonin in aging and Alzheimer's disease. *Journal of Pineal Research*, *38*, 145–152. <https://doi.org/10.1111/j.1600-079X.2004.00196.x>
- Wu, Y. H., Zhou, J. N., Van Heerikhuizen, J., Jockers, R., & Swaab, D. F. (2007). Decreased MT1 melatonin receptor expression in the suprachiasmatic nucleus in aging and Alzheimer's disease. *Neurobiology of Aging*, *28*, 1239–1247. <https://doi.org/10.1016/j.neurobiolaging.2006.06.002>
- Wullschlegel, S., Loewith, R., & Hall, M. N. (2006). TOR signaling in growth and metabolism. *Cell*, *124*, 471–484. <https://doi.org/10.1016/j.cell.2006.01.016>
- Wyse, C. A., & Coogan, A. N. (2010). Impact of aging on diurnal expression patterns of *CLOCK* and *BMAL1* in the mouse brain. *Brain Research*, *1337*, 21–31. <https://doi.org/10.1016/j.brainres.2010.03.113>
- Yamamoto, D., Takahashi, M., Nagashima, S., Wakabayashi, T., Yamazaki, H., Takei, A., Takei, S., Sakai, K., Ebihara, K., Iwasaki, Y., Yada, T., & Ishibashi, S. (2020). Peripheral circadian rhythms in the liver and white adipose tissue of mice are attenuated by constant light and restored by time-restricted feeding. *PLoS One*, *15*, e0234439.
- Yamazaki, S., Numano, R., Abe, M., Hida, A., Takahashi, R.-i., Ueda, M., Block, G. D., Sakaki, Y., Menaker, M., & Tei, H. (2000). Resetting central and peripheral circadian oscillators in transgenic rats. *Science*, *288*, 682–685. <https://doi.org/10.1126/science.288.5466.682>

- Yamazaki, S., Straume, M., Tei, H., Sakaki, Y., Menaker, M., & Block, G. D. (2002). Effects of aging on central and peripheral mammalian clocks. *Proceedings of the National Academy of Sciences of the United States of America*, 99, 10801–10806. <https://doi.org/10.1073/pnas.152318499>
- Yang, S.-B., Tien, A.-C., Boddupalli, G., Xu, A. W., Jan, Y. N., & Jan, L. Y. (2012). Rapamycin ameliorates age-dependent obesity associated with increased mTOR signaling in hypothalamic pomc neurons. *Neuron*, 75, 425–436. <https://doi.org/10.1016/j.neuron.2012.03.043>
- Zeitler, J. M., Daniels, J. E., Duffy, J. F., Klerman, E. B., Shanahan, T. L., Dijk, D.-J., & Czeisler, C. A. (1999). Do plasma melatonin concentrations decline with age? *American Journal of Medicine*, 107, 432–436. [https://doi.org/10.1016/S0002-9343\(99\)00266-1](https://doi.org/10.1016/S0002-9343(99)00266-1)
- Zeitler, J. M., Duffy, J. F., Lockley, S. W., Dijk, D.-J., & Czeisler, C. A. (2007). Plasma melatonin rhythms in young and older humans during sleep, sleep deprivation, and wake. *Sleep*, 30, 1437–1443. <https://doi.org/10.1093/sleep/30.11.1437>
- Zhang, G., Li, J., Purkayastha, S., Tang, Y., Zhang, H., Yin, Y., Li, B., Liu, G., & Cai, D. (2013). Hypothalamic programming of systemic ageing involving IKK- β , NF- κ B and GnRH. *Nature*, 497, 211–216. <https://doi.org/10.1038/nature12143>
- Zhang, R., Lahens, N. F., Ballance, H. I., Hughes, M. E., & Hogenesch, J. B. (2014). A circadian gene expression atlas in mammals: Implications for biology and medicine. *Proceedings of the National Academy of Sciences of the United States of America*, 111, 16219–16224. <https://doi.org/10.1073/pnas.1408886111>
- Zhang, W., Chen, X. Y., Su, S. W., Jia, Q. Z., Ding, T., Zhu, Z. N., & Zhang, T. (2016). Exogenous melatonin for sleep disorders in neurodegenerative diseases: A meta-analysis of randomized clinical trials. *Neurological Sciences*, 37, 57–65.
- Zhou, J.-N., Hofman, M. A., & Swaab, D. F. (1995). VIP neurons in the human SCN in relation to sex, age, and Alzheimer's disease. *Neurobiology of Aging*, 16, 571–576.
- Zisapel, N. (2018). New perspectives on the role of melatonin in human sleep, circadian rhythms and their regulation. *British Journal of Pharmacology*, 175, 3190–3199.

How to cite this article: Buijink MR, Michel S. A multi-level assessment of the bidirectional relationship between aging and the circadian clock. *J Neurochem*. 2021;157:73–94. <https://doi.org/10.1111/jnc.15286>