

Multimodality imaging for myocardial injury in acute myocardial infarction and the assessment of valvular heart disease

Podlesnikar, T.

Citation

Podlesnikar, T. (2022, June 28). Multimodality imaging for myocardial injury in acute myocardial infarction and the assessment of valvular heart disease. Retrieved from https://hdl.handle.net/1887/3420621

Version: Publisher's Version

License: License agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden


Downloaded

https://hdl.handle.net/1887/3420621

from:

Note: To cite this publication please use the final published version (if applicable).

Part I



Cardiovascular Magnetic Resonancederived Left Ventricular Strain in Acute Myocardial Infarction

Effect of early metoprolol during
ST-segment elevation myocardial infarction
on left ventricular strain: feature-tracking
cardiovascular magnetic resonance
substudy from the METOCARD-CNIC trial

Tomaž Podlesnikar, Gonzalo Pizarro, Rodrigo Fernández-Jiménez, Jose M Montero-Cabezas, Javier Sánchez-González, Chiara Bucciarelli-Ducci, Nina Ajmone Marsan, Zlatko Fras, Jeroen J Bax, Valentin Fuster, Borja Ibáñez, Victoria Delgado

JACC Cardiovasc Imaging 2019;12(7 Pt 1):1188-98

ABSTRACT

Background: Early intravenous metoprolol before primary percutaneous coronary intervention (PCI) in ST-segment elevation myocardial infarction (STEMI) portends better outcomes in the METOCARD-CNIC trial. The aim of the present study was to evaluate the effect of early intravenous metoprolol on left ventricular (LV) strain assessed with feature-tracking cardiovascular magnetic resonance (CMR).

Methods: A total of 197 patients with acute anterior STEMI who were enrolled in the METO-CARD-CNIC trial (100 allocated to intravenous metoprolol before primary PCI and 97 control patients) were evaluated. LV global circumferential strain (GCS) and global longitudinal strain (GLS) were measured with feature-tracking CMR at 1 week and 6 months after STEMI and compared between randomization groups.

Results: Patients who received early intravenous metoprolol had significantly more preserved LV strain compared to the control patients at 1 week after STEMI (GCS: -13.9±3.8% versus -12.6±3.9%, respectively; P=0.013; GLS: -11.9±2.8% versus -10.9±3.2%, respectively; P=0.032). In both groups, LV strain significantly improved during follow-up (mean difference between 6-month and 1-week strain for the metoprolol group: GCS: -2.9%, 95% CI: -3.5% to -2.4; GLS: -2.9%, 95% CI: -3.4 to -2.4; both P<0.001; the control group: GCS: -3.4%, 95% CI: -3.9% to -2.8%; GLS: -3.4%, 95% CI: -3.9% to -3.0%; both P<0.001). When dividing the overall cohort of patients in quartiles of GCS and GLS, there were significantly less patients in the first quartile (i.e. the worst LV systolic function) who received early intravenous metoprolol compared to control patients at 1 week and 6 months (P<0.05 for GCS and GLS at both time points).

Conclusions: In patients with anterior STEMI, early administration of intravenous metoprolol before primary PCI was associated with significantly less patients with severely depressed LV GCS and GLS, both at 1 week and 6 months. Feature-tracking CMR represents a complementary tool to evaluate the benefits of cardioprotective therapies.

INTRODUCTION

The long-term treatment with beta-blockers after ST-segment elevation myocardial infarction (STEMI) is well established and the benefit appears to be greatest for patients with myocardial infarction complicated by heart failure, left ventricular (LV) systolic dysfunction, or ventricular arrhythmias, 1,2 Current European and American guidelines recommend initiating oral beta-blockers in the first 24 hours after STEMI.^{1,2} The role of routine early, intravenous beta-blockers administration prior to primary percutaneous coronary intervention (PCI) is less firmly established. In the context of reduced oxygen supply during myocardial infarction, beta-blockers have the potential to reduce ischemic injury when administered prior to PCI, through their effect on slowing of heart rate, decreasing myocardial contractility, and lowering systemic blood pressure. In addition, some beta-blockers have shown to be able to reduce reperfusion-injury by inhibiting neutrophils function.³ The Effect of Metoprolol in Cardioprotection During an Acute Myocardial Infarction (METOCARD-CNIC) trial showed that early administration of intravenous metoprolol before primary PCI significantly reduced infarct size 1 week post-STEMI as evaluated by cardiovascular magnetic resonance (CMR) imaging.4 In addition, early metoprolol administration was associated with improved long-term LV ejection fraction (LVEF), fewer indications for cardioverter-defibrillator implantation, and fewer heart failure readmissions. 5 Accordingly, current European guidelines indicate that intravenous beta-blockers should be considered at the time of presentation in STEMI patients undergoing primary PCI provided that there are no contraindications, no signs of acute heart failure, and the systolic blood pressure is >120 mmHg.1

The impact of early intravenous metoprolol on LV myocardial strain has not yet been evaluated. In contrast to LVEF, LV strain does not rely on geometrical assumptions, shows superior intra- and inter-observer reproducibility and can detect subtle systolic dysfunction in patients with preserved LVEF.^{6,7} Recent development of feature-tracking CMR allows multidirectional myocardial strain assessment from standard cine images without the need for specialized pulse sequences or additional scanning time.⁸ In the METOCARD-CNIC trial population, we evaluated LV global circumferential (GCS) and longitudinal (GLS) strain measured with feature-tracking CMR both at 1 week and 6 months after primary PCI.

METHODS

Patient population

The present study included patients who were enrolled in the METOCARD-CNIC trial and completed 1-week and 6-month CMR study. Briefly, the multicenter randomized METOCARD-CNIC clinical trial recruited patients with first anterior STEMI undergoing primary PCI.⁹ A total of 270 patients were randomized to receive up to 15 mg intravenous metoprolol before reperfusion versus conventional therapy. All patients received oral metoprolol, first dose 12-24 hours after STEMI. Exclusion criteria were Killip class III to IV acute heart failure, systolic blood pressure persistently <120 mmHg, PR interval >240 milliseconds (or type II–III atrioventricular block), heart rate persistently <60 beats/min, or active treatment with any beta-blocker agent. Of the initial population, 202 patients underwent 2 CMR studies, at 1 week (5 to 7 days) and 6 months after STEMI. Conventional CMR parameters of LV dimensions, function and myocardial scar and LV GCS and GLS measured with feature-tracking analysis were evaluated at both time points for the overall population as a single group, and for each randomization treatment arm individually.

The study was approved by the ethical committees and institutional review boards at each participating center. All eligible patients gave written informed consent.

Cardiovascular magnetic resonance

The CMR protocol has been described in detail elsewhere. Data acquisition was performed with 1.5 and 3.0 T CMR scanners. LV 2-, 3- and 4-chamber views and a stack of contiguous short-axis slices covering the whole LV were acquired with steady-state free precession functional cine imaging. Typical acquisition parameters were: voxel size 1.6×2 mm, slice thickness 8 mm, gap 0 mm, cardiac phases 25-30, TR 3.5, TE 1.7, flip angle 40, SENSE 1.5, averages 1, FOV 360 × 360 mm. Subsequently, segmented inversion recovery gradient echo sequence acquired 10-15 minutes after a cumulative dose of 0.2 mmol/kg intravenous gadolinium contrast agent (Magnevist, Schering AG, Berlin, Germany) was employed for myocardial necrosis/fibrosis imaging. LV volumes, LV mass, LVEF and late gadolinium enhancement (LGE) data were analyzed with dedicated software (QMass MR 7.5; Medis, Leiden, the Netherlands) as described previously.

Feature-tracking CMR analysis

Feature-tracking CMR analysis was performed on steady-state free precession cine images with dedicated software (CVI⁴² v5.3, Circle Cardiovascular Imaging, Calgary, Canada) (Figure 1, Videos

1 and 2). First, the LV endo- and epicardium were delineated at end-diastole in the LV 2-, 3- and 4-chamber views and contiguous short-axis slices and the LV reference points were defined: the mitral annulus and the LV apex in long-axis views and the anterior right ventricular insertion point in the short-axis slices. The most basal short-axis slices, in which the image plane showed LV myocardium only at end-diastole but not at end-systole were excluded. The outlined myocardium borders were automatically tracked throughout the cardiac cycle with fully automated feature-tracking analysis. The quality of the myocardium tracking was visually evaluated. Global time-strain curves were obtained and peak LV GCS and GLS values were recorded.




Figure 1: Feature tracking cardiovascular magnetic resonance. (A) Left ventricular (LV) mid-cavity short-axis and 4-chamber long-axis end-diastolic steady-state free precession images. LV endo- and epicardium (red and green lines) were delineated and LV reference points were defined: the anterior right ventricular insertion point in the short-axis view and the mitral annulus and LV apex in the 4-chamber view. The same method was repeated in the remaining long- and short-axis slices. **(B and C)** Visual evaluation of myocardium tracking (Video 1 and 2 in supplementary material). The interventricular septum and LV anterior wall in the short-axis view and the mid-to-apical septum and apex in 4-chamber view (infarcted area) show impaired deformation compared to the other myocardial segments (B = end-diastole, C = end-systole). **(D)** Global time-strain curves were obtained and peak global circumferential strain (-11.9%, top image) and peak global longitudinal strain (-10.2%, bottom image) values were recorded.

A single observer (TP) performed feature-tracking analysis of CMR data. The same observer repeated the analysis of 20 randomly selected CMR scans after 4 weeks to assess the intra-observer variability. A second observer (JMMC), blinded to the results of the first observer, re-measured a different subset of 20 randomly selected CMR scans for the assessment of inter-observer variability.

Statistical analysis

Continuous variables are presented as mean ± standard deviation and categorical variables as frequencies (percentages). Patients were divided into 2 groups according to the treatment received. Comparisons between the early metoprolol group and the control group were performed using independent samples t-test for continuous variables and Pearson's Chi square test or Fischer's exact test for categorical variables. Fischer's exact test was used when the expected value of a categorical variable was <5. Comparisons between 1-week and 6-month CMR data were performed using paired samples t-test. In addition, the study population was divided in quartiles of LV GCS and GLS. The number of patients within the first quartile of LV GCS and GLS (worst LV systolic function) at each randomization treatment arm (early intravenous metoprolol vs control patients) was compared with Pearson's Chi square test at 1 week and 6 months of follow-up. In addition, logistic regression analysis was performed to assess the value of LV GCS and GLS 1 week after STEMI to predict LVEF normalization (≥50%) at 6 months. Odds ratios and 95% confidence intervals (CI) were calculated and adjusted for infarct size (LGE extent) at 1-week CMR, demographic and clinical variables (age, sex, body mass index, presence of hypertension, diabetes, dyslipidemia, smoking status) and treatment randomization arm (early intravenous metoprolol vs. control patients).

The intra- and inter-observer agreement for GCS and GLS measurements were assessed with intraclass correlation coefficients. A two-sided P-value of <0.05 was statistically significant and excellent agreement was defined as an intraclass correlation coefficient >0.9. All statistical analyses were performed using IBM SPSS Statistics 23 (IBM, Armonk, New York).

RESULTS

Of the initial 202 patients who underwent 2 CMR studies, feature-tracking CMR analysis was feasible in 197 patients (early metoprolol group: N=100; control group: N=97) and they formed the population of the present analysis. LV GLS analysis at 6 months was feasible in 195 patients (early metoprolol group: N=99; control group: N=96).

Patients demographics, cardiovascular risk factors, clinical characteristics at recruitment and procedural characteristics of the overall population (mean age 58.1 years, 88% male) and the patients divided according to received randomization treatment (metoprolol vs control) are presented in Table 1. There were no statistically significant differences between both groups. Conventional and feature-tracking CMR parameters of LV structure and function, evaluated at 1 week and 6 months after STEMI for the overall population and for each randomization treatment arm individually, are presented in Table 2.

Table 1: Patients demographics, cardiovascular risk factors, clinical status at recruitment and procedural characteristics.

| | Total (N=197) | Metoprolol (N=100) | Control (N=97) | P-value |
|--|------------------|-----------------------|-------------------|---------|
| Demographics | | | | |
| Age (years) | 58.1±11.3 | 57.8±12.3 | 58.4±10.1 | 0.698 |
| Sex (male) | 173 (88) | 87 (87) | 86 (89) | 0.865 |
| BMI (kg/m²) | 27.6±3.5 | 27.6±3.5 | 27.6±3.5 | 0.900 |
| Cardiovascular risk factors | | | | |
| Hypertension | 74 (38) | 37 (37) | 37 (38) | 0.955 |
| Diabetes mellitus | 39 (20) | 21 (21) | 18 (19) | 0.616 |
| Dyslipidemia | 85 (43) | 43 (43) | 42 (43) | 0.935 |
| Smoking* | 126 (64) | 64 (64) | 62 (64) | 0.839 |
| Clinical status at recruitment | | | | |
| Killip class II† | 19 (10) | 8 (8) | 11 (11) | 0.441 |
| Systolic BP (mmHg) | 142±19 | 142±18 | 142±19 | 0.949 |
| Diastolic BP (mmHg) | 88±16 | 89±16 | 87±15 | 0.266 |
| Heart rate (bpm) | 82±13 | 82±14 | 81±13 | 0.539 |
| Procedural characteristics | | | | |
| Ischemia duration (min) | 194±65 | 197±62 | 191±68 | 0.488 |
| TIMI grade 0-1 flow before primary PCI | 163 (83) | 80 (80) | 83 (86) | 0.373 |
| Successful PCI (TIMI grade 2-3 flow) | 194 (99) | 100 (100) | 94 (97) | 0.117 |

BMI = body mass index; BP = blood pressure; PCI = percutaneous coronary intervention; TIMI = Thrombolysis in Myocardial Infarction. Values are mean ± SD or n (%).

†all other patients were Killip class I (Killip class III to IV were study's exclusion criteria).

Table 2: Effect of treatment randomization on conventional and feature tracking CMR parameters.

| | | 6 months | | | | | | | | |
|----------------|--------------------|-----------------------|-------------------|--------------------------------|---------|--------------------|-----------------------|-------------------|--------------------------------|---------|
| | Overall (N=197) | Metoprolol (N=100) | Control (N=97) | Mean difference [95% CI] | P-value | Overall (N=197) | Metoprolol (N=100) | Control (N=97) | Mean difference [95% CI] | P-value |
| LVEDV (mL) | 172.6±36.2 | 169.8±33.4 | 175.5±38.8 | -5.6 [-15.8 to 4.5] | 0.276 | 191.6±42.6 | 187.0±38.8 | 196.5±45.9 | -9.5 [-21.5 to 2.5] | 0.119 |
| LVESV (mL) | 97.8±31.3 | 92.9±26.6 | 102.8±34.9 | -9.8 [-18.6 to -1.1] | 0.028 | 104.4±40.8 | 98.2±36.1 | 110.8±44.5 | -12.6 [-24.1 to -1.2] | 0.031 |
| LVEF (%) | 44.2±9.4 | 45.8±9.1 | 42.6±9.6 | 3.2 [0.6 to 5.8] | 0.017 | 47.0±10.8 | 48.7±10.0 | 45.3±11.4 | 3.4 [0.4 to 6.4] | 0.028 |
| LV mass (g) | 111.5±25.4 | 109.1±25.2 | 113.9±25.5 | -4.7 [-11.9 to 2.4] | 0.192 | 85.7±17.6 | 84.6±17.4 | 86.8±17.7 | -2.3 [-7.2 to 2.7] | 0.371 |
| LGE (%) | 22.7±12.8 | 20.9±11.6 | 24.7±13.8 | -3.8 [-7.4 to -0.2] | 0.039 | 16.9±9.7 | 15.7±9.5 | 18.0±9.7 | -2.3 [-5.1 to 0.5] | 0.104 |
| LV GCS (%) | -13.3±3.9 | -13.9±3.8 | -12.6±3.9 | -1.4 [-2.4 to -0.3] | 0.013 | -16.4±4.2 | -16.9±4.0 | -15.9±4.4 | -0.9 [-2.1 to 0.3] | 0.122 |
| LV GLS (%) | -11.4±3.0 | -11.9±2.8 | -10.9±3.2 | -0.9 [-1.8 to -0.1] | 0.032 | -14.6±3.0 | -14.8±2.9 | -14.4±3.0 | -0.4 [-1.2 to 0.5] | 0.379 |

CI = confidence interval; CMR = cardiovascular magnetic resonance; GCS = global circumferential strain; GLS = global longitudinal strain; LGE = late gadolinium enhancement; LV = left ventricular; LVEDV = left ventricular end-diastolic volume; LVEF = left ventricular ejection fraction; LVESV = left ventricular end-systolic volume.

Values are mean ± SD unless otherwise indicated.

^{*}smoking was defined as current or quitted <10 years ago.

LV conventional and feature-tracking CMR parameters 1 week after STEMI

One week after intervention (metoprolol or control), patients who received early intravenous metoprolol showed significantly smaller LV end-systolic volumes, higher LVEF and smaller infarct sizes assessed by LGE (Table 2, Figure 2). In addition, patients who received early intravenous metoprolol had more preserved LV GCS and GLS than patients in the control group (GCS: -13.9±3.8% versus -12.6±3.9%, respectively; P=0.013; GLS: -11.9±2.8% versus -10.9±3.2%, respectively; P=0.032).




Figure 2: Time course and effect of treatment randomization on conventional and feature tracking cardiovascular magnetic resonance parameters after ST-segment elevation myocardial infarction. Left ventricular end-diastolic volume (LVEDV) (A), left ventricular end-systolic volume (LVESV) (B), left ventricular ejection fraction (LVEF) (C), late gadolinium enhancement (LGE) (D), peak global circumferential strain (GCS) (E) and peak global longitudinal strain (GLS) (F) in the early intravenous metoprolol and the control group, at 1 week and at 6 months after the acute event. The asterisks represent the mean values and the error bars represent the standard errors of the mean. P-values describe the statistical significance between both treatment arms at each time point.

Changes in LV conventional and feature-tracking CMR parameters between 1-week and 6-month follow-up after STEMI

There were significant changes in conventional CMR parameters and LV strain between 1-week and 6-month follow-up in the overall population and in both study treatment arms (Table 3, Figure 2 and 3). LV end-diastolic and end-systolic volumes significantly increased over time. However, LV dilation was more pronounced for LV end-diastolic volumes than for LV end-systolic volumes, partly explaining the significant improvement of LVEF over time. The percentage of LV myocardium with LGE significantly decreased over the 6 months of follow-up. In addition, LV GCS and GLS significantly improved over the 6-month follow-up (mean difference between 6-month and 1-week strain for the metoprolol group: GCS: -2.9%, 95% CI: -3.5% to -2.4; GLS: -2.9%, 95% CI: -3.4 to -2.4; both P<0.001; the control group: GCS: -3.4%, 95% CI: -3.9% to -2.8%; GLS: -3.4%, 95% CI: -3.9% to -3.0%; both P<0.001).

Table 3: Time course of LV conventional and feature tracking CMR parameters after STEMI.

| | Overall (N=197) | | | Metoprolol (N=100) | | | Control (N=97) | | |
|-------------|--------------------|----------------|---------|--------------------|----------------|---------|--------------------|----------------|---------|
| | Mean difference | 95% CI | P-value | Mean difference | 95% CI | P-value | Mean difference | 95% CI | P-value |
| LVEDV (mL) | 18.9 | 15.3 to 22.5 | <0.001 | 16.4 | 11.6 to 21.2 | <0.001 | 21.5 | 16.2 to 26.8 | <0.001 |
| LVESV (mL) | 6.7 | 3.4 to 9.9 | <0.001 | 4.9 | 0.4 to 9.3 | 0.032 | 8.5 | 3.6 to 13.4 | 0.001 |
| LVEF (%) | 2.7 | 1.8 to 3.6 | <0.001 | 2.9 | 1.5 to 4.2 | <0.001 | 2.6 | 1.3 to 3.9 | <0.001 |
| LV mass (g) | -25.8 | -28.5 to -23.2 | <0.001 | -24.6 | -28.3 to -20.9 | <0.001 | -27.0 | -30.9 to -23.1 | <0.001 |
| LGE (%) | -5.8 | -6.7 to -4.8 | <0.001 | -5.1 | -6.5 to -3.8 | <0.001 | -6.5 | -7.8 to -5.1 | <0.001 |
| LV GCS (%) | -3.2 | -3.5 to -2.8 | <0.001 | -2.9 | -3.5 to -2.4 | <0.001 | -3.4 | -3.9 to -2.8 | <0.001 |
| LV GLS (%) | -3.2 | -3.5 to -2.8 | <0.001 | -2.9 | -3.4 to -2.4 | <0.001 | -3.4 | -3.9 to -3.0 | <0.001 |

 ${\sf STEMI=ST-segment\ elevation\ myocardial\ infarction;\ other\ abbreviations\ as\ in\ Table\ 2.}$




Figure 3: Time course of conventional and feature tracking cardiovascular magnetic resonance parameters after ST-segment elevation myocardial infarction in the overall population. Left ventricular end-diastolic volume (LVEDV) (A), left ventricular end-systolic volume (LVESV) (B), left ventricular ejection fraction (LVEF) (C), late gadolinium enhancement (LGE) (D), peak global circumferential strain (GCS) (E) and peak global longitudinal strain (GLS) (F) in the overall population at 1 week and at 6 months after the acute event. Dots are individual patient data. Blue lines represent the mean ± standard error of the mean. P-values describe the statistical significance between the two time points.

LV GCS and GLS at 1 week after STEMI were significant predictors of LVEF normalization (LVEF ≥50%) at 6-month follow-up (Table 4). Each 1 percent increase in LV GCS was associated with 40.8% higher likelihood of LVEF normalization (P<0.001) and each 1% of increase in LV GLS was associated with 40.9% higher likelihood of LVEF normalization at 6 months after STEMI (P<0.001). Both, LV GCS and GLS, remained significant predictors of LVEF normalization after adjusting for the extent of LGE on 1-week CMR, demographic and clinical variables and treatment randomization arm (P<0.001 for both).

Table 4: Left ventricular ejection fraction (LVEF), left ventricular global circumferential (GCS) and longitudinal (GLS) strain at 1 week after myocardial infarction as predictors of LVEF normalization (LVEF ≥50%) at 6 months after the acute event.

| | Univariate analysis | | | Mult | ivariate an | alysis 1* | Multivariate analysis 2† | | |
|----------|---------------------|-----------------|---------|---------------|-----------------|-----------|--------------------------|-----------------|---------|
| | Odds ratio | 95% CI | P-value | Odds ratio | 95% CI | P-value | Odds ratio | 95% CI | P-value |
| LVEF (%) | 1.289 | 1.203- 1.382 | <0.001 | 1.190 | 1.100- 1.286 | <0.001 | 1.231 | 1.129- 1.342 | <0.001 |
| GCS (%) | 0.592 | 0.513- 0.682 | <0.001 | 0.723 | 0.619- 0.843 | <0.001 | 0.715 | 0.610- 0.839 | <0.001 |
| GLS (%) | 0.591 | 0.505- 0.692 | <0.001 | 0.718 | 0.600- 0.860 | <0.001 | 0.666 | 0.542- 0.819 | <0.001 |

CI = confidence interval.

LV conventional and feature-tracking CMR parameters 6 months after STEMI

The improvements in LV conventional and feature-tracking CMR parameters resulted in non-significant differences in LV end-diastolic volumes, LV mass and LGE between both treatment arms at 6 months (Table 2, Figure 2). However, patients who received early intravenous metoprolol still had significantly smaller LV end-systolic volumes and higher LVEF. In addition, patients who received early intravenous metoprolol showed a non-significant trend for more preserved LV strain compared to patients in the control group (GCS: -16.9±4.0% versus -15.9±4.4%, respectively; P=0.122; GLS: -14.8±2.9% versus -14.4±3.0%, respectively; P=0.379).

The effect of early metoprolol on severe LV systolic dysfunction

When dividing the overall cohort of patients in quartiles of GCS and GLS, there were significantly less number of patients receiving early intravenous metoprolol in the first GCS and GLS quartile (i.e. the worst LV systolic function), both at 1 week and at 6 months (Table 5, Figure 4). At 1 week after STEMI, there were 18 patients who received early intravenous metoprolol versus 31 patients with the conventional treatment in the first GCS quartile group (\geq -10.0%) (P=0.023) and 13 patients who received early metoprolol versus 36 control patients in the first GLS quartile group (\geq -9.3%) (P<0.001). At 6 months after STEMI, there were 17 patients who received early intravenous metoprolol versus 32 patients with the conventional treatment in the first GCS quartile group (\geq -13.1%) (P=0.009) and 18 patients who received early metoprolol versus 31 control patients in the first GLS quartile group (\geq -12.8%) (P=0.023).

^{*}adjusted for the extent of late gadolinium enhancement (LGE) on 1-week cardiovascular magnetic resonance (CMR). †adjusted for the extent of LGE on 1-week CMR, age, sex, body mass index, presence of hypertension, diabetes, dyslipidemia, smoking status and treatment randomization arm (early intravenous metoprolol vs control).

Table 5: Number of patients in GCS and GLS quartiles at 1 week and 6 months after STEMI.

| | LV GCS 1 week | | | | | | | | | |
|------------|---------------------------|------------------------------------|------------------------------------|---------------------------|--|--|--|--|--|--|
| | 1st quartile (≥-10.0%) | 2nd quartile (-10.0% to -13.1%) | 3rd quartile (-13.1% to -16.3%) | 4th quartile (<-16.3%) | | | | | | |
| Metoprolol | 18 | 22 | 34 | 26 | | | | | | |
| Control | 31 | 28 | 15 | 23 | | | | | | |
| | LV GLS 1 week | | | | | | | | | |
| | 1st quartile (≥-9.3%) | 2nd quartile (-9.3% to -11.3%) | 3rd quartile (-11.3% to -13.2%) | 4th quartile (<-13.2%) | | | | | | |
| Metoprolol | 13 | 34 | 25 | 28 | | | | | | |
| Control | 36 | 16 | 25 | 20 | | | | | | |
| | LV GCS 6 months | | | | | | | | | |
| | 1st quartile (≥-13.1%) | 2nd quartile (-13.1% to -16.4%) | 3rd quartile (-16.4% to -19.8%) | 4th quartile (<-19.8%) | | | | | | |
| Metoprolol | 17 | 30 | 27 | 26 | | | | | | |
| Control | 32 | 20 | 22 | 23 | | | | | | |
| | LV GLS 6 months | | | | | | | | | |
| | 1st quartile (≥-12.8%) | 2nd quartile (-12.8% to -15.0%) | 3rd quartile (-15.0% to -16.8%) | 4th quartile (<-16.8%) | | | | | | |
| Metoprolol | 18 | 26 | 29 | 26 | | | | | | |
| Control | 31 | 23 | 20 | 22 | | | | | | |

Abbreviations as in Table 2.




Figure 4: Number of patients within the first quartile of LV GCS and GLS (worst LV systolic function) in the early metoprolol group versus controls at 1-week and 6-month follow-up. Patients in the first global circumferential strain (GCS) and global longitudinal strain (GLS) quartile (worst LV systolic function) were compared according to the treatment received (early intravenous metoprolol vs conventional therapy) at 1 week and at 6 months after acute ST-segment elevation myocardial infarction.

Reproducibility of global left ventricular strain measurements

Excellent intra- and inter-observer variabilities for the feature-tracking CMR analysis of GCS and GLS were obtained. The intra-observer intraclass correlation coefficients (95% CI) for the measurement of LV GCS and GLS were 0.990 (0.975-0.996) and 0.982 (0.955-0.993), respective-

ly. Furthermore, the inter-observer intraclass correlation coefficients (95% CI) for the measurement of LV GCS and GLS were 0.995 (0.987-0.998) and 0.990 (0.976-0.996), respectively.

DISCUSSION

The present study demonstrates that in patients with anterior STEMI treated with primary PCI, early administration of intravenous metoprolol was associated with more preserved LV GCS and GLS at 1 week after myocardial infarction as compared to control patients. In addition, early administration of intravenous metoprolol before primary PCI was associated with significantly less patients with severely depressed GCS and GLS both at 1 week and 6 months. Altogether, these data indicate that early intravenous metoprolol before reperfusion improves short and long-term LV systolic dysfunction as evaluated with feature-tracking CMR.

LV conventional and feature-tracking CMR parameters 1 week after STEMI

Acute myocardial infarction results in myocardial cell necrosis and changes in extracellular collagen matrix that portend adverse consequences on LV structure and function. 10 While early intravenous beta-blocker administration offers physiological rationale for lowering the myocardial infarction burden, their routine use has been disputed over the last decades due to the conflicting data on patients outcome.11 The METOCARD-CNIC trial was the first randomized control trial in the modern era of primary PCI in STEMI patients, showing that early administration of intravenous metoprolol resulted in significant reduction of LV end-systolic volumes, increase in LVEF and smaller LGE-assessed infarct size 1 week after anterior STEMI, as evaluated by CMR imaging.4 The present study provides additional information on the effect of early intravenous metoprolol on LV systolic function by means of circumferential and longitudinal shortening, assessed with novel feature-tracking CMR algorithm. This is important since LV strain with speckle tracking echocardiography has been shown to be a more sensitive marker of LV dysfunction⁷ and to provide incremental prognostic information over LVEF in the STEMI population.¹² Recently, clinical implications of feature-tracking CMR in STEMI have been demonstrated.^{13,14} Our results show that GCS and GLS were more preserved in patients who received early intravenous metoprolol, supporting the rationale to use beta-blocker intravenously in clinically stable STEMI patients before primary PCI.1

Time course of LV structural and functional changes after STEMI

In the healing process of acute myocardial infarction important structural and functional changes take place in both the infarct area and the remote zone. 10 Several studies have focused on LV remodeling after acute myocardial infarction.¹⁵⁻¹⁷ In a large prospective STEMI registry including 507 patients treated with primary PCI and imaged with CMR at 1 week and 6 months, LV end-diastolic volume increased (from 79±21 mL/m² to 81±23 mL/m²; P=0.06) and LV end-systolic volume decreased (from 41±19 mL/m² to 39±21 mL/m²; P=0.02) over time. ¹⁶ This resulted in a significant increase in LVEF (from 50±12% to 54±13%, respectively; P<0.001). In the present study including a homogenous population with anterior STEMI patients treated with primary PCI, LV end-diastolic and LV end-systolic volumes both increased significantly over time in patients receiving early intravenous metoprolol as well as in control patients (Table 3, Figure 2 and 3). However, the increase was proportionally larger for LV end-diastolic volume than for LV end-systolic volume, resulting in an increase in LVEF. Furthermore, several authors have reported a reduction in infarct size, assessed with LGE CMR in STEMI patients treated with primary PCI. 16,18 Engblom et al. 18 showed a progressive decrease of LGE, expressed as the percentage of total LV mass, from days 1, 7, 42 to 182; however, there was no significant additional reduction of hyperenhanced myocardium at 1 year. The LGE reduction occurred predominantly during the first week after infarction (63% of the total 1-year reduction). In addition, Bodi et al. 16 reported significant reduction of LGE from 1 week to 6 months after STEMI (21±14% and 17±12%, respectively; P<0.001). This is in line with the results of the present study, which also demonstrated a decrease in LV hyperenhancement from 1 week to 6 months post-infarction.

In addition, the present study evaluated LV strain with feature-tracking CMR. LV strain has been extensively studied with speckle tracking echocardiography after acute myocardial infarction. On the other hand, global LV strain with CMR after myocardial infarction has been less extensively evaluated, but a few studies investigated the time changes of regional LV strain, using different myocardial tagging techniques. Kidambi *et al.* Showed an improvement of infarct zone peak systolic circumferential strain from day 2 to day 90 in 39 patients after STEMI treated with primary PCI, using complementary spatial modulation of magnetization myocardial tagging technique. Neizel *et al.* demonstrated an improvement in peak systolic circumferential strain in the myocardial segments with >50% transmural LGE (P<0.05) with strain-encoded imaging. The present study is, however, the first to assess the time course of GCS and GLS in a large anterior STEMI population with feature-tracking CMR. We demonstrated an overall improvement of 3.2% of GCS and GLS between 1-week and 6-month follow-up (P<0.001 for both).

The effect of metoprolol on long-term results

The results of the METOCARD-CNIC trial have shown long-term benefit of early intravenous metoprolol after acute anterior STEMI.⁵ Patients who received early intravenous metoprolol had smaller LV end-systolic volumes and more preserved LVEF at 6 months after STEMI, however there were no statistically significant differences in LGE-assessed infarct size between both treatment arms. In the present analysis, GCS and GLS showed a tendency towards more preserved values in the metoprolol group, but the differences did not reach the level of statistical significance. These results suggest that GCS and GLS are more closely related to myocardial infarct size, assessed with LGE CMR, than to changes in LV volumes, described by LVEF. This is in line with the published reports, showing that GLS with echocardiography is a better predictor of LGE-assessed infarct size compared to LVEF, whether measured in the acute phase after revascularization or at follow-up.^{22,23}

The different effects of metoprolol on GCS and GLS between 1-week and 6-month follow-up might be explained by the kinetics of the healing process of myocardial infarction. Edema is a very dynamic process during the first week after myocardial infarction,²⁴ and strain closely associates with its intensity and volume.²⁰ Moreover, cardioprotective therapies may affect the extent and intensity of post-myocardial infarction edema.²⁵ We may reasonably assume that the differences in LV GCS and GLS between both treatment arms were more pronounced in the acute phase because of a blunted edematous reaction in metoprolol treated patients as compared to control patients and have diluted at 6-month follow-up due an overall large resorption of edema and necrotic tissue.^{26,27}

Importantly however, when dividing the overall cohort of patients in quartiles of GCS and GLS, there was a significantly fewer number of patients receiving early intravenous metoprolol in the first GCS and GLS quartile (i.e. the worst LV systolic function), both at 1 week and at 6 months after STEMI (Figure 4). This shows that early metoprolol administration has a long-term beneficial effect on the healing process of STEMI and prevents severe LV systolic dysfunction. Our results support the use of early intravenous metoprolol in STEMI patients without contraindications to beta-blockers undergoing primary PCI.

Study limitations

Feature-tracking is a novel technique to assess LV strain with CMR. Recommendations on how to perform feature-tracking analysis are lacking, there are no accepted standard reference values for LV strain and the agreement between different vendors of feature-tracking software is

largely unknown. ²⁸ However, LV strain with feature-tracking CMR has shown to closely correlate with myocardial tagging and speckle tracking echocardiography and demonstrated superior intra- and inter-observer variability compared to both methods. ^{29,30} Furthermore, evaluation of LV strain was not a predefined study endpoint of the METOCARD-CNIC trial. Of the initial 202 patients who underwent 2 CMR studies in the METOCARD-CNIC trial, 5 patients were excluded from the LV strain analysis (7 patients from the analysis of GLS at 6 months) due to poor CMR cine image quality, which may have influenced our results. However, 98% (97%) feasibility of strain assessment with feature-tracking CMR is similar to what has been described before. ^{29,30}

CONCLUSION

Early intravenous metoprolol is associated with improved LV strain at 1 week after the acute anterior STEMI. Furthermore, early intravenous metoprolol is associated with less patients having worst LV systolic function at 1-week and at 6-month follow-up, compared with control patients. In conclusion, early metoprolol administration before primary PCI reduces the incidence of severe LV systolic dysfunction, both at short- and long-term follow-up as evaluated by feature-tracking CMR.

REFERENCES

- Ibanez B, James S, Agewall S, Antunes MJ, Bucciarelli-Ducci C, Bueno H *et al.* 2017 ESC Guidelines for the management of acute myocardial infarction in patients presenting with ST-segment elevation: The Task Force for the management of acute myocardial infarction in patients presenting with ST-segment elevation of the European Society of Cardiology (ESC). *Eur Heart J* 2018;39(2):119-77.
- O'Gara PT, Kushner FG, Ascheim DD, Casey DE, Jr., Chung MK, de Lemos JA et al. 2013 ACCF/AHA
 guideline for the management of ST-elevation myocardial infarction: executive summary: a report
 of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. J Am Coll Cardiol 2013;61(4):485-510.
- Garcia-Prieto J, Villena-Gutierrez R, Gomez M, Bernardo E, Pun-Garcia A, Garcia-Lunar I et al. Neutrophil stunning by metoprolol reduces infarct size. Nat Commun 2017;8:14780.
- 4. Ibanez B, Macaya C, Sanchez-Brunete V, Pizarro G, Fernandez-Friera L, Mateos A et al. Effect of early metoprolol on infarct size in ST-segment-elevation myocardial infarction patients undergoing primary percutaneous coronary intervention: the Effect of Metoprolol in Cardioprotection During an Acute Myocardial Infarction (METOCARD-CNIC) trial. Circulation 2013;128(14):1495-503.
- 5. Pizarro G, Fernandez-Friera L, Fuster V, Fernandez-Jimenez R, Garcia-Ruiz JM, Garcia-Alvarez A et al. Long-term benefit of early pre-reperfusion metoprolol administration in patients with acute myocardial infarction: results from the METOCARD-CNIC trial (Effect of Metoprolol in Cardioprotection During an Acute Myocardial Infarction). J Am Coll Cardiol 2014;63(22):2356-62.
- Stanton T, Leano R, Marwick TH. Prediction of all-cause mortality from global longitudinal speckle strain:
 comparison with ejection fraction and wall motion scoring. Circ Cardiovasc Imaging 2009;2(5):356-64.
- Kraigher-Krainer E, Shah AM, Gupta DK, Santos A, Claggett B, Pieske B et al. Impaired systolic function by strain imaging in heart failure with preserved ejection fraction. J Am Coll Cardiol 2014;63(5):447-56.
- Schuster A, Hor KN, Kowallick JT, Beerbaum P, Kutty S. Cardiovascular Magnetic Resonance Myocardial Feature Tracking: Concepts and Clinical Applications. Circ Cardiovasc Imaging 2016;9(4):e004077.
- 9. Ibanez B, Fuster V, Macaya C, Sanchez-Brunete V, Pizarro G, Lopez-Romero P et al. Study design for the "effect of METOprolol in CARDioproteCtion during an acute myocardial InfarCtion" (METO-CARD-CNIC): a randomized, controlled parallel-group, observer-blinded clinical trial of early pre-reperfusion metoprolol administration in ST-segment elevation myocardial infarction. Am Heart J 2012;164(4):473-80.e5.
- 10. Jugdutt BI. Ventricular remodeling after infarction and the extracellular collagen matrix: when is enough enough? *Circulation* 2003;108(11):1395-403.

- Chen ZM, Pan HC, Chen YP, Peto R, Collins R, Jiang LX et al. Early intravenous then oral metoprolol in 45,852 patients with acute myocardial infarction: randomised placebo-controlled trial. Lancet 2005;366(9497):1622-32.
- 12. Antoni ML, Mollema SA, Delgado V, Atary JZ, Borleffs CJ, Boersma E *et al.* Prognostic importance of strain and strain rate after acute myocardial infarction. *Eur Heart J* 2010;31(13):1640-7.
- Gavara J, Rodriguez-Palomares JF, Valente F, Monmeneu JV, Lopez-Lereu MP, Bonanad C et al. Prognostic Value of Strain by Tissue Tracking Cardiac Magnetic Resonance After ST-Segment Elevation Myocardial Infarction. *JACC Cardiovasc Imaging* 2018;11(10):1448-57.
- Eitel I, Stiermaier T, Lange T, Rommel KP, Koschalka A, Kowallick JT et al. Cardiac Magnetic Resonance Myocardial Feature Tracking for Optimized Prediction of Cardiovascular Events Following Myocardial Infarction. JACC Cardiovasc Imaging 2018;11(10):1433-44.
- Joyce E, Hoogslag GE, Leong DP, Debonnaire P, Katsanos S, Boden H et al. Association between left ventricular global longitudinal strain and adverse left ventricular dilatation after ST-segment-elevation myocardial infarction. Circ Cardiovasc Imaging 2014;7(1):74-81.
- Bodi V, Monmeneu JV, Ortiz-Perez JT, Lopez-Lereu MP, Bonanad C, Husser O et al. Prediction of Reverse Remodeling at Cardiac MR Imaging Soon after First ST-Segment-Elevation Myocardial Infarction: Results of a Large Prospective Registry. Radiology 2016;278(1):54-63.
- Ganame J, Messalli G, Masci PG, Dymarkowski S, Abbasi K, Van de Werf F et al. Time course of infarct
 healing and left ventricular remodelling in patients with reperfused ST segment elevation myocardial infarction using comprehensive magnetic resonance imaging. Eur Radiol 2011;21(4):693-701.
- Engblom H, Hedstrom E, Heiberg E, Wagner GS, Pahlm O, Arheden H. Rapid initial reduction of hyperenhanced myocardium after reperfused first myocardial infarction suggests recovery of the peri-infarction zone: one-year follow-up by MRI. Circ Cardiovasc Imaging 2009;2(1):47-55.
- 19. Antoni ML, Mollema SA, Atary JZ, Borleffs CJ, Boersma E, van de Veire NR *et al.* Time course of global left ventricular strain after acute myocardial infarction. *Eur Heart J* 2010;31(16):2006-13.
- Kidambi A, Mather AN, Swoboda P, Motwani M, Fairbairn TA, Greenwood JP et al. Relationship between myocardial edema and regional myocardial function after reperfused acute myocardial infarction: an MR imaging study. Radiology 2013;267(3):701-8.
- Neizel M, Korosoglou G, Lossnitzer D, Kuhl H, Hoffmann R, Ocklenburg C et al. Impact of systolic and diastolic deformation indexes assessed by strain-encoded imaging to predict persistent severe myocardial dysfunction in patients after acute myocardial infarction at follow-up. J Am Coll Cardiol 2010;56(13):1056-62.

- Vartdal T, Brunvand H, Pettersen E, Smith HJ, Lyseggen E, Helle-Valle T et al. Early prediction of infarct size by strain Doppler echocardiography after coronary reperfusion. J Am Coll Cardiol 2007;49(16):1715-21.
- 23. Gjesdal O, Helle-Valle T, Hopp E, Lunde K, Vartdal T, Aakhus S *et al.* Noninvasive separation of large, medium, and small myocardial infarcts in survivors of reperfused ST-elevation myocardial infarction: a comprehensive tissue Doppler and speckle-tracking echocardiography study. *Circ Cardiovasc Imaging* 2008;1(3):189-96.
- 24. Fernandez-Jimenez R, Barreiro-Perez M, Martin-Garcia A, Sanchez-Gonzalez J, Aguero J, Galan-Arriola C *et al.* Dynamic Edematous Response of the Human Heart to Myocardial Infarction: Implications for Assessing Myocardial Area at Risk and Salvage. *Circulation* 2017;136(14):1288-300.
- Fernandez-Jimenez R, Galan-Arriola C, Sanchez-Gonzalez J, Aguero J, Lopez-Martin GJ, Gomez-Talavera S et al. Effect of Ischemia Duration and Protective Interventions on the Temporal Dynamics of Tissue Composition After Myocardial Infarction. Circ Res 2017;121(4):439-50.
- Dall'Armellina E, Karia N, Lindsay AC, Karamitsos TD, Ferreira V, Robson MD et al. Dynamic changes
 of edema and late gadolinium enhancement after acute myocardial infarction and their relationship
 to functional recovery and salvage index. Circ Cardiovasc Imaging 2011;4(3):228-36.
- Fieno DS, Hillenbrand HB, Rehwald WG, Harris KR, Decker RS, Parker MA et al. Infarct resorption, compensatory hypertrophy, and differing patterns of ventricular remodeling following myocardial infarctions of varying size. J Am Coll Cardiol 2004;43(11):2124-31.
- Claus P, Omar AM, Pedrizzetti G, Sengupta PP, Nagel E. Tissue Tracking Technology for Assessing Cardiac Mechanics: Principles, Normal Values, and Clinical Applications. *JACC Cardiovasc Imaging* 2015;8(12):1444-60.
- Khan JN, Singh A, Nazir SA, Kanagala P, Gershlick AH, McCann GP. Comparison of cardiovascular magnetic resonance feature tracking and tagging for the assessment of left ventricular systolic strain in acute myocardial infarction. *Eur J Radiol* 2015;84(5):840-8.
- Obokata M, Nagata Y, Wu VC, Kado Y, Kurabayashi M, Otsuji Y et al. Direct comparison of cardiac magnetic resonance feature tracking and 2D/3D echocardiography speckle tracking for evaluation of global left ventricular strain. Eur Heart J Cardiovasc Imaging 2016;17(5):525-32.