



Universiteit  
Leiden  
The Netherlands

## **Do early-life eating habits predict later autistic traits? Results from a population-based study**

van't Hof, M.; Ester, W.A.; Berckelaer-Onnes, I. van; Hillegers, M.H.J.; Hoek, H.W.; Jansen, P.W.

### **Citation**


Van't Hof, M., Ester, W. A., Berckelaer-Onnes, I. van, Hillegers, M. H. J., Hoek, H. W., & Jansen, P. W. (2021). Do early-life eating habits predict later autistic traits?: Results from a population-based study. *Appetite*, 156. doi:10.1016/j.appet.2020.104976

Version: Publisher's Version

License: [Creative Commons CC BY 4.0 license](https://creativecommons.org/licenses/by/4.0/)

Downloaded from: <https://hdl.handle.net/1887/3276789>

**Note:** To cite this publication please use the final published version (if applicable).



## Research report

## Do early-life eating habits predict later autistic traits? Results from a population-based study

Maarten van 't Hof<sup>a,b,c,d</sup>, Wietske A. Ester<sup>c,d,e</sup>, Ina van Berckelaer-Onnes<sup>c,f</sup>, Manon H. J. Hillegers<sup>b</sup>, Hans W. Hoek<sup>d,g,h</sup>, Pauline W. Jansen<sup>b,i,\*</sup><sup>a</sup> Generation R, Erasmus University Medical Center, Rotterdam, the Netherlands<sup>b</sup> Department of Child and Adolescent Psychiatry/Psychology, Erasmus University Medical Center, Rotterdam, the Netherlands<sup>c</sup> Sarr Expert Centre for Autism, Lucertis Child and Adolescent Psychiatry, Rotterdam, the Netherlands<sup>d</sup> Parnassia Psychiatric Institute, The Hague, the Netherlands<sup>e</sup> Department of Child and Adolescent Psychiatry, Curium-LUMC, Leiden University Medical Center, Oegstgeest, the Netherlands<sup>f</sup> Faculty of Social Sciences, Clinical Child and Adolescent Studies, University Leiden, Leiden, the Netherlands<sup>g</sup> Department of Psychiatry, University Medical Center Groningen, University of Groningen, Groningen, the Netherlands<sup>h</sup> Department of Epidemiology, Mailman School of Public Health, Columbia University, New York, United States<sup>i</sup> Department of Psychology, Education and Child Studies, Erasmus University Rotterdam, Rotterdam, the Netherlands

## ARTICLE INFO

## Keywords:

Breastfeeding  
Eating behavior  
Autism spectrum disorder  
Autistic traits  
Infancy  
General population

## ABSTRACT

Eating problems are common among children with Autism Spectrum Disorder (ASD), but it is unknown to what extent infant eating behavior is associated with later autistic traits. As eating behavior is currently not included in ASD screening instruments, it is important to evaluate whether infant eating behavior predicts later autistic traits and might therefore be used to enhance the early detection of ASD. We investigated the association of breastfeeding and eating behavior during infancy with later autistic traits in the population-based Generation R cohort. We included 3546 mother-child dyads with maternal reports on feeding and eating at age two months and autistic traits at six years. Eating behavior was assessed with seven items on specific eating habits and the Social Responsiveness Scale was used to evaluate autistic traits. Covariates included child sex, and maternal psychopathology and autistic traits. Linear regression analyses showed that being formula fed at two months was associated with a higher autistic trait score at six years (adjusted  $B = 0.07$ ; 95% CI: 0.00–0.14). Children who were drinking only small quantities (adjusted  $B = 0.17$ , 95% CI: 0.04–0.30) and were hungry/not satisfied (adjusted  $B = 0.23$ , 95% CI: 0.08–0.39) at age two months also had a higher autistic traits score at age six years. We found no interactions with sex or breastfeeding. This study shows that eating behavior during infancy is related with autistic traits in childhood. Although the associations were fairly small, these findings suggest that early-life eating problems might be relevant for early detection of ASD and a potential addition to ASD-specific screening instruments.

## 1. Introduction

Autism Spectrum Disorder (ASD) is a complex neurodevelopmental disorder that is present from early life onwards. It is characterized by persistent deficiencies in communication and social interaction, and restricted, repetitive patterns of behaviors, interests and activities. Although several symptoms are already present in the early developmental period, some symptoms manifest itself only later in life (American Psychiatric Association, 2013). ASD can be reliably diagnosed as young as 24 months (Johnsson & Myers, 2007), but the current mean

age at diagnosis ranges from 38 to 120 months (Daniels & Mandell, 2014). There is an urgency to identifying children with ASD, as particularly early interventions can improve language and cognitive abilities (Dawson & Burner, 2011) and result in better long-term outcomes across the school age years (Clark et al., 2018) compared to interventions at a later stage. Detection of the earliest signs of ASD can enhance timely identification and diagnosis, which then permits early treatment to achieve the most optimal long-term outcomes for children with ASD.

There are several widely used instruments available for ASD screening. The Modified Checklist for Autism in Toddlers (M-CHAT(R/

\* Corresponding author. Department of Child and Adolescent Psychiatry/Psychology, Erasmus University Medical Center, Rotterdam, the Netherlands.

E-mail address: [p.w.jansen@erasmusmc.nl](mailto:p.w.jansen@erasmusmc.nl) (P.W. Jansen).

<https://doi.org/10.1016/j.appet.2020.104976>

Received 25 February 2020; Received in revised form 17 September 2020; Accepted 17 September 2020

Available online 21 September 2020

0195-6663/© 2020 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

F); Robins et al., 2001; Diana L. Robins et al., 2014) is currently the most commonly used screening instrument (Øien et al., 2019). Other instruments like the Social Communication Questionnaire (SCQ; Rutter et al., 2003) and the Early Screening of Autistic Traits Questionnaire (ESAT/CoSoS in Dutch; Swinkels et al., 2006) are also frequently used. These screening instruments focus primarily on ASD deficiencies described in the DSM-5 criteria, but also include items specific to other ASD-related deficiencies like having an unusual response to sensory stimuli (Robins et al., 2001; Rutter et al., 2003; Swinkels et al., 2006). Although eating problems are often seen among children with ASD, they are currently not included in ASD screening instruments.

Eating problems –like severe or prolonged picky eating and food neophobia (fear of new/unknown foods)– are seen in 46–89% of children with ASD (Ledford & Gast, 2006) compared to 6–50% of children in the general population (Taylor et al., 2015). Food neophobia is two to three times more common in children with ASD than in non-ASD children (Wallace et al., 2018). In previous studies in the general population, we also identified eating problems in early childhood as a possible early sign of ASD (Cardona Cano et al., 2016). While eating problems may already be expressed early in life, to our knowledge there are no studies on the association of eating behavior in infancy with ASD, except for research on breastfeeding (Tseng et al., 2019).

Absence of breastfeeding has been examined as a risk factor for ASD numerous times. A recent meta-analysis indicated that breastfeeding may protect against developing ASD and suggests genetic factors, content of breastmilk, and skin-to-skin contact as potential causal pathways for this protective effect (Tseng et al., 2019). However, eating problems can also complicate breastfeeding, with unsuccessful breastfeeding thus potentially being indicative of other eating problems seen in children with ASD. Current research calls for prospective studies to confirm the suspected protective effect of breastfeeding on the development of ASD (Tseng et al., 2019).

We investigated the association between eating behavior in infancy, differentiated into breastfeeding and feeding habits, with later autistic traits in a prospective, population-based cohort. Knowing whether infant eating behavior predicts later autistic traits could be useful to enhance screening for autism at an early stage. We hypothesize that breastfeeding is associated with fewer autistic traits in later childhood, while problematic eating behaviors in infants is prospectively associated with a higher autistic traits score.

## 2. Method

### 2.1. Data source

This study uses data from the Generation R Study, a population-based, longitudinal cohort study from the fetal stage onwards; the cohort is based in Rotterdam, the Netherlands. In short, all pregnant women living in Rotterdam, with an expected delivery date between April 2002 and January 2006, were invited to participate in the study (participation rate: 61%). The Generation R study design and population has been described in detail (Jaddoe et al., 2012). Written informed consent was obtained from all participants. The study was approved by the Medical Ethical Committee of the Erasmus Medical Center, Rotterdam.

### 2.2. Participants

Our study population includes all parents of 6625 children who gave consent for participation in the postnatal phase of the Generation R study (Fig. 1). We excluded children with missing data on eating behavior ( $n = 1779$ ) or breastfeeding ( $n = 69$ ) at two months and those without information on autistic traits at 6 years ( $n = 1231$ ). This resulted in a study sample of 3546 mother-child dyads.




Fig. 1. Flow diagram of the study population.

### 2.3. Measures

#### 2.3.1. Breastfeeding

Information about feeding mode was obtained by a maternal postal questionnaire when children were two months old. We assessed if infants were breastfed (yes/no) and whether they received breastfeeding only, a combination of breast- and formula feeding, or formula feeding only at the time the questionnaire was filled out.

#### 2.3.2. Eating behavior

Eating behavior at age two months was assessed in the same maternal questionnaire using seven single items related to eating and feeding difficulties. These assessed relatively common difficulties and behavior, and were partly based on previous studies that also used single items on infant feeding (Micali et al., 2009; Wolke et al., 1994). Parents (mostly mothers, 93.7%) responded; they were asked if their infant exhibited one or more of the following feeding habits: 'Drinks slowly', 'Drinks only small quantities', 'Drinks very greedily', 'Is hungry or not satisfied', 'Spits up a lot', 'Regurgitates mouthfuls of food' and 'Refuses breastfeeding'. Response options were Yes or No.

#### 2.3.3. Autistic traits

Autistic traits were assessed using maternal reports of the validated Social Responsiveness Scale (SRS) (Constantino, 2002) when children were six years old. The SRS is an autism screening questionnaire providing a quantitative measure of (sub)clinical autistic traits related to social cognition and communication. The psychometric properties are

good, including a test-retest reliability of 0.88 and a two-year stability correlation of 0.83 (Constantino et al., 2003). The original questionnaire was reduced to an 18-item shortened SRS to minimize subject burden, with selected items encompassing all DSM-IV autism domains. The shortening was done in consultation with the SRS test developer, as described previously (Román et al., 2013). Our SRS short form consists of the Social Cognition (5 items), Social Communication (8 items) and Mannerism (5 items) subscales. Items assessed traits in the past three months and were scored on a 4-point scale (not true, sometimes true, often true, almost always true). The 18-item shortened SRS shows high correlations of 0.93–0.99 with the full scale, as described in Blanken et al. (2015). Diagnostic validity studies comparing the SRS with the Autism Diagnostic Interview- Revised (ADI-R) and Autism Diagnostic Observation Scale (ADOS) found acceptable sensitivity and specificity rates of 70–80% (Hampton & Strand, 2015).

To identify children with an ASD diagnosis, we examined medical records using a multifaceted screening procedure (White et al., 2018). First, cut-off scores for the SRS were used to identify children with elevated ASD traits. Second, to rule out false-negatives, children scoring in the top 15% of the total score of the Child Behavior Checklist (CBCL/1½-5) (Achenbach & Rescorla, 2001) at age six years were then screened using the 40-item Social Communication Questionnaire (SCQ), as reported by a parent (Berument et al., 1999). Additionally, psychiatric diagnoses and treatment were routinely assessed at all contact moments between ages 6–9 years (health center visits and questionnaires). If children scored positive in one or more of these three information sources, the medical records at their general practitioners were examined. If an ASD diagnosis was confirmed through the medical records, the child was considered to be a clinically confirmed case of ASD.

#### 2.3.4. Covariates

We included several child and family covariates based on previous studies (Blanken et al., 2015; Cardona Cano et al., 2016; Jansen et al., 2012). Information about sex, birth weight and gestational age at birth was obtained from midwife- and hospital registries. Maternal age, educational level and ethnicity were obtained by postal questionnaire. We used the Brief Symptom Inventory (BSI) to assess maternal psychopathology during pregnancy. The BSI is a validated 53-item self-reported questionnaire assessing diverse psychiatric problems (Derogatis & Melisaratos, 1983). We also used the Autism-Spectrum Quotient (AQ-short) to assess maternal autistic traits. Although this questionnaire was distributed when the children were 9 years old, we assumed that maternal autistic traits were fairly stable across adulthood. The AQ-short is a 28-item self-report assessing autistic traits on the scales 'social behavioral difficulties' and 'fascination for numbers/patterns', which sum into a total score (minimum = 28, maximum = 112). A cut-off score of >65 is suggested for a quick screening of autistic traits in a clinical setting (Hoekstra et al., 2011).

#### 2.4. Statistical analyses

First, separate linear regressions were conducted to evaluate the relationship between breastfeeding at two months and autistic traits at six years. Three models were evaluated: model 1 was an unadjusted model, model 2 included the covariates maternal age, education and psychopathology, and model 3 additionally included measures of maternal autistic traits. Only covariates that altered the unadjusted beta by more than 10% were included in model 2. Maternal autistic traits were included to demonstrate the possible influence of a familial predisposition for ASD. We also tested sex interactions and finally evaluated the association between breastfeeding at two months and clinically confirmed ASD diagnosis by logistic regression.

In a second set of analyses, linear regressions were conducted to evaluate the relationship between eating behavior at two months and autistic traits at six years, with separate regressions for each type of eating behavior. The same three adjusted and unadjusted models were

evaluated for each eating behavior, using a similar method to that described above. We tested for possible interactions of sex with eating behavior, as well as for breastfeeding with eating behavior.

We estimated missing values on confounders (ranging from 0% missing for sex to 32.1% for maternal autistic traits) using multiple imputation techniques. All variables included in the analyses were used to estimate the missing values (Graham, 2009). Regressions were conducted on the imputed data and reported estimates are the pooled results of 30 imputed datasets. All statistical analyses were performed with SPSS 21.

### 3. Results

#### 3.1. Non-response analyses

We compared the characteristics of participating mother-child dyads ( $n = 3546$ ) with those excluded from the study due to missing data on determinants or outcome ( $n = 3079$ ). Data were more often missing in mothers who were younger, had a lower education level, were of non-Dutch ethnicity, and who had higher psychopathology and autistic traits scores (all  $p$ -values <.01). Excluded children had a lower birth weight, were more often boys, and more often received no breastfeeding than included children (all  $p$ -values <.01).

#### 3.2. Sample characteristics

Child and maternal characteristics of the study sample are presented in Table 1. In total, 51.0% of children were girls and 68.8% received

**Table 1**

General characteristics of parent-child dyads from Generation R cohort ( $n = 3546$ ).

	Percentage or median <sup>a</sup>	% missing
Child characteristics		
Sex		0.0
Girl, %	51.0	
Age at assessment (months)		
Feeding and eating behavior	2.8 (0.4–6.0)	7.4
Social Responsiveness Scale	71.8 (58.7–106.8)	0.0
Gestational age at birth	40.1 (27.1–43.4)	0.0
Birth weight (grams)	3480 (780–5610)	0.0
Breastfeeding at 2 months		0.0
Yes, %	68.8	0.0
Breastfeeding type at 2 months		0.0
Breastfeeding only, %	41.2	
Breastfeeding & formula feeding, %	27.7	
Formula feeding only, %	31.2	
Eating behavior at 2 months		
Drinks slowly, %	13.9	0.0
Drinks only small quantities, %	6.0	0.0
Drinks very greedily, %	39.1	0.0
Is hungry or not satisfied, %	3.9	0.0
Spits up a lot, %	12.2	0.0
Regurgitates mouthfuls of feed, %	32.0	0.0
Refuses to breastfeed, %	1.7	0.0
Autistic traits at 6 years	0.2 (0.0–2.5)	0.0
Maternal characteristics		
Age at intake (years)	31.9 (15.3–45.4)	0.0
Ethnicity		0.2
Dutch, %	68.0	
Other, %	31.9	
Education		2.7
Higher vocational education/university, %	59.5	
Lower vocational education, %	26.0	
Less than high school, %	11.8	
Psychopathology score	0.1 (0.0–2.7)	15.9
Autism traits score	50.0 (28.0–84.0)	32.1

<sup>a</sup> Values are percentages for categorical variables and medians (range) for continuous non-normally distributed variables.

breastfeeding at two months. More than two-thirds (68.0%) of the mothers had a Dutch background. The study sample included 50 children (1.4%) who had a GP-confirmed diagnosis of ASD. Of the mothers, 5.9% showed elevated levels of autistic traits (AQ-short score >65).

3.3. Prospective relationship between breastfeeding and autistic traits

Table 2 shows that receiving formula feeding at two months was associated with a higher autistic traits score at six years (confounder adjusted B = 0.07; 95% CI: 0.00 to 0.14) than when receiving breastfeeding.

Particularly receiving only formula feeding was associated with a higher autistic traits score at six years (confounder adjusted B = 0.08; 95% CI: 0.00 to 0.15), while receiving mixed formula and breastfeeding was not significantly associated with autistic traits score. Adjustment for maternal age, education and psychopathology led to a change in the association between breastfeeding and autistic traits of more than 10%, while the additional inclusion of maternal autistic traits in the model did not notably affect this association. We found no interactions of breastfeeding and sex on autistic traits.

Sensitivity analyses showed that children who were breastfed at two months did not differ in their odds of a later ASD diagnosis, compared with children who were not breastfed (OR = 0.81; 95% CI: 0.44 to 1.48, p = .50).

3.4. Prospective relationship between eating behavior and autistic traits

Table 3 indicates that infants who drank only small quantities at two months had a higher autistic traits score at six years (confounder adjusted B = 0.17; 95% CI: 0.04 to 0.30) compared to infants who did not drink small quantities. Results further show that the infant eating behavior 'being hungry or not being satisfied' at two months was associated with later autistic traits (confounder adjusted B = 0.23; 95% CI: 0.08 to 0.39). The inclusion of maternal autistic traits in the models did not notably affect the association between infant eating behavior and later autistic traits. The other eating behaviors were not associated with later autistic traits. No interaction of sex or breastfeeding with eating behavior was found.

4. Discussion

This study –in a general population sample– shows that infants who were formula fed, drank only small quantities, or who were often hungry or not satisfied at the age of two months had a higher autistic traits score in childhood at age 6 years.

Table 2  
Prospective association between breastfeeding and child autistic traits (n = 3546).

	B for SRS total score at six years (95% CI)		
	Model 1	Model 2	Model 3
Breastfeeding at 2 months (reference response = yes)	0.15 (0.02; 0.22)**	0.07 (0.00; 0.14)*	0.07 (0.00; 0.14)*
Type of feeding at 2 months			
Breastfeeding only	Reference	Reference	Reference
Breastfeeding & formula feeding	0.05 (-0.02; 0.13)	0.03 (-0.05; 0.10)	0.02 (-0.05; 0.09)
Formula feeding only	0.17 (0.10; 0.24)**	0.08 (0.01; 0.16)*	0.08 (0.00; 0.15)*

Model 1: unadjusted.

Model 2: adjusted for maternal age, education and psychopathology.

Model 3: adjusted for maternal age, education, psychopathology and autistic traits.

Note: We found no significant interaction with sex.

\*\*p < .01, \*p < .05.

Table 3

Prospective association between infant eating behavior and child autistic traits (n = 3546).

Eating behavior at age 2 months (reference response = no)	B for SRS total score at six years (95% CI)		
	Model 1	Model 2	Model 3
Drinks slowly	0.12 (0.03; 0.21)*	0.07 (-0.01; 0.16)	0.06 (-0.02; 0.15)
Drinks only small quantities	0.28 (0.15–0.40)**	0.20 (0.07–0.33)**	0.17 (0.04; 0.30)**
Drinks very greedily	0.03 (-0.04; 0.09)	0.00 (-0.06; 0.07)	0.01 (-0.05; 0.07)
Is hungry or not satisfied	0.40 (0.23–0.56)**	0.24 (0.09; 0.40)**	0.23 (0.08; 0.39)**
Spits up a lot	0.08 (-0.01; 0.18)	0.06 (-0.03; 0.16)	0.05 (-0.04; 0.14)
Regurgitates mouthfuls of feed	-0.05 (-0.12; 0.01)	-0.04 (-0.10; 0.03)	-0.03 (-0.10; 0.03)
Refuses to breastfeed	0.05 (-0.20; 0.29)	0.02 (-0.21; 0.26)	-0.00 (-0.23; 0.23)

Model 1: unadjusted.

Model 2: adjusted for maternal age, education and psychopathology.

Model 3: adjusted for maternal age, education, psychopathology and autistic traits.

\*\*p < .01, \*p < .05.

Note: We found no significant interactions between sex or breastfeeding (yes/no) with eating behavior.

4.1. Breastfeeding

Our results on breastfeeding are in line with previous studies suggesting a negative association between breastfeeding and later ASD (Tseng et al., 2019). Our study contributes to the clinical findings that breastfeeding is associated with a lower degree of autistic traits at a subclinical level in a large general population sample. Although in our subgroup of children with an ASD diagnosis the association pointed in the same direction, caution is needed in drawing firm conclusions since this result was not significant, mostly likely due to the small number of children with a clinically confirmed ASD diagnosis (n = 50).

Although it is often suggested that breastfeeding and ASD are causally linked, these potential causal pathways are still quite hypothetical. Genetic factors may well be implicated since autism has a strong genetic basis, as illustrated by a high concordance rate (64–91%) seen in twin studies (Tick et al., 2016). Moreover, literature indicates that other maternal factors may also play a role in the association between ASD and breastfeeding. Multiple studies indicated that general maternal psychopathology is associated with shorter breastfeeding duration (Boucher et al., 2017; Tavoulari et al., 2016). Also, the broader autism phenotype is related to elevated depressive symptomatology (Ingersoll & Hambrick, 2011) and socio-affective impairment (Berthoz et al., 2013) which can negatively affect breastfeeding duration (Falsett et al., 2019; Henderson et al., 2003). Nonetheless, our results show that psychopathology and particularly the level of autistic traits of mothers only explain a small part of the association between breastfeeding and later autistic traits in children. This is in line with a previous study indicating that the broader autism phenotype does not fully explain the association between breastfeeding and ASD (Soke et al., 2019). We should also consider the possibility of residual confounding. For example, maternal under- and over-responsivity to sensory stimuli-which are common in people with ASD (Lai et al., 2011) and influence the duration that women give breastfeeding (Britton et al., 2006; Tharner et al., 2012) - were not evaluated in our study.

Also, multiple child factors may contribute to the association between breastfeeding and ASD. Sensory problems that have been found in infants at high risk for ASD (Bryson et al., 2007; Germani et al., 2014; Mulligan & White, 2012) could affect the initiation and/or continuation

of breastfeeding, suggesting a reverse causal pathway. This may be due to the fact that breastfeeding provides a richer variation in oral sensory stimulation (taste) compared to formula milk (Beauchamp & Mennella, 2011), because the taste of breastmilk (e.g. bitterness) is affected by the mother's consumption (Mastorakou et al., 2019). Also, compared to other feeding methods, breastfeeding provides more close skin-to-skin contact (Liu et al., 2013) and frequent touch by the mother (Kuzela et al., 1990) which are possibly less pleasant for children with sensory problems. Finally, a dysregulated breastfeeding pattern of endless sucking in children with ASD (Lucas & Cutler, 2015) may lead to sore nipples and breastfeeding discontinuation (Ahluwalia et al., 2005; Li et al., 2008).

#### 4.2. Eating behavior

The current study showed that specific infant eating problems –particularly *drinking only small quantities* and *being hungry or not satisfied*– were associated with a higher autistic traits score later in childhood. This is in line with our previous study across middle childhood: in that study we found that autistic traits at six years were prospectively associated with elevated levels of picky eating and food responsiveness at ten years in both boys and girls. Among girls only, autistic traits at six years were also associated with elevated levels of emotional over- and under-eating at ten years (van 't Hof et al., 2019). Our current results add to the literature that the association between eating behavior and autistic traits seems to be present even from very early infancy. It is, however, important to emphasize that, although both studies show a prospective association between eating behavior and autistic traits, based on these findings we cannot draw any conclusions on causality.

There are several possible explanations for the reported association. Drinking only small quantities and being hungry or not satisfied are both linked with appetite self-regulation. Typically developing children show good self-regulation, also in feeding (Saltzman et al., 2018), but self-regulation difficulties have been reported among children with ASD (Gomez & Baird, 2005; Jahromi, 2017). It is possible that infant eating problems are an early expression of the self-regulation difficulties typically seen in ASD.

Delays in the development of early motor skills are also often reported in children with ASD (Chinello et al., 2018) and can be related to eating behavior problems in infancy. Specific impairment in oral motor function can affect eating behavior by limited bolus control, and/or manipulation and transit of liquids and solids (Weir et al., 2007), while impairments in oral and pharyngeal sensory-motor functioning may also inhibit feeding skills (Goday et al., 2019). Finally, sensory processing deficits are another common problem in children with ASD (Leekam et al., 2007) and these have been linked to eating problems in ASD (Nadon et al., 2011).

In contrast to our previous study evaluating the association between eating behavior and autistic traits in middle childhood (van 't Hof et al., 2019), we found no sex differences in the current study. This is most likely because sex differences were previously found solely in the association between emotional eating and autistic traits (van 't Hof et al., 2019). Emotional eating, however, is mostly shaped by shared environmental factors rather than genetic risk (Herle et al., 2018) and develops during childhood. So, potentially emotional eating problems are not yet present in infants, and even if they were present, such eating behavior would be very difficult to assess at a very young age.

#### 4.3. Implications

Information on eating problems in infancy can potentially be used by professionals as part of the screening for ASD in the general population. Although most early signs of ASD can only reliably be observed from age 1–2 years onwards, previous research has indicated several possible biological and behavioral markers of ASD in infancy. Research on biological markers found that early brain development (Hazlett et al., 2017)

may be indicative of ASD as early as six months of age. Furthermore, infant gaze patterns (Shic et al., 2014), reduced looking time (Falck-Ytter et al., 2013), less alternating gaze during interaction with adults (Thorup et al., 2018), and a decline in eye fixation (Jones & Klin, 2013) have all been identified as early behavioral markers of ASD using eye-tracking techniques.

However, these early biological and behavioral markers require expensive and invasive brain imaging and eye-tracking techniques that cannot be easily used for widespread screening of ASD. Our current results suggest that the addition of certain eating behaviors to ASD-specific screening instruments like the M-CHAT, SCQ and the CoSoS might be helpful in providing a more complete ASD phenotype. Such addition might also enhance the validity, and with that the screening properties, of these instruments. In addition, we suggest the standardized monitoring of infant eating behavior in well-baby clinics to increase the identification of eating and feeding problems by using standardized questionnaires such as the BEBQ or by the inclusion of eating behavior items in developmental screening instruments.

#### 4.4. Strengths and limitations

This study has several strengths. First, the large data set enabled us to assess the association between infant eating behavior and autistic traits prospectively and to control for possible confounding factors. We were also able to use validated instruments to assess child (SRS) and maternal (AQ-short) autistic traits in a general population setting, thereby optimizing validity and minimizing measurement bias. However, as infant eating behavior questionnaires, like the Baby Eating Behavior Questionnaire (BEBQ) (Llewellyn et al., 2011), were not available when the children in this study were 2 months old, we assessed a limited number of different eating behaviors using unvalidated, single items. Furthermore, we also did not assess the reasons why mothers stopped breastfeeding their child. More detailed information on eating and feeding behaviors could help us to better understand the underlying mechanisms.

Another limitation of our study was that mothers with a higher psychopathology and autistic traits score were more likely to be lost for follow-up. Considering the genetic component in ASD, children with relatively high levels of autistic traits may also have dropped out selectively. Although no data on autistic traits were available for the excluded sample, we expect the effects of drop-out to be fairly small based on results from another large cohort study (Nilsen et al., 2009). A final limitation was that although the prevalence of children in this study with an ASD diagnosis (1.4%) was comparable to the prevalence found in the United States (one in 59 children, 1.7%) (Baio et al., 2018), the number of children with a confirmed ASD diagnosis in the sample was still small, meaning we had insufficient power to detect statistically significant differences.

### 5. Conclusions

Our study shows that infant eating problems and the absence of breastfeeding at two months of age are associated with a higher autistic traits score in later childhood. Although the associations were fairly small, these findings suggest that efforts to detect autism at an early stage might benefit from assessing early-life eating problems, alongside other factors like social interaction deficits. We recommend complementing ASD-specific screening instruments with an assessment of early-life eating habits.

#### Authors' contributions

This study was conceived and designed by MH, WE, MHJH, HH and PJ. Analyses were performed and interpreted by MH, WE and PJ. MH, WE and PJ prepared the draft manuscript with input from IO, MHJH, HH. All authors read and approved the final manuscript.

## Funding

The first phase of the Generation R Study is made possible by financial support from the Netherlands Organization for Health Research and Development (ZonMW, grant number 10.000.1003).

## Ethical statement

### Ethics approval and consent to participate

Written informed consent was obtained from all participants. The study was approved by the Medical Ethical Committee of the Erasmus Medical Center, Rotterdam.

## Declaration of competing interest

PWJ received support from ZonMW (Mental Health Care Research Program, Fellowship 636320005). The authors declare that they have no further competing interest.

## Acknowledgements

The Generation R Study is run by Erasmus Medical Center in close collaboration with Erasmus University Rotterdam (School of Law and Faculty of Social Sciences), the Municipal Health Service Rotterdam area, the Rotterdam Homecare Foundation, and the Stichting Trombosedienst and Artsenlaboratorium Rijnmond, Rotterdam. We gratefully acknowledge the contribution of general practitioners, hospitals, midwives, and pharmacies in Rotterdam. We thank Jackie Senior for editing the manuscript.

## References

- Achenbach, T., & Rescorla, L. (2001). *Manual for ASEBA school-age forms & profiles. Youth, & Families: University of Vermont, Research Center for Children.*
- Ahluwalia, I. B., Morrow, B., & Hsia, J. (2005). Why do women stop breastfeeding? Findings from the pregnancy risk assessment and monitoring system. *Pediatrics*, *116*(6), 1408–1412. <https://doi.org/10.1542/peds.2005-0013>.
- American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders* (5th ed.). American Psychiatric Association. <https://doi.org/10.1176/appi.books.9780890425596.744053>.
- Baio, J., Wiggins, L., Christensen, D. L., Maenner, M. J., Daniels, J., Warren, Z., Kurzius-Spencer, M., Zahorodny, W., Robinson Rosenberg, C., White, T., Durkin, M. S., Imm, P., Nikolaou, L., Yeargin-Allsopp, M., Lee, L.-C., Harrington, R., Lopez, M., Fitzgerald, R. T., Hewitt, A., ... Dowling, N. F. (2018). Prevalence of autism spectrum disorder among children aged 8 years - autism and developmental disabilities monitoring network, 11 Sites, United States, 2014. *Morbidity and Mortality Weekly Report. Surveillance Summaries*, *67*(6), 1–23. <https://doi.org/10.15585/mmwr.ss6706a1>.
- Beauchamp, G. K., & Mennella, J. A. (2011). Flavor perception in human infants: Development and functional significance. *Digestion*, *83*(SUPPL.1), 1–6. <https://doi.org/10.1159/000323397>.
- Berthoz, S., Lalanne, C., Crane, L., & Hill, E. L. (2013). Investigating emotional impairments in adults with autism spectrum disorders and the broader autism phenotype. *Psychiatry Research*, *208*(3), 257–264. <https://doi.org/10.1016/j.psychres.2013.05.014>.
- Berument, S. K., Rutter, M., Lord, C., Pickles, A., & Bailey, A. J. (1999). Autism screening Questionnaire : Diagnostic validity autism screening questionnaire : Diagnostic validity. *The British Journal of Psychiatry*, *175*(5), 444–451. <https://doi.org/10.1192/bjp.175.5.444>.
- Blanken, L. M. E., Mous, S. E., Ghassabian, A., Muetzel, R. L., Schoemaker, N. K., El Marroun, H., Van Der Lugt, A., Jaddoe, V. W. V., Hofman, A., Verhulst, F. C., Tiemeier, H., & White, T. (2015). Cortical morphology in 6- to 10-year old children with autistic traits: A population-based neuroimaging study. *American Journal of Psychiatry*, *172*(5), 479–486. <https://doi.org/10.1176/appi.ajp.2014.14040482>.
- Boucher, O., Julvez, J., Guxens, M., Arranz, E., Ibarluzea, J., Sánchez De Miguel, M., Fernández-Somoano, A., Tardon, A., Rebagliato, M., Garcia-Esteban, R., O'Connor, G., Ballester, F., & Sunyer, J. (2017). Association between breastfeeding duration and cognitive development, autistic traits and ADHD symptoms: A multicenter study in Spain. *Pediatric Research*, *81*(3), 434–442. <https://doi.org/10.1038/pr.2016.238>.
- Britton, J. R., Britton, H. L., & Gronwaldt, V. (2006). Breastfeeding, sensitivity, and attachment. *Pediatrics*. <https://doi.org/10.1542/peds.2005-2916>.
- Bryson, S. E., Zwaigenbaum, L., Brian, J., Roberts, W., Szatmari, P., Rombough, V., & McDermott, C. (2007). A prospective case series of high-risk infants who developed autism. *Journal of Autism and Developmental Disorders*, *37*(1), 12–24. <https://doi.org/10.1007/s10803-006-0328-2>.
- Cardano Cano, S., Hoek, H. W., van Hoeken, D., de Barse, L. M., Jaddoe, V. W. V., Verhulst, F. C., & Tiemeier, H. (2016). Behavioral outcomes of picky eating in childhood: A prospective study in the general population. *Journal of Child Psychology and Psychiatry*, *57*(11), 1239–1246. <https://doi.org/10.1111/jcpp.12530>.
- Chinello, A., Di Gangi, V., & Valenza, E. (2018). Research in Developmental Disabilities Persistent primary reflexes affect motor acts : Potential implications for autism spectrum disorder. *Research in Developmental Disabilities*, *83*, 287–295. <https://doi.org/10.1016/j.ridd.2016.07.010>.
- Clark, M. L. E., Vinen, Z., Barbaro, J., & Dissanayake, C. (2018). School age outcomes of children diagnosed early and later with autism spectrum disorder. *Journal of Autism and Developmental Disorders*, *48*(1), 92–102. <https://doi.org/10.1007/s10803-017-3279-x>.
- Constantino, J. N. (2002). *Social responsiveness scale (SRS), manual.* Western Psychological services.
- Constantino, J. N., Davis, S. a., Todd, R. D., Schindler, M. K., Gross, M. M., Brophy, S. L., Metzger, L. M., Shoushtari, C. S., Splinter, R., & Reich, W. (2003). Validation of a Brief quantitative measure of autistic traits: Comparison of the social responsiveness scale with the Autism diagnostic Interview -revised. *Journal of Autism and Developmental Disorders*, *33*(4), 427–433. <https://doi.org/10.1023/A:1025014929212>.
- Daniels, A. M., & Mandell, D. S. (2014). Explaining differences in age at autism spectrum disorder diagnosis: A critical review. *Autism*, *18*(5), 583–597. <https://doi.org/10.1177/1362361313480277>.
- Dawson, G., & Burner, K. (2011). Behavioral interventions in children and adolescents with autism spectrum disorder: A review of recent findings. *Current Opinion in Pediatrics*, *23*(6), 616–620. <https://doi.org/10.1097/MOP.0b013e32834cf082>.
- Derogatis, L. R., & Melisaratos, N. (1983). The Brief symptom inventory: An introductory report. *Psychological Medicine*, *13*(3), 595–605. <https://doi.org/10.1017/S0033291700048017>.
- Falck-Ytter, T., Bölte, S., & Gredebäck, G. (2013). Eye tracking in early autism research. *Journal of Neurodevelopmental Disorders*, *5*(1), 28. <https://doi.org/10.1186/1866-1955-5-28>.
- Falsett, C. F., Santos, I. M. M. dos, & Vasconcellos, A. M. (2019). Interfering factors of the breastfeeding process in children bearing various health needs: Contributions to nursing. *Rev Fund Care Online*, *11*(5), 1278–1285. <https://doi.org/10.9789/2175-5361.2019.v11i5.1278-1285>.
- Germani, T., Zwaigenbaum, L., Bryson, S., Brian, J., Smith, I., Roberts, W., Szatmari, P., Roncadin, C., Sacrey, L. A. R., Garon, N., & Vaillancourt, T. (2014). Brief report: Assessment of early sensory processing in infants at high-risk of autism spectrum disorder. *Journal of Autism and Developmental Disorders*, *44*(12), 3264–3270. <https://doi.org/10.1007/s10803-014-2175-x>.
- Goday, P. S., Huh, S. Y., Silverman, A., Lukens, C. T., Dodrill, P., Cohen, S. S., Delaney, A. A. L., Feuling, M. B., Noel, A. A. R. J., Gisel, E., Kessler, D. B., Kraus de Camargo, O., Browne, J., & Phalen, J. A. (2019). Pediatric feeding disorder: Consensus definition and conceptual framework. *Journal of Pediatric Gastroenterology and Nutrition*, *68*(1), 124–129. <https://doi.org/10.1097/MPG.0000000000002188>.
- Gomez, C. R., & Baird, S. (2005). Identifying early indicators for autism in self-regulation difficulties. *Focus on Autism and Other Developmental Disabilities*, *20*(2), 106–116. <https://doi.org/10.1177/10883576050200020101>.
- Graham, J. W. (2009). Missing data analysis: Making it work in the real world. *Annual Review of Psychology*, *60*, 549–576. <https://doi.org/10.1146/annurev.psych.58.110405.085530>.
- Hampton, J., & Strand, P. S. (2015). A review of level 2 parent-report instruments used to screen children aged 1.5-5 for autism: A meta-analytic update. *Journal of Autism and Developmental Disorders*, *45*(8), 2519–2530. <https://doi.org/10.1007/s10803-015-2419-4>.
- Hazlett, H. C., Gu, H., Munsell, B. C., Hyung, S., Styner, M., Wolff, J. J., Elison, J. T., Swanson, M. R., Zhu, H., Botteron, K. N., Collins, D. L., Constantino, J. N., Dager, S. R., Estes, A. M., Evans, A. C., Fonov, V. S., Gerig, G., Kostopoulos, P., McKinstry, R. C., ... Piven, J. (2017). Early brain development in infants at high risk for autism spectrum disorder. *Nature*, *542*(7641), 348–351. <https://doi.org/10.1038/nature21369>.
- Henderson, J. J., Evans, S. F., Straton, J. A. Y., Priest, S. R., & Hagan, R. (2003). Impact of postnatal depression on breastfeeding duration. *Birth*, *30*(3), 175–180. <https://doi.org/10.1046/j.1523-536X.2003.00242.x>.
- Herle, M., Fildes, A., & Llewellyn, C. H. (2018). Emotional eating is learned not inherited in children, regardless of obesity risk. *Pediatric Obesity*, *13*(10), 628–631. <https://doi.org/10.1111/ijpo.12428>.
- Hoekstra, R. A., Vinkhuyzen, A. A. E., & Wheelwright, S. (2011). The construction and validation of an abridged version of the autism-spectrum quotient ( AQ-Short ). *Journal of Autism and Developmental Disorders*, *41*(5), 589–596. <https://doi.org/10.1007/s10803-010-1073-0>.
- Ingersoll, B., & Hambrick, D. Z. (2011). The relationship between the broader autism phenotype, child severity, and stress and depression in parents of children with autism spectrum disorders. *Research in Autism Spectrum Disorders*, *5*(1), 337–344. <https://doi.org/10.1016/j.rasd.2010.04.017>.
- Jaddoe, V. W. V., Van Duijn, C. M., Franco, O. H., Van Der Heijden, A. J., Van IJendoorn, M. H., De Jongste, J. C., Van Der Lugt, A., MacKenbach, J. P., Moll, H. A., Raat, H., Rivadeneira, F., Steegers, E. A. P., Tiemeier, H., Uitterlinden, A. G., Verhulst, F. C., & Hofman, A. (2012). The generation r study: Design and cohort update 2012. *European Journal of Epidemiology*, *27*(9), 739–756. <https://doi.org/10.1007/s10654-012-9735-1>.
- Jahromi, L. B. (2017). Self-regulation in young children with autism spectrum disorder: An interdisciplinary perspective on emotion regulation, executive function, and

- effortful control. In *International review of research in developmental disabilities* (1st ed., Vol. 53, pp. 45–89). Elsevier Inc. <https://doi.org/10.1016/bs.irrdd.2017.07.007>.
- Jansen, P. W., Roza, S. J., Jaddoe, V. W., Mackenbach, J. D., Raat, H., Hofman, A., Verhulst, F. C., & Tiemeier, H. (2012). Children's eating behavior, feeding practices of parents and weight problems in early childhood: Results from the population-based generation R study. *International Journal of Behavioral Nutrition and Physical Activity*, 9(1), 130. <https://doi.org/10.1186/1479-5868-9-130>.
- Johnson, C. P., & Myers, S. M. (2007). Identification and evaluation of children with autism spectrum disorders. *Pediatrics*, 120(5), 1183–1215. <https://doi.org/10.1542/peds.2007-2361>.
- Jones, W., & Klin, A. (2013). Attention to eyes is present but in decline in 2–6-month-old infants later diagnosed with autism. *Nature*, 504(7480), 427–431. <https://doi.org/10.1038/nature12715>.
- Kuzela, A. L., Stifter, C. A., & Worobey, J. (1990). Breastfeeding and mother-infant interactions. *Journal of Reproductive and Infant Psychology*, 8(3), 185–194. <https://doi.org/10.1080/02646839008403623>.
- Ledford, J. R., & Gast, D. L. (2006). Feeding problems in children with autism spectrum disorders: A review. *Focus on Autism and Other Developmental Disabilities*, 21(3), 153–166. <https://doi.org/10.1177/10883576060210030401>.
- Leekam, S. R., Nieto, C., Libby, S. J., Wing, L., & Gould, J. (2007). Describing the sensory abnormalities of children and adults with autism. *Journal of Autism and Developmental Disorders*, 37(5), 894–910. <https://doi.org/10.1007/s10803-006-0218-7>.
- Li, R., Fein, S. B., Chen, J., & Grummer-Strawn, L. M. (2008). Why mothers stop breastfeeding: Mothers' self-reported reasons for stopping during the first year. *Pediatrics*, 122(Supplement 2), S69–S76. <https://doi.org/10.1542/peds.2008-1315i>.
- Liu, J., Leung, P., & Yang, A. (2013). Breastfeeding and active bonding protects against children's internalizing behavior problems. *Nutrients*, 6(1), 76–89. <https://doi.org/10.3390/nu6010076>.
- Llewellyn, C. H., van Jaarsveld, C. H. M., Johnson, L., Carnell, S., & Wardle, J. (2011). Development and factor structure of the baby eating behaviour questionnaire in the gemini birth cohort. *Appetite*, 57(2), 388–396. <https://doi.org/10.1016/j.appet.2011.05.324>.
- Lucas, J., & Cutler. (2015). Autism: Dysregulated breastfeeding behaviors? *The Journal of Perinatal Education*, 24(3), 171–180. <https://doi.org/10.1891/1058-1243.24.3.171>.
- Mastorakou, D., Ruark, A., Weenen, H., Stahl, B., & Stieger, M. (2019). Sensory characteristics of human milk: Association between mothers' diet and milk for bitter taste. *Journal of Dairy Science*, 102(2), 1116–1130. <https://doi.org/10.3168/jds.2018-15339>.
- Micali, N., Simonoff, E., & Treasure, J. (2009). Infant feeding and weight in the first year of life in babies of women with eating disorders. *The Journal of Pediatrics*, 154(1), 55–60. <https://doi.org/10.1016/j.jpeds.2008.07.003>.
- Mulligan, S., & White, B. P. (2012). Sensory and motor behaviors of infant siblings of children with and without autism. *American Journal of Occupational Therapy*, 66(5), 556–566. <https://doi.org/10.5014/ajot.2012.004077>.
- Nadon, G., Feldman, D. E., Dunn, W., & Gisel, E. (2011). Association of sensory processing and eating problems in children with autism spectrum disorders. *Autism Research and Treatment*, 27–29. <https://doi.org/10.1155/2011/541926>.
- Nilsen, R. M., Vollset, S. E., Gjessing, H. K., Skjærven, R., Melve, K. K., Schreuder, P., Alsaker, E. R., Haug, K., Daltveit, A. K., & Magnus, P. (2009). Self-selection and bias in a large prospective pregnancy cohort in Norway. *Paediatric & Perinatal Epidemiology*, 23(6), 597–608. <https://doi.org/10.1111/j.1365-3016.2009.01062.x>.
- Øien, R. A., Nordahl-Hansen, A., & Schjølberg, S. (2019). Screening instruments for ASD. In F. Volkmar (Ed.), *Encyclopedia of autism spectrum disorders*. Springer. [https://doi.org/10.1007/978-1-4614-6435-8\\_102295-1](https://doi.org/10.1007/978-1-4614-6435-8_102295-1).
- Robins, D. L., Casagrande, K., Barton, M., Chen, C. M. A., Dumont-Mathieu, T., & Fein, D. (2014). Validation of the modified checklist for autism in toddlers, revised with follow-up (M-CHAT-R/F). *Pediatrics*, 133(1), 37–45. <https://doi.org/10.1542/peds.2013-1813>.
- Robins, D. L., Fein, D., Barton, M. L., & Green, J. A. (2001). The modified checklist for autism in toddlers: An initial study investigating the early detection of autism and pervasive developmental disorders. *Journal of Autism and Developmental Disorders*, 31(2), 131–144. <https://doi.org/10.1023/A:1010738829569>.
- Román, G. C., Ghassabian, A., Bongers-Schokking, J. J., Jaddoe, V. W. V., Hofman, A., De Rijke, Y. B., Verhulst, F. C., & Tiemeier, H. (2013). Association of gestational maternal hypothyroxinemia and increased autism risk. *Annals of Neurology*, 74(5), 733–742. <https://doi.org/10.1002/ana.23976>.
- Rutter, M., Bailey, A., & Lord, C. (2003). *The social communication questionnaire*. Western Psychological Services.
- Saltzman, J. A., Fiese, B. H., Bost, K. K., & McBride, B. A. (2018). Development of appetite self-regulation: Integrating perspectives from attachment and family systems theory. *Child Development Perspectives*, 12(1), 51–57. <https://doi.org/10.1111/cdep.12254>.
- Shic, F., Macari, S., & Chawarska, K. (2014). Speech disturbs face scanning in 6-month-old infants who develop autism spectrum disorder. *Biological Psychiatry*, 75(3), 231–237. <https://doi.org/10.1016/j.biopsych.2013.07.009>.
- Soke, G. N., Maenner, M., Windham, G., Moody, E., Kaczaniuk, J., DiGiuseppi, C., & Schieve, L. A. (2019). Association between breastfeeding initiation and duration and autism spectrum disorder in preschool children enrolled in the study to explore early development. *Autism Research*, 12(5), 816–829. <https://doi.org/10.1002/aur.2091>.
- Swinkels, S. H. N., Dietz, C., Daalen, E. Van, Kerkhof, I. H. G. M., Engeland, H. Van, & Buitelaar, J. K. (2006). Screening for autistic spectrum in children aged 14 to 15 months. I: The development of the early screening of autistic traits questionnaire (ESAT). *Journal of Autism and Developmental Disorders*, 36(6), 723–732. <https://doi.org/10.1007/s10803-006-0115-0>.
- Tavoulari, E.-F., Benetou, V., Vlastarakos, P. V., Psaltopoulou, T., Chrousos, G., Kreams, G., Gryparis, A., & Linos, A. (2016). Factors affecting breastfeeding duration in Greece: What is important? *World Journal of Clinical Pediatrics*, 5(3), 349. <https://doi.org/10.5409/wjcp.v5.i3.349>.
- Taylor, C. M., Wernimont, S. M., Northstone, K., & Emmett, P. M. (2015). Picky/fussy eating in children: Review of definitions, assessment, prevalence and dietary intakes. *Appetite*, 95, 349–359. <https://doi.org/10.1016/j.appet.2015.07.026>.
- Tharner, A., Luijk, M. P. C. M., Raat, H., Ijzendoorn, M. H., Bakermans-Kranenburg, M. J., Moll, H. A., Jaddoe, V. W. V., Hofman, A., Verhulst, F. C., & Tiemeier, H. (2012). Breastfeeding and its relation to maternal sensitivity and infant attachment. *Journal of Developmental and Behavioral Pediatrics*, 33(5), 396–404. <https://doi.org/10.1097/DBP.0b013e318257fac3>.
- Thorup, E., Nyström, P., Gredebäck, G., Bölte, S., & Falck-Ytter, T. (2018). Reduced alternating gaze during social interaction in infancy is associated with elevated symptoms of autism in toddlerhood. *Journal of Abnormal Child Psychology*, 46(7), 1547–1561. <https://doi.org/10.1007/s10802-017-0388-0>.
- Tick, B., Bolton, P., Happé, F., Rutter, M., & Rijdsdijk, F. (2016). Heritability of autism spectrum disorders: A meta-analysis of twin studies. *The Journal of Child Psychology and Psychiatry and Allied Disciplines*, 57(5), 585–595. <https://doi.org/10.1111/jcpp.12499>.
- Tseng, P. T., Chen, Y. W., Stubbs, B., Carvalho, A. F., Whiteley, P., Tang, C. H., Yang, W., Chen, T. Y., Li, D. J., Chu, C. S., Yang, W.-C., Liang, H.-Y., Wu, C.-K., Yen, C.-F., & Lin, P. Y. (2019). Maternal breastfeeding and autism spectrum disorder in children: A systematic review and meta-analysis. *Nutritional Neuroscience*, 22(5), 354–362. <https://doi.org/10.1080/1028415X.2017.1388598>.
- van 't Hof, M., Ester, W. A., Serdarevic, F., van Berckelaer-Onnes, I., Tiemeier, H., Hoek, H. W., & Jansen, P. W. (2019). The sex-specific association between autistic traits and eating behavior in childhood: An exploratory study in the general population. *Appetite* <https://doi.org/10.1016/j.appet.2019.104519>.
- Wallace, G. L., Llewellyn, C., Fildes, A., & Ronald, A. (2018). Autism spectrum disorder and food neophobia: Clinical and subclinical links. *American Journal of Clinical Nutrition*, 108(4), 701–707. <https://doi.org/10.1093/ajcn/nqy163>.
- Weir, K., McMahon, S., Barry, L., Ware, R., Masters, I. B., & Chang, A. B. (2007). Oropharyngeal aspiration and pneumonia in children. *Pediatric Pulmonology*, 42(11), 1024–1031. <https://doi.org/10.1002/ppul.20687>.
- White, T., Muetzel, R. L., Marroun, H. El, Blanken, L. M. E., Jansen, P., Bolhuis, K., Kocovska, D., & Mous, S. E. (2018). Paediatric population neuroimaging and the generation R Study: The second wave. *European Journal of Epidemiology*, 33(1), 105–131. <https://doi.org/10.1007/s10654-017-0319-y>.
- Wolke, D., Gray, P., & Meyer, R. (1994). Excessive infant crying: A controlled study of mothers helping mothers. *Pediatrics*, 94(3), 322–332.