



Universiteit
Leiden
The Netherlands

Fa(c)t checking: how fatty acids shape metabolism and function of macrophages and dendritic cells

Almeida, L.; Everts, B.

Citation

Almeida, L., & Everts, B. (2021). Fa(c)t checking: how fatty acids shape metabolism and function of macrophages and dendritic cells. *European Journal Of Immunology*, 51(7), 1628-1640. doi:10.1002/eji.202048944

Version: Publisher's Version

License: [Creative Commons CC BY-NC-ND 4.0 license](https://creativecommons.org/licenses/by-nc-nd/4.0/)

Downloaded from: <https://hdl.handle.net/1887/3214691>

Note: To cite this publication please use the final published version (if applicable).

REVIEW**Fa(c)t checking: How fatty acids shape metabolism and function of macrophages and dendritic cells****Luís Almeida**  and **Bart Everts** 

Department of Parasitology, Leiden University Medical Centre, Leiden, The Netherlands

In recent years there have been major advances in our understanding of the role of free fatty acids (FAs) and their metabolism in shaping the functional properties of macrophages and DCs. This review presents the most recent insights into how cell intrinsic FA metabolism controls DC and macrophage function, as well as the current evidence of the importance of various exogenous FAs (such as polyunsaturated FAs and their oxidation products—prostaglandins, leukotrienes, and proresolving lipid mediators) in affecting DC and macrophage biology, by modulating their metabolic properties. Finally, we explore whether targeted modulation of FA metabolism of myeloid cells to steer their function could hold promise in therapeutic settings.

Keywords: Dendritic cells · Fatty Acids · Macrophages · Metabolism · Proresolving mediators**Introduction**

Macrophages are innate immune cells of myeloid origin with a diverse range of functions. They not only play a key role in maintaining tissue homeostasis under noninflammatory conditions, but also in phagocytosis and killing of microbes during an infection and as well as in driving wound healing and tissue regeneration during the resolution phase of an inflammatory response [1, 2]. To be able to acquire these distinct functional traits, macrophages can adopt various polarization states, which are imprinted by the local cues they are exposed to in the tissues where they reside [3–5]. The most well-studied differentiation states are classically activated macrophages (also called M1 macrophages) and alternatively activated macrophages (also referred to as M2 macrophages), and both can be modelled in *in vitro* culture systems. Although these two polarization states fail to capture the full diversity of functional states that macrophages can adopt *in vivo*, they are commonly used as models for studying the proinflammatory and anti-inflammatory properties of these cells, respectively [6].

Like macrophages, DCs also belong to the group of myeloid innate immune cells. DCs sit at the crossroads between the innate and the adaptive immune system, working as special-

ized APCs that are capable of initiating T-cell responses [7]. During steady state, DCs reside in peripheral tissues in a quiescent state. Upon sensing of pathogens or tissue-derived danger signals, DCs undergo a phenotypic and functional change involving enhanced internalization and processing of Ags [8–10] and migration towards tissue draining lymph nodes, where they can induce an adaptive immune response by priming and activating Ag-specific T cells [9, 11]. Furthermore, DCs also play a role in the induction of tolerance during steady state due to exposure to tolerizing signals. These tolerogenic DCs can mediate tolerance by promoting peripheral T-cell anergy and apoptosis, decreasing effector and memory T-cell responses, and inducing the differentiation and activation of Tregs [12–15].

Due to the central role played by DCs and macrophages in the immune response, it is important to understand how their function is regulated and what kind of stimuli are needed to initiate/sustain their activation and polarization in specific scenarios. For instance, there has been a longstanding interest in defining the signaling pathways regulating macrophage and DC function in the context of “classical” immunological cues such as cytokines, chemokines, and PAMPs [16, 17]. More recently, there has been a growing appreciation that metabolic signals and alterations in cellular metabolism can also dictate immune cell function (see [18] for a detailed introduction to immune cell metabolism). Recent research about immunometabolism has contributed to the

Correspondence: Dr. Bart Everts
e-mail: b.everts@lumc.nl




Figure 1. Cellular metabolism of intra- and extracellular fatty acids. A schematic overview of key processes involved in cellular FA uptake and metabolism. Core metabolic pathways connected to FA metabolism are indicated as well as the main processes involved in SPM synthesis from PUFAs such as ω -3 and ω -6 FAs. Specifically, PUFAs, many of which are essential FAs obtained from food, can be metabolized by cyclooxygenases (COX) and lipoxygenases (LOX) to give rise to SPMs. Main PUFAs that serve as substrate for these enzymes are linoleic acid (LA) and arachidonic acid (AA), which are ω -6 PUFAs, and eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA), both ω -3 PUFAs. Several enzymatic reactions lead to synthesis of SPMs (highlighted in green). The metabolism of EPA by COX-2 eventually gives rise to E-series Resolvins (RvE), while with DHA the products are more diverse. The actions of LOX on DHA lead to products from the Maresin (Mar) family, while both LOX and COX-2 can give products from the Protectin family (PD) or D-series Resolvins (RvD). In the case of ω -6 PUFAs, such as AA, COX and LOX, give products.

realization that stimuli and changes in the environment macrophages and DCs are exposed to, eventually converge into alterations in their metabolic properties. It has become clear that reprogramming of metabolic pathways, such as glycolysis, oxidative phosphorylation (OXPHOS), fatty acid (FA) synthesis and oxidation (FAO) are not only associated with, but are also crucial for shaping functional responses of DCs and polarization of macrophages to environmental cues [18].

The notion that FAs (both intracellular as well as extracellular) and their metabolism play a central role in shaping DC and macrophage biology has gained significant traction in recent years [19]. Extracellular FAs can be synthesized *de novo* from carbons derived from other core metabolic pathways such as the TCA cycle, glycolysis and glutaminolysis, hydrolyzed from intracellular lipid stores, or directly obtained from extracellular space (Fig. 1). These FAs play a pivotal structural role when used for incorporation into cellular membranes. Moreover, through their oxidation in mitochondria they serve an important role in generating energy via OXPHOS as well as in generating various TCA cycle intermediates that can act as signaling metabolites or

that can be used for the synthesis of other macromolecules. As a result, metabolism of intracellular FAs is a central regulator of DC and macrophage function. In addition, various FAs present in the extracellular environment, released by other cells, including adipocytes, tumor cells and other immune cells, or obtained through diet [20], have also been shown to have the potential to alter the functional properties of DCs and macrophages [20–22].

The most well-studied exogenous FAs in the context of myeloid cell biology are short-chain fatty acids (SCFAs), saturated fatty acids (SFAs), and unsaturated fatty acids (UFAs), which could be monounsaturated (MUFAs) and polyunsaturated (PUFAs). In addition, specialized proresolving mediators (SPMs) which are products from PUFA metabolism (mainly ω -3 and ω -6 PUFAs; see Fig. 1) [23], can also have strong modulatory effects on myeloid cells including macrophages and DCs [23, 24]. Various mechanisms have been proposed through which these exogenous FAs can affect DC and macrophage function. These include acting as signaling molecules engaging receptors, serving as structural components, and interestingly, altering the metabolism of these cells.

In this review, we will discuss the most recent insights into how intrinsic FA metabolism controls DC and macrophage function, as well as the current evidence showing how various exogenous FAs (such as PUFAs and their oxidation products—prostaglandins, leukotrienes, and SPMs) affect DC and macrophage function, by modulating their metabolic properties.

Role of intrinsic fatty acid metabolism in myeloid cell function

Fatty acid oxidation

Macrophages

Free FAs used for FAO can be acquired either by uptake of dietary fats and subsequent hydrolysis, by lipolysis of stored acylglycerols, or by *de novo* FA synthesis [20]. These FAs can be oxidized either in peroxisomes, and/or be transported into the mitochondria by Carnitine Palmitoyltransferase (CPT) where they will undergo FAO. FAO results in the formation of multiple units of acetyl-CoA, which can serve as substrate in the TCA cycle to fuel OXPHOS. It is a tightly regulated pathway, with the rate limiting step being the transport of acyl-CoA into the mitochondrial matrix by CPT1. The control of CPT1 activity is, therefore, a key checkpoint for regulating mitochondrial FAO (Fig. 1) [25].

Murine macrophages stimulated with IL-4 *in vitro* are characterized by increased FAO activity [26]. Likewise, tumor-associated macrophages (TAMs) which share phenotypic and anti-inflammatory characteristics with *in vitro*-generated M2 macrophages [27] were shown to also have high levels of FAO [28]. Moreover, increased FAO is also correlated with efferocytosis, a process used by M2-like macrophages to remove apoptotic cells to maintain tissue homeostasis [29]. Initial studies have suggested that FAO itself is crucial for M2 polarization, as pharmacological inhibition of CPT1 using etomoxir prevented M2 differentiation [26, 30–32]. However, more recent work has questioned these findings [33]. Using genetic approaches, it was shown that conditional deletion of CPT1a had no effect on acquisition of an M2 phenotype. Moreover, etomoxir was found to have substantial off-target effects at concentrations used in these earlier studies [33]. However, it should be noted that long-term genetic deletion of crucial FAO enzymes, such as CPT1a, may result in metabolic adaptation by usage of compensatory pathways to support cellular processes normally dependent on long-chain FAO, that cells may not be able to resort to upon acute pharmacological inhibition [34]. These issues will need to be resolved to fully understand the importance of FAO in alternative activation of macrophages.

In vitro-generated murine M1 macrophages are characterized by high expression of iNOS leading to the synthesis of NO [35], which is known to impair OXPHOS by inhibiting the electron transport chain in an auto- or paracrine manner (Fig. 2A) [36], and promote the synthesis of ROS that help in

the microbicidal activity of M1 macrophages. As a consequence, FAO is severely compromised in these cells [30]. However, LPS-stimulated peritoneal macrophages (pMacrophages) are characterized by increased OXPHOS [37], which may stem from a lower potential to produce NO by pMacrophages compared to their *in vitro* counterparts [30]. Under certain conditions, FAO can also be important for specific proinflammatory properties of macrophages. NLRP3 inflammasome activation, and synthesis of IL-1 β was impaired following pharmacological inhibition of CPT1a with etomoxir suggesting dependency on FAO [38–40]. However, the exact mechanism by which FAO boosts NLRP3 activations is still not clear, although increased FAO-fuelled mitochondrial ROS production has been implicated [40, 41].

One of the mechanisms by which FA metabolism regulates macrophage polarization involves changes in histone acetylation of IL-4-inducible genes fuelled by an increase in cellular concentrations of acetyl-CoA (Ac-CoA) [31]. It was found that IL-4 receptor (IL-4R) signaling increases ATP citrate lyase (ACLY) expression and activity (see Fig. 2B), resulting in an accumulation of nuclear and cytosolic Ac-CoA to enable efficient histone acetylation. Tracing experiments revealed that of glutamine, glucose and palmitate (PA), the latter was the largest source of carbon for Ac-CoA [31]. This may indicate that FAO in M2 macrophages fuels the TCA cycle to increase ACLY-driven Ac-CoA output to support histone acetylation required for expression of M2 macrophage-associated genes.

Thus, the relationship between macrophage phenotype and FAO does not appear to be as black and white, as initially thought. A picture is emerging that FAO can—depending on the context—support both pro- or anti-inflammatory properties by appropriating specific functions. For instance, in proinflammatory macrophages, FAO may be used to produce ROS to support activation of the NLRP3 inflammasome, as well as potentially fuel the TCA cycle to compensate for cataplerosis of intermediates that are being extracted from the TCA cycle for synthesis of amino acids and other macromolecules needed for proinflammatory activation [38–41]. On the other hand, in M2 macrophages there is evidence that FAO, in addition to being involved in epigenetic remodeling by serving as a source of Ac-CoA required for acquisition of an M2 phenotype, also contributes to maintenance of anti-inflammatory activities, such as efferocytosis [29], by burning through FAs that otherwise accumulate in these cells as a consequence of this process. Moreover, β -oxidation of FAs from apoptotic cells enhanced IL-10 transcription and synthesis, thereby reinforcing their anti-inflammatory phenotype [42].

Dendritic cells

Several studies have highlighted the importance of FAO in regulating the functional properties of DCs in a stimulus- and subset-specific manner [43–45]. Different TLR stimuli engage FAO to a different extent in human moDCs [43]. While TLR4 stimulation was found to induce glycolysis, TLR7/8 stimulation with pRNA increased FAO and OXPHOS in human moDCs (Fig. 3A).




Figure 2. FA metabolism of pro- and anti-inflammatory macrophage Schematic depiction of how FA metabolism and uptake control the function of (A) pro- and (B) anti-inflammatory macrophages. Green lines indicate positive signaling, while red lines indicate an inhibitory effect. (A) Proinflammatory macrophages increase FA synthesis upon TLR signaling. This increase in FA synthesis drives 4-1BBL activity which helps to sustain inflammation. TLR-driven FA synthesis also leads to an increase in lipid droplets which is associated with increased PGE2 and IL-1β synthesis. Moreover, TLR signaling induces iNOS expression and subsequent NO synthesis which inhibits the ETC, thereby reducing FAO, and promoting ROS formation which helps with microbicidal functions. Increased FA synthesis upon TLR stimulation also leads to SFA synthesis, which increases PC levels in the cell membrane, leading to less fluidity and K⁺ efflux, thereby activating NLRP3 and IL-1β synthesis. TLR1/2, TLR7, and TLR9 activation increased FA synthesis and de novo SFA and MUFA synthesis. TLR3 activation leads to the opposite effect, inhibiting FA synthesis, along with MUFA and SFA synthesis. (B) IL-4R signaling activates STAT1, STAT3, and STAT6 which promote the transcription of TAM genes and CD36. IL-4R signaling also promotes de novo FA synthesis which increases Acyl-CoA levels in the cell. Extrinsic FAs can also increase Acyl-CoA levels by being transported intracellularly by CD36 or FATP-1. Increased Acyl-CoA promotes OXPHOS and FAO in a CPT1a-dependent manner. The increased flux in OXPHOS and FAO results in elevated levels of mitochondrial Acetyl-CoA which can be transported to the cytosol in an ACLY-dependent manner or through the Malate-Aspartate shuttle. Cytosolic Acetyl-CoA can participate in histone acetylation of M2-like and TAM genes. PGE2 impairs mitochondrial membrane potential in M2-like macrophages by dysregulation of the Malate-Aspartate shuttle by increasing cAMP-induced Got1 expression.




Figure 3. FA metabolism of immunostimulatory and tolerogenic DCs: Schematic depiction of how FA metabolism and uptake control the function of (A) immunostimulatory and (B) tolerogenic dendritic cells. Green lines indicate positive signaling, while red lines indicate an inhibitory effect. Dashed lines represent effects from exogenous stimuli. (A) In immunostimulatory DCs, TLR signaling increases FA synthesis to promote ER expansion and LD formation, which contributes to upregulation of MHCII and MHCI. PGC1α and PPARγ activation is also associated with increased FA synthesis in immunostimulatory DCs. TLR7/8 stimulation can lead to the phosphorylation of BCKDE1α in a PINK-mediated manner. This results in increased OXPHOS and FAO which in turn is associated with increased activation of cDCs and pDCs. Additionally, extrinsic FAs, such as OA and PA, which are transported intracellularly by MSR1, can boost TLR4 signaling and increase IL-23 synthesis. (B) In tolerogenic DCs, increased OXPHOS and FAO are associated with reduced IL-6 and IL-12 synthesis and increased IDO expression. Tumor-derived FAs can increase OXPHOS and FAO by either being transported by CPT1a or by activating PPARα. This PPARα activation can also increase LD formation, which have a high content of oxidized PUFAs. Extrinsic FAs transported by MSR1 and FABP4 can feed these LDs and increase their oxidized PUFA content. These LDs, and tumor-derived FAs themselves, impair cross-presentation by suppressing MHCI surface expression. External stimuli, such as VitD3 and Wnt5a, can also increase FAO and OXPHOS, with Wnt5a doing so in a PPARγ-dependent manner.

This increase in FAO and OXPHOS was due to branched-chain alpha-keto acid dehydrogenase complex E1-alpha subunit (BCKDE1 α) phosphorylation in a PTEN-induced putative kinase 1 (PINK1)-dependent manner. Interestingly, inducing PINK1 activity in tolerogenic DCs stimulated FAO and rendered these DCs immunostimulatory [43], suggesting FAO can promote proinflammatory functions in DCs. Other studies have also implicated FAO in supporting proinflammatory DC activation by showing that its pharmacological inhibition by etomoxir suppressed both murine cDC and pDC activation stimulated with a TLR9 agonist, as evidenced by decreased expression of costimulatory molecules and decreased synthesis of proinflammatory cytokines [44]. Similarly, murine pDCs increase FAO upon stimulation of TLR9, which was found to be required for efficient type 1 interferon (I-IFN) production [45]. Tumor cells may appropriate this mechanism by secreting α -Fetoprotein, which inhibits FAO and OXPHOS in DCs, leading to impaired stimulation of Ag-specific effector functions [46]. This points towards a link between FAO and triggering of endosomal TLRs, which may be explained by fact that engagement of endosomal TLRs generally lead to strong type 1-IFN production, which can promote FAO in an autocrine manner [41]. Possibly, in this context, FAO may take over the role from glycolysis as main carbon source to fuel the synthesis of TCA cycle intermediates, which in DCs stimulated with cell membrane-associated TLRs is required to support the anabolic demands of immunogenic DC activation [47].

However, increased FAO has also been implicated in supporting tolerogenic properties of DCs (Fig. 2B). DCs rendered tolerogenic *in vitro* using Vitamin-D3 or *in vivo* through Wnt5a signaling in the tumor microenvironment, display increased FAO [48–50] on which they in part rely for induction of Tregs [50, 51]. In the latter study, FAO was mechanistically shown to enhance indolamine 2,3-dioxygenase-1 (IDO) activity and suppress IL-6 and IL-12 cytokine expression by DCs, culminating in Treg generation. Additionally, failure of TLR-stimulated murine CD11c+ DCs or BM-DCs to switch from OXPHOS to glycolysis, due to deficiency in miRNA-142, which normally suppresses CPT1a activity, locked these cells in a tolerogenic state with reduced synthesis of proinflammatory cytokines and reduced ability to activate T cells both *in vitro* and *in vivo* [52].

Taken together, FAO may support, depending on the studied DC subset and/or nature of activating signal, either proinflammatory or anti-inflammatory properties of these cells. How exactly the same metabolic pathway can underpin these divergent immunological properties in DCs remains to be determined.

Fatty acid synthesis

De novo FA synthesis is a process in the cytoplasm whereby acyl chains are generated from Ac-CoA through the action of FA synthases. Most of the Ac-CoA which is converted into FAs is derived from carbohydrates originating from the glycolytic pathway and TCA cycle [15, 46].

Macrophages

There are several studies that have linked de novo FA synthesis to supporting proinflammatory function of macrophages (Fig. 2A). For instance, genetic models to block FA synthesis have shown that NO production and proinflammatory signaling are reduced in both human and murine macrophages upon TLR stimulation [53–56]. Additionally, FA synthesis also appears to support TLR-driven 4-1BBL activity, a member of the TNF superfamily which regulates the sustained production of proinflammatory cytokines in TLR-activated macrophages [57]. Consistent with this latter finding, inhibition of FA synthesis in a murine model of psoriasis, in which prolonged proinflammatory activity of M1 macrophages is a driving factor behind the disease, alleviated the symptoms [57]. However, conditional deletion of ACC1, the enzyme that converts Acetyl-CoA into Malonyl-CoA used for FA synthesis, did not compromise proinflammatory macrophage responses [58]. These discrepancies in outcome might, as alluded to above, arise from the difference in effects of instant pharmacological targeting with potential off-target effects versus long-term deletion. In addition, different TLR ligands appear to have different effects on FA synthesis and the lipidome of macrophages [59]. Stimulation of MyD88-dependent TLRs (i.e. TLR1/2, TLR7, or TLR9) increased de novo SFA and MUFA synthesis, while TLR3 stimulation reduced both SFA and MUFA synthesis. Moreover, inhibition of MUFA synthesis, without affecting SFA synthesis, disturbed the TLR-driven reprogramming of the lipidome, resulting in an increased inflammatory response [59].

Together, this links SFA synthesis to proinflammatory properties of macrophages. Several mechanisms have been proposed through which FA synthesis can support proinflammatory properties of macrophages. First, it may help in changing membrane lipid composition to alter fluidity to facilitate membrane-associated effector functions, such as phagocytosis [60–62], as has been shown following TLR4 stimulation [63]. Here, sterol regulatory element binding protein-1a (SREBP-1a), a key transcriptional regulatory protein of FA metabolism, was activated downstream of TLR4 and increased FA synthesis. Inhibiting this pathway led to defective phagocytosis, resulting from a reduction in the interaction between lipid rafts and the cytoskeleton, presumably due to reduced accumulation of newly synthesized fatty acyl chains within membrane phospholipids [63]. Secondly, synthesized FAs can be used to form lipid droplets (LDs) that have a role in the first line of defense against pathogens, by serving as anchors for immune proteins and as docking sites for phagocytic membranes. This facilitates the encounter between immune proteins and phagocytized pathogens, while also protecting the cell from possible unwanted damage due to the cytotoxic properties of these proteins [64]. Additionally, LDs can change cellular metabolism by uncoupling themselves from mitochondria upon infection, to lower mitochondrial FAO. Finally, it was recently shown that LD development, as a consequence of commitment to triacylglycerol synthesis following TLR stimulation, was needed for increased synthesis of M1-associated inflammatory mediators such as IL-1 β and PGE2 [65].

Dendritic cells

Studies with murine BMDCs have shown that LPS stimulation promotes de novo FA synthesis to support expression of cytokines and costimulatory markers required for potent T-cell activation [47]. Moreover, DC differentiation and subsequent upregulation of MHCII requires FA synthesis [66–68] (Fig. 3A). However, in contrast to pharmacological targeting of FA synthesis as used in the aforementioned study to interrogate the role of de novo FA synthesis, genetic targeting of ACC1 did not appear to affect DC cytokine expression [58].

Mechanistically, FA synthesis-driven DC activation was linked to increased ER and Golgi expansion to allow for efficient translation of these proteins [47]. It is interesting to note that in this study increased LD formation dependent on FA synthesis was also observed. This increase in LD formation has been reported by others as well following LPS or IL-4 stimulation, both in *in vitro* cultured BM-DCs as well as primary CD11c+ murine splenic, and lymph node DCs [69], and was associated with enhanced T-cell activation [47, 69]. This suggests that these LDs serve not just as passive lipid storage organelles but may also be linked to key processes fundamental to DC biology, such as Ag processing and presentation. Indeed inactivation of genes which regulate the assembly of lipid bodies abrogated cross-presentation by DCs [70]. The exact mechanism by which LDs participate in Ag cross-presentation is not yet known, but previous work showed that lipid composition of LDs affected MHC-I expression in DCs [71].

Role of exogenous fatty acids on myeloid cell function

Apart from intracellular FA metabolism, free FAs present in the extracellular space, which can come from the diet or released by other cells, can also affect myeloid cell function [20, 72, 73]. This can be either by serving as structural components, by being used as nutrients to fuel cellular metabolic pathways following uptake, or by acting as signaling molecules through engagement of surface or intracellular receptors. Here, we will specifically focus on their effect on macrophage and DC metabolism and thereby function.

Fatty acid uptake

Macrophages

For many of their biological effects, extracellular FAs first need to be imported into the cell. Macrophages express various receptors and transporters that mediate this process (Fig. 2B). Studies have shown that FA transport protein 1 (FATP1), an important transporter for FA uptake, plays a role in the metabolic reprogramming of macrophages during inflammation [74]. FATP1 overexpression in murine BM-macrophages induced FAO and, in turn reduced an

LPS-induced inflammatory response. Additionally, by inhibiting FATP1, instead a proinflammatory macrophage phenotype could be promoted. CD36, a scavenger receptor, is also involved in the transport of exogenous FAs into the cytosol [32], where they can fuel FAO as shown in IL-4-driven M2 differentiation [32]. This was functionally important as CD36-deficient macrophages were impaired in their M2 polarization. Likewise, human macrophages displayed a reduction in LPS-induced IL-12 and TNF- α synthesis following exposure to FAs [75]. Taken together, this suggests that uptake of exogenous FAs is generally linked to promotion of anti-inflammatory macrophage function.

Dendritic cells

The role for FA uptake in regulating DC function seems to be more complex than in macrophages. On the one hand, activation of both *in vitro*-cultured BMDCs and primary CD11c+ murine DCs (isolated from spleen and lymph nodes) resulted in increased FA accumulation and LD formation (Fig. 3A), which was correlated with increased expression of scavenger receptors, such as Macrophage scavenger receptor 1 (MSR1) [69], suggesting FA uptake may support DC immunogenicity. On the other hand, there is also evidence for a tolerising effect of FA uptake by DCs, especially in the tumor microenvironment [21] (Fig. 3B). Tumor-associated DCs (TADCs) upregulate scavenger receptors, including MSR1, FA binding protein 4 (FABP4), and lipoprotein lipase, which promote exogenous FA uptake, [76–78], and correspondingly display high lipid content and LD accumulation. Functionally, these TADCs showed impaired Ag presentation and subsequent T-cell stimulating ability. Interestingly, inhibiting FA synthesis or MSR1 activity restored their lipid content to normal levels and as well as their T-cell priming abilities [77]. Recently, it was shown that FAs taken up by TADCs can serve as ligands for peroxisome proliferator-activated receptor alpha (PPAR α) which is a member of ligand-activated nuclear transcription factors regulating lipid metabolism. PPAR α binding promoted LD synthesis, as well as increased FAO, which resulted in reduced DC immunogenicity [79]. Interestingly, correspondingly, inhibition of PPAR α activation in this context, restored DCs function and enhanced antitumor immune responses in a therapeutic setting. Even though the anti-inflammatory properties of PPAR α and PPAR δ are well documented (as reviewed in [80]), in some contexts their activity is associated with inflammatory responses. For instance, a recent publication showed that deletion of PPAR δ in CD11c+ cells in mice dampened palmitic acid-induced IL-12 and TNF synthesis, and upregulation of costimulatory molecules, resulting in attenuated development of atherosclerosis [81].

An explanation for why FA uptake and LD formation can, depending on the context, either support or interfere with DC immunogenicity, may come from the nature of the FAs these cells accumulate. LDs in TADCs were shown to contain high levels of oxidized PUFAs [82, 83] compared to non-TADCs, which has been linked to tumor-derived molecules that prompt lipid peroxidation in TADCs [84]. This was found to drive accumulation of

MHC-I-peptide complexes in lysosomes and late endosomes, limiting cross-presentation and, subsequently, cytotoxic T-cell priming. This would be consistent with recent work by Ugolini et al. [85] showing that uptake of oxidized truncated FAs impaired DC Ag cross-presentation in cancer, without affecting direct presentation.

Fatty acids as signaling molecules

Saturated fatty acids

It is well described that in general signaling by exogenous SFAs exert proinflammatory effects on DCs and macrophages [86]. Interestingly, several recent studies have now also revealed an important role for metabolic rewiring in this process. Activation of exogenous SFAs into Acyl-CoA, was shown to activate the NLRP3 inflammasome, driving an M1 type while UFAs prevented this [38] (Fig. 2A). The authors showed that these SFAs promoted the synthesis of phosphatidylcholine, leading to loss of membrane fluidity and K⁺ efflux, enabling subsequent NLRP3 activation. UFAs were able to inhibit this effect by instead redirecting SFAs to triacylglycerol synthesis. Furthermore, exposure of macrophages to palmitic acid (PA), a SFA, was associated with impaired wound healing, a state of low-grade chronic inflammation and increased IL-1 β and IL-23 synthesis [87–89]. In an environment rich in FAs, DCs are also stimulated toward a proinflammatory phenotype. Specifically, accumulation of PA and oleic acid (OA) amplified TLR signaling and led to an increase in IL-23 expression, which in a model of psoriasis worsened disease progression [88] (Fig. 3A). This was linked to PA inhibiting hexokinase activity and perturbing TCA metabolism in TLR-activated cells, leading to an increase in mtROS and proinflammatory cytokines. Nevertheless, the exact mechanisms or receptors by which SFAs can modulate macrophage or DC function are still not fully elucidated. It was previously thought that SFAs could bind TLRs, thus, activating a proinflammatory phenotype in macrophages. However, recent data show that, while TLR4 signaling is needed for SFA-induced inflammation, SFAs do not bind directly to TLR4 [90].

Polyunsaturated fatty acids

One of the most well-studied bioactive FAs known to modulate myeloid cell function is Prostaglandin E2 (PGE2), an oxidation product of AA that can bind specific receptors (Fig. 1). While PGE2 was already reported to inhibit murine BM-macrophage activation and polarization both in vitro and in vivo [91], more recent work elucidated metabolic effects of PGE2 on M2 macrophages. The authors observed that PGE2, alongside a drop in expression of a subset of M2 markers, caused a dissipation of the mitochondrial membrane potential in IL-4-stimulated M2 macrophages [92] (Fig. 2B). This was due to PGE2 affecting the transcription of several genes related to maintenance of

Table 1. SPMs and their association with protection against inflammatory diseases

Disease	SPM	References
Alzheimer's	RvD1, RvE1	[122, 123]
Arthritis	17-HDHA, RvD1, RvE3	[119, 124]
Atherosclerosis	Resolvins and Lipoxins	[125]
Colitis	RvE1	[103]
Diabetes	RvD1, RvE1, Lipoxins	[126]
Parkinson's	RvD1	[127]

mitochondrial membrane potential in a cAMP-mediated manner. PGE2 initiated the transcription of genes that regulate the malate-aspartate shuttle, including Got1. Another PUFA, Leukotriene-B4 (LTB4) has also recently been linked to regulating macrophage metabolism [93]. Type 1 diabetic mice have increased levels of circulating LTB4. Macrophages from these mice displayed increased FAO and CPT2 expression when compared to macrophages from control mice. This was associated with an increased proinflammatory signature. These effects were reduced upon blocking LTB4 signaling using a receptor antagonist.

Specialized proresolving mediators

Recently, SPMs have received considerable attention given the growing evidence for their key role in active resolution of inflammation. There are already some strong correlations between deregulation of SPM metabolism, and certain chronic inflammatory diseases such as Alzheimer's disease, atherosclerosis, arthritis, and type-2 diabetes [23] (Table 1). The proresolving properties of SPMs stem in a large part from their ability to suppress inflammatory properties of macrophages and DCs. SPMs promote the shift from M1-like to M2-like macrophages, increase phagocytic and efferocytotic activities of macrophages, and reduce IL-12 synthesis by DCs [94, 95]. However, until now, little is known about whether SPMs may affect metabolism of these cells or how cellular metabolism of those cells affects SPM synthesis.

Given the known dampening effects of SPMs on proinflammatory DC and macrophage activation and the clear functional link between engagement of certain metabolic pathways and anti-inflammatory properties of these cells, it is tempting to speculate that a mechanism by which SPMs mediate these effects is by modulating DC and macrophage metabolism. One way SPMs might achieve this is by binding to PPARs. PPARs can be activated by many different ligands, including long-chain SFAs and UFAs, eicosanoids or other products of PUFA oxidation such as SPM 15-deoxy- δ -12,14-prostaglandin J2 and Maresin-1 [96–100]. Interestingly, PPAR signaling is known to play a role in attenuating the inflammatory function of macrophages as well as DCs by regulating their metabolism [79, 101]. Additionally, SPMs can bind surface receptors within the family of G-protein-coupled receptors (GPRs) [102]. Although it remains to be established whether signaling through GPRs that bind SPMs (e.g. GPR32, ALX/FPR2, ChemR23 [103, 104]) could drive metabolic reprogramming,

the fact that signaling through other GPRs, such as GPR120 and GPR40, has already been described to affect FA synthesis in adipocytes and hepatocytes, makes it tempting to speculate that SPMs may also alter macrophage and DC metabolism and thereby their function via GPR signaling [105, 106].

Perspectives and outlook

There is a growing body of evidence for a key role of FAs in the regulation of myeloid cell function and the inflammatory response by serving as nutrients, structural components of cells, signaling molecules, and/or epigenetic regulators. Many studies point to the increasing importance of FA metabolism in the function of DCs, such as T-cell priming and Ag-presentation, and that of macrophages such as microbicidal activity, phagocytosis, and efferocytosis. While pro- and anti-inflammatory macrophage and DC phenotypes are often characterized by, and dependent on, engagement of FA synthesis and FAO, respectively, it is becoming increasingly clear that, depending on the context, particularly FAO can support both pro- and anti-inflammatory properties of these cells. How exactly a single metabolic pathway can appropriate these different functions that allow it to support such diverse immunological responses is still poorly understood and it is one of the key outstanding questions that awaits to be addressed. However, one could hypothesize that the activity of metabolic pathways directly connected to FA metabolism, controlled by specific environmental cues and/or the nature of cell subset intrinsic metabolic imprinting, can play a decisive role in how products derived from FA metabolism are being redirected and used, and thereby what the final functional output is of such a pathway.

Many important insights in the effects of FAs on myeloid cell metabolism and function have been gained from *in vitro* studies. However, more *in-depth in vivo* studies will be crucial to fully capture the complexity of this interaction and will be needed further in this field. First, this pertains to the complexity of the FA composition and metabolic changes in the local microenvironment that are not easily reproduced *in vitro*. Metabolic conditions in general, and FA composition in particular—such as those found at sites of inflammation or the tumor microenvironment—are highly complex and are likely fluctuating over time. Second, the diversity in macrophage phenotypes and DC subsets and associated metabolic properties found in tissues cannot be fully modelled *in vitro*. For instance, this is well illustrated by tissue-resident macrophages which have a very different metabolic profile than BM-macrophages generated *in vitro* [107]. Likewise, evidence is accumulating that specific DC subsets found *in vivo*, have distinct metabolic properties and requirements for FA metabolism for their function [108]. To what extent most aforementioned studies, often using *in vitro*-generated BMDCs, faithfully recapitulate the metabolic programs that are engaged and needed for primary DCs *in situ* is questionable. Moreover, what role FA metabolism plays in the biology of the recently discovered new DC subsets is yet to be addressed [109, 110].

An important hurdle in studying FA metabolism of macrophages and DCs *in situ* is that they are generally present at very low frequencies. However, recent advances in single-cell technologies, such as single-cell RNA sequencing and high dimensional flow cytometry [111], are now making it possible to characterize in unprecedented depth phenotypes and metabolic characteristics of rare cell populations at single-cell level in clinical and tissue samples [112]. Moreover, imaging mass cytometry combined with imaging mass spectrometry could provide crucial additional spatial information about local metabolite and FA abundance and myeloid cell phenotype in tissues. These are some of the promising technological advances that will no doubt spur new discoveries in this exciting field of immunometabolism.

As exemplified in this review, there is a growing number of studies that show that extracellular FAs and SPMs can modulate DC and macrophage function by altering their metabolic properties. An aspect that has thus far received less attention is, if and how cellular metabolism shapes the production and release of these lipid mediators. There are first indications that show indeed synthesis of certain PUFAs is dependent on LD formation fuelled by *de novo* FA synthesis [65]. Whether production of anti-inflammatory FAs, such as SPMs, is supported by different metabolic programs than proinflammatory FAs warrants further investigation.

Finally, there is increasing evidence that suggests that deregulated FA metabolism in macrophages and DCs can contribute to development of several inflammatory diseases. This link has been particularly well established in atherosclerosis and type 2 diabetes where dysfunctional lipid handling and metabolism by macrophages have been shown to be an important driver of pathology (i.e. plaque formation and tissue-specific insulin resistance, respectively) [74, 113, 114]. Moreover, altered lipid handling by macrophages and DCs due to hyperlipidemia as in human studies are associated with, and in murine models causally linked to increased chances of developing autoimmune diseases, such as psoriasis, rheumatoid arthritis (RA), and systemic lupus erythematosus (SLE) [57, 81, 115]. Therefore, therapies aimed at targeting FA metabolism have proven their value in treatment of such disorders. For example, PPAR γ agonists are successfully used in the clinic to counteract hyperlipidemia and hyperglycemia. However, the beneficial effects of these drugs have been primarily attributed to alterations in FA metabolism of metabolic tissues rather than myeloid cells [116]. Nonetheless, given the important role of FA metabolism in the regulation of macrophage and DC biology, now also efforts are undertaken to evaluate whether direct manipulation of FA metabolism of those cells could be used to shape their functional properties for therapeutic purposes. In this respect, there are several preclinical mouse studies that have given promising results. For instance, pharmacological inhibition of FAO with etomoxir in myeloid-derived suppressor cells enhanced the effectiveness of cancer therapies in mice [117]. Additionally, etomoxir improved antitumor response following checkpoint blockade treatment *in vivo* [50], which was associated with a switch from a tolerogenic to an immunogenic

TADC phenotype. Apart from targeting core FA metabolism in DCs or macrophages to shape their function for therapeutic purposes, there are also interesting developments aimed at promoting the synthesis of SPMs for treatment of inflammatory disorders. In this respect it is interesting to note that it is already known that aspirin, a common anti-inflammatory drug, triggers the synthesis of several SPMs by modifying COX-2 activity and inhibiting COX-1 [118]. These data suggest the anti-inflammatory properties which have been attributed to aspirin lie, in part, in its ability to promote SPM synthesis. The therapeutic potential of SPMs has also been evaluated more directly. It was shown that RvD1 leads to cartilage protection and better disease outcome in a murine arthritis model [119]. Additionally, through lipidomic analysis of human samples, 17-HDHA was shown to be associated with lower pain in arthritis patients, pointing toward a possible therapeutic application of these compounds in treating inflammatory diseases [120]. Indeed, in a human skin blister model, it was shown that administration of SPMs into the inflamed site promoted resolution [121]. These findings have paved the way for a phase I clinical trial (NCT04308889) that is currently ongoing, in which the effects of dietary supplementation with ω -3 FAs in a human inflammation blister model are assessed, to determine whether SPMs may promote resolution of inflammation. Additional work will be needed to establish to what extent the potential beneficial effects of these treatments are dependent on functional modulation of myeloid cells.

In conclusion, the intricate connection between FA metabolism and myeloid cell function, makes it a highly interesting target for therapeutic intervention to modulate immune responses and to potentially treat diseases marked by a compromised inflammatory response, such as cancer, or by a failure to resolve inflammation, as occurs in various chronic inflammatory disorders.

Acknowledgments: This work was supported by funding from the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie Grant agreement No. 812890.

Conflict of interest: The authors declare no financial or commercial conflict of interest.

References

- 1 Wynn, T. and Barron, L., Master regulators of inflammation and fibrosis. *Semin. Liver Dis.* 2010. **30**: 245–257.
- 2 Gordon, S., Alternative activation of macrophages. *Nat. Rev. Immunol.* 2003. **3**: 23–35.
- 3 Gordon, S. and Taylor, P. R., Monocyte and macrophage heterogeneity. *Nat. Rev. Immunol.* 2005. **5**: 953–964.

- 4 Murray, P. J. and Wynn, T. A., Protective and pathogenic functions of macrophage subsets. *Nat. Rev. Immunol.* 2011. **11**: 723–737.
- 5 Mosser, D. M. and Edwards, J. P., Exploring the full spectrum of macrophage activation. *Nat. Rev. Immunol.* 2008. **8**: 958–969.
- 6 Murray, P. J., Allen, J. E., Biswas, S. K., Fisher, E. A., Gilroy, D. W., Goerdt, S., Gordon, S., et al. Macrophage activation and polarization: nomenclature and experimental guidelines. *Immunity* 2014. **41**: 14–20.
- 7 Sallusto, F. and Lanzavecchia, A., Efficient presentation of soluble antigen by cultured human dendritic cells is maintained by granulocyte/macrophage colony-stimulating factor plus Interleukin 4 and downregulated by tumor necrosis factor α . *J. Exp. Med.* 1994. **179**: 1109–1118.
- 8 Reis, E. and Sousa, C., Toll-like receptors and dendritic cells: for whom the bug tolls. *Semin. Immunol.* 2004. **16**: 27–34.
- 9 Henri, S., Guillemins, M., Poulin, L. F., Tamoutounour, S., Ardouin, L., Dalod, M. and Malissen, B., Disentangling the complexity of the skin dendritic cell network. *Immunol. Cell Biol.* 2010. **88**: 366–375.
- 10 Platt, C. D., Ma, J. K., Chalouni, C., Ebersold, M., Bou-Reslan, H., Carano, R. A. D., Mellman, I., et al., Mature dendritic cells use endocytic receptors to capture and present antigens. *Proc. Natl. Acad. Sci. USA* 2010. **107**: 4287–4292.
- 11 Martín-Fontecha, A., Sebastiani, S., Höpken, U. E., Uguccioni, M., Lipp, M., Lanzavecchia, A. and Sallusto, F., Regulation of dendritic cell migration to the draining lymph node: impact on T lymphocyte traffic and priming. *J. Exp. Med.* 2003. **198**: 615–621.
- 12 Kryczanowski, F., Raker, V., Graulich, E., Domogalla, M. P. and Steinbrink, K., IL-10-modulated human dendritic cells for clinical use: identification of a stable and migratory subset with improved tolerogenic activity. *J. Immunol.* 2016. **197**: 3607–3617.
- 13 Raker, V. K., Domogalla, M. P. and Steinbrink, K., Tolerogenic dendritic cells for regulatory T cell induction in man. *Front. Immunol.* 2015. **6**: 1–11.
- 14 Boks, M. A., Kager-Groenland, J. R., Haasjes, M. S. P., Zwaginga, J. J., van Ham, S. M. and ten Brinke, A., IL-10-generated tolerogenic dendritic cells are optimal for functional regulatory T cell induction: a comparative study of human clinical-applicable DC. *Clin. Immunol.* 2012. **142**: 332–342.
- 15 Li, H. and Shi, B., Tolerogenic dendritic cells and their applications in transplantation. *Cell Mol. Immunol.* 2015. **12**: 24–30.
- 16 Murray, P. J., Macrophage polarization. *Annu. Rev. Physiol.* 2017. **79**: 541–566.
- 17 Kapsenberg, M. L., Dendritic-cell control of pathogen-driven T-cell polarization. *Nat. Rev. Immunol.* 2003. **3**: 984–993.
- 18 O'Neill, L. A. J., Kishton, R. J. and Rathmell, J., A guide to immunometabolism for immunologists. *Nat. Rev. Immunol.* 2016. **16**: 553–565.
- 19 O'Neill, L. A. J. and Pearce, E. J., Immunometabolism governs dendritic cell and macrophage function. *J. Exp. Med.* 2016. **213**: 15–23.
- 20 Galli, C. and Risé, P., Origin of fatty acids in the body: endogenous synthesis versus dietary intakes. *Eur. J. Lipid Sci. Technol.* 2006. **108**: 521–525.
- 21 Koundouros, N. and Poulgiannis, G., Reprogramming of fatty acid metabolism in cancer. *Br. J. Cancer* 2020. **122**: 4–22.
- 22 Wu, H., Han, Y., Rodriguez Sillke, Y., Deng, H., Siddiqui, S., Treese, C., Schmidt, F., et al., Lipid droplet-dependent fatty acid metabolism controls the immune suppressive phenotype of tumor-associated macrophages. *EMBO Mol. Med.* 2019. **11**: 1–17.
- 23 Serhan, C. N. and Levy, B. D., Resolvins in inflammation: emergence of the pro-resolving superfamily of mediators. *J. Clin. Invest.* 2018. **128**: 2657–2669.

- 24 Serhan, C. N., Chiang, N. and Dalli, J., The resolution code of acute inflammation: novel pro-resolving lipid mediators in resolution. *Semin. Immunol.* 2015. 27: 200–215.
- 25 Lehner, R. and Quiroga, A. D., Fatty acid handling in mammalian cells [Internet]. *Biochemistry of lipids, lipoproteins and membranes: Sixth edition*. Elsevier, Amsterdam, The Netherlands, 2016; pp. 149–184. <https://doi.org/10.1016/B978-0-444-63438-2.00005-5>
- 26 Huang, S. C. C., Smith, A. M., Everts, B., Colonna, M., Pearce, E., Schilling, J. D. and Pearce, E. J., Metabolic reprogramming mediated by the mTORC2-IRF4 signaling axis is essential for macrophage alternative activation. *Immunity* 2016. 45: 817–830.
- 27 Vitale, I., Manic, G., Coussens, L. M., Kroemer, G. and Galluzzi, L., Macrophages and metabolism in the tumor microenvironment. *Cell Metab.* 2019. 30: 36–50.
- 28 Su, P., Wang, Q., Bi, E., Ma, X., Liu, L., Yang, M., Qian, J. and Yi, Q., Enhanced lipid accumulation and metabolism are required for the differentiation and activation of tumor-associated macrophages. *Cancer Res.* 2020. 80: 1438–1450.
- 29 Park, D., Han, C. Z., Elliott, M. R., Kinchen, J. M., Tramont, P. C., Das, S., Collins, S., et al. Continued clearance of apoptotic cells critically depends on the phagocyte Ucp2 protein. *Nature* 2011. 477: 220–224.
- 30 Den, B. J. V., Lao, N. and Menon, D., Macrophage immunometabolism: where are we (going)? *Trends Immunol.* 2017. 38: 395–406.
- 31 Govarrubias, A. J., Aksoylar, H. I., Yu, J., Snyder, N. W., Worth, A. J., Iyer, S. S., Wang, J., et al. Akt-mTORC1 signaling regulates ACLY to integrate metabolic input to control of macrophage activation. *Elife* 2016. 5: 1–19.
- 32 Huang, S. C.-C., Everts, B., Ivanova, Y., O'sullivan, D., Nascimento, M., Smith, A. M., Beatty, W., et al., Cell-intrinsic lysosomal lipolysis is essential for alternative activation of macrophages. *Nat. Immunol.* 2014. 15: 846–855.
- 33 Divakaruni, A. S., Hsieh, W. Y., Minarrieta, L., Duong, T. N., Kim, K. K.O., Desousa, B. R., Andreyev, A. Y., et al., Etomoxir inhibits macrophage polarization by disrupting COA homeostasis. *Cell Metab.* 2018. 28: 490–503.
- 34 Van Den Bossche, J. and Van Der Windt, G. J. W., Fatty acid oxidation in macrophages and T cells: time for reassessment? *Cell Metab.* 2018. 28: 538–540.
- 35 Macmicking, J. D., North, R. J., Lacourse, R., Mudgett, J. S., Shah, S. K. and Nathan, C. F., Identification of nitric oxide synthase as a protective locus. *Proc. Natl Acad. Sci.* 1997. 94: 5243–5248.
- 36 Tengan, C. H. and Moraes, C. T., NO control of mitochondrial function in normal and transformed cells. *Biochim. Biophys. Acta-Bioenerg.* 2017. 1858: 573–581.
- 37 Artyomov, M. N., Sergushichev, A. and Schilling, J. D., Integrating immunometabolism and macrophage diversity. *Semin. Immunol. [Internet]*. 2016. 28: 417–424.
- 38 Gianfrancesco, M. A., Dehairs, J., L'homme, L., Herinckx, G., Esser, N., Jansen, O., Habraken, Y., et al., Saturated fatty acids induce NLRP3 activation in human macrophages through K⁺ efflux resulting from phospholipid saturation and Na, K-ATPase disruption. *Biochim. Biophys. Acta—Mol. Cell Biol. Lipids* 2019. 1864: 1017–1030.
- 39 Yang, Q., Liu, R., Yu, Q., Bi, Y. and Liu, G., Metabolic regulation of inflammasomes in inflammation. *Immunology* 2019. 157: 95–109.
- 40 Moon, J. - S., Nakahira, K., Chung, K. - P., Denicola, G. M., Koo, M. J., Pabón, M. A., Rooney, K. T., et al., NOX4-dependent fatty acid oxidation promotes NLRP3 inflammasome activation in macrophages. *Nat. Med.* 2016. 22: 1002–1012.
- 41 Martinon, F., Signaling by ROS drives inflammasome activation. *Eur. J. Immunol.* 2010. 40: 616–619.
- 42 Zhang, S., Weinberg, S., DeBerge, M., Gainullina, A., Schipma, M., Kinchen, J. M., Ben-Sahra, I., et al. Efferocytosis fuels requirements of fatty acid oxidation and the electron transport chain to polarize macrophages for tissue repair. *Cell Metab.* 2019. 29: 443–456.
- 43 Basit, F. and De Vries, I. J. M., Dendritic cells require PINK1-mediated phosphorylation of BCKDE1 α to promote fatty acid oxidation for immune function. *Front. Immunol.* 2019. 10: 1–15.
- 44 Qiu, C. C., Atencio, A. E. and Gallucci, S., Inhibition of fatty acid metabolism by etomoxir or TOFA suppresses murine dendritic cell activation without affecting viability. *Immunopharmacol. Immunotoxicol.* 2019. 41: 361–369.
- 45 Wu, D., Sanin, D. E., Everts, B., Chen, Q., Qiu, J., Buck, M. D., Patterson, A., et al. Type 1 interferons induce changes in core metabolism that are critical for immune function. *Immunity* 2016. 44: 1325–1336.
- 46 Santos, P. M., Menk, A. V., Shi, J., Tsung, A., Delgoffe, G. M. and Butterfield, L. H., Tumor-derived α -fetoprotein suppresses fatty acid metabolism and oxidative phosphorylation in dendritic cells. *Cancer Immunol. Res.* 2019. 7: 1001–1013.
- 47 Everts, B., Amiel, E., Huang, S. C.-C., Smith, A. M., Chang, C. - H., Lam, W. Y., Redmann, V., et al., TLR-driven early glycolytic reprogramming via the kinases TBK1- $IKK\epsilon$ supports the anabolic demands of dendritic cell activation. *Nat. Immunol.* 2014. 15: 323–332.
- 48 Ferreira, G. B., Van Etten, E., Lage, K., Hansen, D. A., Moreau, Y., Workman, C. T., Waer, M., et al., Proteome analysis demonstrates profound alterations in human dendritic cell nature by TX527, an analogue of vitamin D. *Proteomics* 2009. 9: 3752–3764.
- 49 Ferreira, G. B., Kleijwegt, F. S., Waelkens, E., Lage, K., Nikolic, T., Hansen, D. A., Workman, C. T., et al., Differential protein pathways in 1, 25-dihydroxyvitamin D3 and dexamethasone modulated tolerogenic human dendritic cells. *J. Proteome Res.* 2012. 11: 941–971.
- 50 Zhao, F., Xiao, C., Evans, K. S., Theivanthiran, T., Devito, N., Holtzhausen, A., Liu, J., et al., Paracrine Wnt5a- β -catenin signaling triggers a metabolic program that drives dendritic cell tolerization. *Immunity* 2018. 48: 147–160.
- 51 Malinarich, F., Duan, K., Hamid, R. A., Bijin, A., Lin, W. X., Poidinger, M., Fairhurst, A. - M., et al., High mitochondrial respiration and glycolytic capacity represent a metabolic phenotype of human tolerogenic dendritic cells. *J. Immunol.* 2015. 194: 5174–5186.
- 52 Sun, Y., Oravec-Wilson, K., Bridges, S., Mceachin, R., Wu, J., Kim, S. H., Taylor, A., et al., miR-142 controls metabolic reprogramming that regulates dendritic cell activation. *J. Clin. Invest.* 2019. 129: 2029–2042.
- 53 Infantino, V., Iacobazzi, V., Palmieri, F. and Menga, A., ATP-citrate lyase is essential for macrophage inflammatory response. *Biochem. Biophys. Res. Commun.* 2013. 440: 105–111.
- 54 Infantino, V., Convertini, P., Cucci, L., Panaro, M. A., Di Noia, M. A., Calvello, R., Palmieri, F., et al., The mitochondrial citrate carrier: a new player in inflammation. *Biochem. J.* 2011. 438: 433–436.
- 55 Wei, X., Song, H., Yin, L., Rizzo, M. G., Sidhu, R., Covey, D. F., Ory, D. S., et al., Fatty acid synthesis configures the plasma membrane for inflammation in diabetes. *Nature* 2016. 539: 294–298.
- 56 Moon, J.-S., Lee, S., Park, Mi-Ae, Siempos, I. I., Haslip, M., Lee, P. J., Yun, M., et al., UCP2-induced fatty acid synthase promotes NLRP3 inflammasome activation during sepsis. *J. Clin. Invest.* 2015. 125: 665–680.
- 57 Miki, H., Han, K. H., Scott, D., Croft, M. and Kang, Y. J., 4-1BBL regulates the polarization of macrophages, and inhibition of 4-1BBL signaling alleviates imiquimod-induced psoriasis. *J. Immunol.* 2020. 204: 1892–1903.
- 58 Stüve, P., Minarrieta, L., Erdmann, H., Arnold-Schrauf, C., Swallow, M., Guderian, M., Krull, F., et al. De novo fatty acid synthesis during mycobacterial infection is a prerequisite for the function of highly

- proliferative T cells, but not for dendritic cells or macrophages. *Front. Immunol.* 2018. 9: 1–22.
- 59 Hsieh, W.-Y., Zhou, Q. D., York, A. G., Williams, K. J., Scumpia, P. O., Kroenberger, E. B., Hoi, X. P., et al., Toll-like receptors induce signal-specific reprogramming of the macrophage lipidome. *Cell Metab.* 2020. 32: 128–143.
- 60 Araldi, E., Fernández-Fuertes, M., Canfrán-Duque, A., Tang, W., Cline, G. W., Madrigal-Matute, J., Pober, J. S., et al., Lanosterol modulates TLR4-mediated innate immune responses in macrophages. *Cell Rep.* 2017. 19: 2743–2755.
- 61 Blanc, M., Hsieh, W. Y., Robertson, K. A., Kropp, K. A., Forster, T., Shui, G., Lacaze, P., et al., The transcription factor STAT-1 couples macrophage synthesis of 25-hydroxycholesterol to the interferon antiviral response. *Immunity* 2013. 38: 106–118.
- 62 Schumann, J., It is all about fluidity: fatty acids and macrophage phagocytosis. *Eur. J. Pharmacol.* 2016. 785: 18–23.
- 63 Oishi, Y., Spann, N. J., Link, V. M., Muse, E. D., Strid, T., Edillor, C., Kolar, M. J., et al. SREBP1 contributes to resolution of pro-inflammatory TLR4 signaling by reprogramming fatty acid metabolism. *Cell Metab.* 2017. 25: 412–427.
- 64 Bosch, M., Sánchez-Álvarez, M., Fajardo, A., Kapetanovic, R., Steiner, B., Dutra, F., Moreira, L., et al. Mammalian lipid droplets are innate immune hubs integrating cell metabolism and host defense. *Science* 2020. 370: eaay8085.
- 65 Castoldi, A., Monteiro, L. B., van Teijlingen Bakker, N., Sanin, D. E., Rana, N., Corrado, M., Cameron, A. M., et al. Triacylglycerol synthesis enhances macrophage inflammatory function. *Nat. Commun.* 2020. 11: 1–11.
- 66 Ishikawa, F., Niuro, H., Iino, T., Yoshida, S., Saito, N., Onohara, S., Miyamoto, T., et al., The developmental program of human dendritic cells is operated independently of conventional myeloid and lymphoid pathways. *Blood* 2007. 110: 3591–3600.
- 67 Le Naour, F., Hohenkirk, L., Grolleau, A., Misek, D. E., Lescure, P., Geiger, J. D., Hanash, S. and Beretta, L., Profiling changes in gene expression during differentiation and maturation of monocyte-derived dendritic cells using both oligonucleotide microarrays and proteomics. *J. Biol. Chem.* 2001. 276: 17920–17931.
- 68 Rehman, A., Hemmert, K. C., Ochi, A., Jamal, M., Henning, J. R., Barilla, R., Quesada, J. P., et al., Role of fatty-acid synthesis in dendritic cell generation and function. *J. Immunol.* 2013. 190: 4640–4649.
- 69 Maroof, A., English, N. R., Bedford, P. A., Gabrilovich, D. I. and Knight, S. C., Developing dendritic cells become “lacy” cells packed with fat and glycogen. *Immunology* 2005. 115: 473–483.
- 70 Bougnères, L., Helft, J., Tiwari, S., Vargas, P., Chang, B. H.-J., Chan, L., Campisi, L., et al., A role for lipid bodies in the cross-presentation of phagocytosed antigens by MHC class I in dendritic cells. *Immunity* 2009. 31: 232–244.
- 71 Shaikh, S. R., Mitchell, D., Carroll, E., Li, M., Schneck, J. and Edidin, M., Differential effects of a saturated and a monounsaturated fatty acid on MHC class I antigen presentation. *Scand. J. Immunol.* 2008. 68: 30–42.
- 72 Nordestgaard, B. G., Chapman, M. J., Ray, K., Borén, J., Andreotti, F., Watts, G. F., Ginsberg, H., et al., Lipoprotein(a) as a cardiovascular risk factor: current status. *Eur. Heart J.* 2010. 31: 2844–2853.
- 73 Willnow, T. E., Sheng, Z., Ishibashi, S., Herz, J. Inhibition of hepatic chylomicron remnant uptake by gene transfer of a receptor antagonist. *Science* 1994. 264: 1471–1474.
- 74 Johnson, A. R., Qin, Y., Cozzo, A. J., Freerman, A. J., Huang, M. J., Zhao, L., Sampey, B. P., et al. Metabolic reprogramming through fatty acid transport protein 1 (FATP1) regulates macrophage inflammatory potential and adipose inflammation. *Mol. Metab.* 2016. 5: 506–526.
- 75 Klein-Wieringa, I. R., Andersen, S. N., Kwekkeboom, J. C., Giera, M., De Lange-Brokaar, B. J. E., Van Osch, G. J. V. M., Zuurmond, A.-M., et al., Adipocytes modulate the phenotype of human macrophages through secreted lipids. *J. Immunol.* 2013. 191: 1356–1363.
- 76 Gao, F., Liu, C., Guo, J., Sun, W., Xian, L., Bai, D., Liu, H., et al. Radiation-driven lipid accumulation and dendritic cell dysfunction in cancer. *Sci. Rep.* 2015. 5: 1–8.
- 77 Herber, D. L., Cao, W., Nefedova, Y., Novitskiy, S. V., Nagaraj, S., Tyurin, V. A., Corzo, A., et al. Lipid accumulation and dendritic cell dysfunction in cancer. *Nat. Med.* 2010. 16: 880–886.
- 78 Gardner, J. K., Mamotte, C. D. S., Patel, P., Yeoh, T. L., Jackaman, C. and Nelson, D. J., Mesothelioma tumor cells modulate dendritic cell lipid content, phenotype and function. *PLoS One* 2015. 10: 1–29.
- 79 Yin, X., Zeng, W., Wu, B., Wang, L., Wang, Z., Tian, H., Wang, L., et al., PPAR α inhibition overcomes tumor-derived exosomal lipid-induced dendritic cell dysfunction. *Cell Rep.* 2020. 33: 108278.
- 80 Varga, T., Czimmerer, Z. and Nagy, L., PPARs are a unique set of fatty acid regulated transcription factors controlling both lipid metabolism and inflammation. *Biochim. Biophys. Acta—Mol. Basis Dis.* 2011. 1812: 1007–1022.
- 81 Tian, D., Hong, H., Shang, W., Ho, C. C., Dong, J. and Tian, X. Y., Deletion of Ppard in CD11c+ cells attenuates atherosclerosis in ApoE knockout mice. *FASEB J.* 2020. 34: 3367–3378.
- 82 Cao, W., Ramakrishnan, R., Tyurin, V. A., Veglia, F., Condamine, T., Amoscato, A., Mohammadyani, D., et al. Oxidized lipids block antigen cross-presentation by dendritic cells in cancer. *J. Immunol.* 2014. 192: 2920–2931.
- 83 Veglia, F., Tyurin, V. A., Mohammadyani, D., Blasi, M., Duperret, E. K., Donthireddy, L., Hashimoto, A., et al. Lipid bodies containing oxidatively truncated lipids block antigen cross-presentation by dendritic cells in cancer. *Nat. Commun.* 2017. 8: 2122.
- 84 Garris, C. S. and Pittet, M. J., ER stress in dendritic cells promotes cancer. *Cell* 2015. 161: 1492–1493.
- 85 Ugolini, A., Tyurin, V., Tyurina, Y., Tsyganov, E., Donthireddy, L., Kagan, V. E., et al. Polymorphonuclear myeloid-derived suppressor cells limit antigen cross-presentation by dendritic cells in cancer. *JCI Insight.* 2020. 5: e138581.
- 86 Fritsche, K. L., The science of fatty acids and inflammation. *Adv. Nutr.* 2015. 6: 293S–301S.
- 87 Snodgrass, R. G., Bof, M., Zezina, E., Weigert, A., Dehne, N., Fleming, I., Brüne, B. and Namgaladze, D., Hypoxia potentiates palmitate-induced pro-inflammatory activation of primary human macrophages. *J. Biol. Chem.* 2016. 291: 413–424.
- 88 Mogilenko, D. A., Haas, J. T., L’homme, L., Fleury, S., Quemener, S., Levavasseur, M., Becquart, C., et al., Metabolic and innate immune cues merge into a specific inflammatory response via the UPR. *Cell* 2019. 177: 1201–1216.
- 89 Davis, F. M., Dendekker, A., Joshi, A. D., Wolf, S. J., Audu, C., Melvin, W. J., Mangum, K., et al., Palmitate-TLR4 signaling regulates the histone demethylase, JMJD3, in macrophages and impairs diabetic wound healing. *Eur. J. Immunol.* 2020. 50: 1929–1940.
- 90 Lancaster, G. I., Langley, K. G., Berglund, N. A., Kammoun, H. L., Reibe, S., Estevez, E., Weir, J., et al., Evidence that TLR4 is not a receptor for saturated fatty acids but mediates lipid-induced inflammation by reprogramming macrophage metabolism article evidence that TLR4 is not a receptor for saturated fatty acids but mediates lipid-induced inflammation by. *Cell Metab.* 2018. 1–15.
- 91 Zaslona, Z., Serezani, C. H., Okunishi, K., Aronoff, D. M. and Peters-Golden, M., Prostaglandin E2 restrains macrophage maturation via E

- prostanoid receptor 2/protein kinase A signaling. *Blood*. 2012. **119**: 2358–2367.
- 92 Sanin, D. E., Matsushita, M., Klein Geltink, R. I., Grzes, K. M., Van Teijlingen Bakker, N., Corrado, M., Kabat, A. M., et al., Mitochondrial membrane potential regulates nuclear gene expression in macrophages exposed to prostaglandin E2. *Immunity* 2018. **49**: 1021–1033.
- 93 Ramalho, T., Ramalingam, L., Filgueiras, L., Festuccia, W., Jancar, S. and Moustaid-Moussa, N., Leukotriene-B4 modulates macrophage metabolism and fat loss in type 1 diabetic mice. *J. Leukoc. Biol.* 2019. **106**: 665–675.
- 94 Chiu, C. - Y., Gomolka, B., Dierkes, C., Huang, N. R., Schroeder, M., Purschke, M., Manstein, D., et al., Omega-6 docosapentaenoic acid-derived resolvins and 17-hydroxydocosahexaenoic acid modulate macrophage function and alleviate experimental colitis. *Inflamm. Res.* 2012. **61**: 967–976.
- 95 Duffney, P. F., Falsetta, M. L., Rackow, A. R., Thatcher, T. H., Phipps, R. P. and Sime, P. J., Key roles for lipid mediators in the adaptive immune response. *J. Clin. Invest.* 2018. **128**: 2724–2731.
- 96 Zhao, X., Zhang, Y., Strong, R., Grotta, J. C. and Aronowski, J., 15d-Prostaglandin J2 activates peroxisome proliferator-activated receptor- γ , promotes expression of catalase, and reduces inflammation, behavioral dysfunction, and neuronal loss after intracerebral hemorrhage in rats. *J. Cereb. Blood Flow Metab.* 2006. **26**: 811–820.
- 97 Desvergne, B. and Wahli, W., Peroxisome proliferator-activated receptors: nuclear control of metabolism. *Endocr. Rev.* 1999. **20**: 649–688.
- 98 Surh, Y. J., Na, H. K., Park, J. M., Lee, H. N., Kim, W., Yoon, I. S. and Kim, D. - D., 15-Deoxy- Δ 12,14-prostaglandin J2, an electrophilic lipid mediator of anti-inflammatory and pro-resolving signaling. *Biochem. Pharmacol.* 2011. **82**: 1335–1351.
- 99 Bernardo, A., Levi, G. and Minghetti, L., Role of the peroxisome proliferator-activated receptor- γ (PPAR- γ) and its natural ligand 15-deoxy- Δ (12,14)-prostaglandin J2 in the regulation of microglial functions. *Eur. J. Neurosci.* 2000. **12**: 2215–2223.
- 100 Jung, T. W., Park, H. S., Choi, G. H., Kim, D., Ahn, S. H., Kim, D. S., Lee, T. et al. Maresin 1 attenuates pro-inflammatory reactions and ER stress in HUVECs via PPAR α -mediated pathway. *Mol. Cell Biochem.* 2018. **448**: 335–347.
- 101 Vats, D., Mukundan, L., Odegaard, J. I., Zhang, L., Smith, K. L., Morel, C. R., Greaves, D. R., et al., Oxidative metabolism and PGC-1 β attenuate macrophage-mediated inflammation. *Cell Metab.* 2006. **4**: 13–24.
- 102 Spite, M., Clària, J. and Serhan, C. N., Resolvins, specialized proresolving lipid mediators, and their potential roles in metabolic diseases. *Cell Metab.* 2014. **19**: 21–36.
- 103 Arita, M., Bianchini, F., Aliberti, J., Sher, A., Chiang, N., Hong, S., Yang, R., et al., Stereochemical assignment, antiinflammatory properties, and receptor for the omega-3 lipid mediator resolvin E1. *J. Exp. Med.* 2005. **201**: 713–722.
- 104 Krishnamoorthy, S., Recchiuti, A., Chiang, N., Yacoubian, S., Lee, C. - H., Yang, R., Petasis, N. A., et al., Resolvin D1 binds human phagocytes with evidence for proresolving receptors. *Proc. Natl. Acad. Sci. USA* 2010. **107**: 1660–1665.
- 105 Satapati, S., Qian, Y., Wu, M. S., Petrov, A., Dai, Ge, Wang, S. - P., Zhu, Y., et al., GPR120 suppresses adipose tissue lipolysis and synergizes with GPR40 in antidiabetic efficacy. *J. Lipid Res.* 2017. **58**: 1561–1578.
- 106 Ichimura, A., Hirasawa, A., Poulain-Godefroy, O., Bonnefond, A., Hara, T., Yengo, L., Kimura, I., et al., Dysfunction of lipid sensor GPR120 leads to obesity in both mouse and human. *Nature* 2012. **483**: 350–354.
- 107 Davies, L. C., Rice, C. M., Palmieri, E. M., Taylor, P. R., Kuhns, D. B. and McVicar, D. W., Peritoneal tissue-resident macrophages are metabolically poised to engage microbes using tissue-niche fuels. *Nat. Commun.* 2017. **8**: 2074.
- 108 Basit, F., Mathan, T., Sancho, D. and De Vries, J. M., Human dendritic cell subsets undergo distinct metabolic reprogramming for immune response. *Front. Immunol.* 2018. **9**: 1–17.
- 109 Bourdely, P., Anselmi, G., Vaivode, K., Ramos, R. N., Missolo-Koussou, Y., Hidalgo, S., Tosselo, J., et al., Transcriptional and functional analysis of CD1c+ Human dendritic cells identifies a CD163+ subset priming CD8+ CD103+ T cells. *Immunity* 2020. **53**: 335–352.
- 110 Villani, A. - C., Satija, R., Reynolds, G., Sarkizova, S., Shekhar, K., Fletcher, J., Griesbeck, M., et al., Single-cell RNA-seq reveals new types of human blood dendritic cells, monocytes, and progenitors. *Science (80-)* 2017. **356**: eaah4573.
- 111 Galli, E., Friebel, E., Ingelfinger, F., Unger, S., Núñez, N. G. and Becher, B., The end of omics? High dimensional single cell analysis in precision medicine. *Eur. J. Immunol.* 2019. **49**: 212–220.
- 112 Artyomov, M. N. and Van Den Bossche, J., Immunometabolism in the single-cell era. *Cell Metab.* 2020. **32**: 710–725.
- 113 Furuhashi, M., Fucho, R., Görgün, C. Z., Tuncman, G., Cao, H. and Hotamisligil, G. S., Adipocyte/macrophage fatty acid-binding proteins contribute to metabolic deterioration through actions in both macrophages and adipocytes in mice. *J. Clin. Invest.* 2008. **118**: 2640–2650.
- 114 Ménégaut, L., Jalil, A., Thomas, C. and Masson, D., Macrophage fatty acid metabolism and atherosclerosis: the rise of PUFAs. *Atherosclerosis* 2019. **291**: 52–61.
- 115 Ryu, H., Kim, J., Kim, D., Lee, J. - E. and Chung, Y., Cellular and molecular links between autoimmunity and lipid metabolism. *Mol. Cells* 2019. **42**: 747–754.
- 116 Cheng, H. S., Tan, W. R., Low, Z. S., Marvalim, C., Lee, J. Y. H. and Tan, N. S., Exploration and development of PPAR modulators in health and disease: an update of clinical evidence. *Int. J. Mol. Sci.* 2019. **20**: 1–69.
- 117 Hossain, F., Al-Khami, A. A., Wyczechowska, D., Hernandez, C., Zheng, L., Reiss, K., Valle, L. D., et al., Inhibition of fatty acid oxidation modulates immunosuppressive functions of myeloid-derived suppressor cells and enhances cancer therapies. *Cancer Immunol. Res.* 2015. **3**: 1236–1247.
- 118 Serhan, C. N., Clish, C. B., Brannon, J., Colgan, S. P., Chiang, N. and Gronert, K., Novel functional sets of lipid-derived mediators with anti-inflammatory actions generated from omega-3 fatty acids via cyclooxygenase 2-nonsteroidal antiinflammatory drugs and transcellular processing. *J. Exp. Med.* 2000. **192**: 1197–1204.
- 119 Norling, L. V., Headland, S. E., Dalli, J., Arnardottir, H. H., Haworth, O., Jones, H. R., Irimia, D., et al., Proresolving and cartilage-protective actions of resolvin D1 in inflammatory arthritis. *JCI Insight* 2016. **1**: 1–17.
- 120 Valdes, A. M., Ravipati, S., Menni, C., Abhishek, A., Metrusty, S., Harris, J., Nessa, A., et al. Association of the resolvin precursor 17-HDHA, but not D- or E-series resolvins, with heat pain sensitivity and osteoarthritis pain in humans. *Sci. Rep.* 2017. **7**: 1–8.
- 121 Motwani, M. P., Colas, R. A., George, M. J., Flint, J. D., Dalli, J., Richard-Loendt, A., De Maeyer, R. P. H., et al., Pro-resolving mediators promote resolution in a human skin model of UV-killed *Escherichia coli*-driven acute inflammation. *JCI Insight* 2018. **3**: 1–14.
- 122 Emre, C., Hjorth, E., Bharani, K., Carroll, S., Granholm, A. - C., and Schultzberg, M., Receptors for pro-resolving mediators are increased in Alzheimer's disease brain. *Brain Pathol.* 2020. **30**: 614–640.
- 123 Mizwicki, M. T., Liu, G., Fiala, M., Magpantay, L., Sayre, J., Siani, A., Mahanian, M., et al., 1 α ,25-dihydroxyvitamin D3 and resolvin D1 retune the balance between amyloid- β phagocytosis and inflammation in Alzheimer's disease patients. *J. Alzheimer's Dis.* 2013. **34**: 155–170.

- 124 **Arnardottir, H. H., Dalli, J., Norling, L. V., Colas, R. A., Perretti, M. and Serhan, C. N.**, Resolvin D3 is dysregulated in arthritis and reduces arthritic inflammation. *J. Immunol.* 2016. **197**: 2362–2368.
- 125 **Hersberger, M.**, Potential role of the lipoxygenase derived lipid mediators in atherosclerosis: leukotrienes, lipoxins and resolvins. *Clin. Chem. Lab. Med.* 2010. **48**: 1063–1073.
- 126 **De Gaetano, M., McEvoy, C., Andrews, D., Cacace, A., Hunter, J., Brennan, E. and Godson, C.**, Specialized pro-resolving lipid mediators: modulation of diabetes-associated cardio-, reno-, and retino-vascular complications. *Front. Pharmacol.* 2018. **9**: 1488.
- 127 **Krashia, P., Cordella, A., Nobili, A., La Barbera, L., Federici, M., Leuti, A., Campanelli, F., et al.** Blunting neuroinflammation with resolvin D1 prevents early pathology in a rat model of Parkinson's disease. *Nat. Commun.* 2019. **10**: 1–19.

Abbreviations: **Ac-CoA:** acetyl-CoA · **ACLY:** ATP citrate lyase · **CPT:** Carnitine Palmitoyltransferase · **FA:** fatty acid · **FAO:** fatty acid oxidation · **FATP1:** FA transport protein 1 · **GPRs:** G-protein-coupled receptors · **IDO:** indolamine 2,3-dioxygenase-1 · **IL-4R:** IL-4 receptor · **LDs:** lipid droplets · **LTB4:** leukotriene-B4 ·

MSR1: macrophage scavenger receptor 1 · **MUFAs:** monounsaturated · **OA:** oleic acid · **OXPHOS:** oxidative phosphorylation · **PINK1:** PTEN-induced putative kinase 1 · **PGE2:** prostaglandin E2 · **pMacrophages:** peritoneal macrophages · **PPAR α :** peroxisome proliferator-activated receptor alpha · **PUFAs:** polyunsaturated · **RA:** rheumatoid arthritis · **SCFAs:** short-chain fatty acids · **SREBP-1a:** sterol regulatory element binding protein-1a · **SPMs:** specialized proresolving mediators · **SFAs:** saturated fatty acids · **TADCs:** tumour-associated DCs · **TAMs:** tumour-associated macrophages · **UFAs:** unsaturated fatty acids

Full correspondence: Dr. Bart Everts, Department of Parasitology, Leiden University Medical Center, Albinusdreef 2, 2333 ZA Leiden, The Netherlands.
e-mail: b.everts@lumc.nl

Received: 16/11/2020

Revised: 4/3/2021

Accepted: 25/3/2021

Accepted article online: 31/3/2021