



Universiteit
Leiden
The Netherlands

What are mycorrhizal traits?

Chaudhary, V.B.; Holland, E.P.; Charman-Anderson, S.; Guzman, A.; Bell-Dereske, L.; Cheeke, T.E.; ... ; Helgason, T.

Citation

Chaudhary, V. B., Holland, E. P., Charman-Anderson, S., Guzman, A., Bell-Dereske, L., Cheeke, T. E., ... Helgason, T. (2022). What are mycorrhizal traits? *Trends In Ecology & Evolution*, 37(7), 573-581. doi:10.1016/j.tree.2022.04.003

Version: Publisher's Version

License: [Licensed under Article 25fa Copyright Act/Law \(Amendment Taverne\)](#)

Downloaded from: <https://hdl.handle.net/1887/3439723>

Note: To cite this publication please use the final published version (if applicable).

What are mycorrhizal traits?

V. Bala Chaudhary ^{1,*},[@] E. Penelope Holland,² Suw Charman-Anderson,³ Aidee Guzman,⁴ Lukas Bell-Dereske,⁵ Tanya E. Cheeke,⁶ Adriana Corrales,⁷ Jessica Duchicela,⁸ Cameron Egan,⁹ Manju M. Gupta,¹⁰ S. Emilia Hannula,¹¹ Rachel Hestrin,¹² Shabana Hoosein,¹³ Amit Kumar,¹⁴ Genet Mhretu,¹⁵ Lena Neuenkamp,^{16,17} Pushpa Soti,¹⁸ Yichun Xie,¹⁹ and Thorunn Helgason²

Traits are inherent properties of organisms, but how are they defined for organismal networks such as mycorrhizal symbioses? Mycorrhizal symbioses are complex and diverse belowground symbioses between plants and fungi that have proved challenging to fit into a unified and coherent trait framework. We propose an inclusive mycorrhizal trait framework that classifies traits as morphological, physiological, and phenological features that have functional implications for the symbiosis. We further classify mycorrhizal traits by location – plant, fungus, or the symbiosis – which highlights new questions in trait-based mycorrhizal ecology designed to charge and challenge the scientific community. This new framework is an opportunity for researchers to interrogate their data to identify novel insights and gaps in our understanding of mycorrhizal symbioses.

Fitting mycorrhizal symbioses into existing trait-based ecological frameworks

A **trait** (see [Glossary](#)) is defined as a measurable characteristic (morphological, physiological, phenological, behavioral, or cultural) of an individual organism that is measured at either the individual level or another relevant level of organization [1,2]. Plants and animals typically have many distinguishable morphological traits, and, after decades and/or centuries of research, their life histories are generally well described. As a result, conceptual frameworks for trait-based ecology were developed for and are primarily applied to plants [3,4] and animals [5,6], with a proportionate number of plant and animal trait databases emerging to support these efforts [7].

Trait-based approaches are increasingly applied broadly across disciplines within ecology and evolution. Advantages include the ability to make ecological inferences across temporal, spatial, and organizational scales and a predictive understanding of communities and ecosystem processes [8,9]. Commonly used methods employ species traits in order to understand mechanisms behind responses of species to variation in environmental conditions (i.e., response traits) and traits that link species to patterns in ecosystem processes and functioning (i.e., effect traits) [10]. For microbes, a trait-based approach to ecological studies is particularly crucial as many individuals and species are not easily identifiable or culturable for in-depth laboratory studies [11]. Trait-based approaches provide a key to solving ‘big picture’ problems in ecology, such as community responses to anthropogenic global change [5,12] that depend on complex interactions between species and environments, both above and below ground [13,14].

Mycorrhizal symbioses (mycorrhizas) are close associations between roots and certain fungi [15] and, in terrestrial ecosystems, the dominant belowground structures responsible for shuttling the resources (e.g., nutrients, water) that drive primary productivity [16]. In mycorrhizal symbioses, plants provide photosynthetically derived carbon to fungal partners in exchange for increased

Highlights

Applying trait-based approaches to ecological research on mycorrhizal symbioses broadens ecological inferences, but there is no single unified framework to unite disparate language, terminology, and methods across the many multidisciplinary scientists studying mycorrhizas.

We propose an inclusive framework for trait-based mycorrhizal ecology aimed to stimulate scientists around the world to collect and use more mycorrhizal trait data, particularly in understudied areas. This would widen our understanding regarding the ecological role of mycorrhizal symbioses at individual, species, community, and ecosystem scales.

Analyzing how mycorrhizal symbioses fit within existing trait definitions highlights significant theoretical and empirical knowledge gaps, novel questions, and new research directions to improve our understanding of trait-based mycorrhizal ecology.

¹Department of Environmental Studies, Dartmouth College, Hanover, NH 03755, USA

²Department of Biology, University of York, York, YO10 5DD, UK

³Finding Ada Network, Bournemouth BH12 5AG, UK

⁴Department of Environmental Science, Policy, and Management, University of California, Berkeley, Berkeley, CA 94720, USA

⁵Laboratory of Environmental Microbiology, Institute of Microbiology of the Czech Academy of Sciences, Videňská 1083, 142 20 Prague, Czech Republic

⁶School of Biological Sciences, Washington State University, Richland, WA 99354, USA

⁷Centro de Investigaciones en Microbiología y Biotecnología-UR (CIMBIUR), Facultad de Ciencias



Box 1. Wider applications of the framework

Our framework for mycorrhizal traits can be applied to easily incorporate trait-based methods into empirical and theoretical ecological research. Adopting a trait-based framework for mycorrhizal symbioses benefits ecologists from a variety of disciplinary backgrounds.

Plant ecologist

Plant ecologists use existing frameworks for measuring traits and incorporating trait-based methods into ecological studies [32,52,82]. Plant ecologists already measure many plant mycorrhizal traits such as root architecture, photosynthetic pathway, and phenology (see Table 2 in main text). By also including symbiotic mycorrhizal traits such as colonization intensity, plant mycorrhizal response, or resource exchange rates, plant ecologists could further increase their understanding of plant functioning. For instance, examining mycorrhizal colonization intensity in plant roots would facilitate inferences about carbon and nutrient transfer between plant and fungal symbionts, with links to functioning such as plant productivity or pathogen resistance [54,83].

Fungal ecologist

Fungal ecologists have long used traits to categorize fungi according to guilds, and continue to use trait-based perspectives to research the numerous functional roles that fungi play in ecosystems [24]. As methodologies to assess fungi *in situ* continue to improve, we can better measure many fungal mycorrhizal traits such as mycelial traits and enzyme activity (see Table 2 in main text) [11]. Many fungal mycorrhizal traits can be measured using standard laboratory equipment (e.g., centrifuge, filters, microscope) that researchers already have access to. For example, spore size is an indicator of AM fungal aerial dispersal ability, and thus could improve predictions of landscape management impacts on local AM fungal diversity and composition [69]. Just as leaf traits have expanded knowledge of plant life-history strategies [84], the incorporation of important fungal mycorrhizal traits such as spore morphology will expand our understanding of life-history strategies of mycorrhizal fungi.

Data synthesizer

Large-team science to compile and analyze global ecological datasets will increase our understanding of biodiversity and ecosystem functioning. Ecologists examining ecological phenomena across spatial and temporal scales can incorporate mycorrhizal traits to improve our understanding of global trends in mycorrhizal symbioses. For example, merging data on symbiotic mycorrhizal traits such as mycorrhizal type or plant mycorrhizal response from FungalRoot [31] or MycoDB [29] into other ecological synthesis efforts could reveal novel ways to predict global ecological biodiversity and ecosystem function. Furthermore, because many ecological data comprise repeated sampling (e.g., LTER, NEON), they represent an opportunity to monitor understudied phenological mycorrhizal traits, such as shifts in mycorrhizal type, or mycorrhizal influences on plant reproductive phenology.

Naturales, Universidad del Rosario, Bogotá 110151, Colombia

⁸Departamento de Ciencias de la Vida, Universidad de las Fuerzas Armadas ESPE, Sangolquí 171103, Ecuador

⁹Department of Biology, Okanagan College, 1000 KLO Rd, Kelowna, BC, Canada V1Y 4X8

¹⁰Department of Biology, University of Delhi, Sri Aurobindo College, Delhi 110017, India

¹¹Institute of Environmental Sciences, Leiden University, Leiden 2333, The Netherlands

¹²Physical and Life Sciences Directorate, Lawrence Livermore National Laboratory, Livermore, CA 94550, USA

¹³Department of Forest and Rangeland Stewardship, Colorado State University, Fort Collins, CO 80523, USA

¹⁴Institute of Ecology, Faculty of Sustainability, Leuphana University of Lüneburg, 21335 Lüneburg, Germany

¹⁵Department of Biology, Mekelle University, Mekelle 231, Ethiopia

¹⁶University of Bern, Institute of Plant Sciences, Berne 3013, Switzerland

¹⁷Department of Ecology and Multidisciplinary Institute for Environment Studies 'Ramon Margalef', University of Alicante, Alicante 03009, Spain

¹⁸Biology Department, University of Texas Rio Grande Valley, Edinburg, TX 78539, USA

¹⁹School of Life Sciences, The Chinese University of Hong Kong, Hong Kong SAR 999077

access to soil resources such as nitrogen, phosphorus, and water, although the degree of mutual benefit is context dependent [17]. Mycorrhizal symbioses are most well known for their role in nutrient exchange, but are also recognized for their key roles in ecosystems across a range of organizational scales such as promoting plant establishment, plant pathogen protection, plant resistance to heavy metals, drought tolerance, interspecific community interactions, soil aggregation, and global carbon cycling [18]. Mycorrhizal symbioses represent the interface between two different types of modular lifeforms; in nature, most plant roots are associated with more than one **mycorrhizal fungus**. Furthermore, one mycorrhizal fungus can be associated with multiple plants to form nonrandom assemblages of physical networks of **hyphae** that are connected below ground [19,20]. As such, mycorrhizal symbioses are root–mycelial networks of modular lifeforms with varying degrees of complexity ranging from one plant–one fungus to multiple plant–fungal connections.

Applying concepts from, and drawing parallels with, trait-based ecological theory developed for individual organisms with more definable traits can be challenging for symbioses that are inherently defined as associations between multiple organisms. Traits are often defined for the purpose of a specific study, and so terminology, semantics, and interpretations vary across datasets [7,21], even for well-studied and easily identifiable organisms. For organisms with high species-level diversity but few distinguishable morphological traits, such as fungi [22], trait-based ecology often takes a more mechanistic approach, particularly for species that are cryptic or microscopic [23,24]. For symbioses that are not discrete species units but in fact emergent properties of complex root–

*Correspondence: bala.chaudhary@dartmouth.edu (V.B. Chaudhary).
[©]Twitter: @balachaudhary

mycelial networks, such as mycorrhizal symbioses (but see also [25]), trait-based ecology poses an even greater challenge.

Prior morpho–physio–phenological trait definitions for mycorrhizas

Mycorrhizal ecologists, whether invoking the word ‘trait’ or not, have long studied various **mycorrhizal traits** to gain insight into predictors or proxies of mycorrhizal performance [26]. Table 1 lists examples of previously used definitions of morphological, physiological, or phenological traits that have led to considerable advances in our understanding of mycorrhizal ecology. Mycorrhizal type is an emergent property of the plant and fungal taxa involved in the **symbiosis**, and is likely the most commonly studied mycorrhizal trait. The major types of mycorrhizas – **arbuscular mycorrhiza (AM)**, **ectomycorrhiza (EcM)**, **ericoid mycorrhiza (ErM)**, and **orchid mycorrhiza (OrM)** – are similar in that they are all symbiotic root–mycelial networks of fungi and plants with varying degrees of complexity. However, mycorrhizal types vary substantially with respect to plant and fungal taxa involved in the association, morphological form, ecophysiological function, and their comparative roles in biogeochemical cycling [16,27]. Research on mycorrhizal type, among other mycorrhizal traits, is increasingly being conducted with large global databases (e.g., MycoDB, FungalRoot, FUN^{FUN}, FungalTraits) aimed at making broad inferences about the biogeography and functioning of mycorrhizas [24,28–31].

The application of existing ecological conceptual frameworks has also led to advances in trait-based mycorrhizal ecology, in particular the application of Grime’s C–S–R (competitor–stress tolerator–ruderal) framework [32,33]. A fungal-centric perspective characterizes variation in AM fungal traits such as hyphal growth rate, hyphal turnover rate, **spore** phenology, and dispersal ability as alternative competitive strategies for different AM fungal species [34]. Efforts have also been made to classify AM fungi into edaphophilic or rhizophilic guilds related to differential allocation to soil hyphae or root colonization, respectively, and how that relates to mycorrhizal function [35,36]. In EcM fungi, previous work has defined mycorrhizal traits as differences in morphology and physiology of mycelial [37] and reproductive structures [38] that produce differences in species’ capacity for carbon storage, enzymatic activity, nutrient uptake and translocation, dispersal, and habitat colonization [27,39–42]. Alternatively, plant-centric perspectives have shown how mycorrhizal symbioses explain significant variation in plant life history strategies [43] and multivariate root trait space [44], accounting for different C–S–R and resource utilization strategies of plants across the globe.

Methodological limitations (e.g., culturing bias) certainly impair empirically derived knowledge of mycorrhizal traits [45], but disparate definitions across a diversity of trait-based mycorrhizal

Glossary

Arbuscular mycorrhiza (AM): a mycorrhizal association where plant roots display intracellular colonization by fungi of the subphylum Glomeromycotina.

Arbuscule: a specialized mycorrhizal structure present inside plant cells, and the common site of nutrient exchange in arbuscular mycorrhizas (AMs). Other nutrient exchange sites in arbuscular mycorrhizas include hyphal coils.

Dual colonization: colonization of plant roots by two different mycorrhizal types (i.e., AM and EcM), generally demonstrating ontological shifts in particular plant species (e.g., *Quercus* sp., *Salix* sp., *Populus* sp.).

Ectomycorrhiza (EcM): a mycorrhizal association between plant roots and fungi characterized by an intercellular interface consisting of a branched hyphal lattice and mantle.

Ericoid mycorrhiza (ErM): a mycorrhizal association between plants in the family Ericaceae and certain fungi, characterized by intracellular coils.

Functional markers: traits that do not measure a function directly but instead are indicators of functions (e.g., hyphal production by mycorrhizal fungi that influences soil aggregate formation).

Hyphae: the branching filaments of mycorrhizal fungi that make up the mycelium and conjoin to plant roots either intracellularly or extracellularly. Hyphae differ with respect to morphology, environmental persistence, and function (e.g., nutrient absorption vs. transport).

Mantle: sheath of fungal hyphae enveloping plant roots in EcM associations.

Mycorrhizal fungus: the fungal symbiotic partner in a mycorrhizal association.

Mycorrhizal symbioses: symbiotic associations between plant roots and certain fungi. Synonym: mycorrhiza.

Mycorrhizal traits: morphological, physiological, or phenological characteristics of mycorrhizal fungi, plants, and mycorrhizal associations that have functional implications for the symbiosis.

Orchid mycorrhiza (OrM): mycorrhizal association between plants in the family Orchidaceae and certain fungi characterized by intracellular coils called pelotons.

Paris/Arum: alternative root colonization strategies in arbuscular mycorrhizas. The Paris type is

Table 1. Examples of previously published and highly varied definitions of ‘mycorrhizal traits’

Definitions of mycorrhizal traits	Refs
Traits as the type of mycorrhizal symbiosis (e.g., AM, EcM, ErM) or frequency of occurrence (e.g., obligate, facultative) of mycorrhizal symbiosis in a plant species	[31,44,60–64]
Traits as the context-dependent benefits that plants derive from mycorrhizal symbioses	[17,65]
Traits as spore morphology (e.g., size, shape, color) of mycorrhizal fungi	[66–69]
Traits as root and/or soil colonization strategies of mycorrhizal fungi, including fungal biomass allocation and hyphal production	[36,70–73]
Traits as soil aggregation and stabilization capabilities of mycorrhizal symbioses	[74,75]
Traits as C–S–R characteristics of mycorrhizal fungi	[34,76]
Traits as mycorrhizal fungal behaviors such as movement, communication, and decision-making	[77,78]
Traits as mycorrhizal symbiosis properties related to nutrient flux and ecosystem functioning	[79–81]

research efforts also hinder productive scientific discourse. Often, different definitions of mycorrhizal traits are specific to mycorrhizal type, focused on either a plant-centric or a fungal-centric perspective, or are borrowed from existing ecological theories based on distinct unitary organisms, causing confusion for network-based symbioses between modular organisms. A unified language for mycorrhizal traits that spans mycorrhizal types and morpho-physio-phenological characteristics is sorely needed.

Controversy and disagreement in what constitutes a mycorrhizal trait

Thus far, the body of research on trait-based mycorrhizal ecology has used different definitions of traits, with organismal divides stemming from different morphological metrics that are a proxy for functions, direct measures of functions, or measures of mycorrhizal plant and/or mycorrhizal fungal growth that may also approximate function. In many ways, these organismal divides are a result of researchers coming from different disciplinary backgrounds and perspectives [46,47]. A plant-centric perspective can result in studying different types of traits and the use of varied vocabularies that do not easily translate to those using a fungal-centric perspective (and vice versa). Many microbes have relatively few measurable or easily observable features, and thus functional measures are translated into traits. For example, the presence or abundance of saprotrophic fungi may be correlated with litter decomposition rates. Plant traits, on the contrary, are observable, but their relevance for the functioning of the ecosystem may be ambiguous. For example, specific leaf area can be an indication of plant longevity and hence biomass turnover and photosynthetic rates [10]. Geographic region of study can also drive miscommunication as certain regions of the world (particularly the tropics) are comparatively understudied [48], resulting in a greater need to incorporate local terminologies into globally accepted paradigms. Indeed, inconsistencies in terminology surrounding traits are as diverse as trait ecologists, suggesting the need to keep trait definitions broad, malleable, and identified independently from the environment [2].

Divides also exist between researchers who work primarily in EcM-dominated systems and those working in AM- or ErM-dominated systems. Furthermore, certain systems have been studied for longer, as EcM symbioses were identified in the 1880s [15], but the functional significance of the AM symbioses was not discovered until the 1950s [49]. Less is known about AM symbioses compared with other mycorrhizal groups [50], resulting in a lack of the basic biological and taxonomic framework to integrate ecological research with general mycology. This affects the study of mycorrhizal symbioses as there are significant differences in the focus of AM versus EcM studies, hampering the generation of a unified language to describe them [11,47]. Applying trait-based methods to highly context-dependent mycorrhizal symbioses without a standardized vocabulary is challenging and can result in 'locked-in debates' among researchers that hinder scientific advances [51].

An inclusive and unified framework for mycorrhizal traits

It is the opinion of the authors that a common framework and standardized vocabulary will help to further our understanding of the trait-based ecology of mycorrhizal symbioses. As traits are characteristics of organisms, mycorrhizal traits must be inclusive of all organismal components that make up mycorrhizal symbioses, recognizing that the mycorrhizal functions we observe in nature are the imprint of all mycorrhizal traits working together. Therefore, mycorrhizal traits are morphological, physiological, or phenological characteristics of mycorrhizal fungi, plants, and mycorrhizal associations that have functional implications for the mycorrhizal symbiosis. Because this definition is based on symbiotic function and the inherent root-mycelial network nature of all mycorrhizas, it is applicable across all mycorrhizal types. Our definition emphasizes traits that have functional implications for the mycorrhizal symbiosis to further a mechanistic understanding of mycorrhizal performance and fitness. We aim to link trait-based mycorrhizal ecology to the

characterized by coiled hyphae that spread intracellularly from plant cortical cell to cell, while the Arum type spreads in the plant root cortex via intercellular hyphae.

Spore: fungal cells specialized for asexual or sexual reproduction and dispersal. They can be borne on specialized fungal fruiting bodies or directly from mycelial networks.

Symbiosis: an association between organisms that live in close physical contact.

Trait: any measurable characteristic (morphological, physiological, phenological, behavioral, or cultural) of an individual organism that is measured at either the individual or other relevant level of organization.

work of defining mycorrhizal niches and understanding the mechanisms of community assembly [9]. Some mycorrhizal traits are functions (e.g., plant productivity response to mycorrhizal symbiosis), but some are **functional markers**, traits that do not measure a function directly but instead are indicators of mycorrhizal functions (e.g., hyphal production by mycorrhizal fungi that influences soil aggregate formation) [11,52].

Although there are benefits to using inclusive terminology, an overly broad definition of mycorrhizal traits can also cause confusion during scientific discourse when researchers universally refer to 'mycorrhizal traits' but mean different things. We propose to further qualify mycorrhizal traits using language that references their physical location within mycorrhizal networks (Table 2). Therefore, mycorrhizal traits fall into one of three categories: plant mycorrhizal traits, fungal mycorrhizal traits, and symbiotic mycorrhizal traits.

Plant mycorrhizal traits are mycorrhizal traits that are driven largely by the morphological, physiological, or phenological characteristics of the plant partner. Many root traits, for example, represent important plant mycorrhizal traits as they have functional implications for the symbiosis [44]. Fungal mycorrhizal traits are mycorrhizal traits that are dependent on the morphological, physiological, or phenological characteristics of the mycorrhizal fungal partners. Both fungal response and fungal effect traits [23] – particularly physiological traits relating to fungal ecosystem functions – are components of fungal mycorrhizal traits. Finally, symbiotic mycorrhizal traits are morphological, physiological, or phenological characteristics that lie at the intersection of both partners and are dependent on both the plant and fungal partners present. Figure 1 in main

Table 2. An inclusive and unified framework for mycorrhizal traits. Examples given are categorized as morphological, physiological, or phenological traits; the framework is intended to stimulate thought and discussion, so dynamic classifications are encouraged

	Plant mycorrhizal traits	Fungal mycorrhizal traits	Symbiotic mycorrhizal traits
Definition	Mycorrhizal traits dependent on the morphological, physiological, or phenological characteristics of plant partners	Mycorrhizal traits dependent on the morphological, physiological, or phenological characteristics of the fungal partners	Traits that lie at the organismal intersection of mycorrhizal symbioses and are dependent on both plant and fungal partners
Morphological traits (form)	Root characteristics (e.g., diameter, architecture, surface area:volume, root hair density) Root:shoot ratio Growth form (e.g., tree, grass) Resource allocation (e.g., root:shoot) Seed size Phylogenetic history	Fruiting body (e.g., size, shape, color) Spores (e.g., size, color, shape, ornamentation, wall thickness) Mantle (e.g., color, cell morphology) Hyphae (e.g., specific length, architecture) Biomass allocation strategy (e.g., rhizophilic, edaphilic) Culturability	Mycorrhizal type (AM, EcM, ErM, OrM, non-mycorrhizal, dual colonization) Colonization intensity (e.g., abundance of intercellular and intracellular structures) Structures induced by colonization (e.g., Hartig net, arbuscules , vesicles, Paris/Arum form) Species specificity between plant and fungal symbionts Network indices (e.g., nestedness, modularity, connectivity)
Physiological traits (function)	Plant mycorrhizal status (obligate vs. facultative) Photosynthetic pathway Immune responses (e.g., herbivore-induced responses) Growth and transpiration rates Quantity and quality of root exudates Plant nutrient requirements	Hyphal/spore productivity and turnover Nutrient acquisition strategy (e.g., inorganic vs. organic sources, extracellular enzyme production, acid exudation) Melanin content Carbohydrate metabolism and conversion Facilitative/antagonistic interactions with microorganisms	Plant mycorrhizal response (e.g., increased productivity or nutrient status) Exchange rates for resources (e.g., N, P, C, H ₂ O) Gene expression changes induced by symbiosis Plant-fungal influences on metabolic products Functional specificity between plant and fungal symbionts
Phenological traits	Life history (e.g., annual, perennial) Flowering time and seed production Changes in root exudate quality and quantity	Temporal dynamics in production of fruiting bodies, spores, and hyphae Hyphal/spore persistence and longevity Temporal dynamics in fungal community structure	Shifts in mycorrhiza type over plant lifespan Temporal shifts in colonization structures and/or symbiotic exchange




Figure 1. Schematic diagram of plant mycorrhizal traits, fungal mycorrhizal traits, and symbiotic mycorrhizal traits, with graphic depictions of example morphological, physiological, and phenological traits for each.

text shows examples of plant, fungal, and symbiotic mycorrhizal traits across morphological, physiological, and phenological traits.

This framework both accommodates existing trait-based research and identifies gaps in our knowledge due to data limitations. For example, considerably more research has been conducted on morphological mycorrhizal traits than on physiological mycorrhizal traits, and phenological traits are by far the least studied. More research into how plant, fungal, and symbiotic mycorrhizal traits shift with seasons or ontogeny will give greater insight into the range of variation in traits. This trait-based framework also highlights how little we understand about interspecific and intraspecific variation, as well as plasticity in many mycorrhizal traits, particularly at the physiological level. Computational methods linking genes to traits [53] could be employed to explore relationships between plant, fungal, and/or symbiotic mycorrhizal traits and either plant or mycorrhizal fungal gene frequencies (Box 1). Further exploration of relationships between

traits – particularly those that illuminate symbiotic partner resource sharing and connections between mycorrhizal form and function – is warranted, as mycorrhizas are models for studying resource exchange and stability in symbioses [54].

Concluding remarks

There is already consensus among scientists using traits in ecology that standardized definitions and data structures are required to make the most of trait data and to address challenges at the community and ecosystem levels [7,21]. Microorganisms influence almost all ecosystem processes, and a common framework for research into how microbial processes affect ecosystem-level function is crucial for advancing our understanding [55]. Mycorrhizal symbioses occupy a unique and complex position in ecological communities, with a pivotal role in the maintenance of ecosystem function [56], and will be fundamental to meeting United Nations Sustainable Development Goals in the medium to long term [57].

The cha-cha-cha theory suggests that scientific discoveries can be classified as charge, challenge, or chance [58]. Charge problems are obvious to the observer, but require a new way of thinking to devise a solution. Challenge problems require us to devise a new theory to bring unexplained and diverse anomalies together. Chance discoveries require a ‘prepared mind’ to recognize the importance of something that happens by chance. Our framework for mycorrhizal traits raises numerous outstanding questions as charges and challenges to the ecological community in order to be better prepared to recognize future chance discoveries (see [Outstanding questions](#)). By acknowledging how our position of observation flavors our analyses and understanding of mycorrhizal traits through the very language we use to pose research questions [59], we can, as a community of scientists, be better prepared to recognize serendipitous discoveries. A common framework for mycorrhizal traits may engage scientists around the world to collect more trait-based data, especially in understudied areas, generating chance discoveries. It is the authors’ opinion that a common framework for mycorrhizal trait-based ecology will facilitate the next generation of discoveries in this field. This paper describes only a small portion of the exciting work tackling charges at the present time. Shared terminology allows us to better identify synergy between studies approaching similar questions from different angles, and to take on the challenges.

Acknowledgments

Concepts for this paper were conceived during iCOMET, a workshop and international Collaboration on Mycorrhizal Ecological Traits held virtually in March 2020. The authors are grateful to Linh Anh Cat and Natasa Sibanc for contributing to initial discussions, and to Alastair Fitter for providing helpful comments on the manuscript. We thank Pierre-Luc Chagnon, Guillermo Bueno, and two anonymous reviewers for constructive comments that greatly improved the manuscript. We also appreciate and recognize research that could not be included in this paper due to space limitations. The iCOMET workshop was supported by the Natural Environmental Research Council (grant number NE/S008543/1). V.B.C. is supported by the National Science Foundation (award DEB-1844531). L.N. was supported by the Marie Skłodowska Curie Individual Fellowship (MYFUN, Grant agreement ID: 835472).

Declaration of interests

No interests are declared.

References

1. Violle, C. *et al.* (2007) Let the concept of trait be functional! *Oikos* 116, 882–892
2. Dawson, S.K. *et al.* (2022) The traits of “trait ecologists”: an analysis of the use of trait and functional trait terminology. *Ecol. Evol.* 11, 16434–16445
3. Funk, J.L. *et al.* (2008) Restoration through reassembly: plant traits and invasion resistance. *Trends Ecol. Evol.* 23, 695–703
4. Grime, J.P. and Hunt, R. (1975) Relative growth-rate: its range and adaptive significance in a local flora. *J. Ecol.* 63, 393–422
5. Rydning, S. *et al.* (2021) Shape-shifting: changing animal morphologies as a response to climatic warming. *Trends Ecol. Evol.* 36, 1036–1048
6. Stearns, S.C. (1989) Trade-offs in life-history evolution. *Funct. Ecol.* 3, 259–268

Outstanding questions

Where are the research gaps in trait-based mycorrhizal ecology? What new ecological knowledge about mycorrhizal symbioses can be generated by examining multiple mycorrhizal traits across multiple categories, plant, fungal, and symbiotic mycorrhizal traits? Future theoretical and empirical work must consider traits inclusive of all components of mycorrhizal root–mycelial networks that are relevant to the ecological question at hand.

Can a trait-based framework drive novel approaches to linking plant and fungal measurements that are meaningful for the biology of mycorrhizal symbioses? What new experimental systems can be imagined to better measure mycorrhizal traits and understand mycorrhizal ecology *in situ*? What accessible (and affordable) methods can be broadly used across systems to fill knowledge gaps, particularly in understudied regions of the world?

What is the relationship between form and function in mycorrhizal symbioses? Do morphological traits of mycorrhizal plants, mycorrhizal fungi, or the symbiosis predict mycorrhizal functions or behaviors?

Are mycorrhizal traits positively or negatively related to each other? Are tradeoffs more likely to exist between traits belonging to the same mycorrhizal trait category? A trait framework helps to differentiate the origins of trade resources, which can reveal tradeoffs that may exist between traits with shared resource allocation strategies.

Temporal changes in plant traits are well studied, but how do fungal and symbiotic mycorrhizal traits interact with plant mycorrhizal trait phenology? How do relationships between mycorrhizal traits vary temporally? Certain mycorrhizal traits shift phenologically, but temporal patterns are underexplored for most mycorrhizal traits, particularly in long-lived plants, long-lived fungi, and their mycorrhizal associations.

Are plant, fungal, and symbiotic mycorrhizal traits phylogenetically conserved? What is the degree of interspecific and intraspecific variation in mycorrhizal traits, and can mycorrhizal

7. Kearney, M.R. *et al.* (2021) Where do functional traits come from? The role of theory and models. *Funct. Ecol.* 35, 1385–1396
8. Laughlin, D.C. (2014) The intrinsic dimensionality of plant traits and its relevance to community assembly. *J. Ecol.* 102, 186–193
9. McGill, B.J. *et al.* (2006) Rebuilding community ecology from functional traits. *Trends Ecol. Evol.* 21, 178–185
10. Lavorel, S. and Garnier, E. (2002) Predicting changes in community composition and ecosystem functioning from plant traits: revisiting the holy grail. *Funct. Ecol.* 16, 545–556
11. Lekberg, Y. and Helgason, T. (2018) In situ mycorrhizal function—knowledge gaps and future directions. *New Phytol.* 220, 957–962
12. Harvey, B.P. *et al.* (2022) Predicting responses to marine heatwaves using functional traits. *Trends Ecol. Evol.* 37, 20–29
13. Freschet, G.T. *et al.* (2021) Root traits as drivers of plant and ecosystem functioning: current understanding, pitfalls and future research needs. *New Phytol.* 232, 1123–1158
14. Laughlin, D.C. *et al.* (2021) Root traits explain plant species distributions along climatic gradients yet challenge the nature of ecological trade-offs. *Nat. Ecol. Evol.* 5, 1123–1134
15. Frank, B. (2005) On the nutritional dependence of certain trees on root symbiosis with belowground fungi (an English translation of A.B. Frank's classic paper of 1885). *Mycorrhiza* 15, 267–275
16. Smith, S.E. and Read, D.J. (2008) *Mycorrhizal symbiosis*, Elsevier
17. Hoeksema, J.D. *et al.* (2018) Evolutionary history of plant hosts and fungal symbionts predicts the strength of mycorrhizal mutualism. *Commun. Biol.* 1, 116
18. Delavaux, C.S. *et al.* (2017) Beyond nutrients: a meta-analysis of the diverse effects of arbuscular mycorrhizal fungi on plants and soils. *Ecology* 98, 2111–2119
19. van der Heijden, M.G. *et al.* (2015) Mycorrhizal ecology and evolution: the past, the present, and the future. *New Phytol.* 205, 1406–1423
20. Davison, J. *et al.* (2011) Arbuscular mycorrhizal fungal communities in plant roots are not random assemblages. *FEMS Microbiol. Ecol.* 78, 103–115
21. Schneider, F.D. *et al.* (2019) Towards an ecological trait-data standard. *Methods Ecol. Evol.* 10, 2006–2019
22. Hawksworth, D.L. *et al.* (2017) Fungal diversity revisited: 2.2 to 3.8 million species. *Microbiol. Spectr.* 5, 5.4.10
23. Crowther, T.W. *et al.* (2014) Untangling the fungal niche: the trait-based approach. *Front. Microbiol.* 5, 579
24. Zanne, A.E. *et al.* (2019) Fungal functional ecology: bringing a trait-based approach to plant-associated fungi. *Biol. Rev.* 95, 409–433
25. Ellis, C.J. *et al.* (2021) Functional traits in lichen ecology: a review of challenge and opportunity. *Microorganisms* 9, 766
26. Koide, R.T. and Mosse, B. (2004) A history of research on arbuscular mycorrhiza. *Mycorrhiza* 14, 145–163
27. Tedersoo, L. and Bahram, M. (2019) Mycorrhizal types differ in ecophysiology and alter plant nutrition and soil processes. *Biol. Rev.* 94, 1857–1880
28. Bueno, C.G. *et al.* (2017) Plant mycorrhizal status, but not type, shifts with latitude and elevation in Europe. *Glob. Ecol. Biogeogr.* 26, 690–699
29. Chaudhary, V.B. *et al.* (2016) Mycobd, a global database of plant response to mycorrhizal fungi. *Sci. Data* 3, 160028
30. Põlme, S. *et al.* (2020) Fungaltraits: a user-friendly traits database of fungi and fungus-like stramenopiles. *Fungal Divers.* 105, 1–16
31. Soudzilovskaia, N.A. *et al.* (2020) Fungalroot: global online database of plant mycorrhizal associations. *New Phytol.* 227, 955–966
32. Grime, J.P. (1977) Evidence for the existence of three primary strategies in plants and its relevance to ecological and evolutionary theory. *Am. Nat.* 111, 1169–1194
33. Treseder, K.K. *et al.* (2018) Arbuscular mycorrhizal fungi as mediators of ecosystem responses to nitrogen deposition: a trait-based predictive framework. *J. Ecol.* 106, 480–489
34. Chagnon, P.-L. *et al.* (2013) A trait-based framework to understand life history of mycorrhizal fungi. *Trends Plant Sci.* 18, 484–491
35. Phillips, M.L. *et al.* (2019) Fungal community assembly in soils and roots under plant invasion and nitrogen deposition. *Fungal Ecol.* 40, 107–117
36. Weber, S.E. *et al.* (2019) Responses of arbuscular mycorrhizal fungi to multiple coinciding global change drivers. *Fungal Ecol.* 40, 62–71
37. Fernandez, C.W. and Kennedy, P.G. (2018) Melanization of mycorrhizal fungal necromass structures microbial decomposer communities. *J. Ecol.* 106, 468–479
38. Agerer, R. (2006) Fungal relationships and structural identity of their ectomycorrhizae. *Mycol. Prog.* 5, 67–107
39. Courty, P.-E. *et al.* (2010) The role of ectomycorrhizal communities in forest ecosystem processes: new perspectives and emerging concepts. *Soil Biol. Biochem.* 42, 679–698
40. Hobbie, E.A. and Agerer, R. (2010) Nitrogen isotopes in ectomycorrhizal sporocarps correspond to belowground exploration types. *Plant Soil* 327, 71–83
41. Peay, K.G. *et al.* (2012) Measuring ectomycorrhizal fungal dispersal: macroecological patterns driven by microscopic propagules. *Mol. Ecol.* 21, 4122–4136
42. Lilleskov, E.A. *et al.* (2011) Conservation of ectomycorrhizal fungi: exploring the linkages between functional and taxonomic responses to anthropogenic N deposition. *Fungal Ecol.* 4, 174–183
43. Bauer, J.T. *et al.* (2017) Ecology of floristic quality assessment: testing for correlations between coefficients of conservatism, species traits and mycorrhizal responsiveness. *AoB PLANTS* 10, plx073
44. Bergmann, J. *et al.* (2020) The fungal collaboration gradient dominates the root economics space in plants. *Science. Advances* 6, eaba3756
45. Ohsowski, B.M. *et al.* (2014) Where the wild things are: looking for uncultured glomeromycota. *New Phytol.* 204, 171–179
46. Brundrett, M. and Tedersoo, L. (2019) Misdiagnosis of mycorrhizas and inappropriate recycling of data can lead to false conclusions. *New Phytol.* 221, 18–24
47. Bueno, C.G. *et al.* (2019) Misdiagnosis and uncritical use of plant mycorrhizal data are not the only elephants in the room. *New Phytol.* 224, 1415–1418
48. Pagano, M.C. and Lugo, M.A. (2019) *Mycorrhizal fungi in South America*, Springer
49. Mosse, B. (1953) Fructifications associated with mycorrhizal strawberry roots. *Nature* 171, 974
50. Fitter, A.H. *et al.* (2011) Nutritional exchanges in the arbuscular mycorrhizal symbiosis: implications for sustainable agriculture. *Fungal Biol. Rev.* 25, 68–72
51. Norberg, J. *et al.* (2022) Failures to disagree are essential for environmental science to effectively influence policy development. *Ecol. Lett.* 25, 1075–1093. Available from: <https://doi.org/10.1111/ele.13984>
52. Garnier, E. *et al.* (2016) *Plant Functional Diversity: Organism Traits, Community Structure, and Ecosystem Properties*, Oxford University Press
53. Romero-Olivares, A.L. *et al.* (2021) Linking genes to traits in fungi. *Microb. Ecol.* 82, 145–155
54. Kiers, E.T. *et al.* (2011) Reciprocal rewards stabilize cooperation in the mycorrhizal symbiosis. *Science* 333, 880–882
55. Malik, A.A. *et al.* (2020) Defining trait-based microbial strategies with consequences for soil carbon cycling under climate change. *ISME J.* 14, 1–9
56. Leake, J. *et al.* (2004) Networks of power and influence: the role of mycorrhizal mycelium in controlling plant communities and agroecosystem functioning. *Can. J. Bot.* 82, 1016–1045
57. Field, K.J. *et al.* (2021) Mycorrhizal mediation of sustainable development goals. *Plants People Planet* 3, 430–432
58. Koshland, D.E. (2007) The cha-cha-cha theory of scientific discovery. *Science* 317, 761–762
59. Smith, F.A. and Smith, S.E. (2015) How harmonious are arbuscular mycorrhizal symbioses? Inconsistent concepts reflect different mindsets as well as results. *New Phytol.* 205, 1381–1384
60. Wang, B. and Qiu, Y.-L. (2006) Phylogenetic distribution and evolution of mycorrhizas in land plants. *Mycorrhiza* 16, 299–363

function be predicted by plant or fungal partner phylogeny?

Can knowledge of mycorrhizal traits influence the conservation and management of mycorrhizal symbioses in natural and managed ecosystems? Using traits to predict mycorrhizal species distributions, dispersal, and survival will improve our ability to protect and restore these important interaction networks in a changing world.

61. Hempel, S. *et al.* (2013) Mycorrhizas in the central European flora: relationships with plant life history traits and ecology. *Ecology* 94, 1389–1399
62. Moora, M. (2014) Mycorrhizal traits and plant communities: perspectives for integration. *J. Veg. Sci.* 25, 1126–1132
63. Shi, Z. *et al.* (2020) The worldwide leaf economic spectrum traits are closely linked with mycorrhizal traits. *Fungal Ecol.* 43, 100877
64. Bueno, C.G. *et al.* (2021) Towards a consistent benchmark for plant mycorrhizal association databases. *New Phytol.* 231, 913–916
65. Johnson, N.C. *et al.* (2015) Mycorrhizal phenotypes and the law of the minimum. *New Phytol.* 205, 1473–1484
66. Norros, V. *et al.* (2014) Do small spores disperse further than large spores? *Ecology* 95, 1612–1621
67. Pringle, A. *et al.* (2015) The shape of fungal ecology: does spore morphology give clues to a species' niche? *Fungal Ecol.* 17, 213–216
68. Deveautour, C. *et al.* (2019) Biogeography of arbuscular mycorrhizal fungal spore traits along an aridity gradient, and responses to experimental rainfall manipulation. *Fungal Ecol.* 46, 100899
69. Chaudhary, V.B. *et al.* (2020) Trait-based aerial dispersal of arbuscular mycorrhizal fungi. *New Phytol.* 228, 238–252
70. Agerer, R. (2001) Exploration types of ectomycorrhizae. *Mycorrhiza* 11, 107–114
71. Hart, M.M. and Reader, R.J. (2002) Taxonomic basis for variation in the colonization strategy of arbuscular mycorrhizal fungi. *New Phytol.* 153, 335–344
72. Ekblad, A. *et al.* (2013) The production and turnover of extramatrical mycelium of ectomycorrhizal fungi in forest soils: role in carbon cycling. *Plant Soil* 366, 1–27
73. Powell, J.R. *et al.* (2009) Phylogenetic trait conservatism and the evolution of functional trade-offs in arbuscular mycorrhizal fungi. *Proc. R. Soc. Lond. B Biol. Sci.* 276, 4237–4245
74. Rillig, M.C. *et al.* (2015) Plant root and mycorrhizal fungal traits for understanding soil aggregation. *New Phytol.* 205, 1385–1388
75. Lehmann, A. *et al.* (2020) Fungal traits important for soil aggregation. *Front. Microbiol.* 10, 2904
76. Treseder, K.K. and Lennon, J.T. (2015) Fungal traits that drive ecosystem dynamics on land. *Microbiol. Mol. Biol. Rev.* 79, 243–262
77. Bielčik, M. *et al.* (2019) The role of active movement in fungal ecology and community assembly. *Mov. Ecol.* 7, 36
78. Aleklett, K. and Boddy, L. (2021) Fungal behaviour: a new frontier in behavioural ecology. *Trends Ecol. Evol.* 36, 787–796
79. van der Heijden, M.G.A. and Scheublin, T.R. (2007) Functional traits in mycorrhizal ecology: their use for predicting the impact of arbuscular mycorrhizal fungal communities on plant growth and ecosystem functioning. *New Phytol.* 174, 244–250
80. Phillips, R.P. *et al.* (2013) The mycorrhizal-associated nutrient economy: a new framework for predicting carbon–nutrient couplings in temperate forests. *New Phytol.* 199, 41–51
81. Behm, J.E. and Kiers, E.T. (2014) A phenotypic plasticity framework for assessing intraspecific variation in arbuscular mycorrhizal fungal traits. *J. Ecol.* 102, 315–327
82. Kattge, J. *et al.* (2011) TRY – a global database of plant traits. *Glob. Chang. Biol.* 17, 2905–2935
83. Maherali, H. and Klironomos, J.N. (2007) Influence of phylogeny on fungal community assembly and ecosystem functioning. *Science* 316, 1746–1748
84. Wright, I.J. *et al.* (2004) The worldwide leaf economics spectrum. *Nature* 428, 821–827