



Universiteit
Leiden
The Netherlands

Stability and transition of depression subtypes in late life

Veltman, E.; Kok, A.; Lamers, F.; Stek, M.; Mast, R. van der; Rhebergen, D.

Citation


Veltman, E., Kok, A., Lamers, F., Stek, M., Mast, R. van der, & Rhebergen, D. (2020). Stability and transition of depression subtypes in late life. *Journal Of Affective Disorders*, 265, 445-452. doi:10.1016/j.jad.2020.01.049

Version: Publisher's Version

License: [Creative Commons CC BY 4.0 license](https://creativecommons.org/licenses/by/4.0/)

Downloaded from: <https://hdl.handle.net/1887/3184565>

Note: To cite this publication please use the final published version (if applicable).



Research paper

Stability and transition of depression subtypes in late life

Eveline Veltman (MD)^{a,*}, Almar Kok (PhD) (Co-author)^b, Femke Lamers (PhD) (Co-author)^{b,c},
Max Stek (MD PhD) (Co-author)^{b,c}, Roos van der Mast (MD PhD) (Co-author)^{a,d},
Didi Rhebergen (MD PhD)^{b,c}

^a Department of Psychiatry, Leiden University Medical Center, the Netherlands

^b Amsterdam University Medical Center, Vrije Universiteit, Psychiatry, Amsterdam Public Health research institute, the Netherlands

^c GGZ inGeest Specialized Mental Health Care, Amsterdam, the Netherlands

^d Department of Psychiatry, CAPRI-University of Antwerp, Belgium

ARTICLE INFO

Keywords:

Latent transition analysis
Late-life depression
Depression subtypes
Stability of depressive subtypes
Atypical depression
Melancholic depression

ABSTRACT

Background: The heterogeneity of late-life depression hampers diagnosis and treatment. Data-driven methods have identified several subtypes of depression in older persons, but the longitudinal stability of these subtypes remains unknown.

Methods: In total 111 older persons with a major depressive disorder both at baseline and 2-year follow-up from the Netherlands Study of Depression in Older persons (NESDO) were included. Latent class analysis was performed to identify subtypes of depression at baseline and at 2-year follow-up, and latent transition analysis was used to examine the stability of these subtypes over time. Transition rates between subtypes and characteristics of groups were examined.

Results: Two subtypes were identified in both baseline (T0) and follow-up data (T1), including a ‘melancholic’ subtype (prevalence 80.2% (T0) and 62.2% (T1)), and an ‘atypical’ subtype (prevalence 19.8% (T0) and 37.8% (T1)). The melancholic subtype was characterized by decreased appetite and weight and had a stability of 0.86. The atypical subtype was characterized by increased appetite and weight and had a stability of 0.93, although the discriminating power of different symptoms had decreased at T1. Mean age and education differed significantly between stable and transitioning subgroups, other characteristics did not differ between subgroups.

Limitations: Limited sample size might have hampered the analyses.

Conclusions: Subtypes of late-life depression are relatively stable, but symptoms of depression (like weight loss) seem to blur with symptoms of (patho)physiological aging. This underlines the clinical relevance of depression subtyping, but also the importance of further research into subtypes and the influence of aging.

1. Introduction

Late-life depression is a common disorder, with a prevalence ranging from 1–16% (Djernes, 2006). Especially late-life depression is associated with a poorer course (Schaakxs et al., 2018), and comes with a high disease burden for both patients (Gallo et al., 2007) and their caregivers (Scazufta et al., 2002). Better insight into the etiology and treatment options is therefore of great importance. However, to date research is being hindered by the heterogeneity of depression (Lux and Kendler, 2010; Goldberg, 2011), since the diagnosis of major depressive disorder (MDD) does not reflect the wide array of possible depressive symptom combinations. A diagnosis of MDD is made if 5 out of 9 DSM criteria are met, and since several of these symptoms are opposites (e.g.

weight gain or loss), two patients diagnosed with MDD could have zero symptoms overlapping. In addition, previous, longitudinal studies demonstrated low longitudinal stability of depression categories described in the Diagnostic and Statistical Manual of Mental Disorders (DSM) edition 5 (American Psychiatric Association, 2013) such as major depressive disorder, dysthymia, and subthreshold depression (Angst et al., 1997, 2000). Acknowledging this heterogeneity, the DSM 5 contains several specifiers such as depression with atypical or melancholic features, but these have not proved sufficient to predict prognosis and treatment response (Parker et al., 2010; Lojko and Rybakowski, 2017).

A different approach to examine the heterogeneity of depression is through data-driven methods (Lubke and Muthén, 2005), such as latent class analysis (LCA). These techniques cluster patients based on their

* Corresponding author.

E-mail addresses: evelineveltman@gmail.com (E. Veltman), a.kok1@amsterdamumc.nl (A. Kok), f.lamers@ggzingeest.nl (F. Lamers), m.stek@ggzingeest.nl (M. Stek), r.c.van_der_mast@lumc.nl (R. van der Mast), d.rhebergen@ggzingeest.nl (D. Rhebergen).

<https://doi.org/10.1016/j.jad.2020.01.049>

Received 4 July 2019; Received in revised form 8 December 2019; Accepted 12 January 2020

Available online 14 January 2020

0165-0327/ © 2020 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

congregate of different depressive symptoms, without a pre-conceived hypothesis. Studies in younger adults using LCA identified an ‘atypical’ subtype characterized by increased sleep and appetite, and a ‘typical’ or ‘melancholic’ subtype characterized by decreased sleep and appetite, and the presence of psychomotor symptoms (Kendler et al., 1996; Sullivan et al., 1998, 2002; Lamers et al., 2010; Rodgers et al., 2013; Li et al., 2014). In addition, these empirically-derived subtypes and their most distinguishing symptoms as appetite and weight have been linked to distinct biological and genetic correlates, and different neural activity on functional MRI (Lamers et al., 2013; Milaneschi et al., 2017; Simmons et al., 2018). In older adults, earlier data-driven studies have found subtypes mainly based on severity, probably due to inclusion of persons without a formal diagnosis of depression (Hybels et al., 2009, 2013; Lee et al., 2012; Mezuk and Kendler, 2012). Furthermore, not all of these studies distinguished between increase and decrease of sleep, appetite, weight and psychomotor symptoms, while these distinctions appeared to be crucial in identifying subtypes of depression in both younger (Lamers et al., 2010; Alexandrino-Silva et al., 2013; Rodgers et al., 2013). We have performed an LCA on older depressed adults taking this symptom distinction into account, and have found an atypical, melancholic, and moderately severe subtype (Veltman et al., 2017), similar to the aforementioned studies in younger adults. To further examine the validity and clinical usefulness of these subtypes, insight is needed into the stability and the potential transition rate of subjects across subtypes over time.

Latent transition analysis (LTA) is a data-driven method useful for examining the longitudinal stability and transition of subtypes of depression. In younger depressed adults, studies performing LTA have found a relatively high stability of subtypes (48–90%), with the atypical subtype being the most stable (71–79%) (Lamers et al., 2012; Rodgers et al., 2014). In older depressed adults, so far one earlier LTA has been performed (Ni et al., 2017). Their sample consisted of outpatients aged 60 to 96 years, and they demonstrated moderate stability of subtypes during 2-year follow-up (46–61%), with the highest stability in the subtype characterized by a moderate severity and a lack of positive affect. However, they did not differentiate between increasing and decreasing sleep, appetite, weight, and psychomotor symptoms, and as shown before, these symptoms are important in differentiating between subtypes of depression.

The Netherlands Study on Depression in Older persons (NESDO) provides an excellent opportunity to examine the stability of late-life depression subtypes, with recognition of the importance of differentiation between increase and decrease in sleep, appetite and psychomotor symptoms. This paper is a follow-up of an earlier cross-sectional study that identified three subtypes of depression using latent class analysis, including an atypical, a melancholic, and a moderate subtype (Veltman et al., 2017). The atypical subtype was characterized by a high severity, and an increase in sleep, appetite and weight. The melancholic subtype was characterized by a high severity, but a decrease in sleep, appetite and weight. The moderate subtype was characterized by an overall lower prevalence of depressive symptoms. To examine the 2-year longitudinal stability of these subtypes, in this study we perform a latent transition analysis on persons derived from the NESDO-cohort with a major depressive disorder (MDD) both at baseline and at 2-year follow-up. This study population differs from the population in Veltman et al. (2017) in chronicity, since a diagnosis of MDD on both time points was needed for inclusion. Nevertheless, considering the three identified subtypes in our previous study (Veltman et al., 2017), as well as the findings from LTA studies on younger adults identifying similar subtypes (Lamers et al., 2012; Rodgers et al., 2014), we hypothesize to find three subtypes among older depressed people using LTA, including a melancholic, atypical, and moderate subtype, with considerable stability during 2-year follow-up.

2. Methods

2.1. Study population

Data were derived from the baseline and 2-year follow-up measurements of the Netherlands Study of Depression in Older persons (NESDO), a longitudinal multi-site naturalistic cohort study, examining course and consequences of depression in older persons. The NESDO cohort ($n = 510$) consists of persons aged 60–93 years, including 378 persons with a depressive disorder in the previous 6 months at baseline, and 132 non-depressed controls. Depressed older persons were recruited from both mental health care institutes and general practitioners in five regions in the Netherlands, in order to include persons with current late-life depression in various developmental and severity stages. Depression included a 6-month diagnosis of Major Depressive Disorder (MDD) (95%) and/or 6 month dysthymic disorder (26.5%), or minor depression (two to four depressive symptoms lasting at least two weeks, 5.0%) according to DSM-IV-R criteria. Non-depressed older persons were recruited from general practices and were included when no lifetime diagnosis of depression was present. Exclusion criteria were (suspected) dementia, and insufficient command of the Dutch language. The study design of NESDO is described in detail elsewhere (Comijs et al., 2011).

For the present study, persons with a 6-month diagnosis of MDD at baseline (T0), and a 6-month diagnosis of MDD at 2-year follow-up (T1), were included. Persons with dysthymia or minor depression only were excluded. We chose to include only persons with a present diagnosis of MDD on both T0 and T1 because we specifically wanted to examine the longitudinal stability of depression subtypes.

2.2. Symptoms of depression

Ten depressive symptoms were used as indicator variables in the LCA analyses in order to identify depression subtypes. Nine depression key symptoms of the DSM-IV were based on the Dutch version of the Composite International Diagnostic Interview (CIDI) lifetime version 2.1, (World Health Organization 1998; Andrews and Peters 1998) which were used to diagnose depressive and anxiety disorders according to DSM-IV criteria and conducted by specially trained clinical research staff. Changes in appetite and weight were used as two separate variables. All items were coded as ‘not present’ or ‘present’, except for the items regarding changes in appetite, weight, sleep and psychomotor disturbance. Here we created four categories, so for example for weight the categories were: absence of weight changes, weight loss, weight gain, and both gain and loss.

2.3. Covariates

Several socio-demographic, clinical, psychosocial, and physical health indicators were used to characterize subtypes, equal to the covariates used in our baseline paper (Veltman et al., 2017) and mostly overlapping with similar studies in younger adults (Lamers et al., 2012) to enable comparison.

Socio-demographic variables, including age, gender and years of education, were collected during the baseline interview. *Clinical characteristics* such as age of onset of the depressive disorder and comorbid anxiety disorders were assessed by the CIDI. Severity of depressive symptoms was assessed with the 30-item Inventory of Depressive Symptomatology (IDS) (Rush et al., 1996). Further, we assessed several *psychosocial variables*. Overall daily functioning was measured with the World Health Organization Disability Assessment Schedule II (WHODAS) (Chwastiak and Von Korff, 2003; Buist-Bouwman et al., 2008). Apathy was measured using the Apathy Scale (Starkstein et al., 1992), and cognitive functioning by the Mini Mental State Examination (MMSE) (Folstein et al., 1975). *Physical health indicators* were included. Respondents were asked whether they were currently smoking (yes/

no). In addition, their alcohol intake was calculated as number of drinks per week. Pain was measured using a count of pain locations (range 0–7) listed in the Chronic Graded Pain scale (Von Korff et al., 1992). The presence of metabolic syndrome (yes/no) was measured. Cut-off values for metabolic syndrome were based on the adjusted Adult Treatment Panel (ATP III) criteria (Expert panel, 2001), including waist circumference ≥ 88 cm, triglycerides ≥ 150 mg/dl, HDL-cholesterol < 50 mg/dl, systolic blood pressure ≥ 130 mmHg or diastolic blood pressure ≥ 85 mmHg or use of anti-hypertensive drugs, fasting blood glucose ≥ 100 mg/dl or use of a hypoglycemic drug. Persons scoring positive on ≥ 3 criteria were considered to have a metabolic syndrome. Diabetes was defined as fasting plasma glucose level ≥ 7.0 mmol/l. Objective and standardized assessments of height and weight were performed. Body mass index (BMI) was calculated as kilograms divided by meter squared and categorized as underweight-normal ($\text{BMI} < 25$), overweight ($25 \leq \text{BMI} < 30$) and obese ($\text{BMI} \geq 30$). The number of chronic diseases, presence of cardiovascular disease (assessed by self-report supported by appropriate medication use (see Vogelzangs et al. 2010 for a detailed description), the presence of diabetes (based on fasting plasma glucose level ≥ 7.0 mmol/l or use of anti-diabetic medication [ATC-code A10] WHO 2007) was determined. Muscle weakness as an indicator of frailty was measured with a hand-held dynamometer. This was considered present if persons scored below the cut off score after their performance was stratified by gender and BMI, according to Fried et al. (2001).

2.4. Outline of analyses

First, latent class analyses, using 10 CIDI-items as indicator variables, were performed at T0 and T1 separately. The conditional probability of occurrence of each depressive symptom, given class membership, was examined using Cramer's V. After identification of classes, latent transition analysis was performed, to examine the stability of the identified classes. Characteristics of all classes were examined. In univariate analyses, repeated measurement T-tests were used for normally distributed variables, McNemar tests for dichotomous variables, and Wilcoxon paired rank tests for non-parametric variables. In multivariable analyses, the two stable subtypes were compared, and analyses were corrected for variables that significantly changed the bivariate associations, when added separately (change odd's ratio $> 10\%$).

2.5. Latent class analyses

To determine and identify subtypes of depressive disorders, we performed Latent Class Analysis (LCA). Free from any a priori assumptions about the number and characterization of subtypes, data-driven techniques such as LCA cluster persons into unobserved ('latent') classes, based on their values on several observed variables. Analysis starts with one class, suggesting one class fitting for all persons. Then, iteratively, one class is added to the model, and the best fitting number of classes is determined on the basis of several statistical indicators and interpretability. The LCAs were conducted using Mplus version 5 (Muthén and Muthén, 2007).

To determine which model best fitted the data, we examined the sample size adjusted BIC (ssaBIC), entropy (i.e., classification accuracy), the bootstrapped Lo-Mendell-Rubin likelihood ratio test (LMR) (number of bootstraps used = 1000), the bootstrapped likelihood ratio test (BLRT) and the proportion of respondents in each class. Lower ssaBIC values indicate better model fit (Nylund et al., 2007a). Entropy, as a measure of classification accuracy, is presented for models with more than one subtype. Values for entropy can range between 0 and 1 with values closer to 1 indicating greater accuracy. The Lo-Mendell-Rubin test (LMR) provides a p-value, which indicates whether the $k-1$ subtype model is rejected in favor of the k subtype model. Similar to the LMR test, a p-value below 0.05 for the bootstrapped likelihood ratio test (BLRT) indicates that the present model fits the data better than the

model with one class less; a p-value exceeding 0.05 indicates that the model with one class less should be preferred (Nylund et al., 2007a). There is no consensus yet as to which criterion best identifies the best fitting number of classes, but the BLRT seems to be superior to other model fit indices (Nylund et al., 2007a). Finally, we only accepted models in which all classes contained at least five percent of the sample.

2.6. Latent transition analysis

Latent transition analysis is a longitudinal mixture model that explores change in latent classes of individuals over time. Latent class analysis on two time points is used as a measurement model in LTA, from which individual-level change among these classes over time is defined by the transition analyses (Nylund, 2007b). Depending on the outcome of the two separate latent class analyses performed for T0 and T1, the latent transition analysis was modelled according to Nylund (2007b). In case of an equal number of subtypes for both latent classes analyses, these subtypes were compared in order to see to what extent they were similar. The latent transition model, calculating the movement of subjects between subtypes across time points, was modelled with respect to the comparability of subtypes found on T0 and T1. First, the degree of similarity between subtypes was compared through loglikelihood tests. For this purpose, we examined the extent of (in) variance of the subtypes in different models: a model where all conditional probabilities of variables were constrained to be equal across time, a model where all conditional probabilities were estimated freely across time, and models in which all conditional probabilities started out as variant, and were gradually coded as invariant (testing for partial invariance). The higher the number of conditional probabilities that were invariant across time, the higher the similarity of identified classes over time.

3. Results

The total sample consisted of 111 depressed older people, of whom 67% were females, with a mean age of 70.9 (SD ± 7.7) years at baseline and a mean age of depression onset of 44.2 (SD ± 20.1) years.

In Table 1, demographics of the total sample at baseline (T0) and 2-year follow-up (T1) are shown.

Severity significantly decreased over time ($p < 0.01$), as did body mass index ($p = 0.01$), while muscle weakness increased ($p = 0.04$).

Table 2 shows the model fit indices for 1 to 4 classes on both T0 and T1. The best fitting model for the analysis on both baseline and 2-year follow-up was the 2-class model, with a significant Lo-Mendell-Rubin test (LMR) (T0 $p = 0.01$; T1 $p < 0.01$), a significant bootstrapped likelihood ratio test (BLRT) (T0 $p < 0.01$; T1 $p = 0.01$), lowest BIC and highest entropy. There is no consensus yet as to which criterion best identifies the best fitting number of classes, but the BLRT seems to be superior to other model fit indices (Nylund et al., 2007a). From the 3-class model onwards, the BLRT was no longer significant, indicating that adding more than two classes to the model did not improve model fit. Classification accuracy was high, with average posterior probabilities for subgroups ranging between 0.825 – 1.000 ($n = 111$, data not shown), further indicating a reasonable classification quality.

Table 3 shows the estimated symptom profile probabilities for the two-class model on both T0 and T1. Variables are dichotomous (present or not present), except for the variables appetite, weight, sleep, and psychomotor symptoms, which are nominally divided into four categories (increase, decrease, both, or no change). In both subtypes weight and appetite, and to a lesser extent sleep and psychomotor symptoms, are the main significantly differing symptoms between subtypes. The first subtype in both T0 and T1 shows an atypical symptom pattern, with the highest occurrence of both weight gain and increased appetite, and is therefore labeled 'atypical'. The second subtype in both T0 and T1 is characterized by a more typical or melancholic symptom pattern with symptoms such as weight loss and decreased appetite, and is

Table 1
Characteristics of total sample at baseline (T0) and 2-year follow-up (T1) ($N = 111$).

	T0	T1	Overall p-value (df)
Socio-demographics			
Sex, female, %	66.7	*	*
Age, mean (SD), y	70.7(7.6)	*	*
Education, mean (SD), y	10.5(3.8)	*	*
Clinical characteristics			
Age of onset, mean (SD) y	43.8(19.7)	*	*
Severity (IDS), mean (SD)	34.9(13.1)	32.1(12.3)	0.01 (110) ¹
Anxiety dx last year, %	46.8	37.8	0.13 ²
Psychosocial			
Functioning sx, mean (SD)	35.7(16.4)	37.4(16.1)	0.26(105) ¹
Apathy score, mean (SD)	18.3(5.1)	19.7(6.2)	0.09(108) ¹
MMSE score, median (IQR)	28.0(2.0)	28.0(2.0)	0.67 ³
Physical health			
Current smoking, %	27.0	#	#
# alcohol/ week, median (IQR)	0.3(4.9)	#	#
Chronic pain grade, mean (SD)	2.3(1.3)	2.2(1.1)	0.53(100) ¹
Body mass index, median (IQR)	26.6(5.4)	26.3(6.2)	0.01 ³
Cardiovascular disease, yes, %	26.1	27.0	1.00 ²
Diabetes, yes, %	17.1	19.1	0.32 ¹
# chronic diseases, mean(SD)	2.9(1.9)	2.2(1.1)	0.80(109) ¹
Metabolic syndrome, yes, % Muscle weakness, %	21.6	27.9	0.04 ²

Tests used: 1 = paired-samples *t*-test; 2 = McNemar; 3 = Wilcoxon paired rank test.

#, * = demographic like gender and age of onset had no change over time, and from smoking and alcohol we did not have data available on T1.

therefore labeled ‘melancholic’. Although especially the ‘atypical’ label resembles a specific DSM diagnosis, we explicitly do not intend to refer to this DSM diagnosis. Rather, the label is chosen to facilitate comparisons with our previous data-driven study on older depressed adults (Veltman et al., 2017), and with earlier latent class and latent transition analyses in younger depressed adults (Lamers et al., 2012; Rodgers et al., 2014).

We then performed a latent transition analysis to examine the stability of these subtypes between T0 and T1. The best fitting model included appetite as invariant symptom, and all other symptoms as variant. Both subtypes showed high stability (atypical subtype $N = 15$, stability = 0.93; melancholic subtype $N = 62$, stability = 0.87). The estimated transition probability of melancholic to atypical ($N = 27$) was 0.14, and of atypical to melancholic ($N = 7$) was 0.07.

Table 4 shows the characteristics of the stable subjects and transitioning subjects. The group transitioning from atypical to melancholic has the lowest mean age, followed by the stable atypical subtype. The stable melancholic subtype has the lowest years of education.

Table 2
Model fit indices for different classes at T0 and T1.

Model fit indices for different classes, T0											
Classes	Maximum likelihood	BIC	ssaBIC	Entropy	Lo-Mendell-Rubin	BLRT	Proportion per class				
					2LL	p	p	1	2	3	4
1	−680.618	1446.007	1389.123	–	–	–	–	1.000			
2	−649.427	1473.107	1356.180	0.888	62.381	0.0100	0.00	0.198	0.802		
3	−633.110	1529.953	1352.982	0.814	32.635	0.0939	0.1750	0.243	0.568	0.189	
4	−620.831	1594.877	1357.862	0.803	30.650	1.000	0.0900	0.270	0.189	0.432	0.108
Model fit indices for different classes, T1											
Classes	Maximum likelihood	BIC	ssaBIC	Entropy	Lo-Mendell-Rubin	BLRT	Proportion per class				
					2LL	p	p	1	2	3	4
1	−683.285	1451.342	1394.458	–	–	–	–	1.000			
2	−660.989	1496.230	1379.303	0.764	44.593	0.0082	0.0130	0.378	0.622		
3	−648.750	1559.323	1382.352	0.780	26.338	0.9855	0.7130	0.423	0.252	0.324	
4	−643.177	1625.251	1388.237	0.864	26.549	0.8232	0.4140	0.360	0.243	0.297	0.099

Prevalence of diabetes was lowest in the stable atypical group and highest in the group transitioning from atypical to melancholic. Further group differences in demographic characteristics did not reach significance. Multivariable comparison of the stable atypical and the stable melancholic subgroup showed no significant differences (data not shown). Due to low numbers in the transitioning groups (melancholic to atypical, $N = 27$, and atypical to melancholic, $N = 7$), power limited multivariable analyses across all four groups. Next, multivariable comparison of the stable subjects, including both atypical and melancholic subtypes ($N = 77$), versus the transitioning subjects, including persons transitioning from melancholic towards atypical and vice versa ($N = 32$), did not yield significant findings.

4. Discussion

In this study, data-driven subtypes of old age depression and their stability over two-year follow-up were examined. Both at baseline and 2-year follow-up, two depression subtypes were identified. The first, melancholic subtype was mainly characterized by a decrease in appetite and weight. The second, atypical subtype was predominantly characterized by an increase in appetite and weight. Both subtypes had high temporal stability, with a slightly higher stability for the atypical subtype. Whereas transition rates between subtypes were low, subjects were more likely to shift from the melancholic subtype to the atypical subtype than vice versa. The stable atypical subtype had a lower mean age and a lower age of onset than the stable melancholic subtype, but this difference disappeared in multivariable analyses. Diabetes was lowest in the stable atypical group and highest in the group transitioning from atypical to melancholic. Other demographic variables were not significantly correlated to subtypes, neither in univariate, nor in multivariable regression analyses. No variables predicting transitioning, as compared to stable class membership, were discovered in multivariable analyses.

In contrast to our hypothesis and our previously published LCA (Veltman et al., 2017), based on data derived from the same cohort ($N = 359$), we now identified two subtypes (atypical and melancholic) instead of three. The previously identified moderate subtype is now absent, which may be caused by a decline in study population and thereby a decrease in power. Since we specifically wanted to examine the longitudinal stability of depression subtypes, persons were only eligible for inclusion in the current study if they fulfilled DSM-criteria for MDD both at baseline and 2-year follow-up. Hence, a study population of chronic or recurrent MDD was selected, whereas in the former study (Veltman et al., 2017) persons with non-chronic MDD were also included. Since chronic depression has been associated with higher depression severity (Reisinger Walker and Druss, 2015), this selection procedure may have resulted in a study sample with higher mean depression severity, and hence, the absence of a moderately severe

Table 3Estimated symptom profile probabilities of endorsing depressive symptoms from LCA T0 and T1 ($n = 111$).

	T0 Total sample	T0 Class 1 Atypical	T0 Class 2 Melancholic	Cramer's V	P -value T0(df)	T1 Total sample	T1 Class 1 Atypical	T1 Class 2 Melancholic	Cramer's V	P-value	T1(df)
Prevalence		22 (19.8%)	89 (80.2%)				42 (37.8%)	69 (62.2%)			
DSM-IV criterion symptoms	Symptom probabilities					Symptom probabilities					
Depressed mood	99.1	1.00	0.99	0.05	0.62(1)	95.5	1.00	0.93	<0.01	0.07(1)	
Loss of interest	93.7	0.96	0.93	0.04	0.70(1)	91.0	1.00	0.86	<0.01	0.01(1)	
Weight				0.78	<0.01(3)				0.43	<0.01(3)	
No weight change	55.9	0.36	0.61			77.5	0.81	0.75			
Weight loss	31.5	0.00	0.39			16.2	0.05	0.23			
Weight gain	10.8	0.55	0.00			5.4	0.14	0.00			
Both gain and loss	1.8	0.09	0.00			0.9	0.00	0.01			
Appetite				0.80	<0.01(3)				0.47	<0.01(3)	
No change in appetite	26.1	0.09	0.30			37.8	0.57	0.26			
Decreased appetite	55.0	0.14	0.65			43.2	0.00	0.70			
Increased appetite	15.3	0.73	0.01			16.2	0.43	0.00			
Both increase and decrease	3.6	0.05	0.03			2.7	0.00	0.04			
Sleep				0.28	0.03(3)				0.13	0.02(3)	
No change in sleep	9.0	0.05	0.10			13.5	0.24	0.07			
Less sleep	65.8	0.59	0.67			64.0	0.60	0.67			
More sleep	8.1	0.00	0.10			11.7	0.14	0.10			
Both less and more sleep	17.1	0.36	0.12			10.8	0.08	0.16			
Psychomotor				0.25	0.07(3)				0.06	<0.01(3)	
No psychomotor change	22.5	0.14	0.25			27.9	0.29	0.28			
Psychomotor retardation	36.0	0.32	0.37			32.4	0.36	0.31			
Psychomotor agitation	20.7	0.14	0.23			22.5	0.05	0.34			
Both agitation and retardation	20.7	0.41	0.16			16.2	0.31	0.07			
Fatigue/energy loss	94.6	1.00	0.93	0.12	0.21(1)	98.2	1.00	0.97	0.07	0.27(1)	
Guilt/worthlessness	83.8	1.00	0.80	0.22	0.02(1)	70.3	0.88	0.59	0.18	<0.01(1)	
Concentration/ indecisiveness	95.5	1.00	0.94	0.11	0.26(1)	96.4	1.00	0.94	0.10	0.11(1)	
Suicidal ideation	74.8	1.00	0.69	0.29	<0.01(1)	70.3	0.74	0.68	0.03	0.52(1)	

The left half of the table shows the symptom probabilities of the entire cohort and of the two different subtypes at T0 (ranging from 0–1), and the p-value for the difference in probability. The right half of the table shows the symptom probabilities of the two subtypes at T1, again with their corresponding p-values. All symptoms are dichotomous, except for the symptoms weight, appetite, sleep and psychomotor symptoms. These are nominally divided into four categories: no change, increase, decrease, and both increase and decrease.

Table 4

Characteristics of stable and transitioning subtypes at baseline.

	Stable atypical (N = 15)	Stable melancholic (N = 62)	Atypical à Melancholic (N = 7)	Melancholic à Atypical (N = 27)	Overall p-value (df)
Socio-demographics					
Sex, female,%	80.0	64.5	71.4	63.0	0.67(3) ¹
Age, mean (SD), y	67.6(6.7)	71.8(8.0)	65.7(5.0)	71.2(6.8)	0.04(3) ²
Education, mean (SD), y	11.1(3.7)	9.6(3.7)	11.4(4.7)	11.8(3.6)	0.05(3) ²
Clinical characteristics					
Age onset, mean (SD) y	35.1(21.4)	45.8(19.2)	44.3(15.1)	44.1(20.6)	0.28(3) ²
Severity sx, mean (SD)	37.2(11.9)	33.3(14.4)	37.4(7.1)	36.8(11.8)	0.43(3) ²
Anxiety dx last year,%	53.3	48.4	57.1	37.0	0.65(3) ¹
Psychosocial					
Functioning, mean (SD)	37.1(15.4)	36.6(16.1)	41.1(12.6)	31.1(18.3)	0.65(3) ²
Apathy score, mean (SD)	19.6(4.1)	17.5(4.6)	16.7(5.8)	19.8(6.1)	0.17(3) ²
MMSE score, mean (SD)	28.3(1.6)	27.5(1.9)	27.6(1.3)	27.9(2.2)	0.41(3) ²
Physical health					
Current smoking,%	33.3	29.0	28.6%	18.5%	0.70(3) ¹
# alcohol/ week, median (IQR)	1.0(3.7)	0.4(3.7)	0.0(1.0)	0.2(8.2)	0.73(3) ³
Chronic pain grade, mean (SD)	2.5(1.2)	2.2(1.3)	2.3(1.3)	2.2(1.4)	0.95(3) ²
Presence of metabolic syndrome,%	73.3	50.0	57.1	37.0	0.16(3) ¹
Body mass index, median (IQR)	27.8(9.2)	26.6(6.2)	27.4(1.7)	25.5(5.3)	0.22(3) ³
Cardiovascular disease, yes,%	33.3	22.6	57.1	22.2	0.21(3) ¹
Diabetes,%	0.0	22.6	42.9	3.7	0.01(3) ¹
# chronic diseases, mean (SD)	3.1(1.8)	2.8(1.9)	3.6(2.9)	2.7(1.7)	0.76(3) ²
Muscle weakness,%	20.0	27.9	0.0	14.8	0.25(3) ¹

Estimated transition probabilities: atypical to atypical = 0.928; atypical to melancholic = 0.072; melancholic to atypical = 0.135; melancholic to melancholic = 0.865

Statistical tests used: (1) chi-square; (2) Anova; (3) Kruskal-Wallis.

subtype. Severity scores of the different cohorts indeed reveal a higher severity in the current study towards, with an IDS sum score of 34.9 ($SD \pm 13.1$) in the present study and of 30.5 ($SD \pm 13.0$) in our previous LCA study. In post-hoc analyses, we further explored the characteristics of the subjects that were included in our earlier LCA (Veltman et al., 2017), but not in our current LTA. These were subjects with either missing data on T0, or with MDD on T0 but not on T1 ($N = 248$). First, we found that participants in the current study were equally selected with respect to original subtypes ($p = 0.12$); persons from all three LCA-derived classes were equally selected for inclusion in the current study. This suggests that the absence of a moderate severe subtype might be (partly) caused by a relatively small sample ($N = 111$), with less power for complex modeling. Furthermore, included subjects had a higher MDD severity on T0 ($p = 0.02$) compared to excluded subjects. Of the 248 excluded subjects, 64.1% ($N = 159$) did not reach the criteria for MDD, and 35.9% ($N = 89$) had dropped out. Reasons for drop-out were dropout due to unknown reasons between T0 and T1 (8.0%, $N = 20$), refusal to participate at T1 (5.2%, $N = 13$), not eligible due to physical or mental reasons (16.5%, $N = 41$), no contact could be made (2.0%, $N = 5$), or passing away (4.0%, $N = 10$).

Next, the symptom profiles of the identified atypical and melancholic subgroup in both T0 and T1, and their high temporal stabilities correspond to earlier studies on younger depressed adults (Lamers et al., 2012; Rodgers et al., 2014). In line with studies on younger adults, the atypical subtype was characterized by increased appetite and weight, and the melancholic subtype by decreased appetite and weight. In the current study however, the difference in appetite and weight symptoms between both subtypes was smaller at T1 as compared to T0. The reduced item discriminating power and very small number of invariable symptoms (1 out of 10) in the LTA model might imply that the nature of the atypical and melancholic subtypes is not completely stable over time. This might be because old age and aging are often associated with diminishing physical health (e.g. weight loss and muscle weakness) and somatic comorbidity, especially in depressed older persons (Holvast et al., 2017). These (patho)physiological processes might influence the presentation of late-life depression, and decrease the distinguishing power of symptoms such as appetite and weight over time. This hypothesis is supported by our finding that BMI and muscle strength significantly decreased during 2-year follow-up (see Table 1). This indicates the dwindling health of our subjects and may complicate the identification of depressive subtypes in an older population where depressive symptom presentation might be clouded by (patho)physiological aging.

Nevertheless, taking these possible (patho)physiological alterations in aging into account by freeing conditional probabilities across time, the stability of the identified subtypes was high, and several characteristics were in line with earlier studies on subtypes of depression. Age was significantly lower in the stable atypical subgroup and in the subgroup transitioning from atypical to melancholic, compared to the stable melancholic subgroup and the subgroup transitioning from melancholic to atypical. In earlier studies among younger adults identifying a similar atypical subtype, age and age of depression onset were also lowest in the atypical subtype (Lamers et al., 2010; Veltman et al., 2017). In contrast with previous studies, however, in our study prevalence of metabolic syndrome and cardiovascular disease did not differ between the atypical and melancholic subtype, and prevalence of diabetes was lowest in our stable atypical subtype. It was previously demonstrated that chronic depression, in particular in older age, is associated with a wide array of somatic illnesses (Hegeman et al., 2017; Holvast et al., 2017). Likewise, in non-depressed older persons, prevalence of somatic illnesses is higher in older age. Hence, competing pathways, other than pathways associated with depression subtype, may result in onset of diabetes and cardiovascular disease in older persons. To conclude, we hypothesize that in our chronically depressed cohort of older persons, the lack of significant difference in metabolic

and cardiovascular disturbances may rather reflect pathophysiological disturbances associated with either chronicity of depression or aging processes that are not linearly correlated with depression subtypes in late-life. In addition to this, the limited sample size and 2-year follow-up period might have been insufficient to find the true diversity in depression subtypes and their underlying pathophysiological pathways. Future research on the longitudinal stability of late-life depression subtypes with larger sample sizes and longer follow-up is warranted.

4.1. Strengths and limitations

The results of this study should be interpreted in the light of various strengths and limitations. A strength of this study is that it is the first latent transition analysis on older depressed persons taking both increase and decrease of weight, appetite, sleep, and psychomotor changes into account. This differentiation of symptoms has been especially distinguishing in earlier found data-driven subtypes, and it enables comparison with several studies on both younger and older adults. Furthermore, a wide selection of characteristics was examined. The limited sample size is an important limitation of the current study. Due to both selection of patients with chronic or recurrent MDD and to dropout due to morbidity and mortality in this older population, only a small number of the original population in our earlier study was included. This may decrease the reliability of our LTA, and may hamper our univariate and multivariable analyses. Another limitation is the possible violation of the local independence assumption (correlation between variables in a class is accounted for by the latent variable). This could occur if separate variables are included that measure different expressions of the same underlying symptom. Post-hoc we have examined tetrachoric correlations on bivariate symptoms, presented in **Supplement 1**. Several symptoms showed moderately high correlations, which might mean a violation of the local independence. However, there is still no consensus on how strict this interpretation should be. Our current approach enables comparison with the body of literature on latent categorical modeling of depressive subtypes. Our findings are largely in line with previous findings, which strengthens their plausibility and clinical relevance.

5. Conclusion

To conclude, this is the first study that examined the temporal stability of data-driven subtypes of late-life depression taking into account both increase and decrease in sleep, appetite, weight, and psychomotor symptoms. An atypical and a melancholic subtype were found, both with a high stability over 2-year follow-up. Appetite and weight were main distinguishing symptoms, but their discriminating ability decreased over time. Since the overall cohort significantly declined in weight and physical health during 2-year follow-up, the process of (patho) physiological aging might blur the differentiation of depression subtypes in older age. These findings stress the importance of taking the pathophysiological tangle of aging into account when searching for a better approach to diagnosis, treatment and prevention of late-life depression.

6. Author agreement

6.1. Contributors

We hereby state that all authors have seen and approved the final version of the manuscript 'Stability and transition of depression subtypes in late life' being submitted, and that there are no other persons who satisfied the criteria for authorship but are not listed. We further confirm that the order of authors listed in the manuscript has been approved by all of us. We warrant that the article is the authors' original work, hasn't received prior publication, and isn't under consideration for publication elsewhere.

Funding

The infrastructure for NESDO is funded through the Fonds NutsOhra, Stichting tot Steun VCVGZ, NARSAD The Brain and Behaviour Research Fund, and the participating universities and mental health care organizations (VU University Medical Center, Leiden University Medical Center, University Medical Center Groningen, Radboud University Nijmegen Medical Center, and GGZ inGeest, GGZ Nijmegen, GGZ Rivierduinen, Lentis, and Parnassia).

Declaration of Competing Interest

We wish to confirm that there are no known conflicts of interest associated with this publication and there has been no further significant financial support for this work that could have influenced its outcome.

We confirm that the manuscript has been read and approved by all named authors and that there are no other persons who satisfied the criteria for authorship but are not listed. We further confirm that the order of authors listed in the manuscript has been approved by all of us.

We confirm that we have given due consideration to the protection of intellectual property associated with this work and that there are no impediments to publication, including the timing of publication, with respect to intellectual property. In so doing we confirm that we have followed the regulations of our institutions concerning intellectual property.

Acknowledgments

We confirm that we have given due consideration to the protection of intellectual property associated with this work and that there are no impediments to publication, including the timing of publication, with respect to intellectual property. In so doing we confirm that we have followed the regulations of our institutions concerning intellectual property.

We understand that the Corresponding Author is the sole contact for the Editorial process (including Editorial Manager and direct communications with the office). She is responsible for communicating with the other authors about progress, submissions of revisions and final approval of proofs. We confirm that we have provided a current, correct email address which is accessible by the Corresponding Author and which has been configured to accept email from evelineveltman@gmail.com.

Supplementary materials

Supplementary material associated with this article can be found, in the online version, at [doi:10.1016/j.jad.2020.01.049](https://doi.org/10.1016/j.jad.2020.01.049).

References

- Alexandrino-Silva, C., Wang, Y., Viana, M.C., Bulhões, R.S., Martins, S.S., Andrade, L.H., 2013. Gender differences in symptomatic profiles of depression: results from the São Paulo megacity mental health survey. *J. Affect. Disord.* 147, 355–364. <https://doi.org/10.1016/j.jad.2012.11.041>.
- Andrews, G., Peters, L., 1998. The psychometric properties of the composite international diagnostic interview. *Soc. Psychiatry Psychiatr. Epidemiol.* 33, 80–88. <https://doi.org/10.1007/s001270050026>.
- Angst, J., Merikangas, K.R., 1997. The depressive spectrum: diagnostic classification and course. *J. Affect. Disord.* 45, 31–40. [https://doi.org/10.1016/S0165-0327\(97\)00057-8](https://doi.org/10.1016/S0165-0327(97)00057-8).
- Angst, J., Sellaro, R., Merikangas, K.R., 2000. Depressive spectrum diagnoses. *Compr. Psych.* 41, 39–47. [https://doi.org/10.1016/S0010-440X\(00\)80007-3](https://doi.org/10.1016/S0010-440X(00)80007-3).
- Buist-Bouwman, M.A., Ormel, J., De Graaf, R., Vilagut, G., Alonso, J., Van Sonderen, E., Vollebergh, W.A., ESEMeD/MHEDEA 2000 Investigators, 2008. Psychometric properties of the world health organization disability assessment schedule used in the European study of the epidemiology of mental disorders. *Int. J. Methods Psychiatr. Res.* 17, 185–197. <https://doi.org/10.1002/mpr.261>.
- Chwastiak, L.A., Von Korff, M., 2003. Disability in depression and back pain: evaluation of the world health organization disability assessment schedule (WHO das II) in a primary care setting. *J. Clin. Epidemiol.* 56, 507–514. [https://doi.org/10.1016/S0895-4356\(03\)00051-9](https://doi.org/10.1016/S0895-4356(03)00051-9).
- Comijs, H.C., Van Marwijk, H.W., Van der Mast, R.C., Naarding, P., Oude Voshaar, R.C., Beekman, A.T.F., Boshuizen, M., Dekker, J., Kok, R., De Waal, M.W.M., Penninx, B.W., Stek, M.L., Smit, J.H., 2011. The Netherlands study of depression in older persons (NESDO): a prospective cohort study. *BMC Res. Notes* 5, 524. <https://doi.org/10.1186/1756-0500-4-524>.
- Diagnostic and Statistical Manual of Mental Disorders, 5 ed. DC: American Psychiatric Association, Washington 2013. Print.
- Djernes, K., 2006. Prevalence and predictors of depression in populations of elderly: a review. *Acta Psychiatr. Scand.* 113, 372–387. <https://doi.org/10.1111/j.1600-0447.2006.00770.x>.
- Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in adults, 2001. Executive summary of the third report of the national cholesterol education program (NCEP) expert panel on detection, evaluation, and treatment of high blood cholesterol in adults (Adults treatment panel III). *JAMA* 285, 2486–2497. <https://doi.org/10.1001/jama.285.19.2486>.
- Folstein, M.F., Folstein, S.E., McHugh, P.R., 1975. ‘Mini-mental state.’ a practical method for grading the cognitive state of patients for the clinician. *J. Psychiatr. Res.* 129–138. [https://doi.org/10.1016/0022-3956\(75\)90026-6](https://doi.org/10.1016/0022-3956(75)90026-6).
- Fried, L.P., Tangen, C.M., Walston, J., Newman, A.B., Hirsch, C., Gottdiener, J., Seeman, T., Tracy, R., Kop, W.J., Burke, G., McBurnie, M.A., Cardiovascular Health Study Collaborative Research Group, 2001. Frailty in older adults: evidence for a phenotype. *J. Gerontol. A Biol. Sci. Med. Sci.* 256, 146–156. <https://doi.org/10.1093/gerona/56.3.m146>.
- Goldberg, D., 2011. The heterogeneity of “major depression”. *World Psych.* 10, 226–228. <https://doi.org/10.1002/j.2051-5545.2011.tb00061.x>.
- Hegeman, J.M., Van Fenema, E.M., Comijs, H.C., Kok, R.M., Van der Mast, R.C., De Waal, M.W.M., 2017. Effect of chronic somatic diseases on the course of late-life depression. *Int. J. Geriatr. Psych.* 32, 779–787. <https://doi.org/10.1002/gps.4523>.
- Holvast, F., Van Hattem, B.A., Sinnige, J., Schellevis, F., Taxis, K., Burger, H., Verhaak, P.F.M., 2017. Late-life depression and the association with multimorbidity and polypharmacy: a cross-sectional study. *Fam. Pract.* 34, 539–545. <https://doi.org/10.1093/fampra/cmx018>.
- Hybels, C.F., Lazer, D.G., Pieper, C.F., Landerman, L.R., Steffens, D.C., 2009. Profiles of depressive symptoms in older adults diagnosed with major depression: latent cluster analysis. *Am. J. Geriatr. Psych.* 7, 387–396. <https://doi.org/10.1097/JGP.0b013e31819431ff>.
- Hybels, C.F., Landerman, L.R., Blazer, D.G., 2013. Latent subtypes of depression in a community sample of older adults: can depression clusters predict future depression trajectories? *J. Psychiatric Res* 47, 1288–1297. <https://doi.org/10.1016/j.jpsychires.2013.05.033>.
- Kendler, K.S., Eaves, L.J., Walters, E.E., Neale, M.C., Heath, A.C., Kessler, R.C., 1996. The identification and validation of distinct depressive symptoms in a population-based sample of female twins. *Arch. Gen. Psych.* 53, 391–399. <https://doi.org/10.1001/archpsyc.1996.01830050025004>.
- Korff von, M., Ormel, J., Keefe, F.J., Dworkin, S.F., 1992. Grading the severity of chronic pain. *Pain* 50, 133–149. [https://doi.org/10.1016/0304-3959\(92\)90154-4](https://doi.org/10.1016/0304-3959(92)90154-4).
- Lamers, F., de Jonge, P., Nolen, W.A., Smit, J.H., Zitman, F.G., Beekman, A.T., Penninx, B.W., 2010. Identifying depressive subtypes in a large cohort study: results from the Netherlands study of depression and anxiety (NESDA). *J. Clin. Psychiatry* 71 <https://doi.org/10.4088/JCP.09m05398blu>. 1582–1289.
- Lamers, F., Rhebergen, D., Merikangas, K.R., de Jonge, P., Beekman, A.T., Penninx, B.W., 2012. Stability and transitions of depressive subtypes over a 2-year follow-up. *Psychol. Med.* 42, 2083–2093. <https://doi.org/10.1017/S0033291712000141>.
- Lamers, F., Vogelzangs, N., Merikangas, K.R., de Jonge, P., Beekman, A.T., Penninx, B.W., 2013. Evidence for a differential role of HPA-axis function, inflammation and metabolic syndrome in melancholic versus atypical depression. *Mol. Psych.* 18, 692–699. <https://doi.org/10.1038/mp.2012.144>.
- Lee, C., Leoutsakos, J., Lyketsos, C.G., Steffens, D.C., Breitner, J.C.S., Norton, M.C., for the Cache County Investigators, 2012. Latent class-derived subgroups of depressive symptoms in a community sample of older adults: the cache county study. *Int. J. Geriatr. Psychiatry* 27, 1061–1069. <https://doi.org/10.1002/gps.2824>.
- Lojko, D., Rybakowski, J.K., 2017. Atypical depression: current perspectives. *Neuropsychiatr. Dis. Treat.* 13, 2447–2456. <https://doi.org/10.2147/NDT.S147317>.
- Lubke, G.H., Muthén, B.O., 2005. Investigating population heterogeneity with factor mixture models. *Psychol. Methods* 10, 21–39. <https://doi.org/10.1037/1082-989X.10.1.21>.
- Lux, V., Kendler, K.S., 2010. Deconstructing major depression: a validation study of the DSM-IV symptomatic criteria. *Psychol. Med.* 40, 1679–1690. <https://doi.org/10.1017/S0033291709992157>.
- Mezuk, B., Kendler, K.S., 2012. Examining variation in depressive symptoms over the life course: a latent class analysis. *Psychol. Med.* 42, 2037–2046. <https://doi.org/10.1017/S003329171200027X>.
- Milaneschi, Y., Lamers, F., Peyrot, W.J., Baune, B.T., Breen, G., Dehghan, A., Forstner, A.J., Grabe, H.J., Homuth, G., Kan, C., Lewis, C., Mullins, N., Nauck, M., Pistis, G., Preisig, M., Rivera, M., Rietschel, M., Streit, F., Strohmaier, J., Teumer, A., Van der Auwera, S., Wray, N.R., Boomsma, D.I., Penninx, B.W.J.H., 2017. Genetic association of major depression with atypical features and obesity-related immunometabolic dysregulations. *JAMA Psychiatry* 74, 1214–1225. <https://doi.org/10.1001/jamapsychiatry.2017.3016>.
- Muthén, L.K., Muthén, B.O., 2007. *Mplus User's Guide. Fifth Edition.* Muthén & Muthén: Los Angeles, California.
- Ni, Y., Tein, J.Y., Zhang, M., Yang, Y., Wu, G., 2017. Changes in depression among older adults in China: a latent transition analysis. *J. Affect. Disord.* 209, 3–9. <https://doi.org/10.1016/j.jad.2016.11.004>.

- Nylund, K.L., Asparouhov, T., Muthén, B.O., 2007a. Deciding on the number of subtypes in latent class analysis and growth mixture modeling: a monte carlo simulation study. *Struct. Equ. Modeling* 14, 535–569. <https://doi.org/10.1080/10705510701575396>.
- Nylund, K., 2007b. Latent Transition Analysis: Modeling extensions and an Application to Peer victimization. Doctoral Dissertation. University of California, Los Angeles. <http://www.statmodel.com/download/nylunddis.pdf> (accessed 11 October 2019).
- Parker, G., Fink, M., Shorter, E., Taylor, M.A., Akiskal, H., Berrios, G., Bolwig, T., Brown, W.A., Carroll, B., Healy, D., Klein, D.F., Koukopoulos, A., Michels, R., Paris, J., Rubin, R.T., Spitzer, R., Swartz, C., 2010. Issues for DSM-5: whither melancholia? the case for its classification as a distinct mood disorder. *Am. J. Psychiatry* 167, 745–747. <https://doi.org/10.1176/appi.ajp.2010.09101525>.
- Reisinger Walker, E., Druss, B.G., 2015. Rate and predictors of persistent major depressive disorder in a nationally representative sample. *Community Ment. Health J.* 51, 701–707. <https://doi.org/10.1007/s10597-014-9793-9>.
- Rodgers, S., Grosse Holtforth, M., Müller, M., Hengartner, M.P., Rössler, W., Ajdacic-Gross, V., 2013. Symptom-based subtypes of depression and their psychosocial correlates: a person-centered approach focusing on the influence of sex. *J. Affect. Disord* 156, 92–103. <https://doi.org/10.1016/j.jad.2013.11.021>.
- Rodgers, S., Ajdacic-Gross, V., Müller, M., Hengartner, M.P., Grosse Holtforth, M., Angst, J., Rössler, W., 2014. The role of sex on stability and change of depression symptom subtypes over 20 years: a latent transition analysis. *Eu. Arch. Psychiatry Clin. Neurosci.* 264, 577–588. <https://doi.org/10.1007/s00406-013-0475-3>.
- Rush, A.J., Gullion, C.M., Basco, M.R., Jarrett, R.B., Trivedi, M.H., 1996. The inventory of depressive symptomatology (IDS): psychometric properties. *Psychol. Med.* 26, 477–486. <https://doi.org/10.1017/s0033291700035558>.
- Scazufo, M., Menezes, P.R., Almeida, O.P., 2002. Caregiver burden in an elderly population with depression in São Paulo. Brazil. *Soc. Psychiatry Psychiatr. Epidemiol* 37, 416–422. <https://doi.org/10.1007/s00127-002-0571-6>.
- Schaakxs, R., Comijs, H.C., Lamers, F., Kok, R.M., Beekman, A.T.F., Penninx, B.W.J.H., 2018. Associations between age and the course of major depressive disorder: a 2-year longitudinal cohort study. *Lancet Psych.* 7, 581–590. [https://doi.org/10.1016/S2215-0366\(18\)30166-4](https://doi.org/10.1016/S2215-0366(18)30166-4).
- Simmons, W.K., Burrows, K., Avery, J.A., Kerr, K.L., Taylor, A., Bodurka, J., Potter, W., Teague, T.K., Drevets, W.C., 2018. Appetite changes reveal depression subgroups with distinct endocrine, metabolic, and immune states. *Mol. Psych.* <https://doi.org/10.1038/s41380-018-0093-6>. (epub ahead of print). 10.1038/s41380-018-0093-6.
- Starkstein, S.E., Mayberg, H.S., Preziosi, T.J., Andrezejewski, P., Leiguarda, R., Robinson, R.G., 1992. Reliability, validity, and clinical correlates of apathy in Parkinson's disease. *J. Neuropsych. Clin. Neurosci.* 4, 134–139. <https://doi.org/10.1176/jnp.4.2.134>.
- Sullivan, P.F., Kessler, R.C., Kendler, K.S., 1998. Latent class analysis of lifetime depressive symptoms in the national comorbidity survey. *Am. J. Psych.* 155, 1398–1406. <https://doi.org/10.1176/ajp.155.10.1398>.
- Sullivan, P.F., Prescott, C.A., Kendler, K.S., 2002. The subtypes of major depression in a twin registry. *J. Affect. Disord.* 68, 273–284. [https://doi.org/10.1016/s0165-0327\(00\)00364-5](https://doi.org/10.1016/s0165-0327(00)00364-5).
- Veltman, E.M., Lamers, F., Comijs, H.C., De Waal, M.W.M., Stek, M.L., Van der Mast, R.C., Rhebergen, D., 2017. Depressive subtypes in an elderly cohort identified using latent class analysis. *J. Affect. Disord.* 218, 123–130. <https://doi.org/10.1016/j.jad.2017.04.059>.
- Vogelzangs, N., Kritchevsky, S.B., Beekman, A.T., Brenes, G.A., Newman, A.B., Satterfield, S., Yaffe, K., Harris, T.B., Penninx, B.W., Study, Health ABC, 2010. Obesity and onset of significant depressive symptoms: results from a prospective community-based cohort study of older men and women. *J. Clin. Psych.* 71, 391–399. <https://doi.org/10.4088/JCP.08m04743blu>.
- World Health Organization, 1998. Composite International Diagnostic Interview (CIDI), Version 2.1. World Health Organization: Geneva.
- World Health Organization, 2007. WHO Model list of Essential medicines, Version 15. World Health Organization: Geneva.