



Universiteit
Leiden
The Netherlands

Influence of age on upper arm cuff blood pressure measurement

Picone, D.S.; Schultz, M.G.; Otahal, P.; Black, J.A.; Bos, W.J.; Chen, C.H.; ... ; Sharman, J.E.

Citation

Picone, D. S., Schultz, M. G., Otahal, P., Black, J. A., Bos, W. J., Chen, C. H., ... Sharman, J. E. (2020). Influence of age on upper arm cuff blood pressure measurement. *Hypertension*, 75(3), 844-850. doi:10.1161/HYPERTENSIONAHA.119.13973

Version: Publisher's Version

License: [Creative Commons CC BY-NC-ND 4.0 license](https://creativecommons.org/licenses/by-nc-nd/4.0/)

Downloaded from: <https://hdl.handle.net/1887/3181303>

Note: To cite this publication please use the final published version (if applicable).

Influence of Age on Upper Arm Cuff Blood Pressure Measurement

Dean S. Picone, Martin G. Schultz, Petr Otahal, J. Andrew Black, Willem J. Bos, Chen-Huan Chen, Hao-Min Cheng, Antoine Cremer, Nathan Dwyer, Ricardo Fonseca, Alun D. Hughes, Hack-Lyoung Kim, Peter S. Lacy, Esben Laugesen, Nobuyuki Ohte, Stefano Omboni, Christian Ott, Telmo Pereira, Giacomo Pucci, Philip Roberts-Thomson, Niklas B. Rossen, Roland E. Schmieder, Daisuke Sueta, Kenji Takazawa, Jiguang Wang, Thomas Weber, Berend E. Westerhof, Bryan Williams, Hirotugu Yamada, Eiichiro Yamamoto, James E. Sharman; for the Invasive Blood Pressure Consortium

Abstract—Blood pressure (BP) is a leading global risk factor. Increasing age is related to changes in cardiovascular physiology that could influence cuff BP measurement, but this has never been examined systematically and was the aim of this study. Cuff BP was compared with invasive aortic BP across decades of age (from 40 to 89 years) using individual-level data from 31 studies (1674 patients undergoing coronary angiography) and 22 different cuff BP devices (19 oscillometric, 1 automated auscultation, 2 mercury sphygmomanometry) from the Invasive Blood Pressure Consortium. Subjects were aged 64±11 years, and 32% female. Cuff systolic BP overestimated invasive aortic systolic BP in those aged 40 to 49 years, but with each older decade of age, there was a progressive shift toward increasing underestimation of aortic systolic BP ($P<0.0001$). Conversely, cuff diastolic BP overestimated invasive aortic diastolic BP, and this progressively increased with increasing age ($P<0.0001$). Thus, there was a progressive increase in cuff pulse pressure underestimation of invasive aortic PP with increasing decades of age ($P<0.0001$). These age-related trends were observed across all categories of BP control. We conclude that cuff BP as an estimate of aortic BP was substantially influenced by increasing age, thus potentially exposing older people to greater chance for misdiagnosis of the true risk related to BP. (*Hypertension*. 2020;75:844-850. DOI: 10.1161/HYPERTENSIONAHA.119.13973.) • [Online Data Supplement](#)

Key Words: aging ■ cardiovascular disease ■ coronary angiography ■ risk factor ■ sphygmomanometers

Cardiovascular disease (CVD) is the leading cause of death worldwide,¹ and the most important CVD risk factor is high blood pressure (BP). Clinical management of BP is based on measurements from upper arm cuff BP devices, either using auscultation or automated oscillometry. Correct identification and lowering of high BP will reduce the risk of

CVD and all-cause mortality.² However, our recent work revealed that cuff BP does not reflect intraarterial BP either at the central aorta or brachial artery, especially in the systolic BP (SBP) range of 120 to 159 mm Hg.³ The reasons for these differences are not fully understood but are related to

Received September 11, 2019; first decision September 24, 2019; revision accepted December 19, 2019.

From the Menzies Institute for Medical Research, University of Tasmania, Hobart, Australia (D.S.P., M.G.S., P.O., J.A.B., N.D., R.F., P.R.-T., J.E.S.); Royal Hobart Hospital, Hobart, Tasmania (J.A.B., P.R.-T., N.D.); St Antonius Hospital, Department of Internal Medicine, Nieuwegein, the Netherlands (W.J.B.); Department of Internal Medicine, Leiden University Medical Center, the Netherlands (W.J.B.); Department of Medicine, National Yang-Ming University School of Medicine, Taipei, Taiwan (C.-H.C., H.-M.C.); Department of Medical Education, Taipei Veterans General Hospital, Taiwan (C.-H.C., H.-M.C.); Department of Cardiology/Hypertension, University Hospital of Bordeaux, France (A.C.); Institute of Cardiovascular Sciences, University College London, United Kingdom (A.D.H.); Division of Cardiology, Seoul National University Boramae Hospital, South Korea (H.-L.K.); Institute of Cardiovascular Sciences University College London (UCL) and National Institute for Health Research (NIHR) UCL/UCL Hospitals Biomedical Research Centre, United Kingdom (P.S.L., B.W.); Department of Endocrinology and Internal Medicine, Aarhus University Hospital, Denmark (E.L.); Department of Cardio-Renal Medicine and Hypertension, Nagoya City University Graduate School of Medical Sciences, Japan (N.O.); Clinical Research Unit, Italian Institute of Telemedicine, Varese (S.O.); Scientific Research Department of Cardiology, Science and Technology Park for Biomedicine, Sechenov First Moscow State Medical University, Russian Federation (S.O.); Department of Nephrology and Hypertension, University Hospital Erlangen, Friedrich-Alexander University Erlangen-Nürnberg, Germany (C.O., R.E.S.); Polytechnic Institute of Coimbra, ESTES, Department of Physiology, General Humberto Delgado Street 102, Lousã, Portugal (T.P.); Unit of Internal Medicine at Terni University Hospital, Department of Medicine, University of Perugia, Italy (G.P.); Department of Cardiovascular Medicine, Graduate School of Medical Sciences, Kumamoto University, Japan (D.S., E.Y.); Center for Health Surveillance and Preventive Medicine, Tokyo Medical University Hospital, Japan (K.T.); Centre for Epidemiological Studies and Clinical Trials, Shanghai Key Laboratory of Hypertension, The Shanghai Institute of Hypertension, Department of Hypertension, Ruijin Hospital, Shanghai Jiao Tong University School of Medicine, China (J.W.); Cardiology Department, Klinikum Wels-Grieskirchen, Wels, Austria (T.W.); Department of Pulmonary Diseases, VU University Medical Center, Amsterdam, the Netherlands (B.E.W.); and Department of Community Medicine for Cardiology, Tokushima Graduate School of Biomedical Sciences, Japan (H.Y.).

The online-only Data Supplement is available with this article at <https://www.ahajournals.org/doi/suppl/10.1161/HYPERTENSIONAHA.119.13973>.

Correspondence to James Sharman, Menzies Institute for Medical Research, University of Tasmania, Private Bag 23, Hobart, 7000 Australia. Email James.Sharman@utas.edu.au

© 2020 American Heart Association, Inc.

Hypertension is available at <https://www.ahajournals.org/journal/hyp>

DOI: 10.1161/HYPERTENSIONAHA.119.13973

pathophysiological changes to the cardiovascular system that occur with increasing age or disease.⁴⁻⁷

Upper arm cuff BP measurement, whether by auscultation or oscillometry, relies on analysis of signals (Korotkoff sounds or cuff pressure oscillations) arising from the brachial artery.⁸ Major changes in cardiovascular hemodynamics could alter these signals to an extent that may affect cuff BP measurement. This could be highly relevant to increasing age because it is typically accompanied by a multitude of cardiovascular changes, such as lower BP amplification,⁶ impaired ventricular-vascular coupling,⁹ increased arterial stiffness,¹⁰ altered arterial geometry,¹¹ and abnormal blood flow dynamics.^{12,13} The influence of age on cuff BP compared with an intraarterial (invasive) BP reference standard has never been determined systematically, which was the aim of this study. We hypothesized that increasing age would be associated with greater differences between cuff BP and invasive aortic BP.

Methods

The data that support the findings of this study are available from the corresponding author upon reasonable request.

Overview

The analysis was conducted from data within an international consortium designed to better understand the level of cuff BP as an estimate of invasive BP (Invasive Blood Pressure Consortium).³ This comprised an individual participant meta-analysis among 59 separate studies (total 3073 subjects) where cuff-measured BP was recorded simultaneously (or sequentially in the immediate time period) with invasive BP, thus providing a means to examine the difference between cuff BP compared with invasive BP. Studies that measured cuff BP in the angiography waiting room prior or post procedure were excluded. This current analysis focuses on the comparison of upper arm cuff-measured BP versus invasive aortic BP as the reference measurement, which was measured using fluid-filled catheter-manometers or solid-state micromanometer catheters (complete data available for 1674 subjects). Rationale for comparison with aortic BP was because cuff BP aims to measure the pressure load at the arterial sites of interaction with the central organs.^{14,15} Importantly, it is this central aortic BP that more strongly relates to organ damage, stroke and heart attack, compared with peripheral BP (ie, brachial artery) which may substantially differ from central aortic BP, especially for SBP and pulse pressure (PP).^{3,16} Although arm cuff BP is not always expected to be equivalent to aortic BP, cuff SBP systematically underestimates the true (invasive) brachial SBP, and thus may approximate aortic SBP.^{3,17} On the contrary, cuff diastolic BP (DBP) is relatively constant through the arterial system.³ For complete assessment, a secondary (sensitivity) analysis was also undertaken to compare cuff BP with invasive brachial BP (complete data available for 520 subjects). The University of Tasmania Health and Medical Human Research Ethics Committee approved the study (reference: H0015048).

Data Handling

Several steps were taken to ensure the quality of the consortium data. First, only studies that measured cuff and invasive BPs simultaneously or within an immediate period (just before or after the invasive BP recording) were included. Full details on the sequence of cuff and invasive BP measurements are in the Expanded Methods in the [online-only Data Supplement](#). Furthermore, any study that recorded data during nonbasal hemodynamic shifts or aimed to assess the effect of different cuff sizes on the relationship between cuff BP and invasive BP was excluded. A quality score was calculated by judging the key study methods that could have affected data accuracy ([online-only Data Supplement](#)). Detailed systematic reviews for each topic were updated on 28 February 2018 using the same protocols previously published.³

Information on the separate studies included in the present analyses are detailed in Tables S1 and S2 in the [online-only Data Supplement](#). The analysis was conducted on subjects who were aged 40 to 89 years (stratified according to decades of age) because subjects aged younger than 40 or 90 years and older accounted for <4% of the data. Cuff BP was assessed by comparison to invasive BP, defined as cuff BP minus invasive BP. Therefore, a positive value for the difference indicated that cuff BP overestimated invasive BP, whereas a negative value indicated that cuff BP underestimated invasive BP. Cuff PP and invasive PP were calculated as SBP minus DBP. Mean arterial pressure (MAP) was calculated using a 40% form factor (DBP+0.4×PP)¹⁸ because the true MAP, which is defined as the average of all points on the BP waveform, was not available.

Statistical Analyses

Sample clinical characteristics were reported as mean±SD (or median and interquartile range for skewed data) or number (%) of total cases. All differences between cuff BP and invasive BP were reported as mean and 95% CI. Linear mixed models were used to analyze the influence of age on the difference between cuff BP and invasive BP. Multivariable mixed models were used to account for variables known or suspected to affect the relationship between age and the difference between cuff BP and invasive BP. These variables included sex (as a potential confounder) and separately invasive MAP, body mass index, and heart rate (as potential mediators). A random effect term coding each individual study was included in the mixed models to account for the within-study clustering of subjects. From the unadjusted and adjusted models, average marginal effects for the difference between cuff and invasive BP were calculated for each decade of age. The same analysis was performed with stratification by the category of cuff BP according to the 2017 American Heart Association/American College of Cardiology arterial hypertension guidelines (normal: SBP <120 and DBP <80 mmHg; elevated: 120–129 and <80 mmHg; stage 1 hypertension: 130–139 or 80–89 mmHg and stage 2 hypertension: ≥140 or ≥90 mmHg).¹⁹ Sensitivity analyses included determining the influence of age on the difference between cuff BP and invasive BP when (1) age was assessed as a continuous variable; (2) a fluid-filled or micromanometer tip catheter was used for invasive BP measurements; (3) studies were analyzed according to a maximum versus nonmaximum rated study quality score; (4) cuff versus invasive brachial BP was analyzed; (5) cuff BP and invasive SBP and PP amplification (calculated as invasive brachial SBP and PP minus the respective invasive aortic values) were available on the same subjects; and (6) the order of BP measurement was accounted for. *P*<0.05 was considered statistically significant. Data were analyzed using R version 3.5.1 (R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>). The linear mixed models and average marginal effects were generated using the lme4 and ggeffects packages respectively.

Results

Subjects

One thousand six hundred seventy-four subjects from 31 studies met the inclusion criteria (Figure S1). Twenty-two different cuff BP devices (19 oscillometric, 1 automated auscultation, 2 mercury sphygmomanometry) were used. In 16 of the studies, the average of multiple cuff BP readings was used in the analysis. Most subjects were patients who were undergoing coronary angiography procedures. The clinical characteristics in the Table are typical of this patient population; subjects were middle-to-older aged, predominately male, overweight according to body mass index, and 67% had evidence of stenosis in at least one coronary artery. In total, 65% of subjects had cuff BP in the hypertensive range, according to the American Heart Association/American College of Cardiology guidelines.

Table. Sample Characteristics and BP Values Across Decades of Age

Variable	40 to 49 y (n=168)	50 to 59 y (n=403)	60 to 69 y (n=550)	70 to 79 y (n=447)	80 to 89 y (n=106)
Sample Characteristics					
Age, y	45.1±2.8	54.8±2.7	64.0 [62.0 to 67.0]	74.0 [72.0 to 77.0]	82 [81 to 84]
Female sex, %*	45 (27)	121 (30)	178 (33)	147 (33)	40 (38)
Height, cm†	170.7±9.6	167.1±9.1	165.4±10.3	162.9±10.2	158.9±10.1
Weight, kg‡	84.4±20.9	78.3±18.6	73.7±17.6	68.1±14.5	61.1±13.0
Body mass index, kg/m²§	28.9±5.9	27.9±5.8	26.8±5.5	25.4±4.4	24.1±4.1
Heart rate, bpm	70±12	69±12	68±12	67±12	66±12
Hypertension defined by cuff BP ≥130/≥80, %	91 (54)	241 (60)	361 (66)	316 (71)	82 (77)
Hypertension defined by invasive aortic BP ≥130/≥80, %	76 (45)	206 (51)	337 (61)	305 (68)	83 (78)
BP					
Cuff systolic BP	128±18	131±21	136±23	139±22	145±23
Cuff diastolic BP	80±11	79±12	77±13	76±12	76±14
Cuff pulse pressure	48±13	52±15	59±18	63±20	69±20
Invasive aortic systolic BP	125±20	130±25	138±25	143±26	150±26
Invasive aortic diastolic BP	75±11	73±12	70±12	67±12	65±13
Invasive aortic pulse pressure	50±15	58±19	68±21	76±22	85±22

Data are mean±SD or median [interquartile range]. All BP units are mm Hg. BP indicates blood pressure.

*n=1647.

†n=1520.

‡n=1532.

§n=1518.

||n=1453.

Influence of Age on Upper Arm Cuff BP Measurement

Systolic BP

Cuff SBP slightly overestimated invasive aortic SBP in those aged 40 to 49 years, but with each increase in decade of age, there was a progressive shift toward increasing underestimation of invasive aortic SBP (Figure 1 and Table S3, $P<0.0001$). In those aged 70 to 79 and 80 to 89 years, cuff SBP clearly underestimated invasive aortic SBP. After adjusting for sex and separately for invasive MAP, heart rate, and body mass

index, the difference between cuff SBP and invasive aortic SBP across the decades of age were slightly attenuated but remained significant (Tables S4 and S5, $P<0.0001$). Sex, invasive MAP, heart rate, and body mass index (Tables S4 and S5) were also related to the difference between cuff SBP and invasive aortic SBP. After stratification of subjects based on cuff BP guideline categories, each increase in decade of age remained related to a progressive increase in the magnitude of underestimation of invasive aortic SBP (Figure 2A, $P<0.05$ for each cuff BP category).




Figure 1. Cuff blood pressure (BP) compared with invasive aortic systolic BP (circles), diastolic BP (triangles), and pulse pressure (squares) measurements across age decades. Data are mean difference and 95% CI (error bars). Data above the solid horizontal zero line indicates cuff BP is higher than invasive aortic BP and vice versa below the zero line. The trends for the age-related differences in cuff BP compared with invasive aortic BP were statistically significant for systolic, diastolic and pulse pressure, $P<0.0001$ all.




Figure 2. Cuff blood pressure (BP) compared with invasive aortic BP measurements across decades of age and stratified according to the category of BP control (according to the 2017 American Heart Association/American College of Cardiology arterial hypertension guidelines).¹⁹ Panel A, systolic BP (SBP), panel B, diastolic BP (DBP) and panel C, pulse pressure (PP). Data are mean difference and 95% confidence interval (error bars). Within each BP category, there were significant trends for the influence of age on cuff BP compared with invasive aortic BP ($P < 0.05$), albeit borderline for DBP in stage 1 hypertension ($P = 0.086$). Circles, normal BP; triangles, elevated BP; squares, stage 1 hypertension; crosses; stage 2 hypertension.

Diastolic BP

Cuff DBP overestimated invasive aortic DBP in all decades of age. Similar to SBP, with each increase in decade of age, there was a progressive increase in the overestimation of aortic DBP (Figure 1 and Table S3, $P < 0.0001$). The trend was unchanged after adjustment for the variables described above (Tables S4 and S5, $P < 0.0001$). Sex and invasive MAP (Tables S4 and S5) were also related to the difference between cuff DBP and invasive aortic DBP in the adjusted models. After additional stratification of subjects based on the cuff BP category, each increase in decade of age remained related to a progressive increase in the magnitude of overestimation of invasive aortic DBP ($P < 0.01$; Figure 2B), albeit stage 1 hypertension was a borderline trend ($P = 0.086$).

Pulse Pressure

For each increase in decade of age, there was a progressive increase in the magnitude of underestimation of invasive aortic PP by cuff measurements (Figure 1 and Table S3, $P < 0.0001$). The trend was unchanged after adjustment for sex or separately for invasive MAP, heart rate, and body mass index, and

all these variables were related to the difference between cuff PP and invasive aortic PP (Tables S4 and S5, $P < 0.0001$). After additional stratification of subjects based on the cuff BP category, each increase in decade of age remained related to a progressive increase in the magnitude of underestimation of invasive aortic PP (Figure 2C, $P < 0.001$ for each BP category).

The unadjusted differences between cuff SBP, DBP, and PP and invasive aortic SBP, DBP, and PP were not different between the entire study dataset ($N = 1674$) and the subpopulations used in the adjusted models for sex ($n = 1547$) and invasive MAP, heart rate, and body mass index ($n = 1382$). Our previous work details the difference between cuff and invasive BP for each individual study.³

Sensitivity Analyses

Age as a Continuous Variable

Increasing age was related to a progressive increase in the magnitude of underestimation of invasive aortic SBP and PP and overestimation of aortic DBP ($P < 0.0001$ all).

Fluid-Filled or Micromanometer Tip Catheter

The influence of age on cuff BP compared to invasive aortic BP was similar irrespective of the type of catheter used (trend $P < 0.0001$ all; Figure S2).

Study Quality Score

The influence of age on cuff BP compared with invasive aortic BP was similar for the maximum and nonmaximum rated studies (Figure S3).

Cuff BP Compared With Invasive Brachial BP

Five hundred twenty subjects (62 ± 11 years of age, 31% female; detailed characteristics in Table S6) met the inclusion criteria for this sensitivity analysis (Figure S4). Similar trends to aortic BP were observed for the influence of age on cuff SBP compared with invasive brachial (Figure S5 and Table S7), but this was less pronounced than for invasive aortic BP. After adjustment for sex and separately for invasive MAP, heart rate, and body mass index, the influence of age on cuff SBP compared with invasive brachial was not significant (Tables S8 and S9). The influence of age on cuff DBP and PP compared with invasive brachial values was similar to the invasive aortic analysis (Figure S5 and Tables S8 and S9). Stratification based on the cuff BP guideline category (Figure S6) was limited due to low subject numbers in several age and BP category combinations (eg, $n=3$ for 80–89 years of age and normal, elevated or stage one hypertension BP categories). The magnitude of difference between cuff and invasive brachial BP was similar when data were stratified according to the type of catheter (Figure S7), and separately, the type of cuff device used (cuff oscillometry or mercury auscultation; Figure S8).

Cuff BP and BP Amplification

In 372 subjects, the influence of age on cuff SBP compared with both invasive aortic and brachial SBP tracks for the 40 to 49 and 50 to 59 age decades, but then SBP amplification does not continue to drop with increasing age (Figure S9). Cuff PP compared with both invasive aortic and brachial PP does not track with PP amplification. The influence of age on the difference between cuff and invasive aortic SBP, DBP, or PP remained after adjustment for BP amplification (Table S10).

Order of BP Measurement

The influence of age was not different whether cuff and invasive aortic BP were measured simultaneously or if cuff BP was measured just before invasive BP or if invasive BP was measured just before cuff BP (Figure S10).

Discussion

Correct measurement of BP is paramount for the appropriate diagnosis and management of CVD risk.²⁰ The key findings from this study were that there were greater differences between cuff BP and invasive aortic BP with increasing age and that this occurred irrespective of the level of BP according to guideline categories. These findings could have implications for the assessment of true risk related to BP across the lifespan and may also be relevant to understanding the true distribution of aortic BP in population-level studies, as well as clinical hypertension thresholds and validation protocols used to test new BP devices.

Pioneering studies in arterial physiology from the 1950s provided critical insights on BP measurement, showing that

brachial SBP and PP were higher than corresponding aortic SBP and PP (termed BP amplification).^{21,22} Inconsequential differences in DBP between the aorta and brachial artery were also reported. Theoretically, if cuff BP was a close proxy of invasive brachial BP, then typically it should be higher than the corresponding invasive aortic SBP and PP and should agree closely with aortic DBP. However, cuff BP measurements systematically underestimate invasive brachial SBP (-5.7 mmHg) and PP (-12.0 mmHg) and systematically overestimate invasive brachial DBP ($+5.5$ mmHg).³ The systematic underestimation of brachial SBP means that cuff and invasive aortic SBP are not different on average, but there is wide variability with substantial overestimation or underestimation of aortic SBP, depending on the individual and the cuff BP device.³ Invasive aortic DBP is systematically overestimated by cuff DBP. The present study extends on these findings and has found that age has a systematic influence on the cuff SBP, DBP, and PP compared with invasive aortic values.

This study was not designed to determine the mechanisms which explain why chronological age influences the capacity of cuff BP to estimate invasive aortic BP. An excellent analog of vascular aging can be derived from measures of arterial stiffness via methods such as pulse wave velocity, and several studies have examined the relationship between stiffness and cuff BP compared with invasive BP.^{4,5,23–25} In a study of elderly people, higher arterial stiffness was associated with overestimation of invasive aortic BP by auscultatory cuff measurements.^{5,24} However, the opposite was observed among patients with chronic kidney disease,⁴ using oscillometric cuff BP methods. It is unclear whether differences in measurement methods or participant characteristics explain the discordance.²⁶ Others have found no association between arterial stiffness and cuff compared with invasive BP.^{23,25} Nevertheless, there is physiological rationale that is supportive of arterial stiffness causing differences between cuff BP and invasive aortic BP by altering blood flow dynamics and the properties of signals detected by the upper arm cuff.¹³ In previous studies, a lower heart rate has also been associated with greater underestimation of SBP and overestimation of DBP, and this relationship may be influenced by the cuff deflation rate.^{27,28} Our data is consistent with these observations, although in multivariable models, the relationship between lower heart rate and cuff DBP overestimation was nonsignificant. Furthermore, although older subjects did have lower heart rate, the influence of age on differences between cuff BP and invasive aortic BP remained similar after adjusting for heart rate.

Seminal epidemiological data reporting population-level characteristics and changes in BP with aging have been recorded using cuff BP measurement methods.^{29,30} These studies report a rise in SBP with increasing age and, that from ≈ 50 to 60 years of age, PP also increases due to concomitant decreases in DBP.^{29,31} Importantly, because these observations are from cuff BP, they may underestimate the relationship between aortic SBP and PP with age (according to our invasive observations). Similarly, the decline in invasive aortic DBP with increasing age after 50 years is also likely to be markedly more rapid than observed from cuff DBP measurements. These differences will influence the estimates of strength of association based on epidemiological studies and are probable underlying contributors to clinical uncertainty and debate

around treatment thresholds for SBP,^{19,32,33} DBP,^{34,35} and PP.^{16,32} Despite these issues, decades of evidence unequivocally support the value of cuff BP for prediction of cardiovascular risk in adults across the age spectrum examined in this study.² Nevertheless, the impact of our findings on these uncertainties warrants closer examination in prospective studies.

The current findings may also be relevant to cuff BP device validation protocols that are used to test new devices by comparison to mercury sphygmomanometry. The current universal standard for the validation of BP devices does not take into consideration the potential influence of age on cuff-measured BP.³⁶ Our findings indicate that BP devices should be evaluated among a minimum number of subjects across different decades of age. However, this would not fully address the problem because the influence of age on the cuff BP is likely to extend to the reference comparator, mercury sphygmomanometry. Taken together, this emphasizes the urgent need to find better ways to measure BP (that reflect true invasive aortic BP) without confounding influences from age or other factors.

Subjects were studied under cardiac catheterization conditions and had an indication for coronary angiography; thus, the results may not reflect those that would be observed in the general population. Despite this, there is no data to suggest that the influence of age on cuff BP in patients undergoing cardiac catheterization is different in other populations. Interarm cuff BP differences were not assessed systematically in each individual study, and we cannot rule out that some participants may have had obstructive arterial disease that could have influenced cuff BP compared with invasive aortic BP. Heart rate may also influence cuff BP measurement,^{27,28} but in some studies included in this current analysis, heart rate may not have been recorded simultaneously to BP measurement. The influence of age on cuff BP compared with invasive aortic BP did not change when adjusted for heart rate. Reassuringly, the associations we observed between heart rate and the difference between cuff BP compared with invasive aortic BP are consistent with previous work.²⁸ We could not separately compare the different types of cuff BP devices (eg, mercury versus oscillometric) with invasive aortic BP due to a small sample of data recorded using mercury sphygmomanometry data (n=21). Oscillometric devices are designed to measure the same values as mercury sphygmomanometry, although age, PP, and arterial stiffness can influence differences between these methods.^{26,37} Nevertheless, we did not observe major differences between oscillometric devices or mercury sphygmomanometry compared with invasive brachial BP (Figure S8). The influence of age on cuff BP versus invasive aortic BP for prediction of clinical outcomes or management of hypertension could not be assessed in the present study. Addressing this question should be a research priority.

Perspectives

This study adds to growing evidence that there are substantive differences between cuff BP and invasive BP.^{3,4,6} Although cuff BP is the cornerstone for hypertension management, it is relatively crude and imprecise. In an era of rapid advances in technology and analytics, it is imperative that more personalized methods of BP measurement are developed. Ultimately, better measurement of BP should improve clinical care and lead to a reduction in preventable CVD events.

Acknowledgments

Additional members of the Invasive Blood Pressure Consortium are Ahmed M. Al-Jumaily: Institute of Biomedical Technologies, Auckland University of Technology, Auckland, New Zealand; Brian A. Gould: BMI Hospital Blackheath, London, United Kingdom; Fuyou Liang: School of Naval Architecture, Ocean and Civil Engineering, Shanghai Jiao Tong University, Shanghai, China; Sandy Muecke: Department of Critical Care Medicine, Flinders University, Adelaide, Australia; Ronak Rajani: Cardiology Department, Guy's and St Thomas' Hospitals, London, United Kingdom; Ralph Stewart: Green Lane Cardiovascular Service, Auckland City Hospital, University of Auckland, Auckland, New Zealand; George A. Stouffer: Division of Cardiology, University of North Carolina at Chapel Hill, Chapel Hill, United States; Manish D. Sinha: Department of Clinical Pharmacology and Department of Paediatric Nephrology, Kings College London, Evelina London Children's Hospital, Guy's and St. Thomas' NHS Foundation Trust, London, United Kingdom.

Sources of Funding

This work was supported by a Vanguard Grant from the National Heart Foundation of Australia (reference 101836) and Royal Hobart Hospital Research Foundation project grant (reference 19-202).

Disclosures

J.E. Sharman reports the following: his university has received equipment and research funding from manufacturers of BP devices, including AtCor Medical, IEM, and Pulsecor (Uscom). He has no personal commercial interests related to BP companies. The other authors report no conflicts.

References

- Roth GA, Johnson C, Abajobir A, Abd-Allah F, Abera SF, Abyu G, Ahmed M, Aksut B, Alam T, Alam K, et al. Global, regional, and national burden of cardiovascular diseases for 10 causes, 1990 to 2015. *J Am Coll Cardiol*. 2017;70:1–25. doi: 10.1016/j.jacc.2017.04.052
- Ettehad D, Emdin CA, Kiran A, Anderson SG, Callender T, Emberson J, Chalmers J, Rodgers A, Rahimi K. Blood pressure lowering for prevention of cardiovascular disease and death: a systematic review and meta-analysis. *Lancet*. 2016;387:957–967. doi: 10.1016/S0140-6736(15)01225-8
- Picone DS, Schultz MG, Otahal P, Aakhus S, Al-Jumaily AM, Black JA, Bos WJ, Chambers JB, Chen CH, Cheng HM, et al. Accuracy of cuff-measured blood pressure: systematic reviews and meta-analyses. *J Am Coll Cardiol*. 2017;70:572–586. doi: 10.1016/j.jacc.2017.05.064
- Carlsen RK, Peters CD, Khatir DS, Laugesen E, Bøtker HE, Winther S, Buus NH. Estimated aortic blood pressure based on radial artery tonometry underestimates directly measured aortic blood pressure in patients with advancing chronic kidney disease staging and increasing arterial stiffness. *Kidney Int*. 2016;90:869–877. doi: 10.1016/j.kint.2016.05.014
- Messerli FH, Ventura HO, Amodeo C. Osler's maneuver and pseudohypertension. *N Engl J Med*. 1985;312:1548–1551. doi: 10.1056/NEJM198506133122405
- Picone DS, Schultz MG, Peng X, Black JA, Dwyer N, Roberts-Thomson P, Chen CH, Cheng HM, Pucci G, Wang JG, et al. Discovery of new blood pressure phenotypes and relation to accuracy of cuff devices used in daily clinical practice. *Hypertension*. 2018;71:1239–1247. doi: 10.1161/HYPERTENSIONAHA.117.10696
- Boutouyrie P, London GM, Sharman JE. Estimating central blood pressure in the extreme vascular phenotype of advanced kidney disease. *Kidney Int*. 2016;90:736–739. doi: 10.1016/j.kint.2016.06.039
- Benmira A, Perez-Martin A, Schuster I, Aichoun I, Coudray S, Bereksi-Reguig F, Dauzat M. From korotkoff and marey to automatic non-invasive oscillometric blood pressure measurement: does easiness come with reliability? *Expert Rev Med Devices*. 2016;13:179–189. doi: 10.1586/17434440.2016.1128821
- Chen CH, Nakayama M, Nevo E, Fetis B, Maughan WL, Kass DA. Coupled systolic-ventricular and vascular stiffening with age: implications for pressure regulation and cardiac reserve in the elderly. *J Am Coll Cardiol*. 1998;32:1221–1227. doi: 10.1016/s0735-1097(98)00374-x
- Reference Values for Arterial Stiffness Collaboration. Determinants of pulse wave velocity in healthy people and in the presence of cardiovascular risk factors: 'Establishing normal and reference values'. *Eur Heart J*. 2010;31:2338–2350. doi: 10.1093/eurheartj/ehq165

11. Redheuil A, Yu WC, Mousseaux E, Harouni AA, Kachenoura N, Wu CO, Bluemke D, Lima JA. Age-related changes in aortic arch geometry: relationship with proximal aortic function and left ventricular mass and remodeling. *J Am Coll Cardiol*. 2011;58:1262–1270. doi: 10.1016/j.jacc.2011.06.012
12. Bensalah MZ, Bollache E, Kachenoura N, Giron A, De Cesare A, Macron L, Lefort M, Redheuil A, Redheuil A, Mousseaux E. Geometry is a major determinant of flow reversal in proximal aorta. *Am J Physiol Heart Circ Physiol*. 2014;306:H1408–H1416. doi: 10.1152/ajpheart.00647.2013
13. Hashimoto J, Ito S. Aortic stiffness determines diastolic blood flow reversal in the descending thoracic aorta: potential implication for retrograde embolic stroke in hypertension. *Hypertension*. 2013;62:542–549. doi: 10.1161/HYPERTENSIONAHA.113.01318
14. Booth J. A short history of blood pressure measurement. *Proc R Soc Med*. 1977;70:793–799.
15. Karamanou M, Papaioannou TG, Tsooucalas G, Tousoulis D, Stefanadis C, Androutsos G. Blood pressure measurement: lessons learned from our ancestors. *Curr Pharm Des*. 2015;21:700–704. doi: 10.2174/1381612820666141023163313
16. Sharman JE, Marwick TH. Accuracy of blood pressure monitoring devices: a critical need for improvement that could resolve discrepancy in hypertension guidelines. *J Hum Hypertens*. 2019;33:89–93. doi: 10.1038/s41371-018-0122-6
17. Narayan O, Casan J, Szarski M, Dart AM, Meredith IT, Cameron JD. Estimation of central aortic blood pressure: a systematic meta-analysis of available techniques. *J Hypertens*. 2014;32:1727–1740. doi: 10.1097/HJH.0000000000000249
18. Bos WJ, Verrij E, Vincent HH, Westerhof BE, Parati G, van Montfrans GA. How to assess mean blood pressure properly at the brachial artery level. *J Hypertens*. 2007;25:751–755. doi: 10.1097/HJH.0b013e32803fb621
19. Whelton PK, Carey RM, Aronow WS, Casey DE Jr, Collins KJ, Dennison Himmelfarb C, DePalma SM, Gidding S, Jamerson KA, Jones DW, et al. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APhA/ASH/ASPC/NMA/PCNA guideline for the prevention, detection, evaluation, and management of high blood pressure in adults: a report of the American College of Cardiology/American Heart Association task force on clinical practice guidelines. *J Am Coll Cardiol*. 2018;71:e127–e248. doi: 10.1016/j.jacc.2017.11.006
20. Pickering TG, Hall JE, Appel LJ, Falkner BE, Graves J, Hill MN, Jones DW, Kurtz T, Sheps SG, Roccella EJ; Subcommittee of Professional and Public Education of the American Heart Association Council on High Blood Pressure Research. Recommendations for blood pressure measurement in humans and experimental animals: Part 1: blood pressure measurement in humans: a statement for professionals from the Subcommittee of Professional and Public Education of the American Heart Association Council on High Blood Pressure Research. *Hypertension*. 2005;45:142–161. doi: 10.1161/01.HYP.0000150859.47929.8e
21. Kroeker EJ, Wood EH. Comparison of simultaneously recorded central and peripheral arterial pressure pulses during rest, exercise and tilted position in man. *Circ Res*. 1955;3:623–632. doi: 10.1161/01.res.3.6.623
22. Rowell LB, Brengelmann GL, Blackmon JR, Bruce RA, Murray JA. Disparities between aortic and peripheral pulse pressures induced by upright exercise and vasomotor changes in man. *Circulation*. 1968;37:954–964. doi: 10.1161/01.cir.37.6.954
23. Kuwajima I, Hoh E, Suzuki Y, Matsushita S, Kuramoto K. Pseudohypertension in the elderly. *J Hypertens*. 1990;8:429–432. doi: 10.1097/00004872-199005000-00006
24. Finnegan TP, Spence JD, Wong DG, Wells GA. Blood pressure measurement in the elderly: correlation of arterial stiffness with difference between intra-arterial and cuff pressures. *J Hypertens*. 1985;3:231–235. doi: 10.1097/00004872-198506000-00006
25. Bos WJ, van Goudoever J, Wesseling KH, Rongen GA, Hoedemaker G, Lenders JW, van Montfrans GA. Pseudohypertension and the measurement of blood pressure. *Hypertension*. 1992;20:26–31. doi: 10.1161/01.hyp.20.1.26
26. van Poppel NM, Bos WJ, de Beer NA, van Der Kuip DA, Hofman A, Grobbee DE, Wittman JC. Arterial stiffness as underlying mechanism of disagreement between an oscillometric blood pressure monitor and a sphygmomanometer. *Hypertension*. 2000;36:484–488. doi: 10.1161/01.hyp.36.4.484
27. Zheng D, Amoores JN, Mieke S, Murray A. How important is the recommended slow cuff pressure deflation rate for blood pressure measurement? *Ann Biomed Eng*. 2011;39:2584–2591. doi: 10.1007/s10439-011-0347-9
28. Yong PG, Geddes LA. The effect of cuff pressure deflation rate on accuracy in indirect measurement of blood pressure with the auscultatory method. *J Clin Monit*. 1987;3:155–159. doi: 10.1007/bf01695937
29. Franklin SS, Gustin W 4th, Wong ND, Larson MG, Weber MA, Kannel WB, Levy D. Hemodynamic patterns of age-related changes in blood pressure. The Framingham Heart Study. *Circulation*. 1997;96:308–315. doi: 10.1161/01.cir.96.1.308
30. Wolf-Maier K, Cooper RS, Banegas JR, Giampaoli S, Hense HW, Joffres M, Kastarinen M, Poulter N, Primatesta P, Rodríguez-Artalejo F, et al. Hypertension prevalence and blood pressure levels in 6 European countries, Canada, and the United States. *JAMA*. 2003;289:2363–2369. doi: 10.1001/jama.289.18.2363
31. Scuteri A, Morrell CH, Orrù M, Strait JB, Tarasov KV, Ferrelì LA, Loi F, Pilia MG, Delitala A, Spurgeon H, et al. Longitudinal perspective on the conundrum of central arterial stiffness, blood pressure, and aging. *Hypertension*. 2014;64:1219–1227. doi: 10.1161/HYPERTENSIONAHA.114.04127
32. Williams B, Mancia G, Spiering W, Agabiti Rosei E, Azizi M, Burnier M, Clement DL, Coca A, de Simone G, Dominiczak A, et al; Authors/Task Force Members. 2018 ESC/ESH guidelines for the management of arterial hypertension: the task force for the management of arterial hypertension of the European Society of Cardiology and the European Society of Hypertension: the task force for the management of arterial hypertension of the European Society of Cardiology and the European Society of Hypertension. *J Hypertens*. 2018;36:1953–2041. doi: 10.1097/HJH.0000000000001940
33. Messerli FH, Bangalore S. The blood pressure landscape: schism among guidelines, confusion among physicians, and anxiety among patients. *J Am Coll Cardiol*. 2018;72:1313–1316. doi: 10.1016/j.jacc.2018.07.026
34. Mancia G, Grassi G. Aggressive blood pressure lowering is dangerous: the J-curve: pro side of the argument. *Hypertension*. 2014;63:29–36. doi: 10.1161/01.hyp.0000441190.09494.e9
35. Verdecchia P, Angeli F, Mazzotta G, Garofoli M, Reboldi G. Aggressive blood pressure lowering is dangerous: the J-curve: con side of the argument. *Hypertension*. 2014;63:37–40. doi: 10.1161/01.hyp.0000439102.43479.43
36. Stergiou GS, Palatini P, Asmar R, Ioannidis JP, Kollias A, Lacy P, McManus RJ, Myers MG, Parati G, Shennan A, et al; European Society of Hypertension Working Group on Blood Pressure Monitoring. Recommendations and practical guidance for performing and reporting validation studies according to the universal standard for the validation of blood pressure measuring devices by the Association for the Advancement of Medical Instrumentation/European Society of Hypertension/International Organization for Standardization (AAMI/ESH/ISO). *J Hypertens*. 2019;37:459–466. doi: 10.1097/HJH.0000000000002039
37. Stergiou GS, Lourida P, Tzamouranis D, Baibas NM. Unreliable oscillometric blood pressure measurement: prevalence, repeatability and characteristics of the phenomenon. *J Hum Hypertens*. 2009;23:794–800. doi: 10.1038/jhh.2009.20

Novelty and Significance

What Is New?

- Cuff blood pressure (BP) is influenced by increasing age, whereby invasive systolic BP and PP are progressively underestimated, but invasive diastolic BP is progressively overestimated.
- Age-related trends were independent of BP control and similar for comparisons of cuff BP and invasive brachial BP.

What Is Relevant?

- The findings may have implications for BP management with increasing age, population-level studies of BP, hypertension guideline thresholds, and validation protocols that test new BP devices.

Summary

This study has shown that the difference between cuff BP and invasive aortic BP is substantially influenced by increasing age. Altogether, the data underline the need to improve the quality of BP measurement devices for people of all ages.