



Universiteit
Leiden
The Netherlands

Digenic inheritance of MSH6 and MUTYH variants in familial colorectal cancer

Schubert, S.A.; Ruano, D.; Tiersma, Y.; Drost, M.; Wind, N. de; Nielsen, M.; ... ; Wezel, T. van

Citation






Schubert, S. A., Ruano, D., Tiersma, Y., Drost, M., Wind, N. de, Nielsen, M., ... Wezel, T. van. (2020). Digenic inheritance of MSH6 and MUTYH variants in familial colorectal cancer. *Genes, Chromosomes And Cancer*, 59(12), 697-701. doi:10.1002/gcc.22883

Version: Publisher's Version
License: [Creative Commons CC BY 4.0 license](#)
Downloaded from: <https://hdl.handle.net/1887/3181895>

Note: To cite this publication please use the final published version (if applicable).

BRIEF REPORT

Digenic inheritance of *MSH6* and *MUTYH* variants in familial colorectal cancer

Stephanie A. Schubert¹  | Dina Ruano¹ | Yvonne Tiersma² | Mark Drost² | Niels de Wind²  | Maartje Nielsen³  | Liselotte P. van Hest⁴ | Hans Morreau¹ | Noel F. C. C. de Miranda¹  | Tom van Wezel¹ 

¹Department of Pathology, Leiden University Medical Center, Leiden, The Netherlands

²Department of Human Genetics, Leiden University Medical Center, Leiden, The Netherlands

³Department of Clinical Genetics, Leiden University Medical Center, Leiden, The Netherlands

⁴Department of Clinical Genetics, Amsterdam UMC, Vrije Universiteit Amsterdam, Amsterdam, The Netherlands

Correspondence

Tom van Wezel, Department of Pathology, Leiden University Medical Center, Leiden University, Albinusdreef 2, 2333 ZA Leiden, The Netherlands.
Email: t.van_wezel@lumc.nl

Funding information

KWF Kankerbestrijding, Grant/Award Number: 2015-7664; Maag Lever Darm Stichting, Grant/Award Number: MLDS FP13-13; Stichting Sacha Swarttouw-Hijmans; ZonMw Veni, Grant/Award Number: 016.176.144; European Cooperation in Science and Technology (COST); Leiden University Fund/Nypels-van der Zee Fonds

Abstract

We describe a family severely affected by colorectal cancer (CRC) where whole-exome sequencing identified the coinheritance of the germline variants encoding *MSH6* p.Thr1100Met and *MUTYH* p.Tyr179Cys in, at least, three CRC patients diagnosed before 60 years of age. Digenic inheritance of monoallelic *MSH6* variants of uncertain significance and *MUTYH* variants has been suggested to predispose to Lynch syndrome-associated cancers; however, cosegregation with disease in the familial setting has not yet been established. The identification of individuals carrying multiple potential cancer risk variants is expected to rise with the increased application of whole-genome sequencing and large multigene panel testing in clinical genetic counseling of familial cancer patients. Here we demonstrate the coinheritance of monoallelic variants in *MSH6* and *MUTYH* consistent with cosegregation with CRC, further supporting a role for digenic inheritance in cancer predisposition.

KEYWORDS

digenic inheritance, familial colorectal cancer, Lynch syndrome, *MSH6*, *MUTYH*, whole-exome sequencing

1 | INTRODUCTION

Approximately 25% of colorectal cancers (CRCs) are diagnosed in patients with a family history of CRC. However, the majority of familial CRC cannot be explained by clear-cut genetic defects, which hampers appropriate genetic counselling.¹ The most frequent form of hereditary CRC is Lynch syndrome (OMIM#120435), which predisposes to cancers that develop in a context of DNA mismatch repair (MMR) deficiency, including CRC and endometrial cancer. It is caused by heterozygous, pathogenic variants affecting the DNA MMR genes,

MLH1, *MSH2*, *MSH6*, or *PMS2*. *MUTYH*-associated polyposis (MAP; OMIM#608456) is a recessively inherited CRC syndrome caused by biallelic variants in the base-excision repair gene *MUTYH*. The potential of monoallelic, pathogenic *MUTYH* variants to predispose to CRC remains debatable.¹ Some *MUTYH* variants confer greater functional defects in vitro and are associated with more severe clinical phenotypes, such as the variant encoding p.Tyr179Cys compared to p.Gly396Asp.^{2,3}

Digenic inheritance of monoallelic *MSH6* and *MUTYH* variants has been suggested to predispose to Lynch syndrome-associated cancers;

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2020 The Authors. *Genes, Chromosomes & Cancer* published by Wiley Periodicals LLC

however, cosegregation of both variants within CRC families has not yet been demonstrated.⁴⁻⁹ Here, we demonstrate, for the first time, the coinheritance of monoallelic variants in *MSH6* and *MUTYH* consistent with the cosegregation with CRC, further supporting a role for digenic inheritance in cancer predisposition.

2 | MATERIALS AND METHODS

2.1 | Patients

Clinicopathological data of family members was obtained during consultations at the department of Clinical Genetics of the Amsterdam University Medical Centre, Vrije Universiteit Amsterdam. DNA was extracted from peripheral blood and formalin-fixed paraffin-embedded tissues using standard techniques. All patients provided written informed consent. The study was approved by the Medical Ethical Committee of the Leiden University Medical Center, The Netherlands (protocol P01.019).

2.2 | Whole-exome sequencing

Whole-exome sequencing was outsourced to BGI (BGI-Shenzhen, Shenzhen, China); exome libraries were constructed with the BGI capture kit, followed by sequencing on the Complete Genomics' Sequencing Platform (Complete Genomics Inc., San Jose, California). Filtering and variant prioritization was performed as previously described.¹⁰ All variants were selected based on a maximum population frequency <0.01 (in 1000 Genomes phase 3, ExAC 1.0, ESP6500SI-V2 or GoNL release 5).

2.3 | Variant screening

The *MSH6* (p.Thr1100Met) and *MUTYH* (p.Tyr179Cys) variants were validated and investigated in additional family members by using Sanger sequencing of PCR products obtained under standard PCR conditions. The following M13-tailed primer sets were used: 5'-TGT AAA ACG ACG GCC AGT AAA ACC CCC AAA CGA TGA A-3' and 5'-CAG GAA ACA GCT ATG ACC TGC TCC TCT TCC TCA CAG-3' for *MSH6*, and 5'-GAC GTT GTA AAA CGA CGG CCA GTC CCT AGG GTA GGG GAA ATA GG-3' and 5'-CAG GAA ACA GCT ATG ACC ATG AGT TCC TAC CCT CCT GCC ATC-3' for *MUTYH* (M13-tails are underlined).

2.4 | Tumor analysis

MMR deficiency in tumor samples was assessed by microsatellite instability analysis and immunohistochemical detection of the four MMR proteins (MLH1, MSH2, MSH6, and PMS2).¹¹ KRAS codon 12/13 mutations were screened with Sanger sequencing.¹²

2.5 | Functional MMR assay

In vitro MMR activity assay was performed as previously described.¹³

3 | RESULTS

We performed germline whole-exome sequencing on three CRC patients diagnosed before 60 years of age (III-1, III-7, III-8, Figure 1A)




FIGURE 1 The digenic inheritance of *MSH6* and *MUTYH* variants. A, The pedigree shows the coinheritance of the monoallelic variants which encode *MSH6* p.Thr1100Met and *MUTYH* p.Tyr179Cys in a family affected by colorectal cancer. All spouses were unrelated and unaffected by cancer. Genotypes: *MSH6* p.Thr1100Met (T1100M; blue); *MUTYH* p.Tyr179Cys (Y179C; green); -, wild type. E, whole-exome sequencing analysis; T, tumor analysis; ?, unknown phenotype; numbers in symbols, number of unaffected relatives merged for clarity; filled symbols, cancer patients; C, colorectal cancer; E, endometrial cancer; O, ovarian cancer; d., age at death; followed by the age at diagnosis or death. B, in vitro mismatch repair (MMR) activity assay shows wild-type MMR activity of *MSH6* p.Thr1100Met, compared to wild-type *MSH6* (p.Gly529Gly) and a pathogenic *MSH6* mutant (p.Gly1139Ser). Data are shown as mean \pm SEM of three independent experiments [Color figure can be viewed at wileyonlinelibrary.com]

TABLE 1 All rare variants shared by the three individuals from whole-exome sequencing data

Chr	Gene	RefSeq accession number	mRNA change	Protein change	Population frequency ^a	ClinVar classification ^b	Franklin classification ^c	Cancer gene census
1	EBNA1BP2	NM_001159936	c.1034A > T	p.Asn345Ile	0.006009	—	Benign	—
1	MUTYH	NM_001128425	c.536A > G	p.Tyr179Cys	0.001538	Pathogenic	Pathogenic	Yes
1	TESK2	NM_007170	c.983A > G	p.Gln328Arg	0.0006052	—	VUS	—
1	CAPN9	NM_006615	c.55G > T	p.Ala19Ser	0.00006365	—	VUS	—
2	MSH6	NM_000179	c.3299C > T	p.Thr1100Met	0.00004243	Uncertain	VUS	Yes
3	C3orf20	NM_032137	c.1746C > G	p.Phe582Leu	0.005847	—	Likely benign	—
5	DNAH5	NM_001369	c.1781A > G	p.Glu594Gly	—	—	VUS	—
7	KIAA1324L	NM_001142749	c.2369 T > C	p.Val1790Ala	0.0006585	—	VUS	—
7	TRIP6	NM_003302	c.822G > C	p.Glu274Asp	0.0009893	—	VUS	—
7	CUX1	NM_001202543	c.1438A > G	p.Ser480Gly	0.001128	—	Likely benign	Yes
7	ZNF783	NM_001195220	c.46A > G	p.Thr16Ala	0.001083	—	VUS	—
8	PDP1	NM_018444	c.283A > C	p.Ser95Arg	—	—	VUS	—
9	NMRK1	NM_017881	c.304C > G	p.Leu102Val	0.001419	—	VUS	—
9	GAPVD1	NM_015635	c.850G > A	p.Val284Met	0.003596	—	Benign	—
11	INTS5	NM_030628	c.1436A > G	p.Asn479Ser	0.00004607	—	VUS	—
11	GAL3ST3	NM_033036	c.326G > A	p.Arg109His	0.00004731	—	VUS	—
11	SORL1	NM_003105	c.3346A > G	p.Ile1116Val	0.005308	—	VUS	—
14	LTPB2	NM_000428	c.1226G > A	p.Arg409His	0.0000203	—	VUS	—
15	RYR3	NM_001036	c.7812C > G	p.Asn2604Lys	0.002144	Likely benign	Likely benign	—
15	DAPK2	NM_014326	c.179G > A	p.Arg60Gln	0.003725	—	Likely benign	—
16	NLRC5	NM_032206	c.1219G > A	p.Ala407Thr	0.000003542	—	VUS	—
20	C20orf85	NM_178456	c.101G > A	p.Arg34Gln	0.00192	—	Likely benign	—

Abbreviations: Chr, chromosome; VUS, variant of uncertain significance.

^aPopulation frequency (gnomAD 2.1.1).^bClinVar clinical significance (ClinVar database version August 5, 2019).^cFranklin by Genoox (accessed on May 20, 2020).

and who belonged to a CRC family comprising of seven cancer patients divided over two generations. Twenty-two rare variants were shared by the three patients (Tables 1 and S1), including variants in the *MSH6* (NM_000179.2: c.3299C > T, p.Thr1100Met) and *MUTYH* (NM_001128425.1: c.536A > G, p.Tyr179Cys) genes, while the other 20 genes could not be clearly linked to cancer predisposition. The identified *MSH6* variant was classified as a variant of uncertain significance (VUS) in the Leiden Open Variant Database and the InSiGHT DNA Variant Database.^{14,15} The *MUTYH* variant is the most common pathogenic variant found in the Netherlands.²

Fourteen relatives, all unaffected by cancer or polyposis, were genotyped for these *MSH6* and *MUTYH* variants, identifying one additional carrier of both variants, five *MSH6*-only carriers and four *MUTYH*-only carriers. In all probability, the mothers of the sequenced patients, II-1 and II-2, who were affected by ovarian cancer bellow age 74 and CRC at 38 years old respectively, were obligate carriers of both variants; however, DNA was unavailable for testing and, formally, inheritance through the fathers to the sequenced individuals (III-1, III-7, III-8) cannot be excluded. MMR deficiency was not detected in the colorectal carcinoma of patient III-1, which also lacked the *KRAS* mutation typical for MAP tumors (c.34C > T; Table S2). Functional analysis of the *MSH6* p.Thr1100Met variant showed retained MMR function in vitro (Figure 1B).

4 | DISCUSSION

Digenic inheritance of monoallelic *MSH6* and *MUTYH* variants has been suggested to predispose to Lynch syndrome-associated cancers. The involvement of both *MSH6* and *MUTYH* in oxidative DNA damage repair and their physical interaction enhancing *MUTYH*'s repair activity, substantiates the association of variants in these genes.¹⁶ From earlier studies, the inheritance of monoallelic *MUTYH* variants seemed primarily relevant in patients carrying *MSH6* VUSs, which are less strongly associated with MMR deficiency than pathogenic *MSH6* variants (Table S2).⁴⁻⁹ Furthermore, a digenic inheritance model was proposed once before for CRC predisposition in a carrier of variants in the oxidative DNA damage repair genes *MUTYH* and *OGG1*.¹⁷ Although the functional evidence of combined defects in oxidative DNA damage repair genes is still lacking, the coinheritance of *MSH6* and *MUTYH* variants in at least three, but likely five cancer cases within one family warrants further mechanistic and clinical studies. The absence of cancer and numerous polyps in nondigenic carriers further substantiates this association. Tumor analysis of the tumor of one of the digenic carriers and the in vitro MMR activity assay indicated retention of MMR function of *MSH6* p.Thr1100Met protein. In addition, the genetic marker for MAP-tumors (*KRAS* c.34G > T) was absent in this tumor, which points toward retained *MUTYH* repair activity. The combined inheritance of both genetic variants could still result in impaired repair of oxidative DNA damage. More extensive somatic mutation analysis to assess this was, however, not possible, because of low quality of the DNA sample and the unavailability of additional tumor material.

Next to *MSH6* and *MUTYH*, *CUX1* has been described as a cancer-driving gene.¹⁸ *CUX1* is implicated in inflammatory bowel disease and various cancer types, although primarily due to loss-of-function somatic mutations.^{18,19} This gene codes for several isoforms, including the ubiquitously expressed p200 *CUX1*, which, among other functions, has been shown to stimulate the repair of oxidized DNA bases by *OGG1*.²⁰ The identified *CUX1* (NM_001202543: c.1438A > G, p.Ser480Gly) variant, however, was classified as likely benign by the Franklin variant classification tool.²¹ Additional gene reportedly linked to tumorigenesis include *RYR3*,²² *EBNA1BP2*,²³ *TRIP6*,²⁴ and *CAPN9*.²⁵ The *RYR3* (NM_001036: c.7812C > G, p. Asn2604Lys) and *EBNA1BP2* (NM_001159936: c.1034A > T, p. Asn345Ile) variants were classified as likely benign and benign, respectively, while the *TRIP6* (NM_003302: c.822G > C, p.Glu274Asp) and the *CAPN9* (NM_006615: c.55G > T, p.Ala19Ser) variants were classified as VUS.²¹ *TRIP6* promotes cell migration and invasion through Wnt/ β -catenin signaling and was shown to be upregulated in colorectal tumors.²⁴ Therefore, *TRIP6* variants that increase protein stability or expression could potentially stimulate colorectal tumorigenesis. In addition, lost-of-function variants in *CAPN9* might promote tumor formation, as Calpain-9 induces cell cycle arrest and apoptosis, and low expression predicts a poorer prognosis in gastric cancer patients.²⁵ The contribution of the genetic variants, other than *MSH6* and *MUTYH*, to cancer risk cannot be completely excluded. However, none of these variants have been functionally investigated and especially the variants predicted as benign or likely benign are less likely to contribute to an increased cancer risk. Besides, none of these genes have, to date, been associated with a genetic predisposition to any types of cancer.

In conclusion, with the increased application of whole-genome sequencing or large multigene panel testing in clinical genetic counseling, the number of identified individuals carrying multiple potential risk variants is expected to rise. Here, we demonstrate the coinheritance of *MSH6* and *MUTYH* variants consistent with the cosegregation with cancer, further supporting a role for digenic inheritance in CRC predisposition. Our results reiterate that digenic inheritance should be considered as cause of genetic diseases.

ACKNOWLEDGMENTS

The authors thank Juul T. Wijnen for the collection of clinicopathological data and samples. The authors also thank Julia van Hees and Anniek van Veen for their technical support. The authors are thankful to the Leiden University Fund/Nypels-van der Zee Fonds. This project was funded by research grants from the Dutch Digestive Foundation (MLDS FP13-13) and Stichting Sacha Swarttouw-Hijmans awarded to T.v.W. N.F.C.C.d.M. is supported by the KWF Bas Mulder Award UL (2015-7664) and the ZonMw Veni grant (016.176.I44). This article is based upon work from COST Action CA17118, supported by European Cooperation in Science and Technology (COST).

CONFLICT OF INTEREST

The authors declare no conflicts of interest.

AUTHOR CONTRIBUTIONS

Tom van Wezel, Noel F. C. C. de Miranda, and Hans Morreau conceived and designed the study. Dina Ruano performed next-generation sequencing analyses. Noel F. C. C. de Miranda and Stephanie A. Schubert performed analysis and interpretation of whole-exome sequencing data. Mark Drost and Yvonne Tiersma performed functional analysis. Maartje Nielsen and Liselotte P. van Hest performed patient counseling and clinical data acquisition. Hans Morreau performed the pathology review of the samples. Tom van Wezel, Noel F. C. C. de Miranda, Mark Drost, and Niels de Wind supervised the work. Stephanie A. Schubert, Noel F. C. C. de Miranda, and Tom van Wezel wrote the manuscript. All authors read and approved the manuscript.

DATA AVAILABILITY STATEMENT

The data that support the findings of this study are available from the corresponding author upon reasonable request. The data are not publicly available due to privacy restrictions.

ORCID

Stephanie A. Schubert  <https://orcid.org/0000-0002-8655-1350>

Niels de Wind  <https://orcid.org/0000-0002-3143-5061>

Maartje Nielsen  <https://orcid.org/0000-0002-5351-1870>

Noel F. C. C. de Miranda  <https://orcid.org/0000-0001-6122-1024>

Tom van Wezel  <https://orcid.org/0000-0001-5773-7730>

REFERENCES

- Schubert SA, Morreau H, de Miranda NFCC, van Wezel T. The missing heritability of familial colorectal cancer. *Mutagenesis*. 2020;35(3):221-231.
- Nielsen M, Joerink-van de Beld MC, Jones N, et al. Analysis of MUTYH genotypes and colorectal phenotypes in patients with MUTYH-associated polyposis. *Gastroenterology*. 2009;136(2):471-476.
- Komine K, Shimodaira H, Takao M, et al. Functional complementation assay for 47 MUTYH variants in a MutY-disrupted *Escherichia coli* strain. *Hum Mutat*. 2015;36(7):704-711.
- Niessen RC, Sijmons RH, Ou J, et al. MUTYH and the mismatch repair system: partners in crime? *Hum Genet*. 2006;119(1-2):206-211.
- Steinke V, Rahner N, Morak M, et al. No association between MUTYH and MSH6 germline mutations in 64 HNPCC patients. *Eur J Human Genet*. 2008;16(5):587-592.
- Giraldez MD, Balaguer F, Caldes T, et al. Association of MUTYH and MSH6 germline mutations in colorectal cancer patients. *Fam Cancer*. 2009;8(4):525-531.
- Giraldez MD, Balaguer F, Bujanda L, et al. MSH6 and MUTYH deficiency is a frequent event in early-onset colorectal cancer. *Clin Cancer Res*. 2010;16(22):5402-5413.
- van Puijenbroek M, Nielsen M, Reinards THCM, et al. The natural history of a combined defect in MSH6 and MUTYH in a HNPCC family. *Fam Cancer*. 2007;6(1):43-51.
- Win AK, Reece JC, Buchanan DD, et al. Risk of colorectal cancer for people with a mutation in both a MUTYH and a DNA mismatch repair gene. *Fam Cancer*. 2015;14(4):575-583.
- Schubert SA, Ruano D, Elsayed FA, et al. Evidence for genetic association between chromosome 1q loci and predisposition to colorectal neoplasia. *Br J Cancer*. 2017;117(6):1215-1223.
- de Jong AE, van Puijenbroek M, Hendriks Y, et al. Microsatellite instability, immunohistochemistry, and additional PMS2 staining in suspected hereditary nonpolyposis colorectal cancer. *Clin Cancer Res*. 2004;10(3):972-980.
- Nielsen M, Poley JW, Verhoef S, et al. Duodenal carcinoma in MUTYH-associated polyposis. *J Clin Pathol*. 2006;59(11):1212-1215.
- Drost M, Tiersma Y, Glubb D, et al. Two integrated and highly predictive functional analysis-based procedures for the classification of MSH6 variants in Lynch syndrome. *Genet Med*. 2020;22(5):847-856.
- LOVD. *Leiden Open Variant Database v.3.0 Build 22*. <https://www.lovd.nl>. Accessed May 11, 2020.
- InSIGHT. *DNA Variant Database*. <http://insight-database.org/>. Accessed May 11, 2020.
- Gu Y, Parker A, Wilson TM, Bai H, Chang DY, Lu AL. Human MutY homolog, a DNA glycosylase involved in base excision repair, physically and functionally interacts with mismatch repair proteins human MutS homolog 2/human MutS homolog 6. *J Biol Chem*. 2002;277(13):11135-11142.
- Morak M, Massdorf T, Sykora H, Kerscher M, Holinski-Feder E. First evidence for digenic inheritance in hereditary colorectal cancer by mutations in the base excision repair genes. *Eur J Cancer*. 2011;47(7):1046-1055.
- Wong CC, Martincorena I, Rust AG, et al. Inactivating CUX1 mutations promote tumorigenesis. *Nat Genet*. 2014;46(1):33-38.
- Darsigny M, St-Jean S, Boudreau F. Cux1 transcription factor is induced in inflammatory bowel disease and protects against experimental colitis. *Inflamm Bowel Dis*. 2010;16(10):1739-1750.
- Ramdzan ZM, Vadnais C, Pal R, et al. RAS transformation requires CUX1-dependent repair of oxidative DNA damage. *PLoS Biol*. 2014;12(3):e1001807.
- Franklin. franklin.genoox.com. Accessed May 20, 2020.
- Chae YS, Kim JG, Kang BW, et al. Functional polymorphism in the MicroRNA-367 binding site as a prognostic factor for colonic cancer. *Anticancer Res*. 2013;33(2):513-519.
- Liao P, Wang W, Shen M, et al. A positive feedback loop between EBP2 and c-Myc regulates rDNA transcription, cell proliferation, and tumorigenesis. *Cell Death Dis*. 2014;5(1):e1032.
- Chastre E, Abdessamad M, Kruglov A, et al. TRIP6, a novel molecular partner of the MAGI-1 scaffolding molecule, promotes invasiveness. *FASEB J*. 2009;23(3):916-928.
- Peng P, Wu W, Zhao J, et al. Decreased expression of Calpain-9 predicts unfavorable prognosis in patients with gastric cancer. *Sci Rep*. 2016;6:29604.

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of this article.

How to cite this article: Schubert SA, Ruano D, Tiersma Y, et al. Digenic inheritance of *MSH6* and *MUTYH* variants in familial colorectal cancer. *Genes Chromosomes Cancer*. 2020; 59:697-701. <https://doi.org/10.1002/gcc.22883>