



Universiteit
Leiden
The Netherlands

Galactic substructures as tracers of dark matter and stellar evolution

Reino, S.

Citation

Reino, S. (2022, September 27). *Galactic substructures as tracers of dark matter and stellar evolution*. Retrieved from <https://hdl.handle.net/1887/3464660>

Version: Publisher's Version

[Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

License: <https://hdl.handle.net/1887/3464660>

Note: To cite this publication please use the final published version (if applicable).

*Galactic substructures as tracers of
dark matter and stellar evolution*

*Galactic substructures as tracers of
dark matter and stellar evolution*

Proefschrift

ter verkrijging van
de graad van doctor aan de Universiteit Leiden,
op gezag van rector magnificus prof. dr. ir. H. Bijl,
volgens besluit van het college voor promoties
te verdedigen op dinsdag 27 september 2022
klokke 13:45 uur

door

Stella Reino

geboren te Tallinn
in 1991

Promotor: Prof. dr. K. H. Kuijken
Co-promotores: Dr. E. M. Rossi
Dr. R. E. Sanderson (University of Pennsylvania)

Promotiecommissie: Prof. dr. H. J. A. Röttgering
Prof. dr. H. Hoekstra
Dr. A. G. A. Brown
Prof. dr. D. W. Hogg (New York University)
Prof. dr. K. V. Johnston (Columbia University)
Prof. dr. D. Zaritsky (University of Arizona)

ISBN 978-94-6469-015-6
Cover design by Janno Jõulu
Printed by ProefschriftMaken

*“Behind all seen things lies something vaster;
everything is but a path, a portal or a window
opening on something other than itself.”*

– Wind, Sand and Stars by Antoine de Saint-Exupéry

Contents

1	Introduction	1
1.1	The structure of the Galaxy	3
1.1.1	Open clusters	6
1.1.2	Stellar streams and halo substructure	7
1.2	Galactic potential	11
1.2.1	Extracting the Galactic potential with stellar streams	14
1.2.2	The action-clustering method	16
1.3	Thesis outline	19
2	Galactic potential constraints from streams	23
2.1	Introduction	24
2.2	Method	26
2.2.1	Stäckel potential	27
2.2.2	Actions	30
2.2.3	Determination of the best-fit potential	31
2.2.4	Determination of confidence intervals	34
2.3	Stream catalogue	36
2.3.1	The GD-1 stream	38
2.3.2	Orphan Stream	39
2.3.3	The Palomar 5 stream	40
2.3.4	The Helmi stream	41
2.4	Results for a single-component potential	42
2.4.1	Results for the individual stream data sets	42
2.4.2	Results for the combined data set	44
2.5	Results for a two-component potential	47
2.5.1	Results for individual stream data sets	47
2.5.2	Results for the combined data set	51
2.6	Validation	52

2.6.1	Tests of fitting assumptions	53
2.6.2	Predicted orbits	55
2.6.3	Action space and stream orbital phase	55
2.7	Comparison and discussion	60
2.7.1	Enclosed mass estimates	60
2.7.2	The Galactic rotation curve	62
2.8	Conclusions	65
2.A	Data collection	66
2.A.1	GD-1 data	66
2.A.2	Orphan Stream data	68
2.A.3	Palomar 5 stream data	70
2.B	KLD1 example	72
3	Systematics in constraints from stellar streams	77
3.1	Introduction	78
3.2	Method	80
3.2.1	Stäckel potential	80
3.2.2	Actions	83
3.2.3	Clustering measurement	83
3.3	Simulation data	86
3.4	Full stream results	91
3.5	Sections of streams	98
3.6	Orbital phase	99
3.7	Bias dependence on other stream properties	102
3.8	Conclusions and Discussion	106
3.A	Stream velocities	110
3.B	Bias as a function of further stream properties	110
4	Galactic potential constraints from H3	113
4.1	Introduction	114
4.2	Method	116
4.2.1	Measuring action-space clustering	116
4.2.2	Trial potentials	118
4.3	Data	120
4.4	Results	123
4.5	Simulations	126
4.5.1	Interlopers	127
4.5.2	Measurement errors	128
4.5.3	Leave-one-out cross-validation	129
4.6	Leave-one-out cross-validation in the H3 sample	130

4.7	Conclusions and Discussion	135
5	Gaia study of the Hyades	141
5.1	Introduction	142
5.2	Data	143
5.2.1	Astrometry	143
5.2.2	Radial velocities	144
5.3	Membership selection	144
5.3.1	Centre of the cluster	144
5.3.2	Velocity of the cluster	145
5.3.3	Probability of membership	146
5.3.4	Final distance and velocity	148
5.3.5	Systematic astrometric errors	148
5.4	Validation and discussion of membership results	149
5.4.1	Comparison with earlier studies	149
5.4.2	Stellar density of the galactic field	151
5.4.3	Colour-absolute magnitude diagram	153
5.4.4	Spatial distribution of members	156
5.4.5	Velocity distribution of members	159
5.5	Kinematic modelling	162
5.5.1	Overview	162
5.5.2	Initial conditions	163
5.5.3	Stopping criterion	164
5.6	Results	165
5.6.1	Velocity distribution	166
5.6.2	Velocity dispersion	167
5.6.3	Parallax comparison	168
5.6.4	Systematic errors and correlations	168
5.6.5	Astrometric radial velocities	171
5.6.6	Colour-absolute magnitude diagram	172
5.7	Discussion and conclusions	176
5.A	Radial velocity data	177
5.A.1	Perryman et al. (1998)	177
5.A.2	Griffin (2012)	177
5.A.3	Griffin (2013)	178
5.A.4	Mermilliod et al. (2009)	178
5.A.5	White et al. (2007)	178
5.A.6	Tabernero et al. (2012)	178
5.A.7	Debernardi et al. (2000)	178
5.A.8	HADES	178

5.A.9 XHIP	179
5.A.10 RAVE	179
5.A.11 APOGEE	179
5.A.12 GALAH	179
5.A.13 Epoch radial velocities	179
5.A.14 Creating a final, merged table	180
5.B Johnson photometry	181
5.C Double stars	181
5.D Isochrones	182
5.E Expansion and rotation	183
5.F Extension of Lindegren et al. (2000)	184
Bibliography	187
List of publications	197
English Summary	199
Nederlandse Samenvatting	205
Eestikeelne kokkuvõte	211
Curriculum Vitae	217
Acknowledgements	219