

Immune checkpoint inhibitors in mesothelioma Disselhorst, M.J.

Citation

Disselhorst, M. J. (2022, October 25). *Immune checkpoint inhibitors in mesothelioma*. Retrieved from https://hdl.handle.net/1887/3483978

Version:	Publisher's Version
License:	<u>Licence agreement concerning inclusion of doctoral</u> <u>thesis in the Institutional Repository of the University</u> <u>of Leiden</u>
Downloaded from:	https://hdl.handle.net/1887/3483978

Note: To cite this publication please use the final published version (if applicable).



PART III

Translational research



Chapter 6

Efficacy of nivolumab and ipilimumab in patients with malignant pleural mesothelioma is related to a subtype of effector memory cytotoxic T cells: translational evidence from two clinical trials.

Joanne M. Mankor ^{*a,b}, Maria J. Disselhorst ^{*c}, Myrthe Poncin ^a, Paul Baas ^c, Joachim G.J.V. Aerts ^{a,b}, Heleen Vroman ^{a,b}
^a Department of Pulmonary Medicine, Erasmus Medical Center Rotterdam, Doctor Molewaterplein 40, 3015 GD Rotterdam, the Netherlands.
^b Erasmus MC Cancer Institute, Erasmus Medical Center Rotterdam, Doctor Molewaterplein 40, 3015 GD Rotterdam, the Netherlands.
^c Department of Thoracic Oncology, NKI-AVL, Amsterdam, the Netherlands ^{*}these authors contributed equally

EBioMedicine. 2020 Dec;62:103040

Abstract

Background

Combined immune checkpoint inhibitor (ICI) treatment targeting PD-1 and CTLA-4 was suggested to yield clinical benefit over chemotherapy in malignant pleural mesothelioma (MPM), whereas aPD-1 monotherapy failed to provide benefit in phase-III trials. Success of ICI depends on the presence and activation of tumor-specific T cells. Therefore, we investigated whether T-cell characteristics are underlying clinical efficacy of ICI treatment in MPM.

Methods

Comprehensive immune cell profiling was performed on screening and on treatment peripheral blood samples of mesothelioma patients treated with nivolumab (aPD-1) monotherapy (NCT02497508), or a combination of nivolumab and ipilimumab (aCTLA-4) (NCT03048474).

Findings

aPD-1/aCTLA-4 combination treatment induced a profound increase in proliferation and activation of T cells, which was not observed upon aPD-1 monotherapy. Moreover, patients that responded to combination treatment had low frequencies of naive CD8 T cells and high frequencies of effector memory CD8 T cells that re-expressed RA (TEMRA) at screening. The frequency of Granzyme-B and Interferon-g producing TEMRAs was also higher in responding patients.

Interpretation

High proportions of TEMRAs and cytokine production by TEMRAs before treatment, was associated with a better clinical outcome. TEMRAs, which likely comprise tumor-specific T cells, tend to require blockage of both aPD-1 and aCTLA-4 to be reactivated. In conclusion, peripheral blood TEMRAs can play a key role in explaining and predicting clinical benefit upon aPD-1/aCTLA-4 combination treatment.

Funding

Bristol-Myers Squibb sponsored NivoMes and INITIATE clinical trials and provided study drugs. No external funding was applicable for the flow cytometric analyses of peripheral blood samples described in this manuscript.

Research in context

Evidence before this study

Immune monitoring, the assessment of peripheral blood immune cell subsets, yielded valuable insight into peripheral blood T-cell responses to immune checkpoint inhibitors (ICI) in non-small cell lung cancer (NSCLC) and melanoma patients. We searched Pubmed for scientific literature published between Jan 1st 2010 and June 15th 2020 with the following terms: "mesothelioma" AND ("PD-1" OR "PD-L1" OR "CTLA-4" OR "checkpoint") AND ("peripheral blood" OR "immune monitoring"). No previous studies have assessed the peripheral blood immune cell compartment upon ICI treatment in malignant pleural mesothelioma (MPM).

Added value of this study

To our knowledge, we are the first to perform extensive immune monitoring in MPM patients treated with both aPD-1 monotherapy and aPD-1/aCTLA-4 combination therapy. Recently, promising results of Checkmate-743 (NCT02899299) demonstrated that treatment of MPM patients with nivolumab and ipilimumab yielded a statistically significant and clinically meaningful improvement in overall survival, compared to platinum-based chemotherapy plus pemetrexed. These results are in contrast to the lack of benefit seen earlier in the PROMISE-meso trial (NCT02991482) that investigated nivolumab monotherapy as compared to chemotherapy in MPM. We here provide a rationale for the benefit observed upon aPD-1/aCTLA-4 combination treatment in MPM by indicating differences in the peripheral blood T-cell compartment in two phase II clinical trials that assessed aPD-1 monotherapy and aPD-1/aCTLA-4 combination therapy.

Implications of all the available evidence

Combination checkpoint inhibition appears to be more effective than their use alone in MPM, which was already shown in the MAPS2 phase II randomized trial. Preliminary results of the Checkmate-743 support this statement. These findings, combined with our peripheral blood analyses, warrant further research into aPD-1/aCTLA-4 combination in MPM with in-depth peripheral blood and intratumoral T-cell characterization.

Introduction

Malignant pleural mesothelioma (MPM) is a malignancy arising from the mesothelial cells in the pleural cavity, primarily caused by asbestos exposure. Treatment options for MPM are very limited, as platinum-based chemotherapy combined with an antifolate and the optional addition of bevacizumab, are the only approved first-line treatment for MPM. This treatment leads to a median overall survival OS) of 12 -16 months (1,2). Currently, no registered second-line treatments are available, illustrating the urgent need for new treatment options.

Immunotherapies aim for activation of the immune system, leading to efficient tumorspecific immune responses. In current clinical practice, these therapies include monoclonal antibodies that block inhibitory checkpoint receptors, i.e. programmed death 1 (PD-1), programmed death ligand 1 (PD-L1) and cytotoxic T lymphocyte associated antigen 4 (CTLA-4), thereby reinvigorating anti-tumor immune responses (3). So-called immune checkpoint inhibitor (ICI) treatments have transformed the treatment landscape for various malignancies, such as non-small cell lung cancer (NSCLC) and melanoma (4,5).

Unfortunately, ICI treatments are less effective in MPM as compared to other malignancies. The DETERMINE trial showed no survival benefit of ipilimumab (anti-CTLA-4, aCTLA-4) monotherapy over placebo (6) and pembrolizumab and nivolumab, both anti-PD-1 (aPD-1) monotherapies, demonstrated objective response rates (ORR) of 21% and 26% in the KEYNOTE-028 and NivoMes trials respectively (7,8). Recently, the PROMISE-meso phase III randomized trial (NCT02991482) failed to show improvement in PFS (progression-free survival) and OS upon second line aPD-1 treatment (pembrolizumab), as compared to single agent chemotherapy (institutional choice of gencitabine or vinorelbine) (9). The lack of effective ICI treatment in MPM is thought to be dependent on the small number of tumor-infiltrating lymphocytes (TILs) in MPM (10,11) and the immunosuppressive tumor microenvironment (12,13).

Combining aPD-1 and aCTLA-4 therapy has been shown to induce synergistic effects in both preclinical and clinical studies (14,15). Phase II trials in MPM also suggest improved clinical responses upon combination ICI treatment, as the MAPS2 trial (nivolumab plus ipilimumab), the NIBIT-MESO trial (durvalumab (aPD-L1) plus tremelimumab (aCTLA-4)) and the INITIATE trial (nivolumab plus ipilimumab) reported better clinical responses upon combination ICI treatment than reported by trials that investigated monotherapy (nivolumab or pembrolizumab) (16 18). Recently, the first positive results were announced for the Checkmate-743 (19), a phase III trial that combined aPD-1 (nivolumab) with aCTLA-4 (ipilimumab) treatment in previously untreated MPM patients. These results are very promising, although the magnitude of the benefit is still awaited.

Success of aPD-1 treatment in NSCLC and melanoma is thought to depend on pre-existing T-cell infiltration of the tumor (20), proliferation of peripheral PD-1-expressing CD8 T cells (21) and the ratio between T-cell reinvigoration and tumor burden (22). It remains unclear whether the enhanced efficacy observed in ICI combination treatment trials is due to an additive effect of the respective therapies or truly depends on a novel immunological mechanism that is engaged by targeting both PD-1 and CTLA-4 (23).

In order to dissect the immunological mechanisms responsible for the clinical benefit from aPD-1 and aCTLA-4 therapy in MPM, we aimed to investigate the characteristics of lymphocytes present in peripheral blood of MPM patients treated with aPD-1 monotherapy (nivolumab) in the NivoMes trial (8) and aPD-1 and aCTLA-4 combination therapy (nivolumab/ipilimumab) in the INITIATE trial (16). We specifically aimed to evaluate the T- and NK-cell compartment of the peripheral blood, since prior studies established the value of this compartment in the context of aPD-1 and aCTLA-4 treatment (21,22,24).

Methods

Study population

Patients in this study were enrolled in either the NivoMes study (NCT02497508) or the INITIATE study (NCT03048474). Both studies were approved by the institutional review board of the Netherlands Cancer Institute and in accordance with the Declaration of Helsinki. All patients provided written informed consent before enrolment. Collection and analysis of immune cell subsets in peripheral blood were planned a priori as part of the two trials. Clinical results of the NivoMes and INITIATE were previously published (8,16). In summary, in the NivoMes trial, 34 MPM patients progressing after at least one cycle of platinum based chemotherapy, were treated with nivolumab 3 mg/kg every 2 weeks. In the INITIATE trial, 35 MPM patients progressing after at least one cycle of platinum based chemotherapy were treated with nivolumab (240 mg flat dose every 2 weeks) plus ipilimumab (1 mg/kg every 6 weeks up to four times). Peripheral blood was collected from patients on the day of the first ICI treatment and after six weeks of treatment. These samples correspond to the 'screening' and 'on treatment' time points. Response to treatment was assessed according to modified RECIST criteria for mesothelioma (25). For comparison purposes, we decided to define responding patients as having a complete response (CR), partial response (PR) or stable disease (SD) at six months of follow up and non-responding patients as having progressive disease (PD) at six months of follow up. All patients in the 'responder' group experienced a PFS of six months or longer and all patients in in the 'nonresponder' group progressed within six months.

Processing of peripheral blood

Fifty milliliters of blood was drawn at screening and on treatment time points in EDTA tubes and processed. Peripheral blood mononuclear cells (PBMC) were isolated via standard density-gradient centrifugation using Ficoll-Hypaque (GE Healthcare, Chicago, IL, USA). Cells were cryopreserved in 10% dimethylsulfoxide (Sigma-Aldrich, Saint Louis, MO, USA), 40% FCS (Gibco, ThermoFisher, Waltham, MA, USA) and RPMI (Invitrogen, ThermoFisher, Waltham, MA, USA) until further use.

Flow cytometry

Flow cytometry staining was performed on the cryopreserved PBMC samples. After thawing of the PBMCs, cells were stimulated for 4 hours with phorbol 12-myristate 13-acetate and ionomycin (both from Sigma-Aldrich, Saint Louis, MO, USA) and GolgiStop (BD Biosciences, Franklin Lakes, NJ, USA), prior to continuation of the cytokine staining. Supplementary table 1 lists the antibodies used for the different stainings. First, extracellular markers were stained for 30 min at 4 °C. Secondly, the cells were stained with LIVE/DEAD Fixable Aqua Dead Cell Stain Kit (Invitrogen, ThermoFisher, Waltham, MA, USA) for 10 min at 4 °C in order to identify dead cells. Next, FoxP3 transcription factor fixation/permeabilisation mix (eBioscience, ThermoFisher, Waltham, MA, USA) was used to fixate the cells. Subsequently, intracellular markers were stained for 60 min at 4 °C. Data were acquired using an LSR II flow cytometer equipped with three lasers. We used FlowJo v10 (BD Biosciences, Franklin Lakes, NJ, USA) to analyze the data. Fig. 1A, C, D, F and H show the gating strategy. Specific maturation subsets of T cells were identified by the cell surface markers CD45RA and CCR7. Fractions of CD45RA+CCR7+ naive (N) T cells, CD45RA CCR7+ central memory (CM) T cells, CD45RA CCR7 effector memory (EM) T cells and CD45RA+CCR7 effector memory re-expressing RA (EMRA) T cells were identified in both the CD4 and CD8 T-cell compartments.

Statistical analysis

Statistical analyses were performed in R version 4.0.2 and GraphPad V8.0 (GraphPad, San Diego, CA, USA). P < 0.05 was considered statistically significant. Significant differences between the groups were determined with Mann Whitney U tests (non-parametric, non-paired data) and Wilcoxon signed rank tests (non-parametric, paired data). P values were corrected for multiple testing, using the Benjamini and Hochberg False Discovery Rate (26). Log rank test was used to compare Kaplan-Meier curves for PFS and OS. To stratify PFS and OS for proportions of T-cell subsets, the median was used as a cut off for high vs low proportions.

Role of funding sources

Bristol-Myers Squibb sponsored the clinical studies and provided the study drugs in both the NivoMes and INITIATE clinical trials. The analyses of peripheral blood mononuclear

cells (PBMCs) by flow cytometry, described in this manuscript, were not sponsored by any external funding.

		Nivomes	Initiate
Patients screened	38	38	
Included, received at least	34	35	
At least 1 CT for response	33	34	
At least 1 PBMC sample for	31	38	
treatment time point			
PBMC sample at screening	24	38	
PBMC sample at screening	23	32	
evaluation available			
Baseline characteristics			
Ν	23	32	
Age (years) (range)		67 (62 - 73)	65 (62 - 72)
Gender (%)	Male	19 (82.6%)	24 (75%)
	Female	4 (17.4%)	8 (25%)
Histological subtype (%)	Epithelioid	21 (91.3%)	28 (87.5%)
	Sarcomatoid	2 (8.7%)	2 (6.2%)
	Mixed	0 (0%)	2 (6.2%)
WHO (%)	0	10 (43.5%)	11 (34.4%)
	1-2	13 (56.5%)	21 (65.6%)
6 months response (%)	CR	0 (0%)	0 (0%)
	PR	6 (26.1%)	12 (37.5%)
	Epithelioid	6 (100%)	11 (91.7%)
	Sarcomatoid	0 (0%)	1 (8.3%)
	Mixed	0 (0%)	0 (0%)
	SD	1 (4.3%)	4 (12.5%)
	Epithelioid	1 (100%)	2 (50%)
	Sarcomatoid	0 (0%)	1 (25%)
	Mixed	0 (0%)	1 (25%)
	PD	16 (69.6%)	16 (50%)
	Epithelioid	14 (87.5%)	15 (93.8%)
	Sarcomatoid	2 (12.5%)	0 (0%)
	Mixed	0 (0%)	1 (6.2%)
PFS (months) (95% CI)		2.44 (1.3 - 10.0)	6.25 (4.1 - 11.0)
OS (months) (95% CI)		11.5 (5.1 - 21.6)	23.0 (12.5-not
			reached)

Table 1 Characteristics of patients included in translational analysis.

Results

Patient characteristics

Table 1 demonstrates the numbers of peripheral blood samples available from the two clinical trials. Baseline characteristics are shown for the patients of whom PBMCs were collected at screening and at least 1 CT-scan for response evaluation was available.

Monotherapy with aPD-1 treatment does not induce T-cell proliferation

In both NSCLC and melanoma, it was shown that aPD-1 treatment increased proliferation of CD8 T cells in peripheral blood, and the majority of these proliferating CD8 T cells were PD-1 positive (21,22). We therefore analyzed whether aPD-1 monotherapy induced similar changes in T- or NK cell subsets of MPM patients. No significant differences were observed in the frequencies of T cells (Fig. 1B), T-cell subsets (Fig. 1E, G, I), NK cells and NK T cells (Fig. 1B) between screening and 6 weeks after start of treatment. Surprisingly, aPD-1 monotherapy also induced no increase in proliferation of T-cell subsets, as assessed by Ki-67 expression, a cell cycle marker expressed by cycling or recently divided cells (Fig. 1J-L).





Fig. 1. T- and NK-cell characteristics before and during aPD-1 monotherapy (a, c, d, f, h).

Gating strategy for NK-cells (a), T-cells (c), CD4 T-cells subsets (d), CD8 T-cells subsets (f) and Treg subsets (h) respectively. (b, e, g, i) Percentage of T-and NK-cell subsets (b), CD4 T-cell subsets (e), CD8 T-cells subsets (g) and Treg subsets (l) respectively, at screening and on-treatment time points. (j, k, l) Percentage of Ki67+ CD 4 T-cell subsets (j), Tregs subsets (k) and CD8 T-cell subsets (l) respectively, at screening and on-treatment time points. (m, n, o). Paired samples are shown connected by black lines. Percentage of CD4 T-cell subsets (m), Treg subsets (n) and CD8 T-cell subsets (o) respectively, at the screening time point in responding and non-responding patients. Bars depict mean values with standard error of the mean.

Next, we examined whether differences in the frequencies and phenotype of T cells prior to treatment, could help identify patients that responded to aPD-1 monotherapy. We found that MPM patients with a response upon aPD-1 had slightly higher frequency of CM CD4 T cells, whereas all other T-cell frequencies were similar between responding and non-responding MPM patients (Fig. 1M - O). No changes were found in the proportions of proliferating T- and NK cells, assessed by Ki67 expression (data not shown).

Furthermore, no changes in the frequencies of PD-1, CD28, 4-1BB, HLA-DR, inducible T-cell costimulator (ICOS), CD39, lymphocyte-activation gene 3 (LAG-3), T-cell immunoglobulin and mucin-domain containing-3 (TIM-3) and CTLA-4 expressing T-cell subsets induced by aPD-1 treatment or between responding and non-responding patients were observed (data not shown).

In conclusion, aPD-1 treatment did not induce changes in the proportion and proliferation of T-cell and NK cell subsets in MPM patients. No major differences were found between responding and non-responding patients prior to treatment.

aPD-1 and aCTLA-4 combination therapy promotes proliferation of memory T-cell subsets

Secondly, we examined whether aPD-1 and aCTLA-4 combination treatment induced proliferation and activation of T cells. We found that combination treatment increased the proliferation of CM, EM and EMRA CD4 T-cells and in naive and CM CD8 T-cells (Fig. 2E - G). This increase in proliferation was independent of clinical response (Fig. 2 H - J). Furthermore, the frequency of CM, EM and EMRA CD4 T-cell subsets, and CM and EM CD8 T cells that expressed ICOS increased upon combination therapy, indicating that combination therapy induced T-cell activation (Fig. 3A - C). In the CD4 T-cell compartment, this activation was most prominent in non-responding patients (Fig. 3D). Combination treatment did not induce differences in the frequency of the activation and inhibitory markers CD28, 4-1BB, HLA-DR, PD-1, LAG-3, TIM-3, CD39 and CTLA-4 in both CD4 and CD8 T-cell subsets (data not shown).

In conclusion, combining aPD-1 and aCTLA-4 treatment induced proliferation and activation of memory T-cell subsets, however, this proliferation was independent of clinical response.





Fig. 2. T- and NK-cell characteristics before and during aPD-1/CTLA-4 combination therapy(a, b, c, d) Percentage of T-and NK-cell subsets (a), CD4 T-cell subsets (b), Treg subsets (c) and CD8 T-cells subsets (d) respectively, at screening and on-treatment time points. (e, f, g) Percentage of Ki67+ CD 4 T-cell subsets, (TCM p = 0.003, TEM p = 0.007, TEMRA p = 0.028) (e), Tregs subsets (f) and CD8 T-cell subsets (TN p = 0.036, TCM p = 0.03,) (g) respectively, at screening and on-treatment time points. (h, i, j) Comparison between responding (R) and non-responding (NR) patients for the percentage of Ki67+ CD 4 T-cell subsets (TCM R p = 0.01, TCM NR p = 0.04, TEM R p = 0.01) (h), Tregs subsets (i) and CD8 T-cell subsets (j) respectively, at screening and on-treatment time points. Paired samples are shown connected by black lines in each graph. Significance (Wilcoxon signed-rank test for paired analysis of screening and on-treatment samples and Mann-Whitney U test for comparison of response groups) is shown in each graph, with * p < 0.05 and ** p < 0.01. P values were corrected for multiple testing, using the Benjamini and Hochberg False Discovery Rate.



Fig. 3. Percentage of ICOS+ T cell subsets before and during aPD-1/CTLA-4 combination therapy (a, b, c) Percentage of ICOS+ CD 4 T-cell subsets (TCM p = 0.002, TEM p = 0.003, TEMRA p = 0.004) (a), Tregs subsets (b) and CD8 T-cell subsets (TCM p = 0.003, TEM p = 0.012) (c) respectively, at screening and on-treatment time points. (d, e, f) Comparison between responding (R) and non-responding (NR) patients for the percentage of ICOS+ CD 4 T-cell subsets (TN NR p = 0.01, TCM NR p = 0.02, TEM NR p = 0.03, TEMRA NR p = 0.01 (d), Tregs subsets (nTreg NR p = 0.01) (e) and CD8 T-cell subsets (TCM R p = 0.03) (f) respectively, at screening and on-treatment time points. Paired samples are shown connected by black lines in each graph. Significance (Wilcoxon signed-rank test) is shown in each graph, with *p < 0.05 and ** p < 0.01. P values were corrected for multiple testing, using the Benjamini and Hochberg False Discovery Rate

MPM patients responding to combined aPD-1 and aCTLA-4 treatment showed an altered distribution of CD8 T-cell subsets prior to treatment

We investigated whether the frequency or phenotype of T-cell subsets was different prior to treatment in patients that responded, compared to patients that did not respond to aPD-1 and aCTLA-4 combination treatment. MPM patients that responded had a different distribution of their T-cell compartment prior to treatment, with significantly lower

frequencies of naive and CM CD8 T cells and a higher frequency of EMRA CD8 T cells (Fig. 4A - C). Log rank test revealed that patients with a high EMRA CD8 T-cell proportion (cutoff based on the median proportion) at screening, had a significantly longer PFS upon combination treatment (median PFS of 13.1 vs 3.5 months, p = 0.045). Although the OS curves also appeared to differ (median OS of 25.9 vs 10.2 months), this difference was not statistically significant (Fig. 4D and E). Upon further characterization of these EMRA CD8 T cells, we found that the frequency of Granzyme-B and IFNg-expressing EMRA CD8 T cells was increased in responding patients (Fig. 5A and B). Increased cytokine expression was also observed in CM CD8 T cells and EM CD8 T cells (Fig. 5A and B). High or low proportion of Granzyme-B positive EMRA CD8 T cells (cut-off based on the median proportion) prior to treatment was used to stratify PFS and OS. Median PFS was 10.8 months vs 3.5 months for the high vs low groups and median OS was 32.6 vs 10.2 months. Log rank test did not reveal any significant differences between the two curves for both PFS and OS, although a clear trend was seen in the OS curves.



Fig. 4. Comparison of T-cell characteristics before aPD-1/CTLA-4 combination therapy in responding and non-responding patients (a, b, c) Percentage of CD4 T-cell subsets (a), Treg subsets (b) and CD8 T-cell subsets (TN p = 0.017, TCM p = 0.008, TEMRA p = 0.028) (c) respectively, at the screening time point in responding and non-responding patients. Bars depict mean values with standard error of the mean. Significance (MannWhitney U test) is shown in each graph, with * p < 0.05 and ** p < 0.01. P values were corrected for multiple testing, using the Benjamini and Hochberg False Discovery Rate. (d, e) EMRA CD8 T-cells proportions prior to treatment were used to stratify progression-free survival (PFS) (d) and overall survival (OS) (e). Median proportion of EMRA CD8 T cells was used as a cut off between the 'high' vs 'low' group. Statistical significance of the difference between the two

KaplanMeier curves was tested by log rank test with p = 0.045 for PFS (median PFS of 3.5 vs 13.1 months) and p = 0.086 for OS (median OS of 10.2 vs 25.9 months).

In conclusion, patients that responded to combined treatment with aPD-1 and aCTLA-4 had a different T-cell distribution, in particular more EMRA CD8 T cells and less naive CD8 T cells, prior to treatment. The frequency of cytokine-expressing memory CD8 T cells was increased in responding patients, indicating that these memory CD8 T cells are more functionally active.



Fig. 5. Comparison of cytokine frequencies in CD8 T-cell subsets before aPD-1/CTLA-4 combination therapy in responding and non-responding patients (a, b) Percentage of IFNg+ CD8 T-cell subsets (TEM p = 0.008, TEMRA p = 0.006) (a) and Granzyme-B+ CD8 T-cell subsets (TN p = 0.02, TCM p = 0.032, TEMRA p = 0.02) (b) respectively, at the screening time point in responding and non-responding patients. Bars depict mean values with standard error of the mean. Significance (Mann-Whitney U test) is shown in each graph, with * p < 0.05 and ** p < 0.01. P values were corrected for multiple testing, using the Benjamini and Hochberg False Discovery Rate. (c, d) Proportions of Granzyme-B+ EMRA CD8 T-cells prior to treatment were used to stratify progression-free survival (PFS) (d) and overall survival (OS) (e). Median proportion of Granzyme-B+ EMRA CD 8 T cells was used as a cut off between the 'high' vs 'low' group. Statistical significance of the difference between the two KaplanMeier curves was tested by log rank test with p = 0.14 for PFS (not significant, median PFS of 3.5 vs 10.8 months) and p = 0.051 for OS (not significant, median OS of 10.2 vs 32.6 months).

Discussion

Recently, the first positive results were announced for the Checkmate-743 trial, demonstrating that combining aPD-1 and aCTLA-4 therapy led to improved OS in MPM, as compared to chemotherapy (19). In contrast, aPD-1 monotherapy failed to improve PFS and OS (9). Understanding the immunological mechanisms explaining why combination therapy of aPD-1 and aCTLA-4 is effective and monotherapy is not, is thus vital to select effective treatment options for MPM. To the best of our knowledge, we are the first to investigate T-cell characteristics of MPM patients treated with either aPD-1 monotherapy or aPD-1/aCTLA-4 combination therapy, treated during two ICI trials (8,16).

Using comprehensive immune monitoring, we demonstrate that combining aPD-1 with aCTLA-4 treatment strongly induces memory T-cell proliferation and activation of both CD4 and CD8 T cells. Higher frequencies of ICOS-expressing CD4 T cells were only observed in the combination therapy. Since this proliferation and activation was irrespective of clinical response, these results could indicate that aPD-1/aCTLA-4 treatment induces proliferation and activation of bystander, non-tumor specific T cells, which lack the ability to respond to tumor antigens and do not result in a successful anti-tumor immune response. However, the distribution of T-cell subsets prior to treatment was different in MPM patients with a clinical response to combined aPD-1 and aCTLA-4 treatment. Herein, we found increased frequencies of EMRA CD8 T cells (TEMRAs) at the cost of naive CD8 T cells. Survival analysis also showed that PFS was significantly longer in patients with high frequencies of TEMRAs prior to treatment. Furthermore, in responding patients, we found higher frequencies of TEMRAs expressing Granzyme-B and IFNg. Thus, combined aPD-1/aCTLA-4 treatment was associated with the activation and proliferation of memory T cells, but only MPM patients with high frequencies of TEMRAs prior to start of treatment, did benefit. The beneficial presence of TEMRAs could indicate that TEMRAs in particular comprise tumor-specific memory T cells that can be reinvigorated by combination treatment, but not by aPD-1 monotherapy, as these associations were not found in the aPD-1 monotherapy study. Our results are supported by several studies investigating memory CD8 T-cell biology, both in general and in relation to ICI treatment. Characterization of TILs in melanoma patients treated with combined aPD-1/ aCTLA-4 therapy revealed that tumors of responding patients harbored an effector memory T-cell population (CD8+ EOMES+CD69+CD45RO+) that was less abundant in non-responding patients (27). Wei et al. revealed that dual blockade of PD-1 and CTLA-4 engages biological pathways partly different from aPD-1 monotherapy (28). Combined aCTLA-1/ aPD-1 treatment increased the frequencies of a terminally differentiated TBET+EOMES+ CD8 T-cell subset in peripheral blood of melanoma patients, whereas aPD-1 monotherapy did not. Therefore, the authors speculated that combination therapy may be sufficient to attenuate or even reverse T-cell exhaustion. Both studies demonstrated that the combination of aPD-1/aCTLA-4 has a

distinct effect on borderline terminally differentiated memory T-cells, which was not observed upon aPD-1 monotherapy.

Our findings indicate that combination ICI treatment, in contrast to aPD-1 monotherapy, is able to reactivate these crucial TEMRA cells. Further research should provide mechanistic insight in how combined aPD-1 and aCTLA-4 treatment reactivates TEMRAs and should indicate their specificity.

In contrast to the observations of others in NSCLC and melanoma patients, we did not observe increases in T-cell proliferation upon aPD-1 monotherapy in MPM patients. These studies reported that the increase in proliferation peaked 3 weeks after start of treatment, and declined afterwards (21,22). As we evaluated immunological differences 6 weeks after start of treatment, we were most likely too late to assess the effects of aPD-1 monotherapy. However, these differences could also be dependent on tumor type, as aPD-1 therapy depends on pre-existing tumor-specific PD-1-expressing cells, which could be more frequent in NSCLC and melanoma as compared to MPM. Moreover, it has been described earlier that aPD-1 and aCTLA-4 therapy induced longer lasting transcriptional alterations as compared to aPD-1 monotherapy (29), potentially enabling us to detect changes in T-cell characteristics in combination ICI treatment in peripheral blood at a later point in time.

It is important to highlight that the immunological differences found in the two treatment modalities, although they clearly seem to fit response observations, could still be of a phenomenological nature. Thus, our results do not warrant any general conclusions on differences in ICI monotherapy and combination therapy in tumor types other than MPM. Given the limited number of patients analyzed in these studies and the limited number of responding patients, especially in the aPD-1 monotherapy study, our findings need to be validated in a larger and independent MPM patient cohort. Investigating the immunological changes induced by ICI treatment on multiple time points after start of treatment will also provide insight into the duration of these immunological changes upon different ICI treatments, and enable the comparison between MPM and other malignancies. Furthermore, it is not known whether changes in peripheral T-cell subsets reflect changes in the tumor microenvironment (TME) in MPM, and whether tumor specific T cells migrated from the peripheral blood into the TME or vice versa. We are also aware of the fact that nivolumab was administered in a weight dependent dose of 3 mg/kg every 3 weeks in NivoMes, thus modestly differing from the fixed dose of 240 mg/kg every 3 weeks that was administered in INITIATE. However, since Selby et al. (15) demonstrated that no significant alterations in lymphocyte subsets were seen upon different dosing regimens of nivolumab in macagues, we believe that the immune cell alterations described in this manuscript are most likely not caused by dosing differences. At last, it is important to keep in mind that the presumed similarity between pembrolizumab and nivolumab is subject to an ongoing debate in MPM, especially since several studies in non-Caucasian populations demonstrated ORRs to nivolumab that appear to be higher than what was seen in studies performed in Europe and the United States (30,31).

In conclusion, the combined treatment of aPD-1 and aCTLA-4 induced a robust T-cell proliferation and activation in MPM patients, whereas aPD-1 monotherapy did not. The absence of a correlation to clinical response could indicate that these are bystander T-cells. unable to react to tumorantigens. High proportions of TEMRAs that expressed cytokines, prior to treatment, were associated with a better clinical outcome to combination therapy. likely because TEMRAs comprise tumor-specific T cells. This also suggests that TEMRAs can only be reactivated upon combined blockade of both aPD-1 and aCTLA-4. These findings have important implications for future clinical trial design. First, it provides an explanation for the discouraging results of aPD-1 mono-therapy in MPM, since aPD-1 monotherapy appears unable to reinvigorate tumor-specific terminally differentiated memory CD8 T cells in MPM. Second, it grants directions for future research, since aPD-1/aCTLA-4 appears to be a promising treatment modality for MPM, especially now that we are able to select patients up front that are likely to respond. And, finally, it provides a rationale for studying the efficacy of combining these treatments with vaccination strategies like dendritic cell vaccines in non-responding patients, since these vaccines have been shown to induce tumor specific T cells (32).

References

- 1. Vogelzang NJ, Rusthoven JJ, Symanowski J, Denham C, Kaukel E, Ruffie P, et al. Phase III study of pemetrexed in combination with cisplatin versus cisplatin alone in patients with malignant pleural mesothelioma. J Clin Oncol 2003;21:2636-44.
- 2. Zalcman G, Mazieres J, Margery J, Greillier L, Audigier-Valette C, Moro-Sibilot D, et al. Bevacizumab for newly diagnosed pleural mesothelioma in the Mesothelioma Avastin Cisplatin Pemetrexed Study (MAPS): a randomised, controlled, open-label, phase 3 trial. Lancet 2016;387:1405-14.
- 3. Ribas A, Wolchok JD. Cancer immunotherapy using checkpoint blockade. Science 2018.
- 4. Herbst RS, Morgensztern D, Boshoff C. The biology and management of non-small cell lung cancer. Nature 2018;553:7689. 446.
- Michielin O, Van Akkooi ACJ, Ascierto PA, Dummer R, Keilholz U. Cutaneous melanoma: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. Ann Oncol 2019;30:1884-901.
- Maio M, Scherpereel A, Calabro L, Aerts J, Perez SC, Bearz A, et al. Tremelimumab as secondline or third-line treatment in relapsed malignant mesothelioma (DETERMINE): a multicentre, international, randomised, double-blind, placebo-controlled phase 2b trial. Lancet Oncol 2017;18:1261-73.
- 7. Alley EW, Lopez J, Santoro A, Morosky A, Saraf S, Piperdi B, et al. Clinical safety and activity of pembrolizumab in patients with malignant pleural mesothelioma (KEYNOTE-028): preliminary results from a non-randomised, open-label, phase 1b trial. Lancet Oncol 2017;18:623-30.
- Quispel-Janssen J, van der Noort V, de Vries JF, Zimmerman M, Lalezari F, Thunnissen E, et al. Programmed death 1 blockade with nivolumab in patients with recurrent malignant pleural mesothelioma. J Thorac Oncol 2018;13:1569-76.
- Popat S, Curioni-Fontecedro A, Polydoropoulou V, Shah R, O'Brien M, Pope A, et al. A multicentre randomized phase III trial comparing pembrolizumab (P) vs single agent chemotherapy (CT) for advanced pre-treated malignant pleural mesothelioma (MPM): results from the European Thoracic Oncology Platform (ETOP 9-15) PROMISE-meso trial. Ann Oncol 2019;30:v931.
- 10. Marcq E, Siozopoulou V, de Waele J, Van Audenaerde J, Zwaenepoel K, Santermans E, et al. Prognostic and predictive aspects of the tumor immune microenvironment and immune checkpoints in malignant pleural mesothelioma. Oncolmmunology 2017;6.
- 11. Coussens LM, Zitvogel L, Palucka AK. Neutralizing tumor-promoting chronic inflammation: a magic bullet? Science 2013;339:286-91.
- 12. Minnema-Luiting J, Vroman H, Aerts J, Cornelissen R. Heterogeneity in immune cell content in malignant pleural mesothelioma. Int J Mol Sci 2018;19.
- 13. Yap TA, Aerts JG, Popat S, Fennell DA. Novel insights into mesothelioma biology and implications for therapy. Nat Rev Cancer 2017;17:475-88.
- 14. Das R, Verma R, Sznol M, Boddupalli CS, Gettinger SN, Kluger H, et al. Combination therapy with anti-CTLA-4 and anti-PD-1 leads to distinct immunologic changes in vivo. J Immunol 2015;194:950-9.

- 15. Selby MJ, Engelhardt JJ, Johnston RJ, Lu LS, Han M, Thudium K, et al. Preclinical development of ipilimumab and nivolumab combination immunotherapy: mouse tumor models, In vitro functional studies, and cynomolgus macaque toxicology. PLoS One 2016;11.
- 16. Disselhorst MJ, Quispel-Janssen J, Lalezari F, Monkhorst K, de Vries JF, van der Noort V, et al. Ipilimumab and nivolumab in the treatment of recurrent malignant pleural mesothelioma (INITIATE): results of a prospective, single-arm, phase 2 trial. Lancet Respir Med 2019;7:260-70.
- 17. Calabro L, Morra A, Giannarelli D, Amato G, D'Incecco A, Covre A, et al. Tremelimumab combined with durvalumab in patients with mesothelioma (NIBIT-MESO-1): an open-label, non-randomised, phase 2 study. Lancet Respir Med 2018;6:451-60.
- Scherpereel A, Mazieres J, Greillier L, Lantuejoul S, Do P, Bylicki O, et al. Nivolumab or nivolumab plus ipilimumab in patients with relapsed malignant pleural mesothelioma (IFCT-1501 MAPS2): a multicentre, open-label, randomised, non-comparative, phase 2 trial. Lancet Oncol 2019;20:239-53.
- 19. Bratman SV, Yang SYC, lafolla MAJ, Liu Z, Hansen AR, Bedard PL, et al. Personalized circulating tumor DNA analysis as a predictive biomarker in solid tumor patients treated with pembrolizumab. Nat Cancer 2020.
- 20. Tumeh PC, Harview CL, Yearley JH, Shintaku IP, Taylor EJ, Robert L, et al. PD-1 blockade induces responses by inhibiting adaptive immune resistance. Nature 2014;515(7528):568-71.
- Kamphorst AO, Pillai RN, Yang S, Nasti TH, Akondy RS, Wieland A, et al. Proliferation of PD-1+ CD8 T cells in peripheral blood after PD-1-targeted therapy in lung cancer patients. Proc Natl Acad Sci 2017.
- 22. Huang AC, Postow MA, Orlowski RJ, Mick R, Bengsch B, Manne S, et al. T-cell invigoration to tumour burden ratio associated with anti-PD-1 response. Nature 2017.
- 23. Fundamental mechanisms of immune checkpoint blockade therapy, (2018).
- 24. Wei SC, Levine JH, Cogdill AP, Zhao Y, Anang N-AAS, Andrews MC, et al. Distinct cellular mechanisms underlie anti-CTLA-4 and anti-PD-1 checkpoint blockade. Cell 2017;170:1120-33 e17.
- 25. Byrne MJ, Nowak AK. Modified RECIST criteria for assessment of response in malignant pleural mesothelioma. Ann Oncol 2004;15:257-60.
- 26. Benjamini Y, Hochberg Y. Controlling the false discovery rate: a practical and powerful approach to multiple testing on JSTOR. J R Stat Soc Ser B 1995;57:289-300.
- 27. Gide TN, Quek C, Menzies AM, Tasker AT, Shang P, Holst J, et al. Distinct immune cell populations define response to anti-PD-1 monotherapy and anti-PD-1/anti-CTLA-4 combined therapy. Cancer Cell 2019;35:238-55 e6.
- 28. Wei SC, Levine JH, Cogdill AP, Zhao Y, Anang N-AAS, Andrews MC, et al. Distinct cellular mechanisms underlie anti-CTLA-4 and anti-PD-1 checkpoint blockade. Cell 2017;170:1120-33 e17.
- 29. Fairfax BP, Taylor CA, Watson RA, Nassiri I, Danielli S, Fang H, et al. Peripheral CD8 (+) T cell characteristics associated with durable responses to immune checkpoint blockade in patients with metastatic melanoma. Nat Med 2020;26(2):193-9.

- 30. Nakamura A, Kondo N, Nakamichi T, Kuroda A, Hashimoto M, Matsumoto S, et al. Initial evaluation of nivolumab in patients with post-operative recurrence of malignant pleural mesothelioma. Jpn J Clin Oncol 2020;50(8):920-5.
- 31. Okada M, Kijima T, Aoe K, Kato T, Fujimoto N, Nakagawa K, et al. Clinical efficacy and safety of nivolumab: results of a multicenter, open-label, single-arm, Japanese phase II study in malignant pleural mesothelioma (MERIT). Clin Cancer Res 2019;25(18):5485-92.
- 32. Garg AD, Coulie PG, Van den Eynde BJ, Agostinis P. Integrating next-generation dendritic cell vaccines into the current cancer immunotherapy landscape. Trends Immunol 2017;38(8):577-93.

Supplementary Data

Antibody	Fluorochrome	Intracellular/ extracellular	Manufacturer	Clone	CAT number				
T/NK cell staining w	/ith costimulatory r	narkers							
Ki67	FITC	Intracellular	Ebioscience	20Raj1	11-5699-42				
FOXP3	PE	Intracellular	Ebioscience	236A/E7	12-4777-42				
CD45RA	PE Texas Red	Extracellular	Ebioscience	MEM-56	MHCD45RA17				
CD28	PE Cy7	Extracellular	Biolegend	CD28.2	302926				
CD137 (4-1BB)	PerCP Cy5.5	Extracellular	BD	4B4-1	309813				
PD-1	APC	Extracellular	Biolegend	EH12.2H7	329907/329908				
CD3	APC Cy7	Extracellular	Thermofisher (Invitrogen)	UCHT1	557832				
CD8	AF700	Extracellular	Biolegend	SK1	344724				
CCR7	BV421	Extracellular	Biolegend	G043H7	353208				
CD56	BV605	Extracellular	BD	NCAM16.2	562780				
ICOS	BV650	Extracellular	BD	DX29	563832				
HLA-DR	BV711	Extracellular	BD	G46-6	563696				
CD4	BV786	Extracellular	BD	SK3	563877				
LIVE/DEAD stain	BV510	Extracellular	Thermofisher (Invitrogen)	-	L34966				
T/NK cell staining with coinhibitory markers									
Ki67	FITC	Intracellular	Ebioscience	20Raj1	11-5699-42				
FOXP3	PE	Intracellular	Ebioscience	236A/E7	12-4777-42				
CD45RA	PE Texas Red	Extracellular	Ebioscience	MEM-56	MHCD45RA17				
LAG-3	PE Cy7	Extracellular	Biolegend	11C3C65	369309				
CTLA-4	PerCP Cy5.5	Extracellular	Thermofisher (Invitrogen)	14D3	14-1529-82				
PD-1	APC	Extracellular	Biolegend	EH12.2H7	329907/329908				
CD3	APC Cy7	Extracellular	Thermofisher (Invitrogen)	UCHT1	557832				
CD8	AF700	Extracellular	Biolegend	SK1	344724				
CCR7	BV421	Extracellular	Biolegend	G043H7	353208				
CD56	BV605	Extracellular	BD	NCAM16.2	562780				
TIM-3	BV650	Extracellular	BD	7D3	565565				
CD39	BV711	Extracellular	BD	TU66	563680				
CD4	BV786	Extracellular	BD	SK3	563877				
LIVE/DEAD stain	BV510	Extracellular	Thermofisher (Invitrogen)	-	L34966				
T/NK cell staining with intracellular cytokine markers									
Granzyme-B	FITC	Intracellular	Biolegend	QA16A02	372206				
FOXP3	PE	Intracellular	Ebioscience	236A/E7	12-4777-42				
CD45RA	PE Texas Red	Extracellular	Ebioscience	MEM-56	MHCD45RA17				
IL-10	PE Cy7	Intracellular	Biolegend	JES3-9D7	501420				
TNFα	PerCP Cy5.5	Intracellular	eBioscience	MAb11	560679				
PD-1	APC	Extracellular	Biolegend	EH12.2H7	329907/329908				
CD3	APC Cy7	Extracellular	Thermofisher (Invitrogen)	UCHT1	557832				
CD8	AF700	Extracellular	Biolegend	SK1	344724				
CCR7	BV412	Extracellular	Biolegend	G043H7	353208				
CD56	BV605	Extracellular	BD	NCAM16.2	562780				
IL-2	BV650	Intracellular	BD	5344.111	563467				
IFNγ	BV711	Intracellular	BD	B27	564039				
CD4	BV786	Extracellular	BD	SK3	563877				
LIVE/DEAD stain	BV510	Extracellular	Thermofisher (Invitrogen)	-	L34966				

Supplementary Table S1. Antibodies used for flow cytometry staining.