



Universiteit  
Leiden  
The Netherlands

## **DNA mismatch repair-dependent DNA damage responses and cancer**

IJsselsteijn, R.; Jansen, J.G.; Wind, N. de

### **Citation**

IJsselsteijn, R., Jansen, J. G., & Wind, N. de. (2020). DNA mismatch repair-dependent DNA damage responses and cancer. *Dna Repair*, 93. doi:10.1016/j.dnarep.2020.102923

Version: Publisher's Version

License: [Creative Commons CC BY 4.0 license](#)

Downloaded from: <https://hdl.handle.net/1887/3184393>

**Note:** To cite this publication please use the final published version (if applicable).



## Graphical Review

## DNA mismatch repair-dependent DNA damage responses and cancer

Robbert Ijsselsteijn, Jacob G. Jansen, Niels de Wind\*

Department of Human Genetics, Leiden University Medical Center, PO Box 9600, 2300 RC Leiden, the Netherlands



## ARTICLE INFO

## Keywords:

DNA mismatch repair  
DNA damage responses  
Mutagenesis  
Colorectal cancer

## ABSTRACT

Canonical DNA mismatch repair (MMR) excises base-base mismatches to increase the fidelity of DNA replication. Thus, loss of MMR leads to increased spontaneous mutagenesis. MMR genes also are involved in the suppression of mutagenic, and the induction of protective, responses to various types of DNA damage. In this review we describe these non-canonical roles of MMR at different lesion types. Loss of non-canonical MMR gene functions may have important ramifications for the prevention, development and treatment of colorectal cancer associated with inherited MMR gene defects in Lynch syndrome. This graphical review pays tribute to Samuel H. Wilson. Sam not only made seminal contributions to understanding base excision repair, particularly with respect to structure-function relationships in DNA polymerase  $\beta$  but also, as Editor of DNA Repair, has maintained a high standard of the journal.

High-fidelity DNA replication is crucial to preserve the genomic integrity of eukaryotic cells and, consequently, is beneficial for organismal health. Although DNA replication by the processive DNA polymerases  $\delta$  and  $\epsilon$  is highly accurate, inadvertent nucleotide misincorporations occasionally do occur. Polymerase selectivity and proofreading are the first lines of defense against such misincorporations [1]. When a misincorporation escapes proofreading, DNA mismatch repair (MMR) provides a last line of defense. Consequently, inactivation of any of the MMR genes causes a spontaneous mutator phenotype [2].

In addition to correcting base-base mismatches, MMR plays multiple roles in responses to a wide spectrum of DNA damage-induced mutagenic insults (Fig. 1), including methylated nucleotides, oxidative DNA lesions, interstrand crosslinks (ICLs) and helix-distorting nucleotide lesions. Helix-distorting DNA lesions include a variety of nucleotide lesion types, including intrastrand crosslinks induced by ultraviolet (UV) light and bulky nucleotide adducts induced by dietary compounds such as the heterocyclic amines 2-Amino-1-methyl-6-phenylimidazo [4,5-*b*]pyridine (PhIP) and the polycyclic aromatic hydrocarbon (PAH) benzo(a)pyrene (BaP) [3,4].


The role of MMR in DNA damage responses has been described most comprehensively for monofunctional methylating agents that, amongst others, methylate the O<sup>6</sup> position of guanine. When unrepaired by O<sup>6</sup>-Methylguanine-DNA methyltransferase (MGMT) (Fig. 2A), O<sup>6</sup>-methylguanine can be replicated by either the replicative polymerases  $\delta$  and  $\epsilon$  or by the translesion synthesis (TLS) polymerase  $\eta$  [5] (Fig. 2B). The methylation of guanine alters its tautomeric state, allowing it to base-

pair with both cytosine and thymine during replication [6]. The MMR machinery excises the thymidine in the daughter strand of O<sup>6</sup>-methylguanine:thymine mismatches, together with a tract of adjacent nucleotides. However, during gap filling, O<sup>6</sup>-methylguanine again may direct misincorporation of thymine, which results in recurrent cycling between excision and error-prone gap filling (Fig. 2C). During the next cell cycle the excision tracts lead to double stranded DNA breaks by replicative run-off [7]. These double stranded DNA breaks induce apoptosis or genomic deletions and rearrangements when mis-repaired (Fig. 2D). Thus, MMR-induced damage responses ultimately prevent the accumulation of G > A transitions and, when the damage load exceeds a threshold, protects the cell from genomic instability by inducing delayed apoptosis.

Cells are under constant assault from oxidative DNA damage, induced by reactive oxygen species (ROS) [8]. Again, guanines seem to be a preferred target, with 8-oxoguanine being a predominant oxidative DNA lesion [9]. Similar to O<sup>6</sup>-methylguanine, replication of this damaged base can lead to a misincorporation, most frequently adenine [10] that can be excised by MMR and MUTYH [11] to reduce the mutagenicity of 8-oxoguanine (Fig. 3A). In addition, MMR-deficient cells retain a higher amount of oxidative lesions in the DNA than MMR-proficient cells [12]. Overexpression of *MTH1*, which removes 8-oxoguanines from the nucleotide pool, reduces both spontaneous and oxidatively induced mutagenesis in MMR-deficient cells. These data indicate that MMR also excises 8-oxoguanine, incorporated from the nucleotide pool opposite adenine during replication. In contrast, removal of the template adenine by MUTYH results in mutagenesis

\* Corresponding author.

E-mail address: [N.de\\_wind@lumc.nl](mailto:N.de_wind@lumc.nl) (N. de Wind).



**Fig. 1.** Replication of damaged DNA.


A: DNA is continuously threatened by multiple sources of DNA damage, including diet (e.g. red meat), radical oxygen species (dysfunctional mitochondria), UV radiation (sunlight) and various (cytostatic) drugs leading to methylation (CH<sub>3</sub>), oxidation (O\*) or helix distorting DNA damage (=), as well as interstrand crosslinks (ICL). B: DNA damages are replicated by either replicative polymerases or TLS polymerases, leading to different types of compound mismatches. C: When mismatches remain in the DNA, mutation fixation occurs during successive rounds of replication. Mismatch repair/post-TLS repair may remove misincorporations, preventing DNA damage-induced mutagenesis.

(Fig. 3B) [13]. Moreover, they suggest that oxidative DNA damage derived from the pool may significantly contribute to the spontaneous mutator phenotype of MMR-deficient cells.

In contrast to methylation and oxidative base damage, helix-distorting nucleotide lesions are poorly instructive and therefore arrest the processive DNA replication machinery. To counteract the toxicity of such lesions the cell employs TLS DNA polymerases that can replicate across the damaged nucleotides, albeit in an error-prone fashion, resulting in DNA damage-directed mutagenesis [14]. Interestingly, MMR can suppress this mutagenesis and increases checkpoint activation and apoptosis induced by UV, PhIP and BaP [(4, 15), Ijsselsteijn et al. manuscript in preparation]. In support, the MMR heterodimer MSH2-MSH6 is recruited to sites of localized UV damage [16] and selectively binds “mismatched” but not “matched” nucleotides opposite photolyses [17]. The mechanistic basis for the involvement of MMR proteins in responses to such DNA lesions is not yet fully elucidated, but it has been suggested that a non-canonical MMR pathway, dubbed “post-TLS repair”, excises the nucleotide that was mis-incorporated by TLS opposite the damaged nucleotide. Thereby, post-TLS repair decreases DNA damage-induced mutagenesis while the resulting lesion-containing single-stranded DNA tracts activate DNA damage signaling (Fig. 4) [15]. Another model to explain the increased mutagenesis by photolyses suggests that MMR plays a role in the recruitment of the relatively error-free TLS polymerase  $\eta$ , via Proliferating Cell Nuclear Antigen

(PCNA) mono-ubiquitination [18,19]. However, knocking out both Polymerase  $\eta$  and *Msh6* in mouse embryonic stem cells shows a synergistic, rather than an epistatic, response to UV radiation in terms of mutagenesis (Ijsselsteijn et al. manuscript in preparation), supporting the post-TLS repair model.


One of the most difficult to repair DNA damages are ICLs that require a multi-step process where multiple repair pathways converge to repair the covalent linking of two opposite DNA strands. ICLs can result from aldehydes, derived from ethanol or from lipid peroxidation, from some vegetables and they are also induced by several chemotherapeutics, such as Cisplatin, used in the treatment of (amongst others) breast cancer. The repair of ICLs involves Fanconi Anemia (FA)-associated proteins, but also requires homologous recombination, TLS and in some cases nucleotide excision repair [20]. It has been suggested that MMR-deficient cells display reduced sensitivity to ICL-inducing agents, indicating a role for MMR in responses to ICLs [21]. ‘Misincorporations’, generated during gap-filling by TLS opposite the unhooked ICL might be a substrate for MMR, comparable to post-TLS repair of UV-induced intra-strand crosslinks [15]. For specific ICLs, base excision repair polymerase  $\beta$  (Pol $\beta$ ) and MMR play an epistatic role. Both the toxicity and repair of ICLs are diminished in cells, knocked down for MSH2, MLH1 or Pol $\beta$ , but is not further reduced in a double knock-down of Pol $\beta$  and MSH2 or MLH1 [21], suggesting that the role of MMR is post-replicative and linked to Pol $\beta$ .



**Fig. 2.** Roles of MMR in responses to O<sup>6</sup>-methylguanine.  
 A: O<sup>6</sup>-methylguanine (CH<sub>3</sub>) can be repaired by O<sup>6</sup>-Methylguanine-DNA methyltransferase (MGMT) leading to damage reversal. B: Replication of O<sup>6</sup>-methylguanine can lead to a O<sup>6</sup>-methylguanine.T mismatch, which can be removed by MMR resulting in a transient single stranded DNA tract. C: Gap filling can either lead to incorporation of a cytosine or to another misincorporation, that again is a substrate for MMR. D: Futile cycles of excision and misincorporation can lead to recurrent ssDNA tracts. In a subsequent round of replication these can lead to double stranded DNA breaks and consequent rearrangements or apoptosis.

Absence of MMR/post-TLS repair causes both a spontaneous mutator phenotype and increased DNA damage-induced mutagenesis, as well as aberrant DNA damage responses. It therefore is not surprising that loss of MMR/post-TLS repair is strongly associated with carcinogenesis [22]. As mentioned before, MMR/post-TLS repair deficiency results in tolerance of various types of DNA damage and thus MMR/post-TLS repair-deficient cancers might benefit from a targeted treatment. For instance, the tolerance of MMR/post-TLS repair-deficient cells to chemotherapeutic drugs that induce O<sup>6</sup>-methylguanines, such as Dacarbazine and Temozolomide, indicates that these agents may be ineffective for the treatment of MMR/post-TLS repair-deficient tumors. Furthermore, exposure of MMR-proficient cells to DNA-damaging drugs

might lead to selection for MMR/post-TLS repair-deficiency. For instance, glioblastomas have been shown to acquire tolerance of Temozolomide by inactivating MMR/post-TLS repair [23]. Another example is provided by acute myeloid leukemia (AML), which can be a late consequence of immunosuppression by Azathioprine after organ transplantation [24]. Its S<sup>6</sup>-(thio)guanine metabolite is methylated endogenously and induces MMR-dependent cytotoxicity, which leads to selection of MMR/post-TLS repair-deficient cells. Also, recurring AML in non-transplant patients is often MMR/post-TLS repair deficient, which might be associated with acquired tolerance to treatment of the primary AML with 6-thioguanine [25,26]. Melanomas frequently display hallmarks of MMR/post-TLS repair deficiency too [27]. Sunlight,



**Fig. 3.** Processing of 8-oxoguanine. Adenine 'mispairs' by MMR and MUTYH.

A: Guanines in the DNA can be oxidized by reactive oxygen species (ROS) to 8-oxoguanines (O\*). Replication can lead to misincorporations of mostly adenines opposite 8-oxoguanines. When not removed by MMR and MUTYH, these misincorporations result in G > T transversions. B: Guanines that are oxidized (O\*) in the free dNTP pool can be incorporated in the DNA opposite adenines. MMR recognizes these mismatches and removes the 8-oxoguanines from the DNA. In contrast, inadvertent repair of the nucleotide opposite the 8-oxoguanine by MUTYH leads to a mutation.


which is the major etiological agent of these tumors, induces not only intrastrand DNA crosslinks but also oxidative DNA damage [28]. As described above, MMR/Post-TLS repair is associated with protective responses to both types of DNA lesions, which might provide a rationale for its loss in melanoma. Finally, MMR is lost in 15 % of the sporadic colorectal cancers (CRC) [29]. The mechanism of action of most chemotherapeutic treatments of CRC is based on the induction of DNA damage, leading to senescence or death of the rapidly cycling tumor cells. For instance, 5-fluorouracil (5-FU) is incorporated in RNA but also causes nucleotide pool imbalances, while its metabolite 5-FdU can be incorporated in DNA instead of thymidine. MMR-deficient cells display a mild tolerance to 5-FU, suggesting a role for MMR in processing misincorporations due to nucleotide pool imbalances as well as misincorporations opposite 5-FdU residues [30]. In support, treatment with 5-FU provides no benefit to MMR-deficient CRC [31].

Apart from the causal involvement of MMR deficiency in sporadic cancer, inheritance of a germline mutation in one allele of any of the four MMR genes is responsible for the common cancer predisposition Lynch syndrome, historically called Hereditary Non-Polyposis Colorectal Cancer (HNPCC). Carriers have an approximate 40–60 % risk of developing gastrointestinal tumors and, for females, approximately 40 % risk of endometrial tumors, depending on which MMR gene is mutated [32]. Inheritance of a bi-allelic germline mutation in one of the MMR genes leads to constitutional MMR deficiency (cMMRd), a childhood cancer syndrome. In contrast to Lynch syndrome, cMMRd involves a wide spectrum of tumors that is dominated by brain, hematological and intestinal cancers [33]. The prevalence of sporadic MMR deficiency in CRC and the restricted cancer tropism in Lynch syndrome, as compared with cMMRd, suggest that roles of MMR/post-

TLS repair in the intestine, beyond the correction of simple base-base mismatches are involved in the suppression of these cancers.

Cancer development requires an array of mutagenic (tumor initiation and progression) and non-mutagenic (tumor promotion and progression) events. The exposure of intestinal epithelial cells to endogenous and food-derived mutagenic compounds is pertinent to all stages of oncogenesis [34], but might be even more so in the development of MMR/post-TLS repair-deficient cancer. For instance, in gastrointestinal stem cells from individuals with Lynch syndrome, these compounds can promote the mutational inactivation of the remaining wild-type allele of the germ-line mutated MMR gene (Fig. 5). The resulting MMR/post-TLS repair-deficient cells may be prone to oncogenic derailment in different ways. For example, exposure to butyrate produced by the intestinal microbiome from carbohydrates, may provide a specific growth advantage to MMR-deficient crypts [35]. Furthermore, exposure of these crypts to DNA methylating agents from the diet or bile [36,37] may promote their expansion to adjacent crypts [38]. Also, the signatures of spontaneous mutations caused by the MMR deficiency and those induced by helix-distorting nucleotide lesions are different [39]. This increases the number of "hittable" oncogenic targets and therefore both mutagenic processes may synergize in cancer development. Furthermore, as described above, MMR/post-TLS repair-deficient cells may be intrinsically hypermutable by agents that induce helix-distorting nucleotide lesions while cell cycle arrests and apoptosis may be impaired [3,15]. These factors might further contribute to CRC development in Lynch syndrome. Moreover, intestinal mutagens such as PhIP might damage mitochondria, leading to the generation of ROS, which is another source of DNA damage [40]. Finally, neoepitopes resulting from the mutator phenotypes caused by loss of canonical MMR





**Fig. 5.** Etiology of cancer of the gastro-intestinal tract in Lynch syndrome.

**A:** All gastro-intestinal crypt cells of Lynch syndrome carriers are heterozygous for an MMR gene (light red) and are continuously exposed to agents that induce methylating and helix-distorting DNA lesions. **B:** Methylating and helix-distorting DNA lesions can induce inadvertent loss of the wild type allele of the germ-line mutated MMR gene (bright red). **C:** Loss of MMR/post-TLS repair leads to hypermutagenesis, which results in the accumulation of oncogenic mutations (dark red). In addition, neopeptides in these cells trigger the recruitment of myeloid immune cells that excrete reactive oxygen species, which may further increase mutagenesis in MMR/post-TLS repair-deficient cells. **D:** MMR/post-TLS repair-deficient cells also are more tolerant of methylated and to helix distorting DNA damage, which leads to their expansion. Ultimately, these selective 'advantages' can result in rapid carcinogenesis.

## Acknowledgements

NdW is supported by grants from the Netherlands Organization for Scientific Research (ALWOP.131), the Dutch Cancer Society (KWF 11358) and the Dutch Cancer Foundation and the Dutch Digestive Foundation (FP16-02).

## References

- Johansson E. Rais, DNA replication—a matter of fidelity, *Mol. Cell* 62 (5) (2016) 745–755.
- Y. Zhang, F. Yuan, S.R. Presnell, K. Tian, Y. Gao, A.E. Tomkinson, et al., Reconstitution of 5'-directed human mismatch repair in a purified system, *Cell* 122 (5) (2005) 693–705.
- S.L. Smith-Roe, D.C. Hegan, P.M. Glazer, A.B. Buermeyer, Mlh1-dependent suppression of specific mutations induced in vivo by the food-borne carcinogen 2-amino-1-methyl-6-phenylimidazo [4,5-b] pyridine (PhIP), *Mutat. Res.* 594 (1–2) (2006) 101–112.
- W.E. Glaab, T.R. Skopek, Cytotoxic and mutagenic response of mismatch repair-defective human cancer cells exposed to a food-associated heterocyclic amine, *Carcinogenesis* 20 (3) (1999) 391–394.
- L. Haracska, S. Prakash, L. Prakash, Replication past O6-methylguanine by yeast and human DNA polymerase  $\eta$ , *Mol. Cell. Biol.* 20 (21) (2000) 8001–8007.
- M.D. Wyatt, D.L. Pittman, Methylating agents and DNA repair responses: methylated bases and sources of strand breaks, *Chem. Res. Toxicol.* 19 (12) (2006) 1580–1594.
- D. Fu, J.A. Calvo, L.D. Samson, Balancing repair and tolerance of DNA damage caused by alkylating agents, *Nat. Rev. Cancer* 12 (2) (2012) 104–120.
- B. Halliwell, Free radicals, antioxidants, and human disease: curiosity, cause, or consequence? *Lancet* 344 (8924) (1994) 721–724.
- S. Kanvah, J. Joseph, G.B. Schuster, R.N. Barnett, C.L. Cleveland, U. Landman, Oxidation of DNA: damage to nucleobases, *Acc. Chem. Res.* 43 (2) (2010) 280–287.
- B. Van Loon, E. Markkanen, U. Hübscher, Oxygen as a friend and enemy: how to combat the mutational potential of 8-oxo-guanine, *DNA Repair* 9 (6) (2010) 604–616.
- A. Mazurek, M. Berardini, R. Fishel, Activation of human MutS homologs by 8-oxo-guanine DNA damage, *J. Biol. Chem.* 277 (10) (2002) 8260–8266.
- T.L. Deweese, J.M. Shipman, N.A. Larrier, N.M. Buckley, L.R. Kidd, J.D. Groopman, et al., Mouse embryonic stem cells carrying one or two defective Msh2 alleles respond abnormally to oxidative stress inflicted by low-level radiation, *Proc. Natl. Acad. Sci. U. S. A.* 95 (20) (1998) 11915–11920.
- M.T. Russo, M.F. Blasi, F. Chiera, P. Fortini, P. Degan, P. Macpherson, et al., The oxidized deoxynucleoside triphosphate pool is a significant contributor to genetic instability in mismatch repair-deficient cells, *Mol. Cell. Biol.* 24 (1) (2004) 465–474.
- B. Pilzecker, O.A. Buoninfante, H. Jacobs, DNA damage tolerance in stem cells, ageing, mutagenesis, disease and cancer therapy, *Nucleic Acids Res.* 47 (14) (2019) 7163–7181.
- A. Tsaalbi-Shtylik, C. Ferras, B. Pauw, G. Hendriks, P. Temviriyankul, L. Carlee, et al., Excision of translesion synthesis errors orchestrates responses to helix-distorting DNA lesions, *J. Cell Biol.* 209 (1) (2015) 33–46.
- Z. Hong, J. Jiang, K. Hashiguchi, M. Hoshi, L. Lan, A. Yasui, Recruitment of mismatch repair proteins to the site of DNA damage in human cells, *J. Cell. Sci.* 121 (19) (2008) 3146–3154.
- H. Wang, C.W. Lawrence, G.M. Li, J.B. Hays, Specific binding of human MSH2.MSH6 mismatch-repair protein heterodimers to DNA incorporating thymine- or uracil-containing UV lig.M.hT photoproducts opposite mismatched bases, *J. Biol. Chem.* 274 (24) (1999) 16894–16900.
- A. Zlatanou, E. Despras, T. Braz-Petta, I. Boubakour-Azzouz, C. Pouvelle, Grant, et al., The hMsh2-hMsh6 complex acts in concert with monoubiquitinated PCNA and pol  $\eta$  in response to oxidative DNA damage in human cells, *Mol. Cell* 43 (4) (2011) 649–662.
- J. Peña-Díaz, S. Bregenhorn, M. Ghodgaonkar, C. Follonier, M. Artola-Borán, D. Castor, et al., Noncanonical mismatch repair as a source of genomic instability in human cells, *Mol. Cell* 47 (5) (2012) 669–680.
- A.J. Deans, S.C. West, DNA interstrand crosslink repair and cancer, *Nat. Rev. Cancer* 11 (7) (2011) 467–480.
- A. Sawant, A. Kothandapani, A. Zhitkovich, R.W. Sobol, S.M. Patrick, Role of mismatch repair proteins in the processing of cisplatin interstrand cross-links, *DNA Repair* 35 (2015) 126–136.
- D. Liu, G. Keijzers, L.J. Rasmussen, DNA mismatch repair and its many roles in eukaryotic cells, *Mutat. Res. Mutat. Res.* 773 (2017) 174–187.
- S. Yip, J. Miao, D.P. Cahill, A.J. Iafrate, K. Aldape, C.L. Nutt, et al., MSH6 mutations

- arise in Glioblastomas during temozolomide therapy and mediate temozolomide resistance, *Clin. Cancer Res.* 15 (14) (2009) 4622–4629.
- [24] J. Offman, G. Opelz, B. Doehler, D. Cummins, O. Halil, N.R. Banner, et al., Defective DNA mismatch repair in acute myeloid leukemia/myelodysplastic syndrome after organ transplantation, *Blood* 104 (3) (2004) 822–828.
- [25] P.N. Munshi, M. Lubin, J.R. Bertino, 6-Thioguanine: a drug with unrealized potential for cancer therapy, *Oncologist* 19 (7) (2014) 760–765.
- [26] G. Mao, F. Yuan, K. Absher, C.D. Jennings, D.S. Howard, C.T. Jordan, et al., Preferential loss of mismatch repair function in refractory and relapsed acute myeloid leukemia: potential contribution to AML progression, *Cell Res.* 18 (2) (2008) 281–289.
- [27] O. Kubeček, J. Kopecký, Microsatellite instability in melanoma, *Melanoma Res.* 26 (6) (2016) 545–550.
- [28] A.P. Schuch, N.C. Moreno, N.J. Schuch, C.F.M. Menck, C.C.M. Garcia, Sunlight damage to cellular DNA: focus on oxidatively generated lesions, *Free Radic. Biol. Med.* (107) (2017) 110–124.
- [29] H. Brenner, M. Kloor, C.P. Pox, Colorectal cancer, *Lancet* 383 (9927) (2014) 1490–1502.
- [30] M. Meyers, M.W. Wagner, A. Mazurek, C. Schmutte, R. Fishel, D.A. Boothman, DNA mismatch repair-dependent response to fluoropyrimidine-generated damage, *J. Biol. Chem.* 280 (7) (2005) 5516–5526.
- [31] D.J. Sargent, S. Marsoni, G. Monges, S.N. Thibodeau, R. Labianca, S.R. Hamilton, et al., Defective mismatch repair as a predictive marker for lack of efficacy of fluorouracil-based adjuvant therapy in colon cancer, *J. Clin. Oncol.* 28 (20) (2010) 3219–3226.
- [32] E. Stoffel, B. Mukherjee, V.M. Raymond, N. Tayob, F. Kastrinos, J. Sparr, et al., Calculation of risk of colorectal and endometrial cancer among patients with lynch syndrome, *Gastroenterology* 137 (5) (2009) 1621–1627.
- [33] N. Lavoine, C. Colas, M. Muleris, S. Bodo, A. Duval, N. Entz-Werle, et al., Constitutional mismatch repair deficiency syndrome: clinical description in a French cohort, *J. Med. Genet.* 52 (11) (2015) 770–778.
- [34] A.J. Cross, R. Sinha, Meat-related mutagens/carcinogens in the etiology of colorectal cancer, *Environ. Mol. Mutagen.* 44 (1) (2004) 44–55.
- [35] A. Belcheva, T. Irrazabal, Susan, C. Streutker, H. Maughan, S. Rubino, et al., Gut microbial metabolism drives transformation of Msh2-deficient colon epithelial cells, *Cell* 158 (2) (2014) 288–299.
- [36] W. Jia, G. Xie, W. Jia, Bile acid-microbiota crosstalk in gastrointestinal inflammation and carcinogenesis, *Nat. Rev. Gastroenterol. Hepatol.* 15 (2) (2018) 111–128.
- [37] M. Cantwell, C. Elliott, Nitrates, nitrites and nitrosamines from processed meat intake and colorectal cancer risk, *J. Clin. Nutr. Diet.* 3 (2017) 4–27.
- [38] K. Wojciechowicz, E. Cantelli, B. Van Gerwen, M. Plug, A. Van Der Wal, E. Delzenne-Goette, et al., Temozolomide increases the number of mismatch repair-deficient intestinal crypts and accelerates tumorigenesis in a mouse model of lynch syndrome, *Gastroenterology* 147 (5) (2014) 1064–1072 e5.
- [39] J.E. Kucab, X. Zou, S. Morganella, M. Joel, A.S. Nanda, E. Nagy, et al., A compendium of mutational signatures of environmental agents, *Cell* 177 (4) (2019) 821–836 e16.
- [40] E.F. Fang, M. Scheibye-Knudsen, K.F. Chua, M.P. Mattson, D.L. Croteau, V.A. Bohr, Nuclear DNA damage signalling to mitochondria in ageing, *Nat. Rev. Mol. Cell Biol.* 17 (5) (2016) 308–321.
- [41] J.C. Dudley, M.T. Lin, D.T. Le, J.R. Eshleman, Microsatellite instability as a biomarker for PD-1 blockade, *Clin. Cancer Res.* 22 (4) (2016) 813–820.
- [42] C.C. Winterbourn, A.J. Kettle, M.B. Hampton, Reactive oxygen species and neutrophil function, *Annu. Rev. Biochem.* 85 (1) (2016) 765–792.