



Universiteit
Leiden
The Netherlands

Insight in the role of lipids and other systemic factors in hand and knee osteoarthritis: lessons from clinical studies

Loef, M.

Citation

Loef, M. (2022, November 15). *Insight in the role of lipids and other systemic factors in hand and knee osteoarthritis: lessons from clinical studies*. Retrieved from <https://hdl.handle.net/1887/3485903>

Version: Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/3485903>

Note: To cite this publication please use the final published version (if applicable).

6

The lipid profile for the prediction of prednisolone treatment response in patients with inflammatory hand osteoarthritis: The HOPE study

Marieke Loef, Tariq Faquih, Joost von Hegedus, Mohan Ghorasaini, Andreea Ioan-Facsinay, Dennis Mook-Kanamori, Féline Kroon, Martin Giera, Margreet Kloppenburg

Osteoarthritis and Cartilage OPEN. 2021



Abstract

Objective To explore the use of lipidomics for prediction of prednisolone treatment response in patients with inflammatory hand osteoarthritis.

Design The Hand Osteoarthritis Prednisolone Efficacy (HOPE) study included patients (n=92) with symptomatic inflammatory hand osteoarthritis, fulfilling the ACR criteria. The present analyses comprised only patients randomized to prednisolone treatment (10 mg daily, n=40). Response to prednisolone treatment was defined according to the OARSI-OMERACT responder criteria at six weeks. Baseline blood samples were obtained non-fasted. Lipid species were quantified in erythrocytes with the Lipidizer™ platform (Sciex). Oxylipins were analysed in plasma using an in-house LC-MS/MS platform. Elastic net regularized regression was used to predict prednisolone treatment response based on common patient characteristics alone and including the patients' lipid profile. ROC analyses with 1,000 bootstrapped area under the curve (AUC) was used to determine the discriminatory accuracy of the models.

Results Among included patients, 78% fulfilled the OARSI-OMERACT responder criteria. From the general patient characteristics, elastic net selected baseline hand function as only predictor of treatment response, with an AUC of 0.78 (0.56; 0.97). Addition of lipidomics resulted in an AUC of 0.92 (0.78; 0.99) and 0.85 (0.65; 0.98) for inclusion of the Lipidizer™ platform and oxylipin platform, respectively.

Conclusion Our results suggest that the patients' lipid profile may improve the discriminative accuracy of the prediction of prednisolone treatment response in patients with inflammatory hand osteoarthritis compared to prediction by commonly measured patient characteristics alone. Hence, lipidomics may be a promising field for biomarker discovery for prediction of anti-inflammatory treatment response.

Introduction

Hand osteoarthritis (OA) is one of the most prevalent OA phenotypes, and it is associated with pain, stiffness, functional impairment and a loss in quality of life [1–4]. Currently, there is a high unmet need for disease modifying drugs for the treatment of osteoarthritis (OA). The role of inflammation in hand OA and its association with pain [5,6] has sparked increasing interest for targeting inflammation in therapeutic research. To this regard, the Hand Osteoarthritis Prednisolone Efficacy (HOPE) study was set up. The HOPE study is a blinded, randomized placebo-controlled trial, that investigated the effect of prednisolone treatment in patients with painful, inflammatory hand OA. The HOPE study showed a clinically relevant decrease in pain in patients using prednisolone [7]. Since pharmacological treatments usually show marked variation in treatment response, it is important to carefully select patients who will most likely benefit from treatment, to maximize the desired therapeutic effect, and minimize overtreatment and potential adverse effects. Metabolomics may aid the identification of biomarkers of therapeutic responsiveness [8].

Lipids are essential for joint physiology [9,10]. However, to maintain normal physiology, a tight control of lipid species is warranted. In addition, various lipids and their metabolites are involved in pathophysiological settings, in particular in inflammation. Moreover, they have been shown to play an important role in inflammation in auto-immune diseases [11], as well as in OA [12,13]. Therefore, lipidomics, involving the identification and quantification of lipid metabolites, may be particularly relevant as biomarker of therapeutic responsiveness to anti-inflammatory medication. In addition, previous lipid profiling studies have suggested an altered lipid metabolism in patients with OA [14–16]. In particular, associations between differing levels of phospholipids and OA have been observed [16–18]. Hence, the patients' lipid profile may be predictive of response to anti-inflammatory treatment in patients with inflammatory hand OA. To our knowledge, the use of lipidomics for prediction of treatment response in patients with OA has not previously been studied.

Therefore, we explored the patients' lipid profile for the prediction of prednisolone treatment response in patients with inflammatory hand OA.

Methods

Study design

The HOPE study included patients with symptomatic hand OA, fulfilling the American College of Rheumatology criteria [19] and presenting signs of inflammation in the distal and proximal interphalangeal (DIP/PIP) joints. Full description of patient inclusion and procedures can be found elsewhere [7]. Briefly, patients were required to have: finger pain of ≥ 30 mm on a 100 mm visual analogue scale (VAS) and flaring upon 48-hour NSAID washout (defined as ≥ 20 mm worsening), ≥ 4 DIP/PIP joints with osteoarthritic nodes, ≥ 1 DIP/PIP joints with soft swelling or erythema, and ≥ 1 DIP/PIP joints with positive power Doppler signal or synovitis grade ≥ 2 on ultrasound. Patients were excluded from participation in case of chronic inflammatory rheumatic diseases, psoriasis, uncontrolled serious comorbidities, malignancy, infectious disease, and immune modulating drug use within 90 days before baseline. Patients ($n = 92$) were randomly assigned (1:1) to receive 10 mg prednisolone daily, or placebo, for six weeks. The present study comprised of patients randomized to prednisolone treatment only ($n = 40$). Treatment adherence has been reported previously [7]. The HOPE study (Netherlands Trial

Registry (NTR5263)) was approved by the local medical ethics committees and conducted in accordance with Good Clinical Practice guidelines and Declaration of Helsinki. All patients provided written informed consent.

Patient reported outcomes

At baseline and week six, patients completed a VAS for finger pain and VAS global assessment on a 0-100 mm scale, and the Australian/Canadian Hand Osteoarthritis Index (AUSCAN) pain (scored as 0-20) and function (scored as 0-36) subscales (higher scores are worse). At week six, fulfilment of the OMERACT-OARSI responder criteria was assessed, which was defined as a relative improvement $\geq 50\%$ and absolute change $\geq 20/100$ in AUSCAN pain or function, or a relative improvement $\geq 20\%$ and absolute change $\geq 10/100$ in ≥ 2 of the following: AUSCAN pain, AUSCAN function or VAS patient global assessment [20]. In the OMERACT-OARSI criteria, the AUSCAN pain and function subscale scores are used on a 0-100 scale. The AUSCAN pain and function subscale scores were rescaled from 0-20 and 0-36, respectively, to 0-100. We calculated absolute change as the baseline score minus the follow-up score, and relative change as the absolute change divided by the baseline score.

Baseline imaging

All interphalangeal and metacarpophalangeal joints were assessed on baseline radiographs of both hands (30 joints). Radiographic OA severity was investigated using the Kellgren and Lawrence (KL) grading system on a 0-4 scale [21]. Erosive OA was defined as having ≥ 1 joint in the erosive or remodelling phase according to the Verbruggen-Veys score [22]. Synovial thickening was assessed on ultrasound on a 0-3 scale [6]. A sum score adding the scores of all investigated joints was calculated for KL (0-120) and synovitis (0-90). The reliability of all scoring methods was good [7].

Lipidomics measurements

Blood samples were obtained non-fasted at baseline at various time points during the day in EDTA-tubes, following a standardized protocol. The blood samples were centrifuged for 10 minutes at $2200 \times g$ to separate plasma from the cellular fraction. Erythrocytes were isolated by ficoll density gradient centrifugation and washed 3x with PBS. Plasma samples were quenched using $600\mu\text{L}$ MeOH (Honeywell, 349661L), and $8\mu\text{L}$ IS was added (containing: 500pg/mL PGE2-d4, 5ng/mL DHA-d5, 500pg/mL LTB4-d4 and 500pg/mL 15S-HETE-d8). Samples were stored at -80°C topped with argon until further analyses [23].

The Lipidyzer™ platform (Sciex) was used to quantify total lipid content in erythrocytes (nmol/mL). Lipid extraction was performed using methyl-tert-butylether as described by Matyash et al., with some modifications [24]. To $25\mu\text{L}$ of erythrocyte sample the following was added: $160\mu\text{L}$ MeOH, $50\mu\text{L}$ internal standard solution (Lipidyzer™ internal standard kit, containing >50 labeled internal standards for 13 lipid classes), and $550\mu\text{L}$ methyl-tert-butylether. Samples were vortexed and left at room temperature for 30 minutes. Subsequently, $200\mu\text{L}$ water was added for phase separation and the samples were centrifuged at $13.100 \times g$. The upper layer was transferred to a glass vial and lipid extraction was repeated by adding $300\mu\text{L}$ methyl-tert-butylether, $100\mu\text{L}$ MeOH and $100\mu\text{L}$ water. The organic extracts were combined and dried under a gentle stream of nitrogen. Lipidyzer running buffer ($250\mu\text{L}$) was added and samples were transferred to a glass vial with insert for injection. Briefly, the Lipidyzer platform is a flow-injection-based ion-mobility triple quadrupole system consisting of a Sciex 5500 QTrap equipped with Selexion technology coupled to a Shimadzu Nexera series UHPLC

system used for injection and delivering running buffer at 7 $\mu\text{L}/\text{min}$. Two methods were used for the injection of a total of 50 μL of the resuspended samples. First, PC, PE, (L)PC, (L)PE, and SM lipid classes were analyzed using method 1, operating with active DMS separation under the following conditions: DMS temperature low, modifier (propanol) composition low, separation voltage 3500 V, DMS resolution enhancement low. Next, FFA, TAG, DAG, CER, dihydroceramide (DCER), lactosylceramide (LCER), hexosylceramide (HCER), and CE lipids were analyzed applying method 2, for which the DMS cell was not activated. The MS operated under the following conditions: curtain gas 17, CAD gas medium, ion spray voltage 4100 V in ESI+ mode and -2500V in ESI- mode, temperature $200\text{ }^\circ\text{C}$, nebulizing gas 17, and heater gas 25. Further technical detail can be found elsewhere [25–27]. Lipid concentrations were corrected for the erythrocyte protein pellet content, which was quantified using a Micro BCA Protein Assay Kit (Thermo Scientific, Waltham, MA, USA). Samples were measured in a randomized batch controlled fashion. The lipid concentrations were corrected for the erythrocyte protein pellet content. After preprocessing of the Lipidizer™ data (Supplementary file, figure S1), 286 lipid species were available for further analyses (Supplementary file, table S1).

Oxylipins were measured in plasma, using liquid-chromatography combined with mass spectrometry (LC-MS/MS) analysis in negative electrospray ionization mode as described previously [28]. A QTrap 6500 mass spectrometer in negative ESI mode (Sciex, Nieuwerkerk aan den IJssel, The Netherlands) was used, coupled to a LC system employing LC-30AD pumps, a SIL-30AC auto sampler, and a CTO-20AC column oven (Shimadzu, 's-Hertogenbosch, The Netherlands). A Kinetex C18 $50 \times 2.1\text{ mm}$, $1.7\text{ }\mu\text{m}$ column, combined with a C8 pre column (Phenomenex, Utrecht, The Netherlands) was used, kept at $50\text{ }^\circ\text{C}$. A gradient of water and Methanol with 0.01% acetic acid was used. An injection volume of 40 μL was used, with a flow rate of 400 $\mu\text{L}/\text{min}$ (6). Oxylipins were identified using characteristic mass transitions and relative retention times. Only peaks with a signal to noise > 10 were included, resulting in identification of 25 oxylipins. For a subset of these, synthetic standards were available, allowing for quantification (ng/mL). Area ratios were calculated for all other oxylipins.

Statistical analyses

Descriptive statistics were used for baseline patient characteristics. Two-sample t-tests and Chi-square tests were used as appropriate to assess differences in baseline general patient characteristics. We used elastic net (EN) regularized regression for selection of predictors [29]. EN uses an additional tuning parameter (α) to combine the properties of ridge regression and lasso by applying both L1 and L2 penalties. Thereby, it simultaneously performs automatic variable selection and continuous shrinkage, while also dealing with high correlations amongst predictors. Prior to fitting the model, lipid measurements below the detection limit were imputed with the minimum measured value divided by two, all lipid variables were logarithmically transformed due to a non-normal distribution, and were mean scaled to ensure comparability by giving the metabolites equal weight. We performed EN regularization with a logit model, defining the OARSI-OMERACT responder status as the outcome. Prior to fitting the EN models, we performed a 10-fold cross-validation (CV) for selection of the optimal tuning parameters based on the smallest CV mean prediction error. In addition, we used manual α selection based on the out-of-sample deviance ratio and CV mean deviance to investigate the performance of more comprehensive models. First, a model was fit with commonly assessed patient characteristics and patient reported outcomes, measured at baseline (model 1). Second, we fitted model 2 by adding the Lipidizer™ platform lipids to model 1. Third, we fitted model 3 by adding the oxylipins to model 1. Fourth, we

combined the general patient characteristics with both lipid platforms in model 4. Lastly, we fitted a model with the predictors selected by model 2 and 3. We used the Stata command: *elasticnet logit depvar othervars, alpha(0.1(0.1)1) selection(cv, fold(10) alllambdas)*. The discriminatory accuracy of the model was estimated by receiver operating characteristic (ROC) analyses (Stata command: *rocreg*). The area under the curve (AUC) and corresponding 95% confidence intervals (CI) were calculated using 1,000 bootstrap replications. Additionally, we performed sensitivity analyses investigating the association between the lipid predictors and treatment response using univariable logistic regression. Stata V16.1 (StataCorp LP, TX, USA) was used for all analyses.

Availability of data and materials

The data underlying this article cannot be shared publicly due to the privacy of the participants of the HOPE study and legal reasons (HOPE study participants did not sign informed consent to make their data publicly available). The data is available upon request to interested qualified researchers. Data requests should be sent to the corresponding author.

Results

Study population

Baseline lipid measurements and the OARS-OMERACT responder status at week six were available in 40 prednisolone-treated patients. Figure 1 shows a flowchart of included patients. Of these patients, 31 (78%) fulfilled the OARS-OMERACT responder criteria. The percentage of patient fulfilling either the major criteria or a particular combination of minor criteria is presented in supplementary figure 2. Patients responding to prednisolone treatment showed statistically worse baseline AUSCAN function scores (19.6 ± 6.6) than non-responders (11 ± 7.5). None of the other general characteristics differed between responders and non-responders (table 1).




Figure 1. Flowchart of patient numbers

The present analyses included only patients randomized to prednisolone treatment. Of the 46 patients assigned, 2 discontinued the study due to poor efficacy or an adverse event. Four patients were excluded due to missing lipid measurements at baseline.

Table 1. Baseline characteristics of prednisolone-treated patients in the HOPE study

	All prednisolone treated n = 40	Responders n = 31 (78%)	Non-responders n = 9 (23%)
General characteristics			
Age, year	62.4 (9.3)	62.9 (9.4)	60.8 (9.4)
Sex, % women	85	84	89
BMI, kg/m ²	27.4 (4.4)	27.8 (4.2)	26.2 (5.0)
Education, % high	46	42	56
Disease duration	6.7 (7.1)	7.2 (7.4)	4.9 (5.8)
Erosive OA, %	71	74	56
Kellgren-Lawrence sum score, 0-120	35.1 (16.4)	34.1 (16.5)	37.5 (14.7)
Ultrasound synovitis sum score, 0-90	16.2 (6.6)	15.5 (6.4)	18.7 (7.2)
VAS global assessment, 0-100	52.3 (20.6)	54.2 (16.8)	45.6 (30.8)
AUSCAN pain, 0-20	11.0 (3.3)	11.3 (2.4)	10 (5.4)
AUSCAN function, 0-36	17.7 (7.6)	19.6 (6.6)	11 (7.5)

Numbers represent mean (SD) unless otherwise specified. Abbreviations: AUSCAN = Australian/Canadian Hand Osteoarthritis Index, BMI = body mass index, VAS = visual analogue scale

Prediction of treatment response using general patient characteristics

The general characteristics presented in table 1 were entered in model 1 as predictors of OARSI-OMERACT responder status. Only AUSCAN function was selected in the model (worse function associated with response), resulting in an AUC with 95% CI of 0.78 (0.56; 0.94). Predictors entered in the model, predictors selected by EN, and corresponding ROC curves of the models are shown in figure 2. Table 2 presents the baseline concentrations of the selected lipids. Tuning parameters and model deviances of all models are provided in table 3.

Table 2. Baseline levels of selected lipids

	All prednisolone treated n = 40	Responders n = 31 (78%)	Non-responders n = 9 (23%)
Levels selected Lipidizer™ lipids			
DAG(16:0/16:0), nmol/mL	0.28 (0.12)	0.30 (0.12)	0.18 (0.084)
PE(O-18:0/20:4), nmol/mL	66.01 (12.20)	63.26 (10.65)	75.48 (13.04)
Levels selected oxylipins			
9-HOTrE, area ratio	0.12 (0.09)	0.093 (0.059)	0.20 (0.14)
5-HEPE, area ratio	0.011 (0.015)	0.014 (0.016)	0.0043 (0.0032)
10-HDHA, ng/mL	0.0039 (0.0044)	0.0046 (0.0048)	0.0019 (0.0020)

Numbers represent mean (SD). Abbreviations: DAG = diacylglycerol, PE = phosphatidylethanolamine, 9-HOTrE = 9-hydroxy-octadecatrienoic acid, 5-HEPE = 5-hydroxy-eicosapentaenoic acid, 10-HDHA = 10-hydroxy-docosahexaenoic acid.

Added value of lipidomics for prediction of treatment response - Lipidizer™

In model 2, we added the 286 Lipidizer™ platform lipid species to model 1. Cross-validated parameter tuning selected an alpha of 0, resulting in the inclusion of all predictors in the model with an AUC of 0.95 (0.85; 0.99). With only minor increase in deviance (CV mean deviance 1.096 vs 1.095), a model (2a) with an alpha of 1 resulted in the selection of three variables: AUSCAN function and two lipids: diacylglycerol(DAG)(16:0/16:0) (higher levels associated with response), and phosphatidylethanolamine(PE)(O-18:0/20:4) (lower levels associated with response), with an AUC of 0.92 (0.78; 0.99).




Figure 2. Prediction model characteristics.

A) shows the variables included for model fitting of the three prediction models, colours correspond to the lines of the ROC curves in B). Of model 2, only the lipid classes are shown. Variables in bold font were selected in the final models. Model 1: General patient characteristics, model 2: model 1 + Lipidizer™ platform, model 3: model 1 + oxylipin platform, model 4: all variables included. Abbreviations: AUC = area under the curve, AUSCAN = Australian/Canadian Hand Osteoarthritis Index, CE = cholesteryl ester, CER = ceramide, DAG = diacylglycerol, DCER = dihydroceramide, FFA = free fatty acid, HCEr = hexosylceramide, KL = Kellgren-Lawrence, LCER = lactosylceramide, (L)PC = (lyso)phosphatidylcholines, (L)PE = (lyso)phosphatidylethanolamine, OA = osteoarthritis, SM = sphingomyelin, TAG = triacylglycerol, WAS = visual analogue scale, 9-HOTrE = 9-hydroxy-octadecatrienoic acid, 5-HEPE = 5-hydroxy-eicosapentaenoic acid, 10-HDHA = 10-hydroxy-docosahexaenoic acid.

Added value of lipidomics for prediction of treatment response – oxylipins

In model 3, the 25 identified oxylipins were added to model 1. With automated parameter tuning an alpha of 0 was used, selecting all variables for the model, resulting in an AUC of 0.88 (0.73; 0.97). However, with only marginal inflation of the CV mean deviance (1.186 vs 1.184) a more comprehensible model (3a) could be fit, which included AUSCAN function and three oxylipin predictors: 9-hydroxy-octadecatrienoic acid (HOTrE) (lower levels associated with response), 5-hydroxy-eicosapentaenoic acid (HEPE) and 10-hydroxy-docosahexaenoic acid (HDHA) (higher levels associated with response), with an AUC of 0.85 (0.65; 0.98).

Combining all predictors

Lastly, we combined the general patient characteristics with both lipid platforms in model 4. Again, automated parameter tuning resulting in an alpha of 0. Including all 326 variables in the model resulted in an AUC of 0.97 (0.90; 1). A more comprehensive model (4a) could be fit using an alpha of 0.2, resulting in the selection of 27 predictors. This model included all previously selected predictors from models 2 and 3, as well as 21 additional higher order (Lipidizer™) lipids (table 3), resulting in a model with an AUC of 0.99 (0.93; 1). In addition, we ran model 5 in which we included only the 6 predictors previously selected by EN in models 2 and 3. The discriminative ability of this model was only slightly less compared to the full model, with an AUC of 0.95 (0.81; 1), and significantly improved the prediction compared to a model based on general patient characteristics alone (model 1 vs model 5, $p=0.03$).

Table 3. Selected predictors and prediction model parameters

	Selected predictors	Tuning parameters		Out-of-sample deviance ratio	CV mean deviance	AUC (95% CI)
		Alpha	Lambda			
Model 1 General characteristics	1 AUSCAN function	1.00	0.113	0.0838	0.9770132	0.78 (0.56; 0.94)
Model 2 Model 1 + Lipidizer™	301 All variables*	0	23.327	-0.0269	1.095027	0.95 (0.85; 0.99)
Model 2a Model 1 + Lipidizer™ Manual alpha selection	AUSCAN function 3 DAG(16:0/16:0) PE(O-18:0/20:4)	1.00	0.150	-0.0275	1.095633	0.92 (0.78; 0.99)
Model 3 Model 1 + oxylipins	40 All variables*	0	2.193	-0.0689	1.170265	0.88 (0.73; 0.97)
Model 3a Model 1 + oxylipins Manual alpha selection	AUSCAN function 4 9-HOTrE 5-HEPE 10-HDHA	0.60	0.182	-0.0835	1.186297	0.85 (0.65; 0.98)
Model 4 All variables combined	326 All variables* AUSCAN function DAG(16:0/16:0) DAG(18:1/20:4) FFA(22:4) LPE(22:5) PC(16:0/20:1) PC(18:0/20:5) PE(16:0/18:2) PE(O-16:0/22:4) PE(O-18:0/20:4) PE(O-18:0/22:4) SM(24:0)	0	8.853	-0.0505	1.150134	0.97 (0.90; 1)
Model 4a All variables combined Manual alpha selection	27 TAG(45:0)-FA(16:0) TAG(47:1)-FA(16:0) TAG(48:1)-FA(16:1) TAG(49:1)-FA(17:0) TAG(49:2)-FA(16:1) TAG(49:2)-FA(18:1) TAG(50:2)-FA(18:2) TAG(51:1)-FA(16:0) TAG(51:2)-FA(15:0) TAG(54:0)-FA(18:0) TAG(54:2)-FA(16:0) TAG(56:1)-FA(18:1) 9-HOTrE 5-HEPE 10-HDHA AUSCAN function	0.2	0.475	-0.0921	1.195637	0.99 (0.93; 1)
Model 5 Predefined model based on predictor selection of model 2a and 3a	6 DAG(16:0/16:0) PE(O-18:0/20:4) 9-HOTrE 5-HEPE 10-HDHA	0	0.079	0.2993	.7671022	0.95 (0.81; 1)

*See additional file 1, tables A1 and A2 for the included lipids. Abbreviations: AUSCAN = Australian/Canadian Hand Osteoarthritis Index, AUC = area under the curve, CI = confidence interval, CV = cross-validation, DAG = diacylglycerol, FFA = free fatty acid, (L)PE = (lyso)phosphatidylethanolamine, PC = phosphatidylcholine, SM = sphingomyelin, TAG = triacylglycerol, 9-HOTrE = 9-hydroxy-octadecatrienoic acid, 5-HEPE = 5-hydroxy-eicosapentaenoic acid, 10-HDHA = 10-hydroxy-docosahexaenoic acid.

Sensitivity analyses

The univariable associations of baseline lipid levels with prednisolone treatment response are shown in the supplementary file, tables S1 and S2. The lipids included in model 2 and 3 were univariably among the lipids most strongly associated with treatment response, supporting the selection of predictors by the EN models.

Discussion

In this exploratory study we investigated the patients' lipid profile for the prediction of prednisolone treatment response in patients with painful inflammatory hand OA. We showed that lipidomics improved the discriminative accuracy of the prediction, when compared to commonly measured patient outcomes alone. Our results suggest that lipidomics is a promising field for further biomarker discovery for the prediction of anti-inflammatory treatment response.

The added predictive value of lipidomics is an interesting finding. From the Lipidizer™ platform, lipids containing fatty acid chains of palmitic acid (16:0), stearic acid (18:0) and arachidonic acid (20:4) were selected as predictors. Palmitic acid is the most abundant saturated fatty acid (SFA) in humans; under physiological conditions its concentration is tightly controlled by desaturation to palmitoleic acid and oleic acid, or elongation to stearic acid [30]. Pathophysiological conditions may increase SFA content, leading to activation of toll-like receptor (TLR)-4 triggered inflammatory signalling cascades via nuclear factor kappa B (NFκB) and cyclooxygenase (COX)-2, increasing proinflammatory cytokine production [31]. Arachidonic acid, an omega-6 polyunsaturated fatty acid (PUFA), is the main precursor of proinflammatory eicosanoids, although it may also give rise to anti-inflammatory mediators. In addition, hydroxylation of the omega-3 PUFAs eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA) may lead to hydroxyeicosapentaenoic acids (HEPE) and hydroxydocosahexaenoic acids (HDHA), which are precursors of anti-inflammatory and pro-resolving mediators [32]. Possibly, the lipid profile represents an indication of the patients' inflammatory state, and their likelihood to respond to anti-inflammatory treatment. However, we should be careful to avoid causal interpretations of our results since no causal inferences can be drawn from prediction analyses.

Furthermore, our results suggest that amongst other patient characteristics such as pain, radiographic OA severity and synovitis, hand function is the most contributing to the prediction of treatment response. Despite possible influences of the small sample size and patient selection, which likely resulted in a lack of predictive ability of characteristics such as age and sex, as well as regression to the mean, it implies that patients' hand function may be an important outcome to consider when making treatment decisions.

To our knowledge lipidomics for the prediction of treatment response in hand OA has not previously been investigated. A major strength of our study is the use of high-quality trial data. Furthermore, we have used lipidomics data from two different platforms, the standardized and commercially available Lipidizer™ platform for the measurement of a large variety of higher order lipids, and an in-house developed platform for the measurement of oxylipins.

However, there are also limitations to our study. Most notable is the small sample, which has likely resulted in overfitting of the models and a higher degree of uncertainty of the estimations. Also, since no study population with comparable data was available, external validation was not possible. In addition, the analyses have been performed in a specific, carefully selected patient population, therefore results may not be generalizable to other patient populations. The blood samples were obtained non-fasted at variable time points during the day due to differences in scheduled hospital visits. Although this may be viewed as a limitation, this procedure is a good reflection of daily practice and limits patient burden. Moreover, the identification of predictions for treatment response that do not require fasted or strictly scheduled sampling will benefit the feasibility and implementation in clinical practice. However, this may have resulted in additional variability in the lipid measurements. In a recent study by our research group we described intra-day variability (ICC) of (DAG) (16:0/16:0) of 0.62 and of (PE)(O-18:0/20:4) ICC of 0.46 [33], representing moderate to good reproducibility of the lipids selected in model 2a. Furthermore, we cannot exclude *in vitro* auto-oxidation of lipid metabolites. However, as this would have occurred to a similar extent in responders and non-responders, it is unlikely this has influenced our findings. Hence, the use of lipidomics, and in particular the development of a lipid biomarker, for the prediction of prednisolone treatment response warrants further investigation.

In conclusion, this exploratory study suggests that lipidomics may prove valuable in the prediction of prednisolone treatment response in patients with inflammatory hand OA. Prediction of treatment response may aid the selection of patients with a high likelihood of treatment benefit, which is crucial to prevent overtreatment and unnecessary exposure to adverse effects.

References

- [1] Zhang Y, Niu J, Kelly-Hayes M, Chaisson CE, Aliabadi P, Felson DT. Prevalence of symptomatic hand osteoarthritis and its impact on functional status among the elderly: The Framingham Study. *Am J Epidemiol* 2002;156:1021–7. <https://doi.org/10.1093/aje/kwf141>.
- [2] Kwok WY, Vliet Vlieland TPM, Rosendaal FR, Huizinga TWJ, Kloppenburg M. Limitations in daily activities are the major determinant of reduced health-related quality of life in patients with hand osteoarthritis. *Ann Rheum Dis* 2011;70:334–6. <https://doi.org/10.1136/ard.2010.133603>.
- [3] Kloppenburg M, Kwok W-Y. Hand osteoarthritis—a heterogeneous disorder. *Nat Rev Rheumatol* 2012;8:22–31. <https://doi.org/10.1038/nrrheum.2011.170>.
- [4] Loef M, Damman W, Mutsert R de, Rosendaal FR, Kloppenburg M. Health-related quality of life in patients with hand osteoarthritis from the general population and the outpatient clinic. *J Rheumatol* 2019. <https://doi.org/10.3899/jrheum.190781>.
- [5] Kortekaas MC, Kwok W-Y, Reijnen M, Watt I, Huizinga TWJ, Kloppenburg M. Pain in hand osteoarthritis is associated with inflammation: the value of ultrasound. *Ann Rheum Dis* 2010;69:1367–9. <https://doi.org/10.1136/ard.2009.124875>.
- [6] Keen HI, Wakefield RJ, Grainger AJ, Hensor EMA, Emery P, Conaghan PG. An ultrasonographic study of osteoarthritis of the hand: synovitis and its relationship to structural pathology and symptoms. *Arthritis Rheum* 2008;59:1756–63. <https://doi.org/10.1002/art.24312>.
- [7] Kroon FPB, Kortekaas MC, Boonen A, Böhlinger S, Reijnen M, Rosendaal FR, et al. Results of a 6-week treatment with 10 mg prednisolone in patients with hand osteoarthritis (HOPE): a double-blind, randomised, placebo-controlled trial. *The Lancet* 2019;394:1993–2001. [https://doi.org/10.1016/S0140-6736\(19\)32489-4](https://doi.org/10.1016/S0140-6736(19)32489-4).
- [8] Kaddurah-Daouk R, Kristal BS, Weinshilboum RM. Metabolomics: a global biochemical approach to drug response and disease. *Annu Rev Pharmacol Toxicol* 2008;48:653–83. <https://doi.org/10.1146/annurev.pharmtox.48.113006.094715>.
- [9] Villalvilla A, Gómez R, Largo R, Herrero-Beaumont G. Lipid Transport and Metabolism in Healthy and Osteoarthritic Cartilage. *Int J Mol Sci* 2013;14:20793–808. <https://doi.org/10.3390/ijms141020793>.
- [10] During A, Penel G, Hardouin P. Understanding the local actions of lipids in bone physiology. *Prog Lipid Res* 2015;59:126–46. <https://doi.org/10.1016/j.plipres.2015.06.002>.
- [11] Tsuge K, Inazumi T, Shimamoto A, Sugimoto Y. Molecular mechanisms underlying prostaglandin E2-exacerbated inflammation and immune diseases. *Int Immunol* 2019;31:597–606. <https://doi.org/10.1093/intimm/dxz021>.
- [12] Bastiaansen-Jenniskens YM, Siawash M, van de Lest CHA, Verhaar J a. N, Kloppenburg M, Zuurmond A-M, et al. Monounsaturated and Saturated, but Not n-6 Polyunsaturated Fatty Acids Decrease Cartilage Destruction under Inflammatory Conditions: A Preliminary Study. *Cartilage* 2013;4:321–8. <https://doi.org/10.1177/1947603513494401>.
- [13] Alvarez-Garcia O, Rogers NH, Smith RG, Lotz MK. Palmitate has proapoptotic and proinflammatory effects on articular cartilage and synergizes with interleukin-1. *Arthritis Rheumatol Hoboken NJ* 2014;66:1779–88. <https://doi.org/10.1002/art.38399>.
- [14] Castro-Perez JM, Kamphorst J, DeGroot J, Lafeber F, Goshawk J, Yu K, et al. Comprehensive LC-MS E lipidomic analysis using a shotgun approach and its application to biomarker detection and identification in osteoarthritis patients. *J Proteome Res* 2010;9:2377–89. <https://doi.org/10.1021/pr901094j>.
- [15] Zhang Q, Li H, Zhang Z, Yang F, Chen J. Serum metabolites as potential biomarkers for diagnosis of knee osteoarthritis. *Dis Markers* 2015;2015:684794. <https://doi.org/10.1155/2015/684794>.
- [16] Zhang W, Sun G, Aitken D, Likhodii S, Liu M, Martin G, et al. Lysophosphatidylcholines to phosphatidylcholines ratio predicts advanced knee osteoarthritis. *Rheumatol Oxf Engl* 2016;55:1566–74. <https://doi.org/10.1093/rheumatology/kew207>.
- [17] Kosinska MK, Liebisch G, Lochnit G, Wilhelm J, Klein H, Kaesser U, et al. A lipidomic study of phospholipid classes and species in human synovial fluid. *Arthritis Rheum* 2013;65:2323–33. <https://doi.org/10.1002/art.38053>.
- [18] Rockel JS, Zhang W, Shestopaloff K, Likhodii S, Sun G, Furey A, et al. A classification modeling approach for determining metabolite signatures in osteoarthritis. *PLoS ONE* 2018;13. <https://doi.org/10.1371/journal.pone.0199618>.
- [19] Altman R, Alarcón G, Appelrouth D, Bloch D, Borenstein D, Brandt K, et al. The American College of Rheumatology criteria for the classification and reporting of osteoarthritis of the hand. *Arthritis Rheum* 1990;33:1601–10.
- [20] Pham T, van der Heijde D, Altman RD, Anderson JJ, Bellamy N, Hochberg M, et al. OMERACT-OARSI initiative: Osteoarthritis Research Society International set of responder criteria for osteoarthritis clinical trials revisited. *Osteoarthritis Cartilage* 2004;12:389–99. <https://doi.org/10.1016/j.joca.2004.02.001>.
- [21] Kellgren JH, Lawrence JS. Radiological Assessment of Osteo-Arthrosis. *Ann Rheum Dis* 1957;16:494–502. <https://doi.org/10.1136/ard.16.4.494>.
- [22] Verbruggen G, Veys EM. Numerical scoring systems for the anatomic evolution of osteoarthritis of the finger joints. *Arthritis Rheum* 1996;39:308–20. <https://doi.org/10.1002/art.1780390221>.
- [23] Jonasdottir HS, Brouwers H, Toes REM,

Ioan-Facsinay A, Giera M. Effects of anticoagulants and storage conditions on clinical oxylipid levels in human plasma. *Biochim Biophys Acta Mol Cell Biol Lipids* 2018;1863:1511–22. <https://doi.org/10.1016/j.bbalip.2018.10.003>.

[24] Matyash V, Liebisch G, Kurzchalia TV, Shevchenko A, Schwudke D. Lipid extraction by methyl-tert-butyl ether for high-throughput lipidomics. *J Lipid Res* 2008;49:1137–46. <https://doi.org/10.1194/jlr.D700041-JLR200>.

[25] Chouvarine P, Giera M, Kastenmüller G, Artati A, Adamski J, Bertram H, et al. Trans-right ventricle and transpulmonary metabolite gradients in human pulmonary arterial hypertension. *Heart Br Card Soc* 2020;106:1332–41. <https://doi.org/10.1136/heartjnl-2019-315900>.

[26] Alarcon-Barrera JC, von Hegedus JH, Brouwers H, Steenvoorden E, Ioan-Facsinay A, Mayboroda OA, et al. Lipid metabolism of leukocytes in the unstimulated and activated states. *Anal Bioanal Chem* 2020;412:2353–63. <https://doi.org/10.1007/s00216-020-02460-8>.

[27] Contrepois K, Mahmoudi S, Ubhi BK, Papsdorf K, Hornburg D, Brunet A, et al. Cross-Platform Comparison of Untargeted and Targeted Lipidomics Approaches on Aging Mouse Plasma. *Sci Rep* 2018;8:17747. <https://doi.org/10.1038/s41598-018-35807-4>.

[28] Jónasdóttir HS, Ioan-Facsinay A, Kwekkeboom J, Brouwers H, Zuurmond A-M, Toes R, et al. An Advanced LC–MS/MS Platform for the Analysis of Specialized Pro-Resolving Lipid Mediators. *Chromatographia* 2015;78:391–401. <https://doi.org/10.1007/s10337-014-2779-5>.

[29] Zou H, Hastie T. Regularization and variable selection via the elastic net. *J R Stat Soc Ser B Stat Methodol* 2005;67:301–20. <https://doi.org/10.1111/j.1467-9868.2005.00503.x>.

[30] Carta G, Murru E, Banni S, Manca C. Palmitic Acid: Physiological Role, Metabolism and Nutritional Implications. *Front Physiol* 2017;8:902. <https://doi.org/10.3389/fphys.2017.00902>.

[31] Legrand P, Rioux V. The Complex and Important Cellular and Metabolic Functions of Saturated Fatty Acids. *Lipids* 2010;45:941–6. <https://doi.org/10.1007/s11745-010-3444-x>.

[32] Spite M, Clària J, Serhan CN. Resolvins, specialized proresolving lipid mediators, and their potential roles in metabolic diseases. *Cell Metab* 2014;19:21–36. <https://doi.org/10.1016/j.cmet.2013.10.006>.

[33] Loef M, von Hegedus JH, Ghorasaini M, Kroon FPB, Giera M, Ioan-Facsinay A, et al. Reproducibility of Targeted Lipidome Analyses (Lipidyzer) in Plasma and Erythrocytes over a 6-Week Period. *Metabolites* 2020;11. <https://doi.org/10.3390/metabo11010026>.

