



**Universiteit
Leiden**
The Netherlands

Dosage reduction of low weight heparin in patients with renal dysfunction: effects on anti-Xa levels and clinical outcomes

Hornung, P.; Khairoun, M.; Dekker, F.W.; Kaasjager, K.A.H.; Huisman, A.; Jakulj, L.; ... ; Ocak, G.

Citation

Hornung, P., Khairoun, M., Dekker, F. W., Kaasjager, K. A. H., Huisman, A., Jakulj, L., ... Ocak, G. (2020). Dosage reduction of low weight heparin in patients with renal dysfunction: effects on anti-Xa levels and clinical outcomes. *Plos One*, 15(10). doi:10.1371/journal.pone.0239222

Version: Publisher's Version

License: [Creative Commons CC BY 4.0 license](#)

Downloaded from: <https://hdl.handle.net/1887/3184682>

Note: To cite this publication please use the final published version (if applicable).


RESEARCH ARTICLE

Dosage reduction of low weight heparin in patients with renal dysfunction: Effects on anti-Xa levels and clinical outcomes

Paul Hornung¹, Meriem Khairoun¹, Friedo W. Dekker², Karin A. H. Kaasjager³, Albert Huisman⁴, Lily Jakulj⁵, Willem Jan W. Bos^{6,7}, Frits R. Rosendaal², Marianne C. Verhaar¹, Gurbey Ocak^{1,2,6*}

1 Department of Nephrology and Hypertension, University Medical Center Utrecht, Utrecht, the Netherlands, **2** Department of Clinical Epidemiology, Leiden University Medical Center, Leiden, the Netherlands, **3** Department of Internal Medicine, University Medical Center Utrecht, Utrecht, the Netherlands, **4** Department of Clinical Chemistry and Haematology, University Medical Center Utrecht, Utrecht, the Netherlands, **5** Dianet Dialysis Center and Department of Nephrology, Amsterdam University Medical Center, Amsterdam, the Netherlands, **6** Department of Internal Medicine, St Antonius Hospital, Nieuwegein, the Netherlands, **7** Department of Internal Medicine, Leiden University Medical Center, Leiden, the Netherlands

* g.ocak@antoniuziekenhuis.nl



Abstract

Background

To prevent bio-accumulation of low molecular weight heparins (LMWHs) in patients with decreased kidney function, dosage reduction and anti-Xa monitoring has been suggested. The aim of this study was to investigate the effect of pre-emptive dosage reduction of LMWH on anti-Xa levels. Furthermore, we investigated the association between anti-Xa levels and bleeding, thrombotic events and mortality.

Methods

In this single center study, we followed 499 patients with decreased renal function in whom anti-Xa levels were measured. We observed how many patients had anti-Xa levels that fell within the reference range, with a standard protocol of a pre-emptive dosage reduction of LMWH (25% reduction in patients with an estimated glomerular filtration rate (eGFR) between 30 and 60 ml/min/1.73m² and a reduction of 50% in patients with an eGFR below the 30 ml/min/1.73m²). Furthermore, Cox proportional hazard analyses were used to estimate hazard ratios to investigate the association between anti-Xa levels and major bleeding, thrombotic events and mortality within three months of follow-up.

Results

In a cohort of 499 patients (445 dalteparin and 54 nadroparin users), a pre-emptive dosage reduction of LMWH led to adequate levels of anti-Xa in only 19% of the patients (12% for the dalteparin users and 50% for nadroparin users). We did not find an association between anti-Xa levels and bleeding, thrombosis or mortality.

OPEN ACCESS

Citation: Hornung P, Khairoun M, Dekker FW, Kaasjager KAH, Huisman A, Jakulj L, et al. (2020) Dosage reduction of low weight heparin in patients with renal dysfunction: Effects on anti-Xa levels and clinical outcomes. PLoS ONE 15(10): e0239222. <https://doi.org/10.1371/journal.pone.0239222>

Editor: Giuseppe Remuzzi, Istituto Di Ricerche Farmacologiche Mario Negri, ITALY

Received: April 19, 2020

Accepted: September 1, 2020

Published: October 1, 2020

Copyright: © 2020 Hornung et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the manuscript.

Funding: The authors received no specific funding for this work.

Competing interests: The authors have declared that no competing interests exist.

Conclusion

Pre-emptive dosage reduction of LMWH leads to low anti-Xa levels in a large proportion, but this was not associated with bleeding, thrombosis or mortality.

Introduction

Low molecular weight heparins (LMWHs) are derived from unfractionated heparin by depolymerization and have superior pharmacokinetic properties as compared with unfractionated heparin [1–6]. The main anticoagulant pathway of LMWH is through inhibition of activated factor X (factor Xa), which can be monitored by anti-Xa levels in blood [1]. However, monitoring of anti-Xa levels is not routinely performed.

The predominant clearance of LMWH is by the kidneys, and consequently, the biologic half-life is prolonged in patients with impaired renal function [7, 8]. Several studies found increased bleeding risks in patients with renal failure who used LMWHs [9–11]. In a meta-analysis of twelve studies involving 4,971 patients who used LMWHs, patients with an estimated glomerular filtration rate (eGFR) ≤ 30 ml/min/1.73m² had a 2.3-fold increased major bleeding risk as compared with patients with an eGFR > 30 ml/min/1.73m² [9].

In order to prevent bio-accumulation of LMWHs, the American College of Chest Physicians Evidence-Based Clinical Practice Guidelines on antithrombotic therapy suggest a dosage reduction of LMWHs combined with anti-Xa monitoring in patients with decreased renal function [12]. Furthermore, the Dutch Federation of Nephrology guideline for anticoagulation with LMWHs in patients with renal impairment recommends to reduce the dose of LMWHs by 25% in patients with an eGFR between the 30 and 60 ml/min/1.73m² and to decrease the dose of LMWHs by 50% in patients with an eGFR ≤ 30 ml/min/1.73m² [13]. In addition, anti-Xa monitoring is recommended in patients with an eGFR < 60 ml/min/1.73m² with advised peak target ranges of 1.0–2.0 U/ml for once daily dosing and 0.6–1.0 U/ml for twice daily dosing [13].

Although advised in current guidelines, it is not known whether this dosage reduction of LMWHs in patients with an eGFR < 60 ml/min/1.73m² leads to adequate anti-Xa levels. In addition, it is unknown whether the reduced dosage is still effective with regard to clinical endpoints. Therefore, anti-Xa monitoring for safety purposes is still a matter of debate.

The aim of this study was to investigate whether pre-emptive dosage reduction of LMWHs (nadroparin and dalteparin) leads to adequate anti-Xa levels. Furthermore, we aimed to investigate the association between anti-Xa levels and clinical outcomes (major bleeding, thrombotic events and mortality).

Methods

Study design and population

This single center cohort study was conducted at the University Medical Center Utrecht, the Netherlands. Adult patients ≥ 18 years who were admitted to the Department of the Internal Medicine between January 2012 and January 2019 and in whom an anti-Xa measurement was measured, were eligible for this study. We only included patients who used dalteparin or nadroparin in our study with a therapeutic dosage of LMWH and an eGFR below the 60 ml/min/1.73m². This study was approved by the chairs of the different departments of Internal Medicine of the University Medical Center Utrecht. The ethical commission of the University

Medical Center Utrecht approved institutional review board exemption, since this was a retrospective study and participants were not subject to procedures or were not required to follow rules of behaviour. For personal data protection, all data were anonymized before analysis. This study was performed according to the Declaration of Helsinki. We followed patients for three months, until an outcome event (bleeding, thrombosis (venous thrombosis or ischemic stroke) or death) or the end of the follow-up period (June 2019).

Low molecular weight heparin guideline

Our local clinical protocol follows the national Dutch guideline which recommends a dosage reduction of LMWH in patients with an eGFR below 60 ml/min/1.73m², followed by an anti-Xa level measurement when LMWH is continued for more than three days [13]. Dalteparin is the most prescribed LMWH for therapeutic indications in our center followed by nadroparin. A dosage reduction of 25% for patients with an eGFR between 30 and 60 ml/min/1.73m² and of 50% for patients with an eGFR <30 ml/min/1.73m² is recommended after a full first dose of 100%. The recommended dosage per subcutaneous injection is 200 international units (IU) per kg for once daily dalteparin and 100 IU/kg for twice daily. The recommended dosage per subcutaneous injection is 171 IU/kg for once daily nadroparin and 86 IU/kg for twice daily. We categorized patients into three groups (dosage below, according or above the guideline recommendations) based on weight and eGFR according to the guideline and local protocol.

Anti-Xa measurements

Anti-Xa levels were measured four hours after LMWH administration on the third day of LMWH use for both patient with once and twice daily LMWH use. Blood samples were collected in sodium citrate tubes (3.2%, 0.105 M, BD vacutainer, Becton Dickinson, UK). All laboratory measurements were performed in the central clinical ISO15189 certified laboratory of the University Medical Center Utrecht Utrecht. Until June 2018, anti-Xa levels were measured on a STA-Rack evolution coagulation analyser using STA liquid anti-Xa reagent (Diagnostica Stago, Asnières-sur-Seine, France). From June 2018, anti-Xa levels were measured on an ACL Top 750 LAS coagulation analyser using HemoSiL Liquid Anti-Xa reagent (Werfen, Bedford, MA, USA). Both anti-Xa assays were carefully aligned. All assays participated in robust internal and external quality assessment schemes and local performance characteristics were within the pre-defined limits stated by the manufacturers.

The target range for peak anti-Xa levels was between 1.0 U/ml and 2.0 U/ml with once daily LMWH and was between 0.6 U/ml and 1.0 U/ml with twice daily LMWH according to our local protocol and the Dutch guideline. Based on this target range, patients had anti-Xa levels below the range if anti-Xa levels were <1.0 U/ml for once daily LMWH and <0.6 U/ml for twice daily LMWH and above range if anti-Xa levels were >2.0 U/ml for once daily LMWH and >1.0 U/ml for twice daily LMWH.

Demographic and clinical data

Demographic and clinical data were obtained from the electronic medical record. Baseline was set as the day of the first anti-Xa measurement. Data on age, sex, height, weight, LMWH (dalteparin or nadroparin), dosage, treatment frequency, antiplatelet drug use, indication for anticoagulation, prior ischemic stroke, prior bleeding, malignancy, liver cirrhosis, heart failure, hypertension history, peripheral vascular disease, hemoglobin and plasma creatinine concentrations were collected at baseline. The creatinine value preceding the anti-Xa measurement was used to calculate eGFR using the CKD-EPI formula [14]. In 94% of the patients, creatinine measurements were performed within three days before the anti-Xa measurement.

Study outcomes: Bleeding, thrombosis and mortality

Endpoints of the study were major bleeding, thrombotic events (ischemic stroke or venous thrombosis including deep vein thrombosis or pulmonary embolism) and mortality within three months after baseline (date of anti-Xa measurement). Major bleeding events were collected from the medical records and evaluated following the classification of the Scientific and Standardization Committee of the International Society on Thrombosis and Haemostasis [15]. A major bleeding was defined as a fatal bleeding or a symptomatic bleeding in a critical area or organ (intracranial, intraspinal, intraocular, retroperitoneal, intra-articular, pericardial, or intramuscular with compartment syndrome) or bleeding causing a fall in hemoglobin level of more than 20 g/L (1.24 mmol/L) or a bleeding event leading to transfusion of two or more units of red cells. Venous thrombosis was confirmed by compression ultrasound for deep vein thrombosis of the leg or arm. Furthermore, pulmonary embolism had to be diagnosed by computed tomography. Ischemic stroke was confirmed by computed tomography or magnetic resonance imaging.

Statistical analyses

Continuous variables are presented as median and interquartile range (IQR). Categorical variables are presented as counts with corresponding percentages.

To investigate the association between dosage reduction of LMWHs and anti-Xa levels, we calculated the proportions of anti-Xa levels that fell within the range across different groups of guideline adherence (dosage below, within and above guideline recommendation). These analyses were stratified for different eGFR categories (eGFR between 30 and 60 ml/min/1.73m² and eGFR <30 ml/min/1.73m²). Furthermore, we stratified these analyses for dalteparin and nadroparin use.

The observation time in each participant was calculated as the time elapsed between the date of the first anti-Xa level measurement and the outcome event within three months (bleeding, thrombosis or mortality) or censoring date (June 2019). Incidence rates for bleeding, thrombosis or mortality were estimated by dividing the number of patients with an event by the total observation time at risk. Only the first event was taken into account. When estimating the incidence rates for bleeding, we neglected the occurrence of thrombosis and vice versa.

In addition, we calculated crude and adjusted hazard ratios (HRs) with 95% confidence intervals (CIs) to evaluate the association between anti-Xa level categories (below, within and above target range) and outcomes (bleeding, thrombosis and mortality within three months). Hazard ratios were adjusted for age, sex, body mass index, LMWH (dalteparin or nadroparin), daily frequency of LMWH dosing (once or twice daily), antiplatelet drug use, indication for anticoagulation, prior ischemic stroke, prior bleeding, malignancy, liver cirrhosis, heart failure, hypertension history, peripheral vascular disease, hemoglobin concentration and eGFR. Furthermore, we investigated the association between outcomes (bleeding, thrombosis and mortality within three months) and a dosage of LMWH below or above the guideline recommendation as compared with a dosage according to the guideline.

As a sensitivity analysis, a time-dependent cox regression analysis was performed to account for potential influence of changes in anti-Xa levels over time. In addition, we performed a competing risk analysis to calculate hazard ratios for bleeding and thrombosis accounting for death as a competing risk. Furthermore, we also investigated the association between anti-Xa levels (below, within and above target range) and outcomes (major bleeding, thrombosis and mortality) within one month. Finally, we also investigated the association between LMWH dosing and anti-Xa levels and outcomes excluding patients from June 2018

(n = 39), since anti-Xa levels were measured with another analyser. All analyses were done by use of SPSS statistical software version 25.0 (IBM SPSS Statistics).

Results

Baseline characteristics

We included 499 adult patients with an anti-Xa measurement and therapeutic use of dalteparin or nadroparin in our analysis. Patients with an eGFR >60 ml/min/1.73m² (n = 325), who did not use LMWHs (n = 107), with prophylactic dosages of LMWH (n = 73), or patients who used tinzaparin (n = 10) or enoxaparin (n = 5) were excluded (Fig 1).

The median age of patients was 68 years and 39% of the included patients were women (Table 1). Dalteparin was used by 89% of the patients, while nadroparin was used in 11% of the patients. Indication for LMWH use was venous thrombosis in 45%, atrial fibrillation in 37%, artificial heart valves in 7% and 11% of the patient had other reasons. Of the 499 patients, 42% had an eGFR between 30 and 60 ml/min/1.73m² (median eGFR 43 ml/min/1.73m²) and 58% had an eGFR below 30 ml/min/1.73m² (median eGFR 17 ml/min/1.73m²). There were 78 dialysis patients in our study (69 hemodialysis and 9 peritoneal dialysis patients).

Anti-Xa levels

The median anti-Xa levels were 0.44 U/ml (IQR 0.21–0.75) for the total group (n = 499), 0.45 U/ml (IQR 0.17–0.70) for once daily dalteparin users (n = 163), 0.40 U/ml (IQR 0.21–0.64) for twice daily dalteparin users (n = 282), 1.15 U/ml (IQR 0.95–1.48) for once daily nadroparin users (n = 15) and 0.59 U/ml (IQR 0.28–0.92) for twice daily nadroparin users (n = 39).

Table 2 shows the proportion of patients who had anti-Xa levels below, within and above the target range. In the total group, only 21% of the patients had a first anti-Xa measurement within range, while 73% of the patients were below the target range and 6% of the patients were above the target range. Declining eGFR resulted in increasing proportion of patients below the target range. An eGFR between 30 and 60 ml/min/1.73m² resulted in 66% of the patients below target range and an eGFR below 30 ml/min/1.73m² resulted in 78% of the patients below target range. In the dialysis group, only 14% of the patients had a first anti-Xa measurement within range, while 78% of the patients were below the target range and 8% of the patients were above the target range.

Overall, only 19% of the patients with an eGFR <60 ml/min/1.73m² who received a dosage according to the guideline had anti-Xa levels within the target range. The great majority (91%) of patients with a dosage below guideline recommendation had anti-Xa levels below the target range. Moreover, a dosage above the current guideline advice still resulted in anti-Xa levels that were below the target range in 67% of the total cohort.

Table 3 displays differences in the anti-Xa levels between dalteparin and nadroparin users. A dosage according to the guideline resulted in anti-Xa levels within the target range for 12% of the dalteparin users and for 50% of the nadroparin users. A dosage according to the guideline resulted in 2% of the patients within the anti-Xa target range for once daily dalteparin use and 16% for twice daily use. For nadroparin use, 73% of the patients with once daily use and 38% of the patients with twice daily use had an anti-Xa level within target range with a dosage according to the guideline.

A second measurement of anti-Xa levels was done within one week after the first measurement in 171 (47%) of the 363 patients with anti-Xa levels below range in the first measurement. For these 171 patients, the second anti-Xa level was still below range in 71%, within range in 26% and above range in 3%. For the 32 patients who had a first anti-Xa level above range, a




Fig 1. Study outline.

<https://doi.org/10.1371/journal.pone.0239222.g001>

second measurement was done in 12 patients within one week (38%). In these 12 patients, the second anti-Xa level was below range in 17%, within range in 42% and still above range in 42%.

Bleeding

There were 104 first major bleeding events within 3 months of follow-up, of which 26 were intracranial and 78 were extracranial bleeding events. Of the 104 major bleeding events, 20 were fatal bleedings. The major bleeding rate within three months of follow-up was 1238 major bleeding events per 1000 person-years. There was no relation with anti-Xa levels below

Table 1. Baseline characteristics.

		Total (n = 499)
Age, median (IQR), years		68 (60–77)
Sex female (%)		195 (39%)
Low molecular weight heparin		
	Dalteparin once daily	163 (33%)
	Dalteparin twice daily	282 (57%)
	Nadroparin once daily	15 (3%)
	Nadroparin twice daily	39 (8%)
Antiplatelet drug use		87 (17%)
Indication for anticoagulation		
	Atrial fibrillation	223 (45%)
	Venous thrombosis	182 (37%)
	Prosthetic heart valves	37 (7%)
	Other	57 (11%)
Comorbidities		
	Prior ischemic stroke	49 (10%)
	Prior bleeding	169 (34%)
	Malignancy	200 (40%)
	Liver cirrhosis	14 (3%)
	Heart failure	74 (15%)
	Hypertension history	348 (70%)
	Peripheral vascular disease	85 (17%)
Weight, median (IQR), kg		78 (68–90)
Body mass index, median (IQR), kg/m ²		25.6 (22.9–29.4)
Hemoglobin, median (IQR), g/L		9.8 (8.9–11.0)
Renal function (eGFR), median (IQR), ml/min/1.73m ²		27 (15–41)
	30–60 ml/min/1.73m ²	212 (42%)
	<30 ml/min/1.73m ²	287 (58%)

IQR, interquartile range.

<https://doi.org/10.1371/journal.pone.0239222.t001>

or above the target range and bleeding events (Table 4). A dosage below (HR 0.9, 95% CI 0.4–2.0) or above (HR 0.9, 95% CI 0.6–1.5) the guideline recommendation as compared with a dosage according to the guideline was not associated with major bleedings after adjustment.

Thrombosis

There were 27 first thrombotic events (7 patients with ischemic stroke and 20 patients with venous thrombosis) corresponding to an event rate of 281 thrombotic events per 1000 person-years. Of the 27 thrombotic events, 8 were fatal. Anti-Xa level below the target range as compared with adequate anti-Xa levels were associated with a 2.4-fold (95% CI 0.7–7.9) increased thrombotic risk, which did not materially change after adjustment for age, sex, body mass index, LMWH (dalteparin or nadroparin), daily frequency of LMWH dosing (once or twice daily), antiplatelet drug use, indication for anticoagulation, prior ischemic stroke, prior bleeding, malignancy, liver cirrhosis, heart failure, hypertension history, peripheral vascular disease, hemoglobin levels and eGFR (HR 2.5, 95% CI (0.7–8.8)) (Table 4).

The hazard ratio of thrombosis was 2.5 (95% CI 0.7–9.0) for a dosage below the guideline recommendation as compared with a dosage according to the guideline after adjustment. A

Table 2. Distribution of anti-Xa levels.

Anti-Xa levels*		Total (n = 499)	Dosage below guideline (n = 46)	Dosage according guideline (n = 178)	Dosage above guideline (n = 275)
eGFR <60 ml/min/1.73m²					
	Below range	363 (73%)	42 (91%)	138 (78%)	183 (67%)
	Within range	104 (21%)	4 (9%)	33 (19%)	67 (24%)
	Above range	32 (6%)	0 (0%)	7 (4%)	25 (9%)
eGFR 30–60 ml/min/1.73m²					
	Below range	140 (66%)	41 (91%)	67 (70%)	32 (45%)
	Within range	56 (26%)	4 (9%)	23 (24%)	29 (41%)
	Above range	16 (8%)	0 (0%)	6 (6%)	10 (14%)
eGFR <30 ml/min/1.73m²					
	Below range	223 (78%)	1 (100%)	71 (87%)	151 (74%)
	Within range	48 (17%)	0 (0%)	10 (12%)	38 (19%)
	Above range	16 (6%)	0 (0%)	1 (1%)	15 (7%)

*Anti-Xa levels: below range (<1.0 U/ml for once daily low molecular weight heparin and <0.6 U/ml for twice daily low molecular weight heparin), within range (1.0 U/ml—2.0 U/ml for once daily low molecular weight heparin and 0.6 U/ml—1.0 U/ml for twice daily low molecular weight heparin), above range (>2.0 U/ml for once daily low molecular weight heparin and >1.0 U/ml for twice daily low molecular weight heparin).

<https://doi.org/10.1371/journal.pone.0239222.t002>

dosage above the guideline recommendation was not associated with thrombotic events (HR 1.0, 95% CI 0.4–2.7).

Mortality

Of the 499 patients, 146 patients died within three months. The mortality rate was 1471 per 1000 person-years. We did not find an association between anti-Xa levels and mortality (Table 4).

A dosage below (HR 1.0, 95% CI 0.6–1.8) or above (HR 0.8, 95% CI 0.5–1.1) the guideline recommendation as compared with a dosage according to the guideline was not associated with mortality after adjustment.

Sensitivity analyses

The time-dependent cox regression analyses showed similar results as the analysis in which only the first anti-Xa measurement was taken into account. There was no relation with anti-Xa levels above the target range and bleeding events after adjustments (HR 1.0, 95% CI 0.4–2.4). Furthermore, time-dependent cox regression analysis showed an adjusted HR of 1.6 (95% CI 0.6–4.4) of thrombosis for anti-Xa level below the target range as compared with adequate anti-Xa levels. We did not find an association between anti-Xa levels and mortality.

Competing risk models accounting for death showed also similar associations regarding the risk of bleeding and thrombosis. There was also no relation with anti-Xa levels above the target range and bleeding events after adjustments (HR 1.1, 95% CI 0.6–1.8) when accounting death as competing risk. Furthermore, HR of thrombosis was 2.5 (95% CI 0.6–9.6) for anti-Xa level below the target range as compared with adequate anti-Xa levels.

We also investigated the association between anti-Xa levels and outcomes within one month (Table 5). These hazard ratios for one month were comparable with the hazard ratios for three months.

Table 3. Distribution of anti-Xa levels stratified for dalteparin and nadroparin.

anti-Xa levels*	DALTEPARIN			NADROPARIN		
	Dosage below guideline (n = 33)	Dosage according guideline (n = 146)	Dosage above guideline (n = 264)	Dosage below guideline (n = 13)	Dosage according guideline (n = 32)	Dosage above guideline (n = 9)
eGFR <60 ml/min/1.73m²						
Below range	31 (94%)	126 (86%)	182 (68%)	11 (85%)	12 (38%)	1 (11%)
Within range	2 (6%)	17 (12%)	64 (24%)	2 (15%)	16 (50%)	3 (33%)
Above range	0 (0%)	3 (2%)	20 (8%)	0 (0%)	4 (13%)	5 (56%)
eGFR 30–60 ml/min/1.73m²						
Below range	30 (94%)	64 (78%)	32 (48%)	11 (85%)	3 (21%)	0 (0%)
Within range	2 (6%)	16 (20%)	26 (39%)	2 (15%)	7 (50%)	3 (75%)
Above range	0 (0%)	2 (2%)	9 (13%)	0 (0%)	4 (29%)	1 (25%)
eGFR <30 ml/min/1.73m²						
Below range	1 (100%)	62 (97%)	150 (75%)	0 (0%)	9 (50%)	1 (20%)
Within range	0 (0%)	1 (2%)	38 (19%)	0 (0%)	9 (50%)	0 (0%)
Above range	0 (0%)	1 (2%)	11 (6%)	0 (0%)	0 (0%)	4 (80%)

* Anti-Xa levels: below range (<1.0 U/ml for once daily low molecular weight heparin and <0.6 U/ml for twice daily low molecular weight heparin), within range (1.0 U/ml—2.0 U/ml for once daily low molecular weight heparin and 0.6 U/ml—1.0 U/ml for twice daily low molecular weight heparin), above range (>2.0 U/ml for once daily low molecular weight heparin and >1.0 U/ml for twice daily low molecular weight heparin).

<https://doi.org/10.1371/journal.pone.0239222.t003>

Furthermore, we investigated whether results were different when patients were excluded after June 2018 (n = 39), since anti-Xa levels were measured with another analyser. However, this did not change the results. In the total group, a similar percentage of only 21% of the patients had a first anti-Xa measurement within range. In addition, we did not find an association between anti-Xa levels and bleeding, thrombosis or mortality.

Discussion

In this observational cohort study of 499 patients with decreased renal function who received therapeutic dosages of nadroparin or dalteparin, we showed that a pre-emptive dosage reduction of LMWH (25% reduction in patients with an eGFR between 30 and 60 ml/min/1.73m² and a reduction of 50% in patients with an eGFR below the 30 ml/min/1.73m²) leads to adequate levels of anti-Xa in only 19% of the patients. Furthermore, there were large differences in the proportion of patients within the target ranges for dalteparin and nadroparin use. Dosage reduction according to the guideline for patients with eGFR below 60 ml/min/1.73m² resulted in adequate anti-Xa levels in 50% of the nadroparin users and in 12% of the dalteparin users. We did not find an association between anti-Xa levels and bleeding, thrombosis or mortality.

Table 4. Anti-Xa levels and bleeding, thrombosis and mortality within three months.

Anti-Xa levels*	BLEEDING		THROMBOSIS		MORTALITY	
	Crude HR (95% CI)	Adjusted** HR (95% CI)	Crude HR (95% CI)	Adjusted** HR (95% CI)	Crude HR (95% CI)	Adjusted** HR (95% CI)
eGFR <60 ml/min/1.73m ²						
Below range	1.1 (0.7–1.8)	1.0 (0.6–1.8)	2.4 (0.7–7.9)	2.5 (0.7–8.8)	1.2 (0.8–1.8)	1.2(0.8–1.9)
Within range	1 ref	1 ref	1 ref	1 ref	1 Ref	1 ref
Above range	1.2 (0.5–2.8)	1.4 (0.6–3.4)	NA	NA	0.8 (0.4–1.9)	1.0 (0.4–2.4)

NA, not applicable.

* Anti-Xa levels: below range (<1.0 U/ml for once daily low molecular weight heparin and <0.6 U/ml for twice daily low molecular weight heparin), within range (1.0 U/ml–2.0 U/ml for once daily low molecular weight heparin and 0.6 U/ml–1.0 U/ml for twice daily low molecular weight heparin), above range (>2.0 U/ml for once daily low molecular weight heparin and >1.0 U/ml for twice daily low molecular weight heparin).

** Adjusted for age, sex, body mass index, low molecular weight heparin (dalteparin or nadroparin), daily frequency of low molecular weight heparin dosing (once or twice daily), antiplatelet drug use, indication for anticoagulation, prior ischemic stroke, prior bleeding, malignancy, liver cirrhosis, heart failure, hypertension history, peripheral vascular disease, hemoglobin levels and eGFR.

<https://doi.org/10.1371/journal.pone.0239222.t004>

Several previous studies have reported on the association between dosages of LMWHs and anti-Xa levels, mostly focussing on enoxaparin [8, 16–21]. Unadjusted therapeutic dosages of enoxaparin were associated with increased anti-Xa levels in patients with an eGFR <30 ml/min/1.73m² as compared with patients with an eGFR >30 ml/min/1.73m² [8, 20, 21]. Furthermore, several observational studies reported that dosage reduction of enoxaparin in patients with a decreased kidney function led to adequate anti-Xa levels [16–19]. Therefore, these studies recommended pre-emptive dose reductions of enoxaparin in patients with decreased kidney function. In contrast to enoxaparin, previous studies did not find an association between decreased kidney function and increased anti-Xa levels for tinzaparin [22–25].

Few studies have reported on the association between nadroparin or dalteparin use and anti-Xa levels. A previous study showed that anti-Xa activity in patients with eGFR below 60 ml/min/1.73 m² treated with pre-emptive dosage reduction of therapeutic dosages of nadroparin resulted in 51% of the anti-Xa levels within the target range, which is similar to our results [26]. To our knowledge, there are no studies that investigated the association between anti-Xa levels and dalteparin for unadjusted and adjusted dosages in patients with impaired kidney function. Dalteparin (6000 Dalton) has a similar molecular weight as tinzaparin (6500

Table 5. Anti-Xa levels and bleeding, thrombosis and mortality within one month.

Anti-Xa levels	BLEEDING		THROMBOSIS		MORTALITY	
	Adjusted*	Adjusted*	Adjusted*	Adjusted*	Adjusted*	Adjusted*
	HR (95% CI)	HR (95% CI)	HR (95% CI)	HR (95% CI)	HR (95% CI)	HR (95% CI)
eGFR <60 ml/min/1.73m ²						
Below range	1.1 (0.6–1.9)	3.3 (0.7–15.2)	1.1 (0.6–1.9)			
Within range	1 ref	1 ref	1 ref			
Above range	1.7 (0.7–4.6)	NA	1.3 (0.5–3.3)			

NA, not applicable.

* Adjusted for age, sex, body mass index, low molecular weight heparin (dalteparin or nadroparin), daily frequency of low molecular weight heparin dosing (once or twice daily), antiplatelet drug use, indication for anticoagulation, prior ischemic stroke, prior bleeding, malignancy, liver cirrhosis, heart failure, hypertension history, peripheral vascular disease, hemoglobin levels and eGFR.

<https://doi.org/10.1371/journal.pone.0239222.t005>

Dalton) and has a higher molecular weight than nadroparin (4300 Dalton) or enoxaparin (4500 Dalton) [27, 28]. Our finding that a dosage reduction in dalteparin for patients with eGFR below the 60 ml/min/1.73m² resulted in adequate anti-Xa levels in 12% of the patients was in line with the finding of a previous study that showed adequate anti-Xa levels in only 8% of tinzaparin users [25].

The greater proportion of anti-Xa levels within target range for nadroparin (50%) than with dalteparin (12%) after pre-emptive dose reduction in patients with an eGFR <60 ml/min/1.73m² can be explained by the difference in molecular weight (4300 Dalton versus 6000 Dalton, respectively) [29]. LMWHs are cleared through a combination of depolymerization after binding to endothelial cell receptors and macrophages and renal filtration. The degree of renal clearance of LMWHs are dependent on the renal function and on the molecular weight [12]. Therefore, the lower molecular weight of nadroparin than dalteparin leads to a longer biological half-life of nadroparin than dalteparin. Consequently, a strategy of pre-emptive dosage reduction of LMWHs probably leads to lower anti-Xa levels of dalteparin in comparison with nadroparin.

Previous studies also investigated the association between anti-Xa levels and outcome events [30–34]. Low anti-Xa levels (<0.5 IU/ml) as compared with anti-Xa levels in the range of 0.5 to 1.2 IU/ml were found to be associated with an increased 30-day mortality risk [30]. We did not find an association between anti-Xa levels and mortality in our study. Another study in patients after hip replacement reported that low anti-Xa levels were associated with increased post-operative thrombosis risk, while high anti-Xa levels were associated with increased bleeding (wound hematoma) [31]. Yet, another study in patients with acute venous thromboembolism who were treated with dalteparin demonstrated that increased anti-Xa levels (above 0.8 IU/ml) were associated with increased bleeding risks [32]. In line with our study, two previous studies did not show an association between anti-Xa levels and bleeding [33, 34].

The strengths of the present study are the large number of participants with information on anti-Xa levels, kidney function and information on bleeding and thrombotic events. To our knowledge, this is the first study that evaluated the association between dosage reduction of dalteparin and nadroparin, anti-Xa levels and outcomes (bleeding, thrombosis and mortality). This study has also several limitations. First, patients who were treated with tinzaparin and enoxaparin were excluded because of the low number of patients. Second, patients from our tertiary center in our study reflect a high-risk group with increased bleeding, thrombosis and mortality rates in which anti-Xa levels were measured. Therefore, it could be that our findings are not generalizable to all patients with renal dysfunction treated with LMWHs. Third, only in a small proportion of patients a second measurement of anti-Xa levels was performed. This could be a selected group in which a low anti-Xa level was expected explaining the low levels in the second anti-Xa level measuring. Fourth, we had limited power in the group of dialysis patients, patients with a LMWH dosage lower than the recommended dose and in the group of patients who used nadroparin. Furthermore, confidence intervals for thrombotic risk were very wide, because of a limited powered due to the low number of thrombotic events. Fifth, the cause of death was not registered. Sixth, we have no data about the time of LMWH administration and time of blood drawing. Therefore, we cannot exclude the possibility of incorrect drawing leading to low peak target ranges. Finally, we cannot exclude residual confounding and selection bias in the association between anti-Xa levels and outcomes. Selection could have occurred in different stages of our study. First, anti-Xa levels are probably not measured in all patients with an eGFR below 60 ml/min/1.73m² who used LMWH. Second, it is not clear which patients continued oral anticoagulants and which patients were switched to LMWHs during hospitalization. Third, differences in anti-Xa levels between once versus twice

daily dosing of LMWH and nadroparin versus dalteparin use could be due to selection bias rather than pharmacological differences.

The American guideline “Parenteral Anticoagulants: American College of Chest Physicians Evidence Based Clinical Practice Guidelines (9th Edition)” and the Dutch guideline “Anticoagulation with LMWHs in renal impairment” suggests anti-Xa monitoring and dose reductions in patients with renal impairment [12, 13]. The American College of Chest Physicians Evidence Based Clinical Practice Guidelines suggestion for anti-Xa monitoring is made in a summary chapter that does not provide graded recommendations. The Dutch guideline is based on expert opinion. In contrast, the American Society of Hematology (ASH) guideline panel for optimal management of anticoagulation therapy for venous thromboembolism makes a graded recommendation against anti-Xa monitoring in patients with an eGFR below 30 ml/min/1.73m² receiving LMWH for treatment of venous thrombo-embolism (conditional recommendation based on very low certainty in the evidence about effects) [35].

Although the rationale of dosage reductions and anti-Xa monitoring of LMWHs is clear, we showed no association between anti-Xa levels and outcomes. Therefore, our data supports the ASH recommendation against anti-factor Xa monitoring. Moreover, there are some other problems with anti-Xa monitoring. It is not known whether measurements of anti-Xa levels must be performed after each dosage adjustment and whether anti-Xa level monitoring should be continued in patients who are within range. Based on the poor performance of the Dutch Federation of Nephrology guideline recommendation in clinical practice, future studies should investigate whether a strategy without anti-Xa measurements as suggested by the ASH guideline is non-inferior to an anti-Xa guided dosing strategy in patients with an impaired kidney function as suggested by the Dutch and American College of Chest Physicians Evidence Based Clinical Practice guideline.

In conclusion, pre-emptive dosage reduction of LMWHs in patient with an eGFR <60 ml/min/1.73m² leads to a high proportion of patient below target anti-Xa ranges. However, low anti-Xa levels were not associated with clinical outcomes in our study.

Acknowledgments

We would like to thank all patients who have contributed data via their medical records. Furthermore, we gratefully acknowledge M.E. Vianen for the participation in the data acquisition.

Author Contributions

Conceptualization: Paul Hornung, Meriem Khairoun, Gurbey Ocak.

Data curation: Paul Hornung, Meriem Khairoun, Gurbey Ocak.

Formal analysis: Paul Hornung, Meriem Khairoun, Gurbey Ocak.

Investigation: Paul Hornung, Meriem Khairoun, Gurbey Ocak.

Methodology: Paul Hornung, Meriem Khairoun, Friedo W. Dekker, Karin A. H. Kaasjager, Albert Huisman, Lily Jakulj, Willem Jan W. Bos, Frits R. Rosendaal, Marianne C. Verhaar, Gurbey Ocak.

Supervision: Friedo W. Dekker, Karin A. H. Kaasjager, Albert Huisman, Lily Jakulj, Willem Jan W. Bos, Frits R. Rosendaal, Marianne C. Verhaar, Gurbey Ocak.

Validation: Paul Hornung, Meriem Khairoun, Gurbey Ocak.

Writing – original draft: Paul Hornung, Meriem Khairoun, Gurbey Ocak.

Writing – review & editing: Paul Hornung, Meriem Khairoun, Friedo W. Dekker, Karin A. H. Kaasjager, Albert Huisman, Lily Jakulj, Willem Jan W. Bos, Frits R. Rosendaal, Marianne C. Verhaar, Gurbey Ocak.

References

1. Weitz JI. Low-molecular-weight heparins. *N Engl J Med*. 1997; 337: 688–98. <https://doi.org/10.1056/NEJM199709043371007> PMID: 9278467
2. Handeland GF, Abildgaard U, Holm HA, Arnesen KE. Dose adjusted heparin treatment of deep venous thrombosis: a comparison of unfractionated and low molecular weight heparin. *Eur J Clin Pharmacol*. 1990; 39: 107–12. <https://doi.org/10.1007/BF00280041> PMID: 2174783
3. Young E, Wells P, Holloway S, Weitz J, Hirsh J. Ex-vivo and in-vitro evidence that low molecular weight heparins exhibit less binding to plasma proteins than unfractionated heparin. *Thromb Haemost*. 1994; 71: 300–4. PMID: 8029793
4. Young E, Cosmi B, Weitz J, Hirsh J. Comparison of the non-specific binding of unfractionated heparin and low molecular weight heparin (Enoxaparin) to plasma proteins. *Thromb Haemost*. 1993; 70: 625–30. PMID: 8115988
5. Matzsch T, Bergqvist D, Hedner U, Ostergaard P. Effects of an enzymatically depolymerized heparin as compared with conventional heparin in healthy volunteers. *Thromb Haemost*. 1987; 57: 97–101. PMID: 3590085
6. Harenberg J. Pharmacology of low molecular weight heparins. *Semin Thromb Hemost*. 1990; 16 Suppl: 12–8.
7. Sanderink GJ, Guimart CG, Ozoux ML, Jariwala NU, Shukla UA, Boutouyrie BX. Pharmacokinetics and pharmacodynamics of the prophylactic dose of enoxaparin once daily over 4 days in patients with renal impairment. *Thromb Res*. 2002; 105: 225–31. [https://doi.org/10.1016/s0049-3848\(02\)00031-2](https://doi.org/10.1016/s0049-3848(02)00031-2) PMID: 11927128
8. Chow SL, Zammit K, West K, Dannenhoffer M, Lopez-Candales A. Correlation of antifactor Xa concentrations with renal function in patients on enoxaparin. *J Clin Pharmacol*. 2003; 43: 586–90. PMID: 12817521
9. Lim W, Dentali F, Eikelboom JW, Crowther MA. Meta-analysis: low-molecular-weight heparin and bleeding in patients with severe renal insufficiency. *Ann Intern Med*. 2006; 144: 673–84. <https://doi.org/10.7326/0003-4819-144-9-200605020-00011> PMID: 16670137
10. Spinler SA, Inverso SM, Cohen M, Goodman SG, Stringer KA, Antman EM, et al. Safety and efficacy of unfractionated heparin versus enoxaparin in patients who are obese and patients with severe renal impairment: analysis from the ESSENCE and TIMI 11B studies. *Am Heart J*. 2003; 146: 33–41. [https://doi.org/10.1016/S0002-8703\(03\)00121-2](https://doi.org/10.1016/S0002-8703(03)00121-2) PMID: 12851605
11. Macie C, Forbes L, Foster GA, Douketis JD. Dosing practices and risk factors for bleeding in patients receiving enoxaparin for the treatment of an acute coronary syndrome. *Chest*. 2004; 125: 1616–21. <https://doi.org/10.1378/chest.125.5.1616> PMID: 15136367
12. Garcia DA, Baglin TP, Weitz JI, Samama MM. Parenteral anticoagulants: Antithrombotic Therapy and Prevention of Thrombosis, 9th ed: American College of Chest Physicians Evidence-Based Clinical Practice Guidelines. *Chest*. 2012; 141: e24S–e43S. <https://doi.org/10.1378/chest.11-2291> PMID: 22315264
13. Dutch Federation of Nephrology. *Anticoagulation with low molecular weight heparin in renal impairment* <https://wwwnefronl/richtlijnen>. 2012.
14. Levey AS, Stevens LA, Schmid CH, Zhang YL, Castro AF III, Feldman HI, et al. A new equation to estimate glomerular filtration rate. *AnnInternMed*. 2009; 150: 604–12. 150/9/604 [pii].
15. Schulman S, Kearon C, Subcommittee on Control of Anticoagulation of the S, Standardization Committee of the International Society on Thrombosis and Haemostasis. Definition of major bleeding in clinical investigations of antihemostatic medicinal products in non-surgical patients. *J Thromb Haemost*. 2005; 3: 692–4. <https://doi.org/10.1111/j.1538-7836.2005.01204.x> PMID: 15842354
16. Hulot JS, Montalescot G, Lechat P, Collet JP, Ankri A, Urien S. Dosing strategy in patients with renal failure receiving enoxaparin for the treatment of non-ST-segment elevation acute coronary syndrome. *Clin Pharmacol Ther*. 2005; 77: 542–52. <https://doi.org/10.1016/j.clpt.2005.02.012> PMID: 15961985
17. Collet JP, Montalescot G, Choussat R, Lison L, Ankri A. Enoxaparin in unstable angina patients with renal failure. *Int J Cardiol*. 2001; 80: 81–2. [https://doi.org/10.1016/s0167-5273\(01\)00455-7](https://doi.org/10.1016/s0167-5273(01)00455-7) PMID: 11575265
18. Collet JP, Montalescot G, Fine E, Golmard JL, Dalby M, Choussat R, et al. Enoxaparin in unstable angina patients who would have been excluded from randomized pivotal trials. *J Am Coll Cardiol*. 2003; 41: 8–14. [https://doi.org/10.1016/s0735-1097\(02\)02664-5](https://doi.org/10.1016/s0735-1097(02)02664-5) PMID: 12570937

19. Kruse MW, Lee JJ. Retrospective evaluation of a pharmacokinetic program for adjusting enoxaparin in renal impairment. *Am Heart J*. 2004; 148: 582–9. <https://doi.org/10.1016/j.ahj.2004.04.015> PMID: 15459586
20. Becker RC, Spencer FA, Gibson M, Rush JE, Sanderink G, Murphy SA, et al. Influence of patient characteristics and renal function on factor Xa inhibition pharmacokinetics and pharmacodynamics after enoxaparin administration in non-ST-segment elevation acute coronary syndromes. *Am Heart J*. 2002; 143: 753–9. <https://doi.org/10.1067/mhj.2002.120774> PMID: 12040334
21. Bazinet A, Almanric K, Brunet C, Turcotte I, Martineau J, Caron S, et al. Dosage of enoxaparin among obese and renal impairment patients. *Thromb Res*. 2005; 116: 41–50. <https://doi.org/10.1016/j.thromres.2004.10.004> PMID: 15850607
22. Siguret V, Pautas E, Fevrier M, Wipff C, Durand-Gasselien B, Laurent M, et al. Elderly patients treated with tinzaparin (Innohep) administered once daily (175 anti-Xa IU/kg): anti-Xa and anti-IIa activities over 10 days. *Thromb Haemost*. 2000; 84: 800–4. PMID: 11127859
23. Siguret V, Gouin-Thibault I, Pautas E, Leizorovicz A. No accumulation of the peak anti-factor Xa activity of tinzaparin in elderly patients with moderate-to-severe renal impairment: the IRIS substudy. *J Thromb Haemost*. 2011; 9: 1966–72. <https://doi.org/10.1111/j.1538-7836.2011.04458.x> PMID: 21819539
24. Pautas E, Gouin I, Bellot O, Andreux JP, Siguret V. Safety profile of tinzaparin administered once daily at a standard curative dose in two hundred very elderly patients. *Drug Saf*. 2002; 25: 725–33. <https://doi.org/10.2165/00002018-200225100-00005> PMID: 12167068
25. Olie RH, Meertens NEL, Henskens YMC, Ten Cate H. Empirically Reduced Dosages of Tinzaparin in Patients with Moderate-to-Severe Renal Insufficiency Lead to Inadequate Anti-Xa Levels. *Nephron*. 2017; 137: 113–23. <https://doi.org/10.1159/000477474> PMID: 28662505
26. Russcher M, Josephus Jitta N, Kraaijenhagen RJ, Fijnheer R, Pasker-de Jong PC, Gaillard CA. Pre-emptive dosage reduction of nadroparin in patients with renal failure: a retrospective case series. *Clin Kidney J*. 2013; 6: 473–7. <https://doi.org/10.1093/ckj/sft083> PMID: 26064511
27. Atiq F, van den Bemt PM, Leebeek FW, van Gelder T, Versmissen J. A systematic review on the accumulation of prophylactic dosages of low-molecular-weight heparins (LMWHs) in patients with renal insufficiency. *Eur J Clin Pharmacol*. 2015; 71: 921–9. <https://doi.org/10.1007/s00228-015-1880-5> PMID: 26071276
28. Boneu B. Low molecular weight heparins: are they superior to unfractionated heparins to prevent and to treat deep vein thrombosis? *Thromb Res*. 2000; 100: V113–20. [https://doi.org/10.1016/s0049-3848\(00\)00331-5](https://doi.org/10.1016/s0049-3848(00)00331-5) PMID: 11053624
29. Schmid P, Fischer AG, Willemin WA. Low-molecular-weight heparin in patients with renal insufficiency. *Swiss Med Wkly*. 2009; 139: 438–52. smw-11284. <https://doi.org/smw-11284> PMID: 19685350
30. Montalescot G, Collet JP, Tanguy ML, Ankri A, Payot L, Dumaine R, et al. Anti-Xa activity relates to survival and efficacy in unselected acute coronary syndrome patients treated with enoxaparin. *Circulation*. 2004; 110: 392–8. <https://doi.org/10.1161/01.CIR.0000136830.65073.C7> PMID: 15249498
31. Levine MN, Planes A, Hirsh J, Goodyear M, Vochelle N, Gent M. The relationship between anti-factor Xa level and clinical outcome in patients receiving enoxaparin low molecular weight heparin to prevent deep vein thrombosis after hip replacement. *Thromb Haemost*. 1989; 62: 940–4. PMID: 2556813
32. Nieuwenhuis HK, Albada J, Banga JD, Sixma JJ. Identification of risk factors for bleeding during treatment of acute venous thromboembolism with heparin or low molecular weight heparin. *Blood*. 1991; 78: 2337–43. PMID: 1657248
33. Walenga JM, Hoppensteadt D, Fareed J. Laboratory monitoring of the clinical effects of low molecular weight heparins. *Thromb Res Suppl*. 1991; 14: 49–62. [https://doi.org/10.1016/0049-3848\(91\)90403-j](https://doi.org/10.1016/0049-3848(91)90403-j) PMID: 1658970
34. Bara L, Leizorovicz A, Picolet H, Samama M. Correlation between anti-Xa and occurrence of thrombosis and haemorrhage in post-surgical patients treated with either Logiparin (LMWH) or unfractionated heparin. Post-surgery Logiparin Study Group. *Thromb Res*. 1992; 65: 641–50. [https://doi.org/10.1016/0049-3848\(92\)90213-t](https://doi.org/10.1016/0049-3848(92)90213-t) PMID: 1319619
35. Witt DM, Nieuwlaat R, Clark NP, Ansell J, Holbrook A, Skov J, et al. American Society of Hematology 2018 guidelines for management of venous thromboembolism: optimal management of anticoagulation therapy. *Blood Adv*. 2018; 2: 3257–91. <https://doi.org/10.1182/bloodadvances.2018024893> PMID: 30482765