



Universiteit
Leiden
The Netherlands

The influence of perioperative interventions targeting psychological distress on clinical outcome after total knee arthroplasty

Sorel, J.C.; Overvliet, G.M.; Gademan, M.G.J.; Haan, C. den; Honig, A.; Poolman, R.W.

Citation

Sorel, J. C., Overvliet, G. M., Gademan, M. G. J., Haan, C. den, Honig, A., & Poolman, R. W. (2020). The influence of perioperative interventions targeting psychological distress on clinical outcome after total knee arthroplasty. *Rheumatology International*, 40(12), 1961-1986. doi:10.1007/s00296-020-04644-y

Version: Publisher's Version

License: [Creative Commons CC BY 4.0 license](#)

Downloaded from: <https://hdl.handle.net/1887/3206807>

Note: To cite this publication please use the final published version (if applicable).

SYSTEMATIC REVIEW



The influence of perioperative interventions targeting psychological distress on clinical outcome after total knee arthroplasty

Juliette Caroline Sorel^{1,2,6} · Geke Marianne Overvliet³ · Maaïke Gerarda Johanna Gademan^{1,4} · Chantal den Haan⁵ · Adriaan Honig³ · Rudolf Wilhelm Poolman^{1,2,6}

Received: 18 April 2020 / Accepted: 3 July 2020 / Published online: 29 July 2020
© The Author(s) 2020

Abstract

Our aim was to assess the effect of perioperative interventions targeting psychological distress on clinical outcome after total knee arthroplasty (TKA). We searched studies on the effect of perioperative interventions focused on psychological distress used in conjunction with TKA on pain, function, and quality of life (QoL) on PubMed, Embase.com, PsycINFO/OVID, CENTRAL, the Cochrane Database of Systematic Reviews, Scopus, and Web of Science. We included 40 studies (22 RCTs, ten cohort studies, and eight quasi-experimental studies) with a total of 3846 patients. We graded the quality of evidence as low for pain and function and as moderate for QoL. Patients receiving music, education, cognitive behavioural therapy, guided imagery, pain coping skills training, Reiki, occupational therapy with self-monitoring, and biofeedback-assisted progressive muscles relaxing training had lower pain scores or declined opioid prescriptions after TKA. Pain coping skills training, audio recording-guided imagery scripts, video promoting self-confidence, psychological therapies by video, Reiki, music, occupational therapy with self-monitoring, education, and psychotherapy improved postoperative functional outcome. Education through an app improved QoL after TKA. The studies in our systematic review show that perioperative interventions targeting psychological distress for patients receiving TKA seem to have a positive effect on postoperative pain, function, and QoL. RCTs with strict methodological safeguards are still needed to determine if perioperative interventions focused on psychological distress should be used in conjunction with TKA. These studies should also assess which type of intervention will be most effective in improving patient-reported outcome measures and declining opioid prescriptions.

Keywords Total knee arthroplasty · Psychological distress · Pain · Function · Quality of life

Electronic supplementary material The online version of this article (<https://doi.org/10.1007/s00296-020-04644-y>) contains supplementary material, which is available to authorized users.

✉ Juliette Caroline Sorel
juliettesorel@hotmail.com

- ¹ Department of Orthopaedics, Leiden University Medical Center, 2333 ZA Leiden, The Netherlands
- ² Department of Orthopaedic Surgery and Traumatology, Joint Research OLVG, 1091 AC Amsterdam, The Netherlands
- ³ Department of Psychiatry, OLVG, Amsterdam and Amsterdam UMC, Jan Tooropstraat 164, 1105 AZ Amsterdam, The Netherlands
- ⁴ Department of Clinical Epidemiology, Leiden University Medical Center, 2333 ZA Leiden, The Netherlands
- ⁵ Medical Library, Department of Research and Education, OLVG, 1061 AE Amsterdam, The Netherlands
- ⁶ Department of Orthopaedic Surgery and Traumatology, OLVG, Oosterpark 9, 1090 HM Amsterdam, The Netherlands

Introduction

Total knee arthroplasty (TKA) is the treatment of choice for medically operable patients with end-stage osteoarthritis (OA) of the knee joint if non-surgical therapies fail to obtain adequate pain relief and functional improvement [1]. TKA proved to be a cost-effective procedure with excellent postoperative implant-related outcomes, such as radiographic appearance and implant features [2]. Nevertheless, a significant number of patients report pain (8.0–26.5%) on long-term follow-up after TKA [3] and as many as 11–19% of the patients are not satisfied with their procedure [4, 5]. Persistent pain after TKA is commonly treated with opioids after surgery [6]. Currently, increasing misuse and addiction to opioids are a rapidly evolving public health issue [7]. Improving pain scores after surgery by understanding factors influencing postoperative pain may help prevent further expansion of this opioid crisis [7].

Unfavourable outcome after TKA is related to age, gender, level of education, pre-operative function and pain [8], comorbidities [9], social support [9], Body Mass Index (BMI) [10], and surgical factors [11–13]. Preoperative psychological factors such as mental health status, symptoms of anxiety and depression, and poor coping skills have also been examined [13–15]. Systematic reviews [16–18] and meta-analyses [19, 20] on this subject reported that psychological distress might affect the postoperative outcome (pain and function) after TKA. Perioperative interventions targeting these psychological factors may improve clinical outcome after surgery. Previous studies have examined the effect of interventions influencing psychological factors to improve postoperative clinical outcome after TKA [21–24]. We found three previous systematic reviews on psychological interventions in conjunction to orthopaedic surgeries [25–27]. The systematic review of Bay et al. [25] did not support the effectiveness of psychological interventions in improving patient-reported joint outcomes after TKA as the interventions explored by studies were found to be ineffective at the latest follow-up. The results of Szeverenyi et al. [26] and Tong et al. [27] indicated that psychological interventions might improve postoperative outcome of orthopaedic surgery. These previous reviews included several types of orthopaedic procedures (among which TKA, total hip arthroplasty (THA) and spinal procedures) and did not focus on TKA. Besides, the most up-to-date search was performed in January 2018 [27].

To our knowledge, focused systematic reviews of studies on TKA patients with wide search and inclusion criteria investigating the effect of interventions targeting psychological distress on patient-reported outcome measures pain, function and/or quality of life (QoL) after surgery have not yet been reported. The aim of our systematic review was to assess the effect of perioperative interventions focused on psychological distress on pain, function and QoL after primary TKA for OA of the knee.

Methods

Search strategy and study selection

We registered our review protocol at PROSPERO international prospective register of systematic reviews (<https://www.crd.york.ac.uk/PROSPERO/>) with reference number CRD42016052466 (https://www.crd.york.ac.uk/prospero/display_record.php?ID=CRD42016052466). We performed this systematic review according to the Preferred Reporting Items for Systematic Reviews and Meta-analysis (PRISMA) statement criteria [28].

We performed the literature search according to the guidance of Gasparyan et al. [29]. A professional medical librarian (CdH) identified therapeutic studies (published articles and abstracts of major conferences) exploring the influence of any type of perioperative (before TKA, during surgery, or during postoperative rehabilitation) interventions targeting psychological distress on postoperative outcome (pain, function, and/or QoL) after TKA by searching PubMed, Embase.com, PsycINFO/OVID, CENTRAL, the Cochrane Database of Systematic Reviews, Scopus and Web of Science from inception up to May 26, 2020.

The following terms, including synonyms and closely related words, were used as index terms or free-text words: ‘total knee arthroplasty’ and ‘psychological intervention’. Full search strategies for all the databases are available in Supplementary Appendix 1. Duplicate articles were excluded.

Selection of articles was limited to adults > 18 years who had undergone a primary total knee replacement for osteoarthritis of the knee. We included different study designs (RCTs, cohorts, quasi-experimental studies) investigating the effect of any intervention targeting psychological distress on postoperative pain, function and/or QoL. Minimum duration of follow-up was not an inclusion criterion with the aim to create a complete overview of all studies that have investigated the effect of perioperative interventions focused on psychological distress on pain, function and/or QoL. Perioperative interventions influencing psychological factors of patients had to be clearly defined. Full-text availability was required. There were no restrictions with respect to language, age, or publication source of the paper.

Exclusion criteria were studies not meeting domain, determinant, or outcome, case reports, descriptive studies (in which there was no control group), non-primary literature studies (letter to the editor, reviews, thesis, expert opinions) and articles with no separated results of patients after TKA and total hip arthroplasty (THA) or other types of surgery if various surgical procedures were analysed.

Main outcome variables

Two authors (JS & GO) independently screened articles for title and abstract and thereafter full text if the abstract potentially met the inclusion criteria. Subsequently, the authors (JS & GO) individually extracted information regarding study design, baseline patient characteristics, baseline clinical findings, follow-up, number of patients initially included in the study, the number of patients available for follow-up and data regarding the primary outcomes of the systematic review. When there was disagreement with respect to data extraction, a third author (AH or RP) could make the final decision.

Quality assessment

We assessed the risk of bias of the included studies using Cochrane Collaboration's tool for assessing the risk of bias [30]. Using this tool, two authors (JS & GO) independently scored six types of bias (selection bias, performance bias, detection bias, attrition bias, reporting bias, and other types of bias) as low, high, or unclear on potential risk of bias [30].

We used the Grading of Recommendation, Assessment, Development, and Evaluation (GRADE) approach to qualify the overall level of evidence of outcome measures pain, function and/or QoL (<https://www.gradeworkinggroup.org/>). Using the GRADEpro software (McMaster University, 2015, available from www.grade.org), we graded the quality of evidence as high, moderate, low, or very low [31].

Data analysis

We arranged the studies according to the type of perioperative intervention (music, education, psychotherapy, and remaining) and collected data of the effect of perioperative interventions targeting psychological distress on post-operative clinical outcome measures pain, function, and QoL. Initially, our intention was to pool data to perform a meta-analysis.

Results

The search strategy and article selection of articles published from 1964 to 26 May 2020 are shown in the flowchart (Fig. 1). Out of 7835 articles remaining after deduplication,




Fig. 1 Search strategy and article selection

we included 40 studies of which 22 RCTs (one randomised controlled pilot study), 10 cohort studies, and 8 quasi-experimental studies with a total number of 3846 patients.

Interventions

A description of the interventions in the experimental and the control groups and the time at which the interventions were applied are presented in Table 1.

Music

Nine studies examined the effect of perioperative listening to music on postoperative outcome. Eight of these studies [32–34, 36–40] assessed the effect of music on pain and three [35, 36, 40] on function. Music was offered at different time points and different types of music were provided.

Education

The effect of education on postoperative outcome was investigated in fifteen studies in which the time of education varied from 12 weeks before surgery to 3 months after surgery (Table 1).

Psychotherapy

Psychological therapies provided with direct support from a professional were examined by eight studies. The patients in the RCTs of Jacobson et al. [61] and Russo et al. [64], who also received psychological therapy, received their psychological intervention by audio recordings, or watching a video instead of direct contact with a health care professional.

Other/remaining interventions

Four remaining interventions (Reiki, biofeedback relaxing training and enhanced reality analgesia, self-monitoring using a diary), applied to six studies, could not be allocated to the music, educational, or psychological therapy intervention groups and were, therefore, classified as remaining interventions (Table 1).

Outcomes

Outcome measures pain, function, and/or QoL were assessed in 22 RCTs (one randomised controlled pilot study), 10 cohort studies, and 8 quasi-experimental studies. Mean age of the patients ranged from 61.7 to 74.1 years and duration of follow-up ranged between 60 min and 2 years. Due to the heterogeneity of the type of studies, interventions, outcome

measures and follow-up there was no possibility to pool data to perform a meta-analysis.

Pain

34 studies examined the influence of a perioperative intervention targeting psychological distress on clinical outcome pain after the TKA. Many different scoring systems were used to score postoperative pain and eight studies assessed pain medication use as an outcome measure for pain (Table 2).

As shown in Table 2, patients in the intervention groups had significant better postoperative pain scores or declined prescriptions of opioids in 20 studies. Therapies applied in these studies were music during surgery [40] or after surgery [33, 36, 38, 39], education [41, 49, 51–53, 55], cognitive behavioural therapy [57, 58], guided imagery [61], pain coping skills training [62], Reiki therapy [66, 70], occupational therapy in combination with self-monitoring using a diary [68], weight-bearing biofeedback training [67] and biofeedback-assisted progressive muscle relaxing training [71]. The remaining 14 studies did not show a significant effect on any of the pain-related outcome measures or pain medication use at the latest follow-up when using a perioperative intervention focused on psychological distress in conjunction to TKA.

Function

A total of 29 studies examined the effect of an intervention targeting psychological distress on function after the TKA (Table 3).

As shown in Table 3, function was significantly improved by perioperative interventions in 18 studies. Pain coping skills training [62], audiorecording guided imagery scripts [61], video promoting self-confidence and psychological support [64], music [35, 36], occupational therapy in combination with self-monitoring using a diary [68], various types of education [41, 43, 45, 47, 51–53], weight-bearing biofeedback training [67], and psychological therapies (behavioural change intervention [60] and cognitive behavioural therapy [57–59]) positively affected any, but not all, of the functional outcome measures after TKA. In the most recent study by Riddle et al. [63], patients receiving pain coping skills training did not have significantly better scores on WOMAC function and the short physical performance battery. Other types of education [42, 44, 48–50, 55], music during physiotherapy [38], enhanced reality analgesia [69], cognitive behavioural therapy delivered by physiotherapists [56], and psychological support from a professional psychologist [23] did also not affect any of the functional outcome measures after TKA.

Table 1 Overview of included studies

Type of intervention	Study	Description of intervention	When was the intervention applied?
Music	Allred [32] Prospective cohort	I: Easy-listening music with headphones for 20 min C: 20-min quiet rest period	Before and after their first ambulation at the first postoperative day
	Aris [33] RCT	I: Additional relaxing music therapy during recovery (<60 beats per minute) C: Usual care	During recovery
	Chen [34] RCT	I: Five compositions of 30 min soothing piano and Chinese violin music (60–80 beats per minute) C: No music	Ward before surgery, in the waiting area of the surgical room and twice during postoperative recovery
	Hsu [35] Prospective cohort	I: Slow relaxing music with slow tempo, low tone and soft melody C: No music, required to rest in bed	Once a day at the 10 a.m. continuous passive motion (CPM) session on the first and second postoperative day
	Hsu [36] Single-group QES	I: Music for 10 min before receiving CPM until the end of the CPM session C: Rest in bed for 10 min before CPM began	During CPM the first and second days after surgery
	Keshmiri [37] RCT	I1: Isolation of noise by soundproof headphones in conjunction to disposable earplugs I2: Music of patients' choice with headphones C: No isolation of noise or music	During surgery, after the effect of sedative (Propofol) was applied
	Leonard [38] RCT	I: Co-treatment session that used live music to support exercise C: Physiotherapy without music	Postsurgery, after admission to the inpatient rehabilitation unit
	Santhna [39] QES	I: Music for five days post-operatively and analgesics C: No music, only pharmacological intervention	5 days postoperatively
	Simcock [40] RCT	I: Music of patients' choice with headphones C: White noise emanating from the headphones	During surgery, after a spinal-epidural anaesthesia and sedation with propofol
	Atabaki [41] RCT	I: Educational intervention presented as a combination of lecture, group discussion, individual education, questions and answers C: Usual care	Four perioperative stages (one day before surgery, 24 h and 48 h later, upon discharge from the hospital)
Education	Aytekin [42] Prospective cohort	I: Education (about OA, joint protection, home safety, and TKA) and home-based exercise C: No additional training program, usual care	During 12 weeks before the operation
	Chen [43] QES	I: Cognitive-behavioural educational intervention (pamphlet, CD and oral instructions) C Routine care and usual instructions delivered orally	Before surgery after hospitalisation and 1 days postsurgery
	Huang [44] RCT	I: 40-min preoperative home rehabilitation education program by a physiotherapist C: No education program	2–4 weeks prior to admission
	Huang [45] RCT	I: Traditional education, telephone education and mobile education C: Traditional face-to-face and telephone education	Following surgery

Table 1 (continued)

Type of intervention	Study	Description of intervention	When was the intervention applied?
Lee [46] RCPS		I1: Psychoeducation on CPSP and prerecorded hypnotic intervention using audiotapes	One delivered before and another delivered at least 24 h after surgery
		I2: Psychoeducation on CPSP and diaphragmatic breathing relaxation exercise	
		C: Usual care	
Lin [47] QES		I: One-to-one less than 30 min preadmission preoperative teaching*	Preadmission preoperative
		C: Postadmission preoperative teaching and no video	
Louw [48] CCTWAA		I: Education program and an additional 30-min group pain neuroscience education session	Before surgery
		C: Only education program	
Malletschek [49] RCT		I: Additional pain psychoeducation over at least 45 min	3–6 days after TKA
Moulton [50] Prospective cohort		C: Usual care	Preoperative for 2 h
		I: Joint school by members of a multidisciplinary group explaining the process of the surgery	
		C: No joint school	
Piva [51] RCT		I: Interactive education to promote physical activity and healthy eating	During 3 months postoperative: 2 lectures during the first postoperative week and mini-sessions of physical activity promotion in the subsequent weeks
		C: No education	
Reslan [52] QES		I: One to one intervention (30–40 min) including education and exercise training by a nurse	Prior to surgery
		C: Standard hospital care	
Timmers [53] RCT		I: Day-to-day postoperative care information related to topics such as pain, physiotherapy exercises, wound care, and daily self-care activities through an application	During the 28-day period after discharge
		C: Only weekly, basic information	
		I: Usual teaching and preoperative educational intervention**	
Wilson [54] RCT		C: Usual teaching	Teaching session and booklet within 4 weeks prior to surgery Phone call during a week before surgery
Yajnik [55] Retrospective cohort		I: Pain management educational card***	Prior to peripheral nerve block placement on the day of surgery, at the time of ward admission by the bedside nurse and once daily during rounds
		C: Before implementation of pain management educational card	

Table 1 (continued)

Type of intervention	Study	Description of intervention	When was the intervention applied?
Psychotherapy	Birch [56] RCT	I: CBT based pain education of approximately 45 min delivered by 2 physiotherapists C: Usual care	3 sessions preoperatively and 4 sessions postoperatively (2 weeks before surgery until 3 months after surgery)
	Cai [57] RCT	I: CBT C: No CBT	After TKA
	Cai [58] RCT	I: Individually tailored CBT by a physiotherapist and a psychologist C: No CBT	During 4 weeks after surgery
	Das Nair [59] RCT	I: 10 sessions of CBT during hour-long sessions by one or two psychologists C: No CBT	During waiting time for surgery
	Harnirattisai [60] QES	I: 25-min sessions of nurse-patient interaction and discussion**** C: No behavioural change intervention	At the fourth postoperative day and two weeks after surgery
	Jacobson [61] RCT	I: 19- to 21- minute audio recordings of guided imagery\$ scripts designed for TKA patients C: Commercially available 17- to 21-min audio recordings	Every day for two weeks before surgery and three weeks after surgery
	Riddle [62] QES	I: Intervention delivered by trained psychologists# C: No intervention	During 8 weekly sessions from approximately one month prior to surgery to one month after surgery
	Riddle [63] RCT	I1: Eight 50-min sessions of 1-on-1 pain coping skills training I2: Eight 50-min sessions of 1-on-1 arthritis education by registered nurses C: Usual care	Approximately 2 weeks preoperatively to approximately 6 weeks postoperatively
	Russo [64] RCT	I: Video according to the Videoinsight Methods^ principles C: No video	Three times a week during the first 3 months after surgery
	Tristaino [23] Prospective cohort	I: Four psychologist-patient sessions of 30 min focusing on defining the psychological themes and concepts on which to focus the activity C: Standard of care	One before surgery, two during postoperative hospital stay and one during rehabilitation

Table 1 (continued)

Type of intervention	Study	Description of intervention	When was the intervention applied?
Remaining	Baldwin [66] RCT	I1: Three or four 30-min Reiki treatments provided by three expert Reiki professionals I2: Standard of care and three or four sham Reiki session delivered by non-trained people C: Standard of care and sessions of “quiet time”	During the hospital stay
	Christiansen [67] RCT	I: Standard of care rehabilitation plus weight bearing biofeedback training C: Standard of care rehabilitation alone	On the morning before surgery (20 min) and after admission to the post anaesthesia care unit (30 min) and 20 min at the first, second and third postoperative day
	Hiraga [68] NRCT	I: Occupational therapy & self-monitoring using a diary C: Occupational therapy only	From 1 to 2 weeks postoperatively
	Koo [69] RCT	I: Enhanced reality analgesia C: No enhanced reality analgesia	Shortly after physiotherapy for 5 times a week, for 2 weeks Shortly after physiotherapy for 5 times a week, for 1 week
	Notte [70] Prospective cohort	I: Weight bearing (WB) biofeedback-assisted progressive muscle relaxation training sessions using a Nintendo Wii fit Plus game and associated Wii balance board C: Standard of care	Twice weekly at home for 6 weeks after surgery
	Wang [71] QES	I: CPM therapy and 30-min biofeedback relaxation training C: Only CPM therapy	One day before surgery and twice a day on the five first postoperative days, concurrent with CPM therapy

I: intervention group, C: control group, RCT randomised controlled trial, CPM continuous passive motion, QES quasi-experimental study, OA osteoarthritis, TKA total knee arthroplasty, CD compact disk, RCT randomized controlled pilot study, CPSP chronic postsurgical pain, CCTWAA controlled clinical trial with alternating allocation, CBT cognitive behavioural therapy, NRCT non-randomised controlled trial

*Preoperative education about care pathway, knee surgery, pain management, expected discharge goals and in-patient and out-patient arthroplasty rehabilitation by an educational nurse and a booklet

**Preadmission preoperative teaching with an instruction booklet during a preoperative outpatient clinic visit. Upon admission to the hospital, they were presented with an educational videotape

***A booklet containing symptom management after TKA, an individual teaching session, and a follow-up support call by the principal investigator

****25-Min sessions of nurse-patient interaction and discussion regarding specific exercises and physical activity, self-monitoring, goal setting, family support and encouragement, and information prompting

§Guided imagery is a widely used mind–body intervention by the generation of self- or practitioner-guided positive sensory and affective mental images to promote health changes in the body, reducing anxiety and stress, and evoking psychological and physiologic relaxation [61]

#Intervention addressed to the recovery of physical function, the concerns during the recovery period and strategies for coping with pain after the operation delivered by trained therapists

^The video was established to produce positive and therapeutic insight, according to the Videoinsight Methods principles [65]

Table 2 The influence of perioperative interventions targeting psychological distress on pain after the TKA

Type of intervention	Study	Nr TKA	Females (%)	Age mean $\pm$ SD	Follow-up	Outcome score (pain)	I score $\pm$ SD	C score $\pm$ SD	Statistically significance at latest follow-up
Music	Allred 2010	T	56	63.9 (64-84)*	6 hours	VAS	41.2 $\pm$ 25.8	45.1 $\pm$ 31.2	$P = 0.337$
		I	28			MPQ	15.9 $\pm$ 10.6	14.9 $\pm$ 12.3	$P = \text{na}$, no statistical analysis between groups
		C	28			Opioid use (morphine or dilauidid)	na	na	$P = 0.388$ and $P = 0.152$ (regarding which oral medication)
	Aris 2019	T	56		60 minutes	VAS	0 (24.39)**	1.5 (32.61)**	$P = 0.045$
		I	28	63.71 $\pm$ 11.005					
		C	28	64.50 $\pm$ 8.851					
	Chen 2015	T	30	68 (53-85)*	Postoperative days	VAS (recovery)	3.22 $\pm$ 0.22***	3.00 $\pm$ 0.25***	$P = 0.50$
		I	15			VAS (ward)	3.07 $\pm$ 0.26***	2.87 $\pm$ 0.18***	$P = 0.53$
		C	15			Opioid use (parental morphine, meperidine, fentanyl in PO recovery)	7.39 $\pm$ 2.66	6.86 $\pm$ 2.29	$P = 0.57$
						Opioid use (parental morphine, meperidine, fentanyl in the ward)	12.04 $\pm$ 14.43	12.90 $\pm$ 8.05	$P = 0.89$
	Hsu 2019	T	49	73.9 $\pm$ 7.5	2 days	NRS	0.06 $\pm$ 0.24	2.14 $\pm$ 1.10	$P < 0.01$
		I	49						
		C	49						
	Keshmiri 2014	T	83	68.7 $\pm$ 0.96	2-7 days	VAS (day 1-3)	1.33 $\pm$ 0.11 (I1) & 1.44 $\pm$ 0.13 (I2)	1.49 $\pm$ 0.13	$P = 0.718$
		I1	28			VAS (day 4-7)	0.9 $\pm$ 0.15 (I1) & 0.81 $\pm$ 0.13 (I2)	1.23 $\pm$ 0.19	$P = 0.330$
		I2	27			VAS (day 17)	1.09 $\pm$ 0.12 (I1) & 1.08 $\pm$ 0.11 (I2)	1.34 $\pm$ 0.14	$P = 0.435$
		C	28			Days of pain catheter duration (type of pain medication na)	3.43 $\pm$ 0.11 (I1) & 3.48 $\pm$ 0.12 (I2)	3.36 $\pm$ 0.19	$P = 0.452$

Table 2 (continued)

Type of intervention	Study	Nr TKA	Females (%)	Age mean $\pm$ SD	Follow-up	Outcome score (pain)	I score $\pm$ SD	C score $\pm$ SD	Statistically significance at latest follow-up
	Leonard 2019	T 32			Postoperative days	NRS	5.44 $\pm$ 3.2	5.56 $\pm$ 2.52	"No significant difference"
		I 16	11 (68.8)	67.9 (45–87)*		Observational coding for pain	3.06 $\pm$ 3.13	2.31 $\pm$ 2.36	$P = 0.02$
		C 16	12 (75)	67.6 (53–80)*					
	Santhna 2015	T 40	14 (70)	63.80 $\pm$ 5.64	5 (days)	PRI	11.78^	29.23^	$P = 0.00$
		I 20	18 (90)	64.90 $\pm$ 6.94					
		C 20				VAS	14.20^	26.80^	$P = 0.00$
						PPI	15.00^	26.00^	$P = 0.001$
						Paracetamol	16000mg^^	17000 mg^^	$P > 0.05$
						Celecoxib	600 mg^^	1600 mg^^	$P > 0.05$
						Tramadol	125 mg^^	225 mg^^	$P > 0.05$
Education	Simcock 2008	T 30	18 (60)	67.3 $\pm$ 9.1	24 hours	VAS (3 hours PO)	3.87 $\pm$ 3.44	1.47 $\pm$ 1.39	$P = 0.01$
		I 15				VAS (6 hours PO)	5.26 $\pm$ 3.04	3.38 $\pm$ 2.48	$P = 0.075$
		C 15				VAS (24 hours PO)	4.03 $\pm$ 2.89	2.41 $\pm$ 1.67	$P = 0.04$
	Atabaki 2019	T 56			6 (weeks)	WOMAC	40.47 $\pm$ 10.47	57.29 $\pm$ 7.51	$P = 0.001$
		I 48	46 (95.8)	65.39 $\pm$ 5.08					
		C 48	41 (85.4)	63.83 $\pm$ 5.14					
	Aytekin 2019	T 44			6 months	VASpr	0.4 $\pm$ 0.9	0.8 $\pm$ 1.1	"no significant difference between groups"
		I 23	18 (78.3)	67.8 $\pm$ 6.3		VASpa	1.5 $\pm$ 1.5	2.3 $\pm$ 2.3	"no significant difference between groups"
		C 21	18 (85.7)	69.7 $\pm$ 6.4		KOOSpain	87.9 $\pm$ 15.4	92.7 $\pm$ 8.3	"no significant difference between groups"
	Chen 2014	T 92	63 (68.5)	69.26 $\pm$ 9.025	5 days	NRS (worst pain)	4.89 $\pm$ 2.82	5.57 $\pm$ 2.84	$P = 0.308$
Huang 2011		I 42				NRS (average pain)	2.38 $\pm$ 1.97	2.43 $\pm$ 2.03	$P = 0.916$
		C 50				NRS (current pain)	2.46 $\pm$ 2.31	2.57 $\pm$ 2.26	$P = 0.836$
		T 242	174 (71.6)	70.2 $\pm$ 7.3	5 days	VAS	2.4 $\pm$ 0.7	2.5 $\pm$ 0.6	$P = 0.686$
		I 125							
		C 117							

Table 2 (continued)

Type of intervention	Study	Nr TKA	Females (%)	Age mean $\pm$ SD	Follow-up	Outcome score (pain)	I score $\pm$ SD	C score $\pm$ SD	Statistically significance at latest follow-up
Louv 2019	T	103			6 (months)	NRS	na	na	$P = 0.386$
		49	32 (65.3%)	74.1 $\pm$ 9.5		Morfine	2601.62 $\pm$ 1103.90	2734.02 $\pm$ 1324.60	$P = 0.635$
		54	23 (51.9)	69.6 $\pm$ 10.6					
	Malletschek 2019	75	47 (62.7)	59-78*	3 months	KOOSpain	na	na	$P = 0.01$
		37							
		38							
Lee 2019	T	24			6 months	NRS	1.40 $\pm$ 0.89 (I1) & 1.73 $\pm$ 1.40 (I2)	2.23 $\pm$ 1.41	HYP vs. control: $P = 0.134$ and $P = 0.038$ (when controlled for covariates)
	I1	8	7 (87.5)	65.63 $\pm$ 9.27					MET vs. Control: $PP = 0.975$
	I2	8	7 (87.5)	56.25 $\pm$ 11.22					
	C	8	8 (100)	67.88 $\pm$ 10.38					
Moulton 2017	T	563	na	70.1 $\pm$ na	2 years	OKS (6 months PO)	28.71 $\pm$ na	31.60 $\pm$ na	$P = 0.251$
	I	503				OKS (2 years PO)	30.17 $\pm$ na	33.26 $\pm$ na	$P = 0.440$
Piva 2017	T	44			6 months	WOMAC pain	min 1.7 (95% CI -3.0,-4.0) ^^^	min 0.3 (95% CI -1.5, 1.0) ^^^	$P = 0.035$
	I	22	31 (70.5)	68.1 $\pm$ 7.5					
	C	22		68.3 $\pm$ 5.5					
	T	60		na	4 weeks	HSSpain	22.83 $\pm$ 4.78	19.18 $\pm$ 5.14	$PP = 0.001$
Reslan 2018	I	30	19 (63.6)						
	C	30	17 (56.7)						
	T	213			4 weeks	NRS at rest	3.45^	4.59^	$PP = 0.001$
Timmers 2019	I	114	74 (64.9)	64.74 $\pm$ 7.57	after discharge	NRS activity	3.99^	5.08^	$P < 0.001$
	C	99	60 (60.6)	65.63 $\pm$ 7.90		NRS at night	4.18^	5.21^	$P = 0.003$
	T	143	89 (62.6)		3 days	BPI-I	24.4 $\pm$ 14.4	22.4 $\pm$ 15.1	$PP = 0.45$
Wilson 2016	I	73		67 $\pm$ 8		NRS (rest)	2.8 $\pm$ 2.5	2.8 $\pm$ 2.7	$P = 0.70$
	C	70		66 $\pm$ 8		NRS (moving)	5.4 $\pm$ 3.0	6.1 $\pm$ 2.5	$P = 0.20$
						NRS worst pain last 24 hours)	7.0 $\pm$ 2.4	7.0 $\pm$ 2.3	$P = 0.87$
						Opioid use (morphine, hydromorphone, oxycodone, codeine)	40 (45)^**	40 (42)^**	"no difference between groups in daily 24-hours opioid administration"

Table 2 (continued)

Type of intervention	Study	Nr TKA	Females (%)	Age mean $\pm$ SD	Follow-up	Outcome score (pain)	I score $\pm$ SD	C score $\pm$ SD	Statistically significance at latest follow-up
Psychotherapy	Yajnik 2018	T	40	68 (46-80)*	2 days	Opioid use (morphine, MME PO day 1 and 2)	38 (1-117)*	72 (32-285)*	$P = 0.001$
		I	20			Minimum pain (patients' verbal rating 0-10) 1 day PO	0 (0-3)*	0 (0-6)*	$P = 0.151$
		C	20			Maximum pain (patients' verbal rating 0-10) 1 day PO	4 (2-9)*	8 (1-10)*	$P = 0.114$
	Birch 2019	T	60		1 (year)	VAS activity	12 (5-18)^^	9 (3-15)^^	$P = NS$
		I	31	66 $\pm$ 9		VAS rest	7 (1-12)^^	6 (1-12)^^	$P = NS$
		C	29	66 $\pm$ 10					
	Cai 2017	T	108		6 months	KSS	82.61 $\pm$ 6.38	73.30 $\pm$ 8.45	$P < 0.01$
		I	54	62.42 $\pm$ 6.59					
		C	54	63.94 $\pm$ 6.58					
	Cai 2018	T	100	62 (55.9)	6 months	NRS	5.63 $\pm$ 0.73	6.27 $\pm$ 0.86	time effects: $P < .001$; group effects: $P = 0.003$; group-by-time interaction: $P = 0.080$
		I	50	65.26 $\pm$ 8.30					
		C	50	66.18 $\pm$ 7.04					

Table 2 (continued)

Type of intervention	Study	Nr TKA	Females (%)	Age mean $\pm$ SD	Follow-up	Outcome score (pain)	I score $\pm$ SD	C score $\pm$ SD	Statistically significance at latest follow-up
Das Nair 2018	T I C	50 25 25	23 (46)	65.7 $\pm$ 8.6 66.7 $\pm$ 9.9		WOMAC pain	6.5 $\pm$ 3.6	7.5 $\pm$ 2.3	$P = 0.40$
						ICOAP constant pain (item 1-5)	6.4 $\pm$ 4.4	6.2 $\pm$ 3.2	$P = 0.99$
						ICOAP constant pain (item 1, 3, 4, 5)	4.8 $\pm$ 3.7	5.1 $\pm$ 3.0	$P = 0.82$
Jacobson 2016	T I C	58 29 29	51 (62.2)	65 (41-81)*	6 months	ICOAP constant pain (converted rasch score item 1, 3, 4, 5)	5.5 $\pm$ 4.1	6.0 $\pm$ 3.2	$P = 0.75$
						ICOAP intermittent pain (item 6-11)	8.5 $\pm$ 5.6	10.2 $\pm$ 4.5	$P = 0.43$
						ICOAP intermittent pain (item 6, 7, 10, 11)	5.7 $\pm$ 3.8	7.1 $\pm$ 3.3	$P = 0.33$
						ICOAP intermittent pain (converted rasch score item 6, 7, 10 11)	5.5 $\pm$ 3.4	6.7 $\pm$ 3.0	$P = 0.34$
						WOMAC pain VAS daily pain	2.7 $\pm$ 3.1 na	3.5 $\pm$ 3.3 na	$P < 0.001$ P not available at 6 months postoperatively
Riddle 2011	T I C	63 18 45	45 (71.4)	63.8 $\pm$ 11.5 60.8 $\pm$ 9.9	2 months	WOMAC pain	6.0 $\pm$ 4.1	8.6 $\pm$ 3.7	$P = 0.017$
Riddle 2019	T	402			12 months	WOMAC pain	3.3 (95% CI 2.5, 4.2) (I1) & 3.0 (95% CI 2.1, 3.8) (I2)^^	2.9 (95% CI 2.0, 3.8)^^	$P = 0.60$
Tristaino 2015	I1 I2 C T I C	130 135 137 64 33 31	94 (72.3) 85 (63.0) 88 (64.2) 44 (62.0)	62.6 $\pm$ 7.9 64.2 $\pm$ 8.5 62.7 $\pm$ 7.7 64.2 $\pm$ 8.6 66.1 $\pm$ 6.6	4 months	NRS	1.8 (95% CI 1.2, 2.4) (I1) & 2.0 (95% CI 1.3, 2.6) (I2)^^	1.7 (95% CI 1.1, 2.2)^^	$P = na$
						SF-36 bodily pain	70.1 $\pm$ 21.5	67.8 $\pm$ 26.8	$P = 0.715$

Table 2 (continued)

Type of intervention	Study	Nr TKA	Females (%)	Age mean $\pm$ SD	Follow-up	Outcome score (pain)	I score $\pm$ SD	C score $\pm$ SD	Statistically significance at latest follow-up
Remaining	Baldwin 2017	T	56	na	72 hours	VAS	na	na	"Reiki significant pain reduction ($P = 0.003$), Sham Reiki and SOC no significant reduction" $P = na$, not mentioned in significant results)
	Hiraga 2019	I1 I2 C	25 12 19			Opioid use (oxycontin, oxycodone, morphine)	na	na	
		T	41		4 weeks	NRS rest	1.3 $\pm$ 0.4	1.2 $\pm$ 0.4	$P = 0.965$
		I	20	76.4 $\pm$ 7.1		NRS walk	1.3 $\pm$ 0.2	3.2 $\pm$ 0.6	$P = 0.017$
		C	21	76.6 $\pm$ 5.5					
	Koo 2018	T	120		5 weeks	VAS	na (figure)	na (figure)	"No significance was found in VAS analyses between the groups"
		I	60	65.00 $\pm$ 6.97			na	na	$P = 0.000$ (1, 2, 3 days PO)
		C	60	63.71 $\pm$ 5.09			na	na	$P = 0.92$
	Notte 2016	T	43	na	3 days postoperatively	NRS	na	na	
		I	23			Opioid use (type of opioid na)	na	na	
		C	20						
	Wang 2015	T	66	73.5 $\pm$ 9.5	5 days	NRS	3.36 $\pm$ 1.47	4.23 $\pm$ 1.67	$P < 0.001$
		I	33			Opioid use (pethidine PO day 5)	1 (3.2)	0 (0.0)	$P = 0.49$
		C	33			PMU (Acetaminophen or COX-2 inhibitor + pethidine or tramadol PO day 5)	24 (77.4)	21 (63.6)	$P = 0.27$

Nr number; TKA total knee arthroplasty; SD standard deviation; I intervention group; C control group; T total study group; VAS visual analog scale; P value; MPQ short form McGill pain questionnaire; na: not available; PO postoperative; NRS numeric rating score; PRI Pain Rating intensity; PPI Present Pain Intensity; mg milligram; WOMAC Western Ontario and McMaster universities osteoarthritis index; VASpr visual analog scale pain resting; VASpa visual analog scale pain activity; KOOSPain pain subscale of the knee injury and osteoarthritis outcome score; HYP hypnotic intervention; MET minimal-effect treatment; OKS Oxford knee score; 95% CI 95% confidence interval; HSS hospital for special surgery; BPI-I Brief Pain Inventory interference; MME Morphine Milligram Equivalents; NS not significant; KSS knee society score; ICOAP Intermittent and Constant Osteoarthritis Pain scale; SF-36 Short Form-36; SOC stand of care; PMU pain medication use; COX-2 cyclooxygenase-2

Instead of mean and SD: *median (range), **median and mean rank, ***mean and standard error, ^mean rank only, ^^median only, ^^mean estimate with the 95% CI in parentheses, ^^median (interquartile range) instead of mean and SD

Table 3 The influence of perioperative interventions targeting psychological distress on function after the TKA

Type of inter- vention	Study	Nr TKA	Females (%)	Age mean ± SD	Follow-up	Outcome score (function)	I score ± SD	C score ± SD	Statistically sig- nificance at latest follow-up	
Music	Hsu [35]	T	91	67 (73.6)	2 days	CPM angles 1 day PO	24.29 ± 5.00	12.98 ± 4.43	P < 0.01	
		I	49	73.9 ± 7.5		CPM angles 2 days PO	21.22 ± 2.98	16.07 ± 4.49	P < 0.01	
		C	42	71.33 ± 8.45		Active knee flexion ROM 2 days PO	106.22 ± 6.17	95.00 ± 6.80	P < 0.01	
	Hsu [36]	T	49		2 days	Increased degree of knee flexion during CPM	21.22 ± 2.98	10.02 ± 3.03	P < 0.01	
		I	49	34 (69.4)	73.9 ± 7.5					
		C	49	34 (69.4)	73.9 ± 7.5					
	Leonard [38]	T	32			Postopera- tive days	Observational cod- ing for pedalling adherence	7.81 ± 0.40	7.44 ± 1.21	"No significant difference"
		I	16	11 (68.8)	67.9 (45–87)*					
		C	16	12 (75)	67.6 (53–80)*					
		T	96			6 weeks	WOMAC stiffness	19.53 ± 12.34	41.66 ± 10.09	P = 0.001
Education	Atabaki [41]	I	48	46 (95.8)		WOMAC perfor- mance difficulty	43.48 ± 7.96	55.82 ± 4.30	P = 0.001	
		C	48	41 (85.4)						
		T	44			6 months	KOOS total	82.2 ± 16.1	85.5 ± 9.5	"No significant difference between groups"
		I	23	18 (78.3)	67.8 ± 6.3		KOOSdaily living activities	87.2 ± 18.3	91.1 ± 9.2	"No significant difference between groups"
		C	21	18(85.7)	69.7 ± 6.4		KOOSsports	52.8 ± 24.4	56.1 ± 13.1	"No significant difference between groups"
		T	92	63 (68.5)	69.26 ± 9.025	5 days	Overall rating of nine physical func- tion items	12.38 ± 2.806	12.05 ± 3.682	P = 0.625
	Chen [43]	I	42				Ankle pumping	1.55 ± 0.39	1.54 ± 0.44	P = 0.927
		C	50				Quadriceps setting	0.17 ± 0.39	0.23 ± 0.43	P = 0.518
							Knee flexion/exten- sion	0.44 ± 0.53	0.69 ± 0.66	P = 0.062
	Huang 2011	T	242	174 (71.6)	70.2 ± 7.3	5 days	Straight-leg raises MPOAL	1.22 ± 2.58 3.71 ± 0.622	0.64 ± 0.56 3.08 ± 1.090	P = 0.000 P = 0.004
I		125				Ability to walk dur- ing discharge	85.7 ± na	81.2 ± na	P = 0.343	
C		117				ROM	76 ± 22	74 ± 20	P = 0.582	

Table 3 (continued)

Type of intervention	Study	Nr TKA	Females (%)	Age mean $\pm$ SD	Follow-up	Outcome score (function)	I score $\pm$ SD	C score $\pm$ SD	Statistically significance at latest follow-up
Huang [45]	T	150	102 (68.0)		3 months	ROM ITT	110.6 $\pm$ 6.68	105.00 $\pm$ 8.82	$P < 0.001$
		I	75	62.42 $\pm$ 6.59		ROM PP	110.0 $\pm$ 6.33	103.26 $\pm$ 7.57	$P < 0.001$
		C	75	63.94 $\pm$ 6.58					
	Lin [47]	T	60	31 (51.7)		EPC	14.93 $\pm$ na	8.87 $\pm$ na	$P < 0.05$
		I	30	68.6 $\pm$ na		Knee flexion	77.84 $\pm$ na	70.16 $\pm$ na	$P = 0.013$
		C	30			Ambulation ability	na	na	"The differences between groups were not significant"
Louw [48]	T	101			6 months	WOMAC	na	na	$P = 0.222$
		I	49	32 (65.3)					
		C	54	23 (51.9)					
	Mallettschek 2019	T	75	47 (62.7)	3 months	KSS	na	na	$P = 0.08$
		I	37	59 – 78**					
		C	38						
Moulton [50]	T	563	na	70.1 $\pm$ na	2 years	OKS (6 months PO)	28.71 $\pm$ na	31.60 $\pm$ na	$P = 0.251$ (6 months)
		I	503						
		C	60						
	Piva [51]	T	44	31 (70.5)	6 months	OKS (2 years PO)	30.17 $\pm$ na	33.26 $\pm$ na	$P = 0.440$ (2 years)
		I	22			SF-36 PF	76.7 $\pm$ 16.1	70.3 $\pm$ 24.2	$P = 0.017$
		C	22			Single-leg stance test	16.1 $\pm$ 9.6	17.4 $\pm$ 9.8	$P = 0.037$
Reslan [52]	T	60				WOMAC PF	11.8 $\pm$ 6.7	12.8 $\pm$ 10.8	$P = 0.558$
		I	30			Stair-climb	14.3 $\pm$ 4.1	15.6 $\pm$ 7.4	$P = 0.054$
		C	30			Chair-stand	12.2 $\pm$ 2.8	13.7 $\pm$ 7.5	$P = 0.149$
						6-Min walk	472.6 $\pm$ 86.5	518.0 $\pm$ 103.3	$P = 0.638$
						Gait speed	1.14 $\pm$ 0.16	1.18 $\pm$ 0.24	$P = 0.790$
						Daily activity	152.5 $\pm$ 93.3	174.9 $\pm$ 126.1	$P = 0.279$
					4 weeks	HSSfunction	15.73 $\pm$ 3.49	13.92 $\pm$ 3.35	$P = 0.026$
						HSSrom	17.04 $\pm$ 2.55	16.53 $\pm$ 4.20	$P = NS$
						HSSquadriceps muscle strength	9.13 $\pm$ 3.81	8.47 $\pm$ 2.93	$P = NS$
						HSSflexion deformity	10.02 $\pm$ 1.21	8.47 $\pm$ 1.93	$P = 0.007$
						HSSinstability	9.89 $\pm$ 3.41	8.27 $\pm$ 2.89	$P = 0.049$
						LEFS	60.35 $\pm$ 11.22	53.83 $\pm$ 12.98	$P = 0.048$

Table 3 (continued)

Type of intervention	Study	Nr TKA	Females (%)	Age mean $\pm$ SD	Follow-up	Outcome score (function)	I score $\pm$ SD	C score $\pm$ SD	Statistically significance at latest follow-up
Psycho-herapy	Timmers [53]	T	213		4 weeks after discharge	KOOS	37.61 $\pm$ 10.17	43.08 $\pm$ 12.96	$P < 0.001$
		I	114	64.74 (7.57)		Ability to perform physiotherapy	7.50***	6.88***	$P = 0.03$
		C	99	65.63 (7.90)		Ability to perform self-care activities	8.32***	7.64***	$P = 0.004$
	Yajnik 2018	T	40	68 (46–80)*	2 days	Maximum ambulation 1 day PO	20 (0–59)^	12 (0–30)^	$P = 0.069$ (PO1)
		I	20			Maximum ambulation 2 days PO	46 (6–67)^	38 (0–61)^	$P = 0.141$ (PO2)
		C	20						
	Birch 2019	T	60		1 year	OKS	33 (29, 27)^	37 (33, 41)^	$P = NS$
		I	31	66 (9)		6-Min walk	441 (402, 480)^	406 (367, 446)^	$P = NS$
		C	29	66 (10)		Sit to stand	12 (11, 14)^	11 (95% CI 10, 13)^	$P = NS$
	Cai [57]	T	108		6 months	KSS	82.61 $\pm$ 6.38	73.30 $\pm$ 8.45	$P < 0.01$
		I	54	62.42 $\pm$ 6.59		First time out of bed (hours)	22.13 $\pm$ 4.18	36.41 $\pm$ 7.31	$P = < 0.001$
		C	54	63.94 $\pm$ 6.58					
	Cai [58]	T	100	62 (55.9)	6 months	HSS function	80.68 $\pm$ 8.02	68.98 $\pm$ 8.64	$P < 0.001$ (time interaction), $P < 0.001$ (group interaction), $P = 0.003$ (group-by-time interaction)
		I	50	65.26 $\pm$ 8.30					
		C	50	66.18 $\pm$ 7.04					
	Das Nair [59]	T	50		6 months	WOMAC function	20.9 $\pm$ 12.7	32.0 $\pm$ 4.8	$P = 0.009$
		I	25	65.7 $\pm$ 8.6		WOMAC stiffness	3.2 $\pm$ 1.9	4.2 $\pm$ 0.9	$P = 0.11$
		C	25	66.7 $\pm$ 9.9					
	Harnirattisai [60]	T	63	59 (93.7)	6 weeks	PTT total	8.86 $\pm$ 1.89	6.43 $\pm$ 1.66	$P = na$
		I	42	67.88 (60–85)*		PPT standing balance	Δ 2.00 $\pm$ 1.22^	Δ 1.09 $\pm$ 1.22^	$P = 0.016$
		C	21			PPT walking speed	Δ 1.55 $\pm$ 1.02^	Δ 0.76 $\pm$ 0.83^	$P = 0.004$
		T				PPT chair-stand	Δ 2.36 $\pm$ 1.05^	Δ 1.33 $\pm$ 1.02^	$P < 0.001$
		I				ADL and daily requirements exercise activity	na	na	"There were no significant differences in ADL participation"
		C							
		T							
		I							
		C							

Table 3 (continued)

Type of intervention	Study	Nr TKA	Females (%)	Age mean $\pm$ SD	Follow-up	Outcome score (function)	I score $\pm$ SD	C score $\pm$ SD	Statistically significance at latest follow-up
Jacobson [61]	T	58	51 (62.2)	65 (41–81)*	6 months	SF-36 physical	50.4 $\pm$ 6.0	47.3 $\pm$ 7.5	$P = na$
	I	29				WOMAC stiffness	1.9 $\pm$ 1.4	2.1 $\pm$ 1.9	$P = na$
	C	29				WOMAC function	7.2 $\pm$ 7.1	10.2 $\pm$ 10.5	$P = na$
Riddle [62]	T	63	45 (71.4)			Gait velocity	na	na	$P = 0.0154$
	I	18				Timed walk in seconds	7.4 $\pm$ 2.2	8.5 $\pm$ 2.3	(group-by-imaging ability interaction) $P = na$
	C	45		63.8 $\pm$ 11.5 60.8 $\pm$ 9.9	2 months	WOMAC disability	18.3 $\pm$ 12.2	24.1 $\pm$ 10.9	$P = 0.023$ (for differences among discharge scores for the 2 groups after adjusting for baseline differences) $P > 0.05$
Riddle [63]	T	402			12 months	WOMAC function	11.7 (8.6, 14.9) (I1) & 12.2 (9.0, 15.4) (I2)^^^	10.5 (7.4, 13.6)^^^	$P > 0.05$
	I1	130	94 (72.3)	62.6 $\pm$ 7.9		SPPB	8.0 (7.2, 8.7) (I1) & 8.4 (7.6, 9.1) (I2)^^^	8.6 (95% CI 7.8, 9.4)^^^	$P > 0.05$
	I2	135	85 (63.0)	64.2 $\pm$ 8.5					
Russo 2016	C	137	88 (64.2)	62.7 $\pm$ 7.7					
	T	110	na	69.1 $\pm$ na	3 months	SF-36 physical	45.6 $\pm$ 8.3	46.2 $\pm$ 9.9	$P > 0.01$
	I	55				KSS	87.8 $\pm$ 9.6	78.3 $\pm$ 8.2	$P = < 0.005$
Tristaino 2015	C	55				WOMAC	79.9 $\pm$ 13.0	69.7 $\pm$ 9.5	$P = < 0.005$
						VAS functional score	2.8 $\pm$ 1.6	4.0 $\pm$ 1.5	$P = < 0.005$
	T	44	44 (62.0)		4 months	SF-36 PCS	49.5 $\pm$ 6.6	50.9 $\pm$ 9.8	$P = 0.5114$
	I	33		64.2 $\pm$ 8.6		Days until physiotherapy objective reached	8.1 $\pm$ 2.4	8.8 $\pm$ 2.3	$P = 0.2424$
	C	31		66.1 $\pm$ 6.6					

Table 3 (continued)

Type of intervention	Study	Nr TKA	Females (%)	Age mean $\pm$ SD	Follow-up	Outcome score (function)	I score $\pm$ SD	C score $\pm$ SD	Statistically significance at latest follow-up
Remaining	Christiansen 2015	T I C	26 13 13	13 (50)	26 weeks	FTSST Hip moment (Nm/kg) during FTSST Knee moment (Nm/kg) during FTSST Ankle moment (Nm/kg) during FTSST Walking speed (m/s) Hip moment during walking Knee extension moment during walking Ankle moment during walking	9.5 $\pm$ 2.4 0.65 $\pm$ 0.24 1.03 $\pm$ 0.22 0.17 $\pm$ 0.16 1.29 $\pm$ 0.25 0.28 $\pm$ 0.19 0.61 $\pm$ 0.25 0.09 $\pm$ 0.29	9.6 $\pm$ 1.6 0.63 $\pm$ 0.20 0.97 $\pm$ 0.11 0.24 $\pm$ 0.14 1.24 $\pm$ 0.13 0.36 $\pm$ 0.22 0.42 $\pm$ 0.44	<i>P</i> = 0.21 <i>P</i> = 0.686 <i>P</i> = 0.434 <i>P</i> = 0.227 <i>P</i> = 0.68 <i>P</i> = 0.160 <i>P</i> = 0.008
	Hiraga [68]	T I C	41 20 21	16 (80) 19 (90.4)	4 weeks	Daily step count Psychical activity time	3580.5 $\pm$ 1545.2 1741.4 $\pm$ 551.3	2088.4 $\pm$ 2008.3 731.8 $\pm$ 321.1	<i>P</i> = 0.041 <i>P</i> = 0.000
	Koo [69]	T I C	120 60 60	17 (28.3)	5 weeks	WOMAC Graded ambulation distance	14.59 $\pm$ 9.14 na	10.86 $\pm$ 10.84 na	<i>P</i> = 0.398 <i>P</i> = na
						6-Min walk test Timed-stand test	407.00 $\pm$ 83.62 19.29 $\pm$ 2.80	353.35 $\pm$ 82.35 19.00 $\pm$ 6.16	<i>P</i> = 0.163 <i>P</i> = 0.967

Nr Number, TKA total knee arthroplasty, SD standard deviation, I intervention group, C control group, T total study group, CPM continuous passive motion, PO postoperative, P *P* value, ROM range of motion, WOMAC Western Ontario and McMaster Universities osteoarthritis index, KOOS Knee Injury and Osteoarthritis Outcome Score, MPOAL muscle power of the affected leg, ITT intention to treat, PP per protocol, na not available, EPC exercises performance checklist, KSS Knee Society Score, OKS Oxford knee score, SF-36 PF Short Form-36 physical functioning, HSS hospital for special surgery knee score, NS not significant, LEFS lower extremity functional scale, POD postoperative day, PPT physical performance test, ADL activities of daily living, SPFB short physical performance battery, VAS visual analog scale, PCS physical component scale, FTSST five-time sit-to-stand test, Nm/kg Newtonmeter/kilogram, m/s metre per second

*Mean (range)

**Range only

***Mean only

^Median (10th–90th percentiles)

^^Mean estimate with the 95% CI parentheses

^^^Mean change score baseline—6 weeks postoperative

QoL

Two recent studies [49, 53] examined the effect a perioperative intervention on QoL (Table 4). Patients receiving postoperative day-to-day education through an app seemed to report significantly better QoL compared to patients who received usual care [53]. Additional psychoeducation did not significantly improve QoL [49].

Quality assessment

Figure 2 shows our risk of bias assessment of the included studies. Figure 3 represents our judgement about each risk of bias item presented as percentages across all studies. The most prevalent shortcomings regarding the risk of bias were inadequate blinding participants and/or personnel during the study (performance bias) and “other types of bias”. Bias due to inadequate generation of a randomisation sequence or inadequate allocation concealment prior to assignment (selection bias) also caused high scores on the risk of bias (Fig. 3).

The overall level of evidence of the studies using the GRADE approach was qualified as low for pain and for function and as moderate for QoL. Serious uncertainty in the assessment of the risk of bias, inconsistency, and indirectness were the main reasons for downgrading the overall level of evidence (Table 5).

Discussion

In this systematic review, we give an overview of studies that assessed the effect of perioperative interventions targeting psychological distress on pain, function, and QoL applied to patients undergoing TKA for primary OA of the knee. Perioperative music [33, 36, 38–40], education [41, 49, 51–53, 55], cognitive behavioural therapy [57, 58], pain coping skills training [62], guided imagery [61], perioperative Reiki therapy [66, 70], occupational therapy in combination with self-monitoring using a diary [68], and biofeedback-assisted progressive muscle relaxing training [71] seem to improve postoperative pain or to decline opioid prescriptions after TKA. For function, pain coping skills training [62], audio-recording guided imagery scripts [61], video promoting self-confidence and psychological support [64], music [35, 36], occupational therapy in combination with self-monitoring using a diary [68], various types of education [41, 43, 45, 47, 51–53], weight-bearing biofeedback training [67], psychological therapies (behavioural change intervention [60] and cognitive behavioural therapy [57–59]) seem to significantly improve at least one postoperative functional outcome measure. Day-to-day education after TKA using an app might improve postoperative QoL.

Table 4 The influence of perioperative interventions targeting psychological distress on QoL after the TKA

Type of intervention	Study	Nr TKA	Females (%)	Age mean $\pm$ SD	Follow-up	Outcome score (QoL)	I score $\pm$ SD I score $\pm$ SD	C score $\pm$ SD	Statistically significance at latest follow-up
Education	Malletscheck 2019	T	75	59–78*	3 months	KOOS QoL	na	na	P = NS
		I	37						
		C	38						
	Timmers 2019	T	213	64.74 $\pm$ 7.57 65.63 $\pm$ 7.90	4 weeks after discharge	EQ-5D	0.76 $\pm$ 0.16	0.67 $\pm$ 0.25	P < 0.001
		I	114						
		C	99						

Nr Number, TKA total knee arthroplasty, SD standard deviation, QoL quality of life, I intervention group, C control group, T total study group, KOOS Knee Injury and Osteoarthritis Outcome Score, na not available, P P value, NS not significant, EQ-5D EuroQOL Five-Dimensional Questionnaire

Instead of mean and SD

*Range

Fig. 2 Risk of bias summary. Authors' judgements about each risk of bias item for each included study. Green: low risk of bias. Red: high risk of bias. No fill: unclear risk of bias

	Random sequence generation (selection bias)	Allocation concealment (selection bias)	Blinding of participants and personnel (performance bias)	Blinding of outcome assessment (detection bias)	Incomplete outcome data (attrition bias)	Selective reporting (reporting bias)	Other bias
Allred 2010	+	+	+		+		+
Aris 2019	+		+		+	+	+
Atabaki 2019			+	+	+	+	+
Aytekin 2019		+	+	+	+	+	+
Baldwin 2017	+	+	+		+	+	+
Birch 2019	+		+	+	+	+	+
Cai 2017	+				+		+
Cai 2018	+	+	+	+	+		+
Chen 2014	+	+	+	+		+	+
Chen 2015	+	+	+	+	+	+	+
Christiansen 2015	+	+	+	+	+	+	+
Das Nair 2018	+	+	+	+	+	+	+
Harnirattisai 2005	+	+	+	+	+		+
Hiraga 2019	+	+			+	+	+
Hsu 2016	+	+	+	+	+	+	
Hsu 2019	+		+	+	+	+	+
Huang 2011	+	+	+		+		
Huang 2017	+				+		
Jacobson 2016	+	+	+	+	+	+	+
Keshmiri 2014		+	+	+	+		
Koo 2018	+	+		+	+	+	+
Lee 2019	+	+	+	+	+		
Leonard 2019	+	+	+	+	+		+
Lin 1997	+	+	+	+	+	+	
Louw 2019	+	+			+	+	+
Malletschek 2019	+	+	+		+	+	
Moulton 2017	+	+	+	+	+	+	+
Notte 2016	+		+	+	+	+	+
Piva 2017	+	+	+	+	+	+	+
Reslan 2018	+	+	+	+	+		+
Riddle 2011	+	+	+	+	+	+	+
Riddle 2019	+	+	+	+	+	+	+
Russo 2016	+				+		
Santhna 2015	+	+	+	+	+	+	+
Simcock 2008	+	+	+	+	+	+	+
Timmers 2019	+	+			+	+	+
Tristaino 2015	+	+	+		+	+	+
Wang 2015	+	+	+		+	+	+
Wilson 2016	+	+	+	+	+	+	+
Yajnik 2018	+	+	+	+	+		+

This is a methodologically well-conducted systematic review for which a professional medical librarian (CdH) has developed the search strategy to conduct a comprehensive search in several databases to identify eligible studies. Two authors (JS & GO) performed the screening, data extraction, risk of bias assessment, and overall level of evidence grading independently. We have created a complete overview of all studies by minimizing our exclusion criteria regarding study design, minimum follow-up, and language. Studies without significant results on the effect of an intervention

are often refused for publication. Due to the heterogeneity of the outcome measures of the included studies, it was not possible to conduct a funnel plot to assess this type of bias (publication bias) in our systematic review. However, we included multiple studies [32–34, 38, 39, 42, 46, 55, 56, 68] with small sample sizes (smaller than 30 patients) with no significant results on both outcome measures pain and function. Therefore we assume the risk of publication bias to be low.

Fig. 3 Risk of bias graph. Authors' judgements about each risk of bias item presented as percentages across all included studies. Green: low risk of bias. Red: high risk of bias. No fill: unclear risk of bias




Table 5 The overall level of evidence using the GRADE approach

Certainty assessment							No of patients		Certainty
No of studies	Study design	Risk of bias	Inconsistency	Indirectness	Imprecision	Other considerations	ITPD	No ITPD	
Pain (follow up: range 60 min to 6 months; assessed with: Various outcome measures)									
34	19 randomised trials and 15 remaining*	Serious	Serious	Serious	Not serious	all plausible residual confounding would suggest spurious effect, while no effect was observed	1618	996	⊕⊕○○ low
Function (follow up: range 2 days to 2 years; assessed with: Various outcome measures)									
29	16 randomised trials and 13 remaining**	Serious	Serious	Serious	Not serious	all plausible residual confounding would suggest spurious effect, while no effect was observed	1580	1003	⊕⊕○○ low
QoL (follow up: range 24 weeks to 3 months; assessed with: Various outcome measures)									
2	1 randomised trial and one non-randomised trial	Serious	Serious	Not serious	Not serious	all plausible residual confounding would suggest spurious effect, while no effect was observed	151	137	⊕⊕⊕○ moderate

GRADE grading of recommendation, assessment, development, and evaluation, *N* number, ITPD intervention targeting psychological distress, QoL quality of life

* 8 prospective cohort studies, 6 quasi-experimental studies, 1 retrospective cohort study

** 6 prospective cohort studies, 6 quasi-experimental studies, 1 retrospective cohort study

Unfortunately, drawing meaningful conclusions from the included studies was hampered. First of all, there was a substantial heterogeneity with respect to study design, analysis, domain, interventions, and outcome measures, which precluded pooling for a meta-analysis. Second, according to the GRADE approach, we have graded the quality of evidence as low for outcome measures pain and function. Therefore, the true effect of the interventions targeting psychological

distress on postoperative pain and function may be different from our estimate of the effect.

The previous systematic reviews of Szeverenyi et al. [26] and Tong et al. [27] concluded that psychological interventions seem to reduce postoperative side effects and anxiety and to improve recovery and mental components of quality of life after orthopaedic surgeries. However, Szeverenyi et al. [Szeverenyi] did not clarify the type of orthopaedic

procedures (only joint replacement or no joint replacement) and Tong et al. [27] included several orthopaedic procedures (THA, TKA, and spinal procedures) of which only two studies [61, 63] represented separated data of patients undergoing TKA. The findings of our review do not support the earlier systematic review of Bay et al. [25], in which most interventions explored by the included studies were found to be ineffective on patient-reported outcome after THA and TKA. Only three studies with patients receiving TKA were included by Bay et al. [25]. Compared to that review, we included fifteen additional RCTs [33, 34, 37, 38, 41, 44, 45, 49, 53, 54, 56–58, 58, 63]. Second, due to the current lack of RCTs on one specific type of intervention focused on psychological distress (for example only pain coping skills training) applied to patients undergoing TKA, we have decided to also include a wider range of study designs to create a complete overview of the perioperative interventions focused on psychological distress that have been used to decrease pain and improve function and/or QoL after surgery. Besides, ten studies [32, 34, 37, 39, 48, 54, 55, 66, 70, 71] in our systematic review evaluated the degree of postoperative pain not only by measuring pain scores, but also by assessing postoperative prescription of opioids or other types of pain medication. Investigating alternative nonpharmacologic methods to reduce postoperative pain and opioid use may help prevent further expansion of opioid misuse and addiction, which is currently a rapidly evolving public health crisis [7].

To the best of our knowledge, except for the mentioned systematic reviews [25, 26], no other systematic reviews or meta-analysis with comparable objectives have been published. Therefore, this is the first systematic review with wide search and inclusion criteria focused on TKA patients investigating the effect of interventions focused on psychological distress on patient-reported outcome measures pain, function, and QoL after surgery. Unfortunately, our review also highlighted the limitations of current literature on this subject. To avoid heterogeneity of outcome measures between studies, we would discourage the use of different questionnaires to assess patient-reported outcome measures (PROMs) in orthopaedic research. The reliability and reproducibility of the EuroQOL Five-Dimensional Questionnaire (EQ-5D) and the responsiveness of the Patient-Reported Outcomes Measurement Information System (PROMIS) Global Health survey have been well validated for patients undergoing TKA [72]. We would, therefore, recommend the use of the EQ-5D and PROMIS to allow tracking and evaluation of the effectiveness of perioperative interventions for psychological distress in conjunction with TKA in the following studies [72].

Conclusions

The studies included in our systematic review show the positive effect of multiple perioperative interventions targeting psychological distress for patients receiving TKA to improve postoperative pain (or to decline prescriptions of opioids), function, and QoL. RCTs with strict methodological safeguards (such as long-term follow-up, large number of patients participating in the study, low risk of bias) prospectively comparing outcome for patients with and without perioperative support are still needed to determine if perioperative interventions targeting psychological distress should be used in conjunction with primary TKA for OA of the knee. These studies should also assess which type of intervention will be most effective in improving patient-reported outcome measures and declining opioid prescriptions in the future.

Acknowledgements "We are pleased to acknowledge Qiukui Hao, Chinese associate professor from the West China Hospital, Sichuan University, China, for the performance of the data-extraction and quality assessment of two Chinese articles [45, 57]."

Compliance with ethical standard

Conflict of interest Author Juliette Caroline Sorel, Geke Marianne Overvliet, Maaïke Gerarda Johanna Gademan, Chantal den Haan declare that they have no conflicts of interest. Author Adriaan Honig reports personal fees from NOV-Dutch Orthopaedic Society, grants from The Netherlands Organisation for Health Research and Development (in Dutch: ZonMw), other from LINK/LIMA, other from Stryker, personal fees from LINK, personal fees from BMJ, non-financial support from LINK, grants from Achmea Healthcare Foundation (in Dutch Stichting Achmea Gezondheidszorg fonds), grants from Dutch health insurances (Zorgverzekeraars Nederland), grants from Foundation of medical research OLVG, Amsterdam, the Netherlands, grants from Van Rens Foundation, grants from Reuma Nederland, other from McMaster University, outside the submitted work.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Skou ST, Roos EM, Laursen MB et al (2015) A randomized, controlled trial of total knee replacement. *N Engl J Med* 373:1597–1606. <https://doi.org/10.1056/NEJMoa1505467>

2. Jenkins PJ, Clement ND, Hamilton DF, Gaston P, Patton JT, Howie CR (2013) Predicting the cost-effectiveness of total hip and knee replacement: a health economic analysis. *Bone Joint J* 95:115–121. <https://doi.org/10.1302/0301-620X.95B1.29835>
3. Beswick AD, Wylde V, Gooberman-Hill R, Blom A, Dieppe P (2012) What proportion of patients report long-term pain after total hip or knee replacement for osteoarthritis? A systematic review of prospective studies in unselected patients. *BMJ Open*. <https://doi.org/10.1136/bmjopen-2011-000435>
4. Baker P, van der Meulen J, Lewsey J, Gregg P (2007) Nation joint registry for England and Wales. The role of pain and function in determining patient satisfaction after total knee replacement. Data from the National Joint Registry for England and Wales. *J Bone Joint Surg Br* 89:893–900. <https://doi.org/10.1302/0301-620X.89B7.19091>
5. Bourne RB, Chesworth BM, Davis AM, Mahomed NN, Charron KD (2010) Patient satisfaction after total knee arthroplasty: who is satisfied and who is not? *Clin Orthop Relat Res* 468:57–63. <https://doi.org/10.1007/s11999-009-1119-9>
6. Bedard NA, Pugely AJ, Westermann RW, Duchman KR, Glass NA, Callaghan JJ (2017) Opioid use after total knee arthroplasty: trends and risk factors for prolonged use. *J Arthroplasty* 32:2390–2394. <https://doi.org/10.1016/j.arth.2017.03.014>
7. Volkow ND, Collins FS (2017) The Role of science in addressing the opioid crisis. *N Engl J Med* 377:391–394. <https://doi.org/10.1056/NEJMs1706626>
8. Alattas SA, Smith T, Bhatti M, Wilson-Nunn D, Donell S (2017) Greater pre-operative anxiety, pain and poorer function predict a worse outcome of a total knee arthroplasty. *Knee Surg Sports Traumatol Arthrosc* 25:3403–3410. <https://doi.org/10.1007/s00167-016-4314-8>
9. Harmelink KEM, Zeegers AVCM, Hullegie W, Hoogbeem TJ, Nijhuis-van der Sanden MWG, Staal JB (2017) Are there prognostic factors for one-year outcome after total knee arthroplasty? a systematic review. *J Arthroplasty* 32:3840–3853. <https://doi.org/10.1016/j.arth.2017.07.011>
10. Sun K, Li H (2017) Body mass index as a predictor of outcome in total knee replace: A systemic review and meta-analysis. *Knee* 24:917–924. <https://doi.org/10.1016/j.knee.2017.05.022>
11. Núñez M, Núñez E, del Val JL et al (2007) Health-related quality of life in patients with osteoarthritis after total knee replacement: factors influencing outcomes at 36 months of follow-up. *Osteoarthritis Cartilage* 15:1001–1007. <https://doi.org/10.1016/j.joca.2007.02.019>
12. Papakostidou I, Dailiana ZH, Papapolychroniou T et al (2012) Factors affecting the quality of life after total knee arthroplasties: a prospective study. *BMC Musculoskelet Disord* 13:116. <https://doi.org/10.1186/1471-2474-13-116>
13. Heck DA, Robinson RL, Partridge CM, Lubitz RM, Freund DA (1998) Patient outcomes after knee replacement. *Clin Orthop Relat Res* 356:93–110. <https://doi.org/10.1097/00003086-19981000-00015>
14. Lingard EA, Katz JN, Wright EA, Sledge CB (2004) Kinemax Outcomes Group. Predicting the outcome of total knee arthroplasty. *J Bone Joint Surg Am* 86:2179–2186. <https://doi.org/10.2106/00004623-200410000-00008>
15. Roth JS, Buehler KC, Shen J, Naughton M (2013) Patient factors predict functional outcomes after cruciate retaining TKA: a 2-year follow-up analysis. *J Arthroplasty* 28:1321–1326. <https://doi.org/10.1016/j.arth.2013.01.009>
16. Khatib Y, Madan A, Naylor JM, Harris IA (2015) Do psychological factors predict poor outcome in patients undergoing TKA? A systematic review. *Clin Orthop Relat Res* 473:2630–2638. <https://doi.org/10.1007/s11999-015-4234-9>
17. Burns LC, Ritvo SE, Ferguson MK, Clarke H, Seltzer Z, Katz J (2015) Pain catastrophizing as a risk factor for chronic pain after total knee arthroplasty: a systematic review. *J Pain Res* 8:21–32. <https://doi.org/10.2147/JPR.S64730>
18. Hernández C, Díaz-Heredia J, Berraquero ML, Crespo P, Loza E, Ruiz Ibán MÁ (2015) Pre-operative predictive factors of post-operative pain in patients with hip or knee arthroplasty: a systematic review. *Reumatol Clin* 11:361–380. <https://doi.org/10.1016/j.reuma.2014.12.008>
19. Lewis GN, Rice DA, McNair PJ, Kluger M (2015) Predictors of persistent pain after total knee arthroplasty: systematic review and meta-analysis. *Br J Anaesth* 114:551–561. <https://doi.org/10.1093/bja/aeu441>
20. Sorel JC, Veltman ES, Honig A, Poolman RW (2019) The influence of preoperative psychological distress on pain and function after total knee arthroplasty: a systematic review and meta-analysis. *Bone Joint J*. <https://doi.org/10.1302/0301-620X.101B1.BJJ-2018-0672.R1>
21. Clarke HD, Timm VL, Goldberg BR, Hattrup SJ (2012) Preoperative patient education reduces in-hospital falls after total knee arthroplasty. *Clin Orthop Relat Res* 470:244–249. <https://doi.org/10.1007/s11999-011-1951-6>
22. Gordon D, Malhas A, Goubiran A, Subramanian P, Messer C, Houlihan-Burne D (2011) Implementing the rapid recovery program in primary hip and knee arthroplasty in a UK state run hospital. *Eur J Orthop Surg Traumatol* 21:151–158. <https://doi.org/10.1007/s00590-010-0690-9>
23. Tristaino V, Lantieri F, Tornago S, Gramazio M, Carriere E, Camera A (2016) Effectiveness of psychological support in patients undergoing primary total hip or knee arthroplasty: a controlled cohort study. *J Orthopaed Traumatol* 17:137–147. <https://doi.org/10.1007/s10195-015-0368-5>
24. White L, Stockwell T, Hartnell N, Henessy M, Mullan J (2016) Factors preventing kneeling in a group of pre-educated patients post total knee arthroplasty. *J Orthopaed Traumatol* 17:333–338. <https://doi.org/10.1007/s10195-016-0411-1>
25. Bay S, Kuster L, McLean N, Byrnes M, Kuster MS (2018) A systematic review of psychological interventions in total hip and knee arthroplasty. *BMC Musculoskeletal Disorders* 19:201. <https://doi.org/10.1186/s12891-018-2121-8>
26. Szeverenyi C, Kekecs Z, Johnson A, Elkins G, Csernatony Z, Varga K (2018) The use of adjunct psychosocial interventions can decrease postoperative pain and improve the quality of clinical care in orthopedic surgery: A systematic review and meta-analysis of randomized controlled trials. *J Pain* 19(11):1231–1252. <https://doi.org/10.1016/j.jpain.2018.05.006>
27. Tong F, Dannaway J, Enke O, Eslick G (2020) Effect of preoperative psychological interventions on elective orthopaedic surgery outcomes: a systematic review and meta-analysis. *ANZ J Surg* 90(3):230–236. <https://doi.org/10.1111/ans.15332>
28. Moher D, Liberati A, Tetzlaff J, Altman DG, The PRISMA Group (2009) Preferred reporting items for systematic reviews and meta-analyses: The PRISMA statement. *BMJ* 339:b2535. <https://doi.org/10.1371/journal.pmed.1000097>
29. Gasparyan AY, Ayvazyan L, Blackmore H, Kitas GD (2011) Writing a narrative biomedical review: considerations for authors, peer reviewers, and editors. *Rheumatol Int* 31(11):1409–1417. <https://doi.org/10.1007/s00296-011-1999-3>
30. Higgins JPT, Altman DG, Sterne JAC (editors). Chapter 8: Assessing risk of bias in included studies. In: Higgins JPT, Green S (editors). *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.1.0 (updated March 2011). The Cochrane Collaboration, 2011. Available from www.handbook.cochrane.org
31. Guyatt G, Oxman AD, Aki EA et al (2011) GRADE guidelines: 1. Introduction-GRADE evidence profiles and summary of findings tables. *J Clin Epidemiol* 64:383–394. <https://doi.org/10.1016/j.jclinepi.2010.04.026>

32. Allred KD, Byers JF, Sole ML (2010) The effect of music on postoperative pain and anxiety. *Pain Manag Nurs* 11:15–25. <https://doi.org/10.1016/j.pmn.2008.12.002>
33. Aris A, Sulaiman S, Che Hasan MK (2019) The influence of music therapy on mental well-being among postoperative patients of total knee arthroplasty (TKA). *Enferm Clin* 29(2):16–23. <https://doi.org/10.1016/j.enfcli.2019.04.004>
34. Chen HJ, Chen TY, Huang CY, Hsieh YM, Lai HL (2015) Effects of music on psychophysiological responses and opioid dosage in patients undergoing total knee replacement surgery. *Jpn J Nurs Sci* 12:309–319. <https://doi.org/10.1111/jjns.12070>
35. Hsu CC, Chen WM, Chen SR, Tseng YT, Lin PC (2016) Effectiveness of music listening in patients with total knee replacement during CPM rehabilitation. *Biol Res Nurs* 18:68–75. <https://doi.org/10.1177/1099800415572147>
36. Hsu CC, Chen SR, Lee PH, Lin PC (2019) The effect of music listening on pain, heart rate variability, and range of motion in older adults after total knee replacement. *Clin Nurs Res* 28(5):529–547. <https://doi.org/10.1177/1054773817749108>
37. Keshmiri A, Wolf T, Wiech O, Benditz A, Grifka J, Springorum H (2014) Einfluss der intraoperativen Schallprotektion auf postoperative Schmerzen. *Schmerz* 28:82–89. <https://doi.org/10.1007/s00482-013-1368-0>
38. Leonard H (2019) Live music therapy during rehabilitation after total knee arthroplasty: a randomized controlled trial. *J Music Ther* 56:61–89. <https://doi.org/10.1093/jmt/thy022>
39. Santhna LP, Norhamdan MY, Damrudi M (2015) The effectiveness of music therapy for post-operative pain control among total knee replacement patients. *Med & Health* 10(1):66–79
40. Simcock XC, Yoon RS, Chalmers P, Geller JA, Kiernan HA, Macaulay W (2008) Intraoperative music reduces perceived pain after total knee arthroplasty: a blinded, prospective, randomized, placebo-controlled clinical trial. *J Knee Surg* 21:275–278. <https://doi.org/10.1055/s-0030-1247831>
41. Atabaki S, Farahani MA, Haghani S (2019) Effect of rehabilitation education on pain, knee stiffness and performance difficulty in patients undergoing knee replacement surgery: A randomized clinical trial. *J Acute Dis* 8(6):233–238. <https://doi.org/10.4103/2221-6189.272854>
42. Aytekin E, Sukur E, Oz N et al (2019) The effect of a 12 week prehabilitation program on pain and function for patients undergoing total knee arthroplasty: A prospective controlled study. *J Clin Orthop Trauma* 10:345–349. <https://doi.org/10.1016/j.jcot.2018.04.006>
43. Chen SR, Chen CS, Lin PC (2014) The effect of educational intervention on the pain and rehabilitation performance of patients who undergo a total knee replacement. *J Clin Nurs* 23:279–287. <https://doi.org/10.1111/jocn.12466>
44. Huang S-W, Chen P-H, Chou Y-H (2012) Effects of a preoperative simplified home rehabilitation education program on length of stay of total knee arthroplasty patients. *Orthop Traumatol Surg Res* 98:259–264. <https://doi.org/10.1016/j.otsr.2011.12.004>
45. Huang P, He J, Zhang YM (2017) The mobile application of patient management in education and follow-up for patients following total knee arthroplasty. *Zhonghua Yi Xue Za Zhi* 97:1592–1595. <https://doi.org/10.3760/cma.j.issn.0376-2491.2017.20.019>
46. Lee JK, Zubaidah JO, Fadhlil IS, Normala I, Jensen MP (2019) Prerecorded hypnotic peri-surgical intervention to alleviate risk of chronic postsurgical pain in total knee replacement: a randomized controlled pilot study. *Int J Clin Exp Hypn* 67:217–245. <https://doi.org/10.1080/00207144.2019.1580975>
47. Lin PC, Lin LC, Lin JJ (1997) Comparing the effectiveness of different educational programs for patients with total knee arthroplasty. *Orthop Nurs* 16(5):43–49
48. Louw A, Puenteadura EJ, Reed J, Zimney K, Grimm D, Landers MR (2019) A controlled clinical trial of preoperative pain neuroscience education for patients about to undergo total knee arthroplasty. *Clin Rehabil* 33(11):1722–1731. <https://doi.org/10.1177/0269215519857782>
49. Mallettschek A, Tiller D, Wohlrab D (2020) Psychoedukation bei Patienten mit Knieendoprothese : Erweitertes Schmerzmanagement [Psychoeducation in knee arthroplasty patients : Additional pain management]. *Orthopade* 49(1):26–31. <https://doi.org/10.1007/s00132-019-03749-y>
50. Moulton LS, Evans PA, Starks I, Smith T (2017) Preoperative education prior to elective knee arthroplasty surgery does not change patient outcomes. *Musculoskeletal Care* 15:341–344. <https://doi.org/10.1002/msc.1177>
51. Piva SR, Almeida GJ, Gil AB, DiGioia AM, Helsel DL, Sowa GA (2017) Effect of comprehensive behavioral and exercise intervention on physical function and activity participation after total knee replacement: a pilot randomized study. *Arthritis Care Res (Hoboken)* 69(12):1855–1862. <https://doi.org/10.1002/acr.23227>
52. Reslan HA, Moustafa SM, Saghih S, Sharaza ES, Badr LK (2018) Does intervention improve the outcomes of patients after total knee replacement surgery? *Int J Ortho Trauma Nurs* 31:26–31. <https://doi.org/10.1016/j.ijotn.2018.08.001>
53. Timmers T, Janssen L, van der Weegen W, Das D, Marijnissen WJ, Hannink G, van der Zwaard BC, Plat A, Thomassen B, Swen JW, Kool RB, Lambers Heerspink FO (2019) the effect of an app for day-to-day postoperative care education on patients with total knee replacement: randomized controlled trial. *JMIR Mhealth Uhealth* 7(10):e15323. <https://doi.org/10.2196/15323>
54. Wilson RA, Watt-Watson J, Hodnett E, Tranmer J (2016) A randomized controlled trial of an individualized preoperative education intervention for symptom management after total knee arthroplasty. *Orthop Nurs* 35:20–29. <https://doi.org/10.1097/NOR.0000000000000210>
55. Yajnik M, Hill JN, Hunter OO et al (2019) Patient education and engagement in postoperative pain management decreases opioid use following knee replacement surgery. *Patient Educ Couns* 102:383–387. <https://doi.org/10.1016/j.pec.2018.09.001>
56. Birch S, Stilling M, Mechlenburg I, Hansen TB (2020) No effect of cognitive behavioral patient education for patients with pain catastrophizing before total knee arthroplasty: a randomized controlled trial. *Acta Orthop.* 91(1):98–103. <https://doi.org/10.1080/17453674.2019.1694312>
57. Cai LB, Liu YJ, Zhao H, Xu HP, Gao HH, Dong YZ (2017) Cognitive behavior therapy alleviates kinesophobia after total knee arthroplasty. *Zhongguo Zuzhi Gongcheng Yanjiu* 21:3658–3663
58. Cai L, Gao H, Xu H, Wang Y, Liu P, Liu Y (2018) Does a program based on cognitive behavioral therapy affect kinesophobia in patients following total knee arthroplasty? a randomized, controlled trial with a 6-month follow-up. *J Arthroplasty* 33:704–710. <https://doi.org/10.1016/j.arth.2017.10.035>
59. Das Nair R, Mhizha-Murira JR, Anderson P et al (2018) Home-based pre-surgical psychological intervention for knee osteoarthritis (HAPPiKNEES): a feasibility randomized controlled trial. *Clin Rehabil* 32:777–789. <https://doi.org/10.1177/0269215518755426>
60. Harnirattisai T, Johnson RA (2005) Effectiveness of a behavioral change intervention in Thai elders after knee replacement. *Nurs Res* 54:97–107. <https://doi.org/10.1097/00006199-200503000-00004>
61. Jacobson AF, Umberger WA, Palmieri PA, Alexander TS, Myerscough RP, Draucker CB, Steudte-Schmiedgen S, Kirschbaum C (2016) Guided imagery for total knee replacement: a randomized, placebo-controlled pilot study. *J Altern Complement Med* 22:563–575. <https://doi.org/10.1089/acm.2016.0038>
62. Riddle DL, Keefe FJ, Nay WT, McKee D, Attarian DE, Jensen MP (2011) Pain coping skills training for patients with elevated pain catastrophizing who are scheduled for knee arthroplasty: a

- quasi-experimental study. *Arch Phys Med Rehabil* 92:859–865. <https://doi.org/10.1016/j.apmr.2011.01.003>
63. Riddle DL, Keefe FJ, Ang DC et al (2019) Pain coping skills training for patients who catastrophize about pain prior to knee arthroplasty: a multisite randomized clinical trial. *J Bone Joint Surg Am* 6(101):218–227. <https://doi.org/10.2106/JBJS.18.00621>
 64. Russo LR, Benedetti MG, Mariani E, Roberti di Sarsina T, Zaffagnini S (2017) The videoinstight® method: improving early results following total knee arthroplasty. *Knee Surg Sports Traumatol Arthrosc* 25:2967–2971. <https://doi.org/10.1007/s00167-016-4118-x>
 65. Zaffagnini S, Russo LR, Marcheggiani Muccioli GM, Marcacci M (2013) The Videoinstight® method: improving rehabilitation following anterior cruciate ligament reconstruction—a preliminary study. *Knee Surg Sports Traumatol Arthrosc* 21:851–858. <https://doi.org/10.1007/s00167-013-2392-4>
 66. Baldwin AL, Vitale A, Brownell E, Kryak E, Rand W (2017) Effects of reiki on pain, anxiety, and blood pressure in patients undergoing knee replacement: a pilot study. *Holist Nurs Pract* 31:80–89. <https://doi.org/10.1097/HNP.0000000000000195>
 67. Christiansen CL, Bade MJ, Davidson BJ, Dayton MR, Stevens-Lapsley JE (2015) Effects of weight-bearing biofeedback training on functional movement patterns following total knee arthroplasty: a randomized controlled trial. *J Orthop Sports Phys Ther* 45:647–655. <https://doi.org/10.2519/jospt.2015.5593>
 68. Hiraga Y, Hisano S, Nomiyama K, Hirakawa Y (2019) Effects of using activity diary for goal setting in occupational therapy on reducing pain and improving psychological and physical performance in patients after total knee arthroplasty: A non-randomised controlled study. *Hong Kong J Occup Ther*. 32(1):53–61. <https://doi.org/10.1177/1569186119849117>
 69. Koo K, Park DK, Youm YS, Cho SD, Hwang CH (2018) Enhanced reality showing long-lasting analgesia after total knee arthroplasty: prospective. *Random Clin Trial Sci Rep* 8:2343. <https://doi.org/10.1038/s41598-018-20260-0>
 70. Notte BB, Fazzino C, Mooney RA (2016) Reiki's effect on patients with total knee arthroplasty: A pilot study. *Nursing* 46:17–23. <https://doi.org/10.1097/01.NURSE.0000476246.16717.65>
 71. Wang TJ, Chang CF, Lou MF et al (2015) Biofeedback relaxation for pain associated with continuous passive motion in Taiwanese patients after total knee arthroplasty. *Res Nurs Health* 38:39–50. <https://doi.org/10.1002/nur.21633>
 72. Shim J, Hamilton DF (2019) Comparative responsiveness of the PROMIS-10 Global Health and EQ-5D questionnaires in patients undergoing total knee arthroplasty. *Bone Joint J* 101-B:832–837. <https://doi.org/10.1302/0301-620X.101B7.BJJ-2018-1543.R1>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.