



Universiteit  
Leiden  
The Netherlands

## **Anyonic, cosmic, and chaotic: three faces of Majorana fermions**

Cheipesh, Y.I.

### **Citation**

Cheipesh, Y. I. (2022, November 17). *Anyonic, cosmic, and chaotic: three faces of Majorana fermions*. *Casimir PhD Series*. Retrieved from <https://hdl.handle.net/1887/3487143>

Version: Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/3487143>

**Note:** To cite this publication please use the final published version (if applicable).

# Stellingen

behorende bij het proefschrift

*Anyonic, cosmic, and chaotic: three faces of Majorana fermions*

1. A vanishing fermion parity in a superconductor fusion experiment can not become a distinctive signature of an isolated Majorana zero-mode. [chapter 2].
2. The absence of quasiparticles in SYK model manifests itself in a linear in temperature dependence of the tunneling current between it and the fermion bath. [chapter 5].
3. Many-body effects in the nano-device present in the experimental setup for the relic neutrino detection impose fundamental intrinsic limitation on its energy resolution. [chapter 6].
4. The detection of relic neutrinos in the  $\beta$ -decay spectrum of  ${}^3\text{H}$  is fundamentally impossible. The only viable isotope for neutrino detection is  ${}^{171}\text{Tm}$ . [chapter 7].
5. Any gauge-invariant lattice theory of Dirac fermions which preserves chiral symmetry must have a vanishing topological index.
6. Entanglement entropy is not a good measure of the complexity of the fermionic system.
7. There is a  $k$ -body basis independent measure of fermionic complexity  $C(k)$  that is zero for a product state, exponentially decreasing for a Fermi liquid state and  $\delta(k - k^*)$  with  $k^* \propto$  volume of the system for a chaotic state.
8. If one couples two fermionic systems, their internal energies will grow at timescales of the Fermi time  $\tau_F \sim 1/\varepsilon_F$  independently of their initial temperatures.

Yevheniia Cheipesh

27 november 2022