



Universiteit
Leiden

The Netherlands

Traffic accident victims and polytrauma patients: injury patterns, outcome and their influencing factors

Leijdesdorff, H.A.

Citation

Leijdesdorff, H. A. (2022, November 24). *Traffic accident victims and polytrauma patients: injury patterns, outcome and their influencing factors*. Retrieved from <https://hdl.handle.net/1887/3487305>

Version: Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/3487305>

Note: To cite this publication please use the final published version (if applicable).

PART I

INJURY PATTERNS AND OUTCOME



Injury Pattern and Injury Severity of In-Hospital
Deceased Road Traffic Accident Victims in The
Netherlands: Dutch Road Traffic Accidents Fatalities.

H.A. Leijdesdorff
S. Gillissen
I.B. Schipper
P. Krijnen

World Journal of Surgery, 2020.

ABSTRACT

Background

Further reduction of road traffic accident (RTA) fatalities is a key priority in the European Union. Since data on injury patterns related to mortality in RTAs is scarce, this study aimed to analyse injury patterns and injury severity of in-hospital RTA fatalities in the Netherlands.

Methods

All in-hospital deceased RTA victims in the Netherlands during the period 2015 - 2016 were analysed. Data were obtained from the National Trauma Registry. Injury patterns, injury severity, accident- and patient characteristics of road user groups were compared.

Results

A total of 497 deceased RTA victims were analysed, of which most were bicyclists. All analysed motorcyclists had an $ISS \geq 16$. Head trauma was most frequent in pedestrians (73.7%) and bicyclists (71.3%). Thorax trauma was most frequent in motorcyclists and motorists (respectively 60.9% and 65.8%). RTA victims younger than 25 years were more severely injured (median ISS 38, interquartile range [IQR] 29-46) compared to RTA victims aged over 75 years (median ISS 25, IQR 13-30). More than 10% of the severely injured ($ISS \geq 16$) RTA victims were not transported to a level I trauma centre. The majority of this group was older than 75 years.

Conclusions

Further prevention of head trauma is needed to reduce RTA fatalities, especially in bicyclists. Also, under-triage of severe trauma in elderly RTA victims is obvious and should be addressed in the early phases of trauma care, especially during pre-hospital triage and initial care at admission.

INTRODUCTION

Road traffic accidents (RTA) contribute significantly to mortality in both developing and developed countries around the world. RTAs are the main cause of death among the young population (aged 15-29 years). The World Health Organization indicates that the number of RTA fatalities continues to climb, reaching a high of 1.35 million worldwide in 2016 [1]. Despite traffic safety laws and extensive preventive governmental programs, the number of seriously injured RTA victims in the Netherlands has increased by 3% per year since 2006, resulting in 21,300 seriously injured RTA victims in 2015 [2].

The number of RTA fatalities in the Netherlands has declined since the 1990's partly due to the introduction of the systematic approach to trauma patients according to the Advanced Trauma Life Support (ATLS®) [2,3]. However, this number has stabilized at a higher annual average of 620 fatalities in 2015, which was significantly higher than expected based on the trend seen in previous years (2006 till 2014) [2,4,5].

Further decline in RTA fatalities is one of the key priorities of the "Road Safety Program 2011 - 2020" initiated by the European Commission, which aims to reduce the number of RTA victims in the Netherlands to less than 10.600 seriously injured victims and 500 fatalities by 2020 [6]. Therefore, we analysed the injury patterns and injury severity for different types of road users involved in fatal RTAs in The Netherlands. These data may contribute to the awareness of specific high-risk road user groups and potentially lethal injuries during trauma admissions.

METHODS

Study design and patients

This study is a retrospective analysis of all adult (≥ 18 years) RTA victims who died in Dutch hospitals within 30 days after acute admission, during the period 2015 and 2016. All data was obtained from the National Trauma Registry and RTA victims who died at the accident scene or during transport to the hospital could not be included, as they are not registered in the trauma registry. The study was exempted from ethics review board approval because the study used coded data from the existing National Trauma Registry.

Data and definitions

Patient data comprised age, gender, road user group (motorists, motorcyclists, moped riders (incl. mobility scooters), bicyclists (incl. E-bikes) and pedestrians), trauma mechanism (blunt or penetrating trauma), mode of transportation to the hospital (am-

balance, helicopter emergency medical service, self-presenting), hospital of admission (Level I, level-II or level-III trauma centres), season (winter: December-February; spring: March-May; summer: June-August; autumn: September-November) and time (morning: 08.00-12.00; afternoon: 12.00-17.00; evening: 17.00-00.00; night: 00.00-8.00).

Using the Abbreviated Injury Scale update 2008 (AIS 2008) [7], injury characteristics with respect to severity distribution were classified for each anatomic region of the body (head, face, neck, thorax, abdomen, spine, upper extremities, lower extremities, and external) [7]. An AIS ≥ 3 was deemed severe injury for the respective system. Ultimately, the total trauma burden was quantified using the Injury Severity Score (ISS) [8], and ISS ≥ 16 was classified as severe trauma. Vital parameters on arrival included the Revised Trauma Score (RTS) [9] and Glasgow Coma Scale (GCS) [10]. Outcomes of care included length of hospitalization in days and admission to the intensive care unit (ICU).

The continuous variables were categorized using clinically relevant cut off points used by the National Trauma Registry for age (18-24, 25-55, 56-75 and >75 yrs.), GCS (3-8, 9-12, 13-15), respiratory rate (<10, 10-29, >29 per minute) and systolic blood pressure (<75, 75-89, >89 mmHg).

ANALYSIS

The data were statistically analysed using IBM SPSS Statistics for Windows, version 23 (IBM Corp., Armonk, N.Y., USA). Patient groups were compared using the chi-squared test for categorical data, the T-test or ANOVA for normally distributed continuous data and the Mann-Whitney test or Kruskal-Wallis test for skewed continuous data.

RESULTS

A total of 548 RTA victims died in Dutch hospitals between January 1, 2015, and December 31 2016, of which 51 were excluded from analysis due to age restrictions and/or unknown road user group. The average age of the remaining 497 deceased RTA victims was 63.8 years and 68.6% were male.

Patient and accident characteristics

There were statistically significant differences in age and gender between the road user groups (Table 1). Deceased bicyclists (n=209, 42%) were the largest group and had the highest average age (71.1, standard deviation [SD] 16.3 years), with 45.9% older

Table 1. Patient and accident characteristics of 497 in hospital deceased road traffic accident victims per road user group.

Characteristics	Total (n=497)	Motorists (n=149)	Motorcyclists (n=23)	Moped riders (n=59)	Bicyclists (n=209)	Pedestrians (n=57)	p-value
Male, n (%)	341 (68.6)	108 (72.5)	22 (95.7)	47 (79.7)	139 (66.5)	25 (43.9)	<0.0001
Age (years), mean (SD)	63.8 (21.7)	56.2 (25.3)	42.2 (13.9)	63.4 (21.8)	71.1 (16.3)	65.9 (19.0)	<0.0001
By category, n (%)							<0.0001
18-24	42 (8.5)	30 (20.1)	2 (8.7)	4 (6.8)	5 (2.4)	1 (1.8)	
25-55	111 (22.3)	35 (23.5)	16 (69.6)	15 (25.4)	28 (13.4)	17 (29.8)	
56-75	153 (30.8)	32 (21.5)	5 (21.7)	20 (33.9)	80 (38.3)	16 (28.1)	
>75	191 (38.4)	52 (34.9)	0 (0.0)	20 (33.9)	96 (45.9)	23 (40.4)	
Trauma mechanism, n (%)							0.85
Blunt	496 (99.8)	149 (30)	23 (4.6)	59 (11.9)	208 (41.9)	57 (11.5)	
Penetrating	1 (0.2)	0	0	0	1 (100)	0	
Season accident, n (%)							0.001
Winter	132 (26.6)	45 (30.2)	1 (4.3)	12 (20.3)	48 (23.0)	26 (45.6)	
Spring	116 (23.3)	36 (24.2)	7 (30.4)	18 (30.5)	44 (21.1)	11 (19.3)	
Summer	130 (26.2)	30 (20.1)	12 (52.2)	18 (30.5)	59 (28.2)	11 (19.3)	
Autumn	119 (23.9)	38 (25.5)	3 (13.0)	11 (18.6)	58 (27.8)	9 (15.8)	
Time of arrival in hospital, n (%)							0.07
Morning	67 (13.5)	16 (10.8)	4 (17.4)	7 (12.1)	32 (15.3)	8 (14.0)	
Afternoon	189 (38.2)	49 (33.1)	4 (17.4)	20 (34.5)	93 (44.5)	23 (40.4)	
Evening	187 (37.8)	60 (40.5)	12 (52.2)	22 (37.9)	72 (34.4)	21 (36.8)	
Night	52 (10.5)	23 (15.5)	3 (13.0)	9 (15.5)	12 (5.7)	5 (8.8)	
Mode of transportation to hospital, n (%)							0.63
Ambulance	448 (92.4)	133 (92.4)	19 (82.6)	55 (94.8)	190 (92.7)	51 (92.7)	
HEMS	31 (6.4)	9 (6.3)	4 (17.4)	2 (3.4)	13 (6.3)	3 (5.5)	
Self-presenting	6 (1.2)	2 (1.4)	0	1 (1.7)	2 (1.0)	1 (1.8)	

SD: standard deviation; HEMS: helicopter emergency medical service.

than 75 years. Deceased motorcyclists were the youngest road user group (mean 42.2 years, SD 13.9) consisting mostly of men (95.7%). A statistically significant difference between road user groups was also found for the season, as most fatal accidents with motorcyclists occurred in summer (52.2%), whereas nearly half of the fatal accidents with pedestrians took place in winter (45.6%). No differences between road user groups were found for the time of the accident and the mode of transportation to the hospital. All but one of the patients had a blunt trauma mechanism (Table 1).

Table 2. Clinical characteristics of in hospital deceased road traffic accident victims per road user group.

Characteristics	Total (n=497)	Motorists (n=149)	Motorcyclists (n=23)	Moped riders (n=59)	Cyclists (n=209)	Pedestrians (n=57)	p-value
<i>AIS region (AIS≥3), n (%)</i>							
Head	314 (63.2)	74 (49.7)	14 (60.9)	35 (59.3)	149 (71.3)	42 (73.7)	<0.0001
Face	23 (4.6)	6 (4.0)	1 (4.3)	3 (5.1)	9 (4.3)	4 (7.0)	0.92
Neck	13 (2.6)	4 (2.7)	4 (17.4)	1 (1.7)	2 (2.0)	2 (3.5)	<0.0001
Thorax	224 (45.1)	98 (65.8)	18 (78.3)	23 (39.0)	66 (31.6)	19 (33.3)	<0.0001
Abdomen	37 (7.4)	18 (12.1)	8 (34.8)	2 (3.4)	7 (3.3)	2 (3.5)	<0.0001
Spine	48 (9.7)	23 (15.4)	5 (21.7)	5 (8.5)	12 (5.7)	3 (5.3)	0.006
Upper Extremities	4 (0.8)	0 (0.0)	1 (4.3)	0 (0.0)	1 (0.5)	2 (3.5)	0.03
Lower Extremities	110 (22.1)	36 (24.2)	11 (47.8)	8 (13.6)	41 (19.6)	14 (24.6)	0.01
External	19 (3.8)	17 (11.4)	0 (0.0)	1 (1.7)	0 (0.0)	1 (1.8)	<0.0001
ISS, median (IQR)	27 (18-38)	34 (22-43)	43 (34-59)	25 (16-33)	26 (16-33)	29 (21-34)	<0.0001
<i>By ISS category, n (%)</i>							
ISS <16	89 (17.9)	18 (12.1)	0 (0.0)	14 (23.7)	47 (22.5)	10 (17.5)	0.01
ISS ≥ 16	407 (82.1)	131 (87.9)	22 (100.0)	45 (76.3)	162 (77.5)	47 (82.5)	
<i>RTS, n (%)</i>							
12	100 (28.3)	30 (30.0)	2 (14.3)	13 (31.7)	45 (28.8)	10 (23.8)	0.31
11	45 (12.7)	13 (13.0)	0 (0.0)	9 (22.0)	18 (11.5)	5 (11.9)	
<11	208 (58.9)	57 (57.0)	12 (85.7)	19 (46.3)	93 (59.6)	27 (64.3)	
<i>GCS, n (%)</i>							
13-15	142 (30.7)	42 (30.9)	2 (9.5)	21 (37.5)	63 (32.3)	14 (25.9)	0.09
9-12	38 (8.2)	6 (4.4)	1 (4.8)	7 (12.5)	17 (8.7)	7 (13.0)	
3-8	282 (61.0)	88 (64.7)	18 (85.7)	28 (50.0)	115 (59.0)	33 (61.1)	
<i>Respiratory rate, n (%)</i>							
<10	36 (9.4)	21 (18.4)	2 (12.5)	6 (13.6)	5 (3.0)	2 (4.3)	<0.0001
10-29	327 (85.2)	82 (71.9)	14 (87.5)	35 (79.5)	154 (93.9)	42 (91.3)	
>29	21 (5.5)	11 (9.6)	0 (0.0)	3 (6.8)	5 (3.0)	2 (4.3)	
<i>SBP, n (%)</i>							
<75	45 (10.0)	26 (20.2)	3 (15.0)	6 (10.9)	10 (5.2)	0 (0.0)	<0.0001
75-89	34 (7.6)	10 (7.8)	4 (20.0)	2 (3.6)	15 (7.7)	3 (6.0)	
>89	369 (82.4)	93 (72.1)	13 (65.0)	47 (85.5)	169 (87.1)	47 (94.0)	
LOHS (days), median (IQR)	3 (2-8)	2 (1-8)	2 (1-3)	4 (2-10)	3 (2-9)	4 (2-12)	0.01
ICU admittance, n (%)	344 (70.3)	98 (66.2)	17 (73.9)	39 (67.2)	148 (72.9)	42 (73.7)	0.63

AIS: abbreviated injury score; ISS: injury severity score; IQR: interquartile range; RTS: revised trauma score; GCS: Glasgow Coma Scale;

ICU: intensive care unit; SBP: Systolic blood pressure; LOHS: Length of hospital stay.

Clinical characteristics

Injury patterns were strikingly different between the groups of deceased road users (Table 2). Motorcyclists were the most severely injured road user group and all were diagnosed with severe trauma ($ISS \geq 16$), with a median ISS of 43 (interquartile range [IQR] 34-59). This group had the shortest survival (median hospital stay 2 days, IQR 1-13). Severe head injury occurred in the majority of each group of deceased road users but was most prevalent in bicyclists and pedestrians (71.3% and 73.7% respectively). Severe thorax trauma was most seen in motorcyclists and motorists (respectively 78.3% and 65.8%) but was also common in moped riders, bicyclists and pedestrians (30-40%). Severe trauma of the lower extremities, abdomen and spine was most often seen in motorcyclists (47.8%, 34.8% and 21.7%). Significant differences between road user groups were found for respiratory rate and systolic blood pressure upon arrival at the Emergency Department, but no differences were found for GCS, RTS and ICU admission (Table 2).

Table 3. Clinical characteristics of 497 in hospital deceased road traffic accident victims per age category.

Characteristics	Total (n=497)	18-24 yrs. (n=42)	25-55 yrs. (n=111)	56-75 yrs. (n=153)	>75 yrs. (n=191)	p-value
<i>AIS region (AIS\geq3), n (%)</i>						
Head	314 (63.2)	31 (73.8)	83 (74.8)	99 (64.7)	101 (52.9)	0.001
Face	23 (4.6)	6 (14.3)	5 (4.5)	3 (2.0)	9 (4.7)	0.01
Neck	13 (2.6)	2 (4.8)	7 (6.3)	3 (2.0)	1 (0.5)	0.02
Thorax	224 (45.1)	26 (61.9)	58 (52.3)	62 (40.5)	78 (40.8)	0.02
Abdomen	37 (7.4)	8 (19.0)	12 (10.8)	8 (5.2)	9 (4.7)	0.004
Spine	48 (9.7)	4 (9.5)	18 (16.2)	14 (9.2)	12 (6.3)	0.05
Upper Extremities	4 (0.8)	1 (2.4)	1 (0.9)	1 (0.7)	1 (0.5)	0.67
Lower Extremities	110 (22.1)	11 (26.2)	17 (15.3)	34 (22.2)	48 (25.1)	0.22
External	19 (3.8)	7 (16.7)	6 (5.4)	3 (2.0)	3 (1.6)	<0.0001
ISS, Median (IQR)	2718-38)	38 (29-46)	33 (25-43)	26 (17-36)	25 (13-30)	<0.0001
<i>By ISS category, n (%)</i>						
ISS <16	89 (17.9)	1 (2.4)	7 (6.4)	24 (15.7)	57 (29.8)	<0.0001
ISS \geq 16	407 (82.1)	41 (97.6)	103 (93.6)	129 (84.3)	134 (70.2)	
<i>GCS, n (%)</i>						
13-15	142 (30.7)	2 (5.4)	8 (7.6)	37 (26.4)	95 (52.8)	<0.0001
9-12	38 (8.2)	0 (0.0)	3 (2.9)	17 (12.1)	18 (10.0)	
3-8	282 (61.0)	35 (94.6)	94 (89.5)	86 (61.4)	67 (37.2)	
<i>RTS, n (%)</i>						
12	100 (28.3)	0 (0.0)	10 (12.8)	25 (23.1)	65 (44.8)	<0.0001
11	45 (12.7)	0 (0.0)	1 (1.3)	17 (15.7)	27 (18.6)	
<11	208 (58.9)	22 (100.0)	67 (85.9)	66 (61.1)	53 (36.6)	
ICU admission, n (%)	344 (70.3)	28 (66.7)	84 (76.4)	123 (80.9)	109 (58.9)	<0.0001
LOHS in days, median (IQR)	3 (2-8)	2 (1-2)	2 (1-5)	4 (2-10)	4 (2-10)	<0.0001

AIS: abbreviated injury score; ISS: injury severity score; IQR: interquartile range; RTS: revised trauma score; GCS: Glasgow Coma Scale; ICU: intensive care unit; LOHS: Length of hospital stay.

In all age groups, severe head trauma and severe thorax trauma were the most common. A decrease in presence of severe trauma to the head, face, neck, thorax, abdomen and severe external injuries was seen with increasing age (Table 3). Median ISS also significantly decreased with increasing age. Deceased RTA victims aged 18-24 years were the most severely injured age group with a median ISS of 38 (IQR 29-46), had the lowest level of consciousness on hospital admission (GCS 3-8; $p < 0.0001$) and all had $RTS < 11$ ($p < 0.0001$). Deceased RTA victims aged over 75 years had the lowest median ISS of 25 (IQR 13-30). In this group, 29.8% were not diagnosed with severe trauma and 44.8% had the maximum RTS. RTA victims aged 18-24 years had the shortest survival (median length of hospital stay 2 days; IQR 1-2) (Table 3).

Table 4. Characteristics of in hospital deceased road traffic accident victims, by hospital level of trauma care.

Characteristics	Level I Trauma centre (n=394)	Level II or III Trauma centre (n=103)	p-value
<i>Road user group, n (%)</i>			0.23
<i>Motorists</i>	124 (31.5)	25 (24.4)	
<i>Motorcyclists</i>	21 (5.3)	2 (1.9)	
<i>Moped riders</i>	44 (11.2)	15 (14.6)	
<i>Bicyclists</i>	159 (40.4)	50 (48.5)	
<i>Pedestrians</i>	46 (11.7)	11 (10.7)	
<i>ISS, median (IQR)</i>	30 (25-38)	13 (9-25)	<0.0001
<i>By category, n (%)</i>			<0.0001
<i>ISS < 16</i>	34 (8.6)	55 (53.9)	
<i>ISS ≥ 16</i>	360 (91.4)	47 (46.1)	
<i>AIS region (AIS ≥ 3), n (%)</i>			
<i>Head</i>	278 (70.6)	36 (35.0)	<0.0001
<i>Face</i>	20 (5.1)	3 (2.9)	0.35
<i>Neck</i>	13 (3.3)	0 (0.0)	0.06
<i>Thorax</i>	194 (49.2)	30 (29.1)	<0.0001
<i>Abdomen</i>	33 (8.4)	4 (3.9)	0.12
<i>Spine</i>	47 (11.9)	1 (1.0)	0.001
<i>Upper Extremities</i>	4 (1.0)	0 (0.0)	0.31
<i>Lower Extremities</i>	83 (21.1)	27 (26.5)	0.26
<i>External</i>	16 (4.1)	3 (2.9)	0.59
<i>Age (years), mean (SD)</i>	61.1 (22.2)	74.0 (16.2)	<0.0001
<i>By category, n (%)</i>			<0.0001
<i>18-24</i>	39 (9.9)	3 (2.9)	
<i>25-55</i>	103 (26.1)	8 (7.8)	
<i>56-75</i>	127 (32.2)	26 (25.2)	
<i>>75</i>	125 (31.7)	66 (64.1)	

ISS: injury severity score; IQR: interquartile range

AIS: abbreviated injury score; AIS ≥ 3: Severe injury.

Pre-hospital triage

In total, 394 of the 497 fatally injured RTA victims (79.3%) were directly transported to a level-I trauma centre. As expected, patients transported directly to a level-I trauma centre more often had severe trauma (ISS \geq 16) compared to patients transported to level-II and level-III trauma centres (91.4% versus 46.1%; $p<0.0001$; Table 4). Nevertheless, 47 of the 407 (11.5%) patients with severe trauma were not transported to a level-I trauma centre. These RTA victims, of whom 61.7% were 75 years or older, were significantly older than the 360 RTA victims with severe trauma who were transported to a level-I trauma centre hospital (mean 70.7 versus 59.9 years; $p=0.001$). In contrast, no age difference was found between the RTA victims without severe trauma who were transported to a level-I and those transported to a level-II or level-III hospital (mean 74.0 versus 77.6 years; $p=0.30$; Table 5). The patients who were directly transported to a level-I trauma centre had more often severe trauma to the head, thorax and spine ($p<0.001$; Table 4). Fatally injured motorcyclists were mainly transported to a level-I centre, whereas the other fatally injured road users were distributed more evenly between level-I trauma centre and level-II or level-III trauma centre hospitals (Table 4).

Table 5. Characteristics of in hospital deceased road traffic accident victims by severity of trauma and hospital level of trauma care.

Characteristics	ISS < 16		p-value	ISS \geq 16		p-value
	Level I Trauma centre (n = 34)	Level II or III Trauma centre (n = 55)		Level I Trauma centre (n = 360)	Level II or III Trauma centre (n = 47)	
<i>Road user group, n (%)</i>			0.24			0.58
<i>Motorists</i>	10 (29.4)	8 (14.5)		114 (31.7)	17 (36.2)	
<i>Motorcyclists</i>	0	0		21 (5.8)	1 (2.1)	
<i>Moped riders</i>	4 (11.8)	10 (18.2)		40 (11.1)	5 (10.6)	
<i>Bicyclists</i>	18 (52.9)	29 (52.7)		141 (39.2)	21 (44.7)	
<i>Pedestrians</i>	2 (5.9)	8 (14.5)		44 (12.2)	3 (6.4)	
<i>AIS region (AIS\geq3), n (%)</i>						
<i>Head</i>	9 (26.5)	4 (7.3)	0.03	269 (74.7)	32 (68.1)	0.33
<i>Face</i>	0	0	-	20 (5.6)	3 (6.4)	0.74
<i>Neck</i>	0	0	-	13 (3.6)	0	0.38
<i>Thorax</i>	8 (23.5)	9 (16.4)	0.40	186 (51.7)	21 (44.7)	0.37
<i>Abdomen</i>	1 (2.9)	1 (1.8)	1.00	32 (8.9)	3 (6.4)	0.78
<i>Spine</i>	0	0	-	47 (13.1)	1 (2.1)	0.03
<i>Upper Extremities</i>	0	0	-	4 (1.1)	0	1.00
<i>Lower Extremities</i>	6 (17.6)	18 (32.7)	0.12	77 (21.4)	9 (19.1)	0.72
<i>External</i>	1 (2.9)	0	0.38	15 (4.2)	3 (6.4)	0.45
<i>Age (years), mean (SD)</i>	74.0 (17.8)	77.6 (11.0)	0.30	59.9 (22.2)	70.7 (19.5)	0.001
<i>By category, n (%)</i>						
<i>18-24</i>	1 (2.9)	0	0.40	38 (10.6)	3 (6.4)	<0.0001
<i>25-55</i>	4 (11.8)	3 (5.5)		99 (27.5)	4 (8.5)	
<i>56-75</i>	9 (26.5)	15 (27.3)		118 (32.8)	11 (23.4)	
<i>>75</i>	20 (58.8)	37 (67.3)		105 (29.2)	29 (61.7)	

ISS: injury severity score; SD: standard deviation.

AIS: abbreviated injury score; AIS \geq 3: Severe injury.

DISCUSSION

This nationwide study aimed to analyse injury patterns and - the severity of in-hospital deceased adult road traffic accident (RTA) victims in The Netherlands. Two third of the patients over 56 years and all motorcyclists suffered severe trauma ($ISS \geq 16$). Also, of all deceased RTA victims, the severely injured and young patients were mainly transported to a level-I trauma centre hospital, whereas the older severely injured patients and less severely injured patients of all ages were often transported to a level-II or level-III trauma centre hospital.

The majority of the deceased RTA victims (63.2%) had sustained severe head trauma. This prevalence is almost three times higher than that of all RTA victims in the Netherlands (23.0%) [2]. Thus, it can be assumed that the prevention of head trauma may substantially reduce the number of RTA fatalities. In the Netherlands, helmet use is mandatory for motorcyclists and some types of moped vehicles, but not for bicyclists even though bicycling is the most common form of transportation [11]. The highest percentage of fatal RTA accidents in this study involved bicyclists in whom head injuries were the most prevalent type of severe injury ($AIS > 3$). In a previous Dutch study similar results were found as severe traumatic brain injuries were mostly diagnosed in bicyclists, pedestrians and moped riders [12]. Yilmaz et al. found that severe head injuries were more prevalent in bicycle-related trauma admissions in the Netherlands, compared to Australia where helmet use is required by law [13]. Therefore, the implementation of strict nationwide guidelines on helmet use in bicyclists and all moped riders can be considered to help reduce the number of head trauma-related RTA fatalities in the Netherlands. Also, the introduction of new and the improvement of existing governmental protective and preventive measures (traffic education, improved infrastructure design, improved vehicle safety standards and better enforcement of traffic rules) are essential to promote traffic safety in The Netherlands so that the number of RTA fatalities in all road user groups can be decreased.

The second most diagnosed injury in our study was severe thorax trauma, which was found in almost half of all deceased RTA victims. This type of severe injury was most often diagnosed in young victims and motorists and motorcyclists. Our current findings are in line with recent German studies that were retrospectively conducted with autopsy data from deceased RTA victims [14,15]. Both studies found that the majority of RTA victims died from severe head and thorax trauma.

In our study motorcyclists were the youngest group of RTA victims that died in hospital and all were admitted with severe trauma ($ISS \geq 16$). In total, 92 motorcyclists died

during the study period in the Netherlands [16], so 69 of them died on the scene. Compared to other vulnerable RTA victims such as pedestrians and bicyclists (of whom respectively 108 and 374 victims died during the studied period), it can be deduced that motorcyclists are the most vulnerable road users as most of them died on the scene. This conclusion is supported by an earlier Dutch study, which pointed out that young motorcyclists are more vulnerable than other two-wheeled road users [17]. The fact that all but two severely injured motorcyclists were presented in a level-I trauma centre shows that during prehospital triage motorcyclists are recognized as potentially complex patients with severe injuries in multiple anatomic regions. Unfortunately, this is not always the case for other types of road users with life-threatening injuries since many of those were presented in level-II or level-III hospitals with fewer facilities and less experienced staff. Hence, improving adequate prehospital triage and triage on the admission of all RTA victims is essential.

Our analysis shows that the overall injury severity in fatally injured RTA victims is inversely related to age and that younger RTA victims had a higher prevalence of severe ($\text{AIS} \geq 3$) head and thorax injuries. Similar findings were reported by Heinrich et al. and by Osler et al. [15,18]. Both studies found that in-hospital deceased elderly trauma patients showed lower overall injury severity compared to younger deceased trauma patients. This underlines that the elderly are more vulnerable road users, mostly due to pre-existing comorbidities and functional decline in daily life [19,20]. Our study showed that the RTA victims aged over 75 years had better vital signs at initial presentation than RTA victims aged 18-24 years. This may bias the clinicians' interpretation of injury severity during admission and its impact on the chance of survival. Clinicians should be suspicious of (combinations of) potentially lethal injuries to the head and thorax, that do not seem life-threatening at the time of admission.

According to national guidelines set up by the Dutch Trauma Association, trauma patients with an $\text{ISS} \geq 16$ (severe trauma) should directly be transported to a level-I trauma centre, but in the prehospital phase injury severity often is difficult to determine. Although 88% of all deceased RTA victims with severe trauma in this study were transported to a level-I Trauma centre, this percentage decreased with advancing age from 93% in the youngest age group (18-24 years) to 78% in the elderly (75 years and older). This finding is in line with other studies that have shown that old age is a risk factor for under-triage [21,22]. Including age, alongside prehospital RTS, in the pre-hospital triage of RTA victims may improve the clinical outcome of this specific group of trauma patients. To prevent under-triage in the elderly Calland et al. suggested treating all elderly trauma patients (>65 years) with at least one $\text{AIS} > 3$ injury in a Level-I Trauma centre [23]. As RTA victims aged over 75 years in this study were the largest group,

with the lowest mean ISS and the most favourable clinical parameters, more awareness of the vulnerability of elderly RTA victims in prehospital triage is needed. This is even more important as the proportion of elderly RTA victims in both the Netherlands and in the Europe Union has risen during the past decade and will probably continue to rise in the future [2,24].

LIMITATIONS

Results were based on retrospective data from the national trauma registry, which did not include the 702 on-scene deceased RTA victims. These victims represent a group that was probably more severely injured than the in-hospital deceased victims. However, no factual data of the on-scene deceased group regarding injury pattern and severity is available. A second limitation is that the trauma registry includes only a limited and pre-determined set of parameters. In the registry, road users are not classified in detail, so bicycles and e-bikes are combined in one group and mopeds also include different types of vehicles. This lack of detailed specification may have blurred the results in these more heterogeneous road user groups. Also, it was impossible to study the effects of pre-existent vulnerability, speed, helmet use, and other forms of protection as these data are not available.

CONCLUSION

Further prevention of head trauma is needed to reduce the lethality of RTA victims, especially bicyclists, who are the largest group of deceased RTA victims with head trauma as the predominant injury pattern. Elderly form a large part of the deceased RTA victims in the Netherlands, despite the fact that they have the lowest mean ISS and the best clinical parameters on admission. Thus, the risk of under-triage of injury severity in the elderly is obvious and should be addressed in the early phases of trauma care, especially during pre-hospital triage and the initial care at the admission of elderly RTA victims.

REFERENCES

1. Global status report on road safety 2018. Geneva: World Health Organization; 2018. License: CC BYNC-SA 3.0 IGO
2. Weijermars W, Van Schagen I, Goldenbeld C, et al (2016) Monitoring and identifying road safety 2016. (In Dutch) SWOV Institute for Road Safety Research. The Hague, the Netherlands
3. The ATLS Subcommittee; American College of Surgeons' Committee on Trauma; International ATLS working group (2013) Advanced trauma life support (ATLS®): the ninth edition J Trauma Acute Care Surg 74:1363-1366
4. Weijermars W, Bos N, Stipdonk HL (2016) Serious road injuries in The Netherlands dissected. *Traffic Inj Prev* 17:73-79
5. SWOV Institute for Road Safety Research. (2018) SWOV Fact sheet: Road deaths in the Netherlands. The Hague, the Netherlands
6. European Commission (2010) New road safety programme 2011-2020. Luxembourg, Off. J. Eur. Union
7. Gennarelli TA, Wodzin E (2008) The Abbreviated Injury Scale 2005. Update 2008. *Assoc. Adv. Automot. Med. Des Plaines, IL* 2008
8. Baker SP, O'Neill B, Haddon Jr W, et al (1974) The injury severity score: a method for describing patients with multiple injuries and evaluating emergency care. *J Trauma* 14:187-196
9. Champion HR, Sacco WJ, Copes WS et al (1989) A revision of the Trauma Score. *J Trauma* 29:623-629
10. Teasdale G, Jennett B (1974) Assessment of coma and impaired consciousness: A practical scale *Lancet* 81-84
11. SWOV Institute for Road Safety Research (2017) SWOV Fact sheet: Cyclists. The Hague, the Netherlands
12. Leijdesdorff HA, van Dijck JTJM, Krijnen P, et al (2014) Injury pattern, hospital triage, and mortality of 1250 patients with severe traumatic brain injury caused by road traffic accidents. *J Neurotrauma* 31:459-465
13. Yilmaz P, Gabbe BJ, McDermott FT, et al (2013) Comparison of the serious injury pattern of adult bicyclists, between South-West Netherlands and the State of Victoria, Australia 2001-2009. *Injury* 44:848-854
14. Pfeifer R, Schick S, Holzmann C, et al (2017) Analysis of injury and mortality patterns in deceased patients with road traffic injuries: an autopsy study. *World J Surg* 41:3111-3119
15. Heinrich D, Holzmann C, Wagner A, et al (2017) What are the differences in injury patterns of young and elderly traffic accident fatalities considering death on scene and death in hospital? *Int J Legal Med* 131:1023-1037
16. SWOV Institute for Road Safety Research (2017) SWOV Fact sheet: Serious road injuries in the Netherlands. The Hague, the Netherlands.
17. Leijdesdorff HA, Siegerink B, Sier CFM, et al (2012) Injury pattern, injury severity, and mortality in 33,495 hospital-admitted victims of motorized two-wheeled vehicle crashes in The Netherlands. *J. Trauma Acute Care Surg* 72:1363-1368
18. Osler T, Hales K, Baack B, et al (1988) Trauma in the elderly. *Am. J. Surg* 156:537-543
19. Bergeron E, Lavoie A, Clas D, et al (2003) Elderly trauma patients with rib fractures are at greater risk of death and pneumonia. *J Trauma Acute Care Surg* 54:478-485

20. McGwin G, Melton SM, May AK, et al (2000) Long-term survival in the elderly after trauma. *J Trauma Acute Care Surg* 49:470-476
21. van Rein EAJ, van der Sluijs R, Houwert RM, et al (2018) Effectiveness of prehospital trauma triage systems in selecting severely injured patients: Is comparative analysis possible? *Am J Emerg Med* 36:1060-1069
22. Chang DC, Bass RR, Cornwell EE, et al (2008) Undertriage of elderly trauma patients to state-designated trauma centers. *Arch Surg* 143:776-781
23. Calland JF, Ingraham AM, Martin N, et al (2012) Evaluation and management of geriatric trauma: An eastern association for the surgery of trauma practice management guideline. *J Trauma Acute Care Surg* 73: S345-50
24. European Commission (2018) Road Safety In The European Union - Trends, statistics and main challenges 29:359-367. Luxembourg, Off. J. Eur. Union

